



Corrigendum to

“Brief communication: Rethinking the 1998 China floods to prepare for a nonstationary future” published in Nat. Hazards Earth Syst. Sci., 19, 715–719, 2019

Shiqiang Du^{1,2}, Xiaotao Cheng³, Qingxu Huang⁴, Ruishan Chen⁵, Philip J. Ward², and Jeroen C. J. H. Aerts²

¹School of Environmental and Geographical Sciences, Shanghai Normal University, Shanghai, 200234, China

²Institute for Environmental Studies, Vrije Universiteit Amsterdam, Amsterdam, 1081 HV, the Netherlands

³China Institute of Water Resources and Hydropower Research, Beijing, 100038, China

⁴State Key Laboratory of Earth Surface Processes & Resource Ecology, Beijing Normal University, Beijing, 100875, China

⁵School of Geographic Sciences, East China Normal University, Shanghai, 200241, China

Correspondence: Shiqiang Du (shiqiangdu@shnu.edu.cn) and Jeroen C. J. H. Aerts (jeroen.aerts@vu.nl)

Published: 25 June 2019

The reservoir capacity numbers were misrepresented in the originally published version of this article. The error has been corrected, and this may be considered the official version of the data.

The corrected version of the data:

As a result, the lengths of the river banks protected by well-maintained dikes increased from 76 532 km in 1998 to 201 124 km in 2016 and reservoir capacity increased from 493 *billion* to 897 *billion* cubic meters (MWR, 2017).

The originally published version of the data:

As a result, the lengths of the river banks protected by well-maintained dikes increased from 76 532 km in 1998 to 201 124 km in 2016 and reservoir capacity increased from 493 *trillion* to 897 *trillion* cubic meters (MWR, 2017).