



Modelling ignition probability for human- and lightning-caused wildfires in Victoria, Australia

Annalie Dorph^{1,2}, Erica Marshall¹, Kate A. Parkins¹, and Trent D. Penman¹

¹FLARE Wildfire Research, School of Ecosystem and Forest Sciences, University of Melbourne, Creswick, Victoria, Australia

²School of Environmental and Rural Science, University of New England, Armidale, NSW, Australia

Correspondence: Annalie Dorph (annalie.dorph@une.edu.au)

Received: 14 February 2022 – Discussion started: 8 March 2022

Revised: 7 July 2022 – Accepted: 23 September 2022 – Published: 24 October 2022

Abstract. Wildfires pose a significant risk to people and property, which is expected to grow with urban expansion into fire-prone landscapes and climate change causing increases in fire extent, severity and frequency. Identifying spatial patterns associated with wildfire activity is important for assessing the potential impacts of wildfires on human life, property and other values. Here, we model the probability of fire ignitions in vegetation across Victoria, Australia, to determine the key drivers of human- and lightning-caused wildfire ignitions. In particular, we extend previous research to consider the role that fuel moisture has in predicting ignition probability while accounting for environmental and local conditions previously identified as important. We used Random Forests to test the effect of variables measuring infrastructure, topography, climate, fuel and soil moisture, fire history, and local weather conditions to investigate what factors drove ignition probability for human- and lightning-caused ignitions. Human-caused ignitions were predominantly influenced by measures of infrastructure and local weather. Lightning-sourced ignitions were driven by fuel moisture, average annual rainfall and local weather. Both human- and lightning-caused ignitions were influenced by dead fuel moisture with ignitions more likely to occur when dead fuel moisture dropped below 20 %. In future, these models of ignition probability may be used to produce spatial likelihood maps, which will improve our models of future wildfire risk and enable land managers to better allocate resources to areas of increased fire risk during the fire season.

1 Introduction

Wildfires present a significant risk to both people and property, with this risk increasing as urban areas continue to expand into fire-prone landscapes (Syphard et al., 2013). Wildfire-associated risks are likely to increase further with future climate change scenarios predicting increases to fire extent, severity and frequency in fire-prone ecosystems (Bowman et al., 2009; Flannigan et al., 2009). Wildfires require four key factors to start: sufficient biomass, fuel moisture low enough to allow combustion, weather conditions conducive to fire spread and an ignition source (Archibald et al., 2009; Bradstock, 2010). These factors vary in space and time to influence the risk of a wildfire occurring on any day in a particular location. Although weather variables are considered to be determinants of fire annual extent (Bradstock et al., 2014; Penman et al., 2013), the spatial pattern of fire is better predicted by ignitions and fuels (Parisien et al., 2010; Pausas and Paula, 2012). Understanding spatial patterns in fire activity is important for assessing the risks and associated impacts of wildfires with regard to human life, property and other values.

Spatial variation in ignition likelihood has been documented in a number of studies, with different patterns observed depending on the ignition source under examination (Bar Massada et al., 2013; Clarke et al., 2019; Liu et al., 2012). Sources of wildfire ignition can either be caused by humans or natural. Ignitions caused by humans may be either intentional or accidental and are often related to indicators of human settlement, such as the distance to the nearest road or housing density (Bar Massada et al., 2013;

Clarke et al., 2019). Natural ignition sources include lightning strikes, which can account for up to 90 % of recorded wildfire ignitions (Clarke et al., 2019; Keeley and Syphard, 2018). The spatial pattern of lightning ignitions differs from human-caused ignitions (Bar Massada et al., 2013) with variables such as local weather or topography driving the probability of lightning ignition (Clarke et al., 2019; Liu et al., 2012). Further, lightning storms may be responsible for multiple ignitions at a time, potentially resulting in more concurrent fires or larger fires when these ignitions converge (Read et al., 2018). The proximity of human-caused ignitions to population-dense areas means that early detection of fires is common. However, the potential for lightning-caused ignitions to occur in remote or hard-to-access locations means that detection of these fires can be difficult. Improved models of lightning ignition probability may therefore aid the early detection of and response to wildfires.

Fuels are composed of both live and dead vegetation. Fuel moisture is a critical factor affecting how fire interacts and moves through fuels (Chandler et al., 1983). Fuel moisture in both live and dead fuel contributes to fire ignition, spread and severity in several ecosystems (Chuvienco et al., 2004, 2009; Dennison et al., 2009; Nolan et al., 2016a). For example, fuel moisture content thresholds have been associated with wildfire occurrence in Australian forests and woodlands (Nolan et al., 2016a). Forest fuel moisture content can shift across these thresholds in very short periods of time (e.g. from hours to a week; Dennison et al., 2009; Nolan et al., 2016a), causing a forest to shift from low to high flammability rapidly. It is therefore important to consider the influence of fuel moisture on fire ignitions at fine temporal scales.

Fuel moisture has previously only been tested in studies of fire ignitions, based on calculations from meteorological data (Dowdy and Mills, 2012; Liu et al., 2012; Miranda et al., 2012; Wotton and Martell, 2005). These methods may lead to uncertain estimates of fuel moisture in areas with highly heterogeneous topography and vegetation (Nieto et al., 2010). The recent development of fuel moisture estimates at regular temporal intervals from remotely sensed MODIS data may provide a solution (Nolan et al., 2016a, b). These methods integrate remotely sensed data with climate modelling inputs to generate layers estimating both live and dead fuel moisture across large spatial and temporal scales. Using these inputs as predictors of fire ignitions may improve our estimates of ignition probability for both human-caused and lightning-ignited fires.

Wildfires in south-eastern Australia have resulted in significant loss of human life and property (Blanchi et al., 2010; Filkov et al., 2020; Haynes et al., 2010). A better understanding of both human-caused and lightning-caused ignitions and the associated risks to human life and property is therefore important for this area. This research was aimed at modelling the probability of fire ignitions across Victoria, Australia. Specifically, we ask whether key drivers of human- and lightning-caused fire ignitions are consistent with previously

reported global patterns. In addition, we extend this previous research to determine to what extent fuel moisture influences ignition probability relative to topographic, human and climatic variables.

2 Methods

2.1 Study area

Our study area was the state of Victoria in south-eastern Australia. The population in Victoria is ~ 6.6 million, of which the majority lives in Melbourne (<https://www.abs.gov.au/>, last access: 7 January 2021). Remnant native vegetation in Victoria covers approximately 46 % of the state owing to previous land clearing for agriculture and human settlements (Fig. 1). There is a climatic gradient across the state, with average annual rainfall in the north-west averaging ~ 300 mm, and in the south-east ranging from 1000 to 1500 mm (<http://www.bom.gov.au/>, last access: 7 January 2021). Average daily maximum temperatures in summer also vary across the state, ranging from 27–30 °C in the north-west to 18–24 °C in the south-east (<http://www.bom.gov.au/>, last access: 7 January 2021).

2.2 Data compilation

Historical fire ignition data were obtained from the Victorian Country Fire Authority (CFA) and the Department of Environment, Land, Water and Planning (DELWP) for the period between 2000 and 2019 ($n = 67927$). These databases have approximately 20 different ignition causes. For this study, ignition causes were reclassified broadly into human-caused ($n = 59146$; from arson or accidental sources including burn-offs, campfires, electrical, fireworks, heat or cutting equipment, power transmission lines, re-light, vehicles, and waste disposal) and lightning-ignited fires ($n = 8781$) as previous work found consistent patterns in the drivers of the different types of human ignitions in the study area (Clarke et al., 2019).

The analytical pathways for human and lightning ignitions were necessarily different. For the analysis of human-caused ignitions, a set of random points were generated across Victoria from a uniform distribution ($n = 75281$). Each random point was assigned a random date and time within the date range of the ignitions data. These random points were used as absence data in the statistical model, providing a random sample of points where fire ignitions did not occur. For the lightning ignitions model, data of all lightning events in Victoria over a certain time period were obtained from the Global Position and Tracking System Pty. Ltd. (GPATS) Australia. These lightning events were each assigned a probability of starting a fire. Therefore, absence data were abundant within the dataset and random points were not required.

Data from raster layers representing a range of natural and built environments were extracted for each ignition and ran-




Figure 1. Study area showing the distribution of (a) recorded human ignitions (presence data) in remnant native vegetation and areas of cleared or modified (i.e. matrix) vegetation (from the Country Fire Authority [CFA] and Department of Land, Water and Planning [DELWP] ignitions data), (b) randomly generated points (human ignitions absence data), (c) recorded lightning ignitions in native and matrix vegetation (from CFA and DELWP ignitions data), and (d) a kernel density estimate for all lightning strikes detected in Victoria for the study period (from the Global Position and Tracking System Pty Ltd dataset). Native vegetation and matrix layers sourced from: Department of Environment, Land, Water and Planning (2017, <https://www.data.vic.gov.au/>, last access: 13 November 2020).

dom point. Details of the different environmental variables used, and descriptions of the layers and their data sources are listed in Table 1. Variables were selected based on those identified as being important from previous studies. For example, human-caused ignitions have a strong relationship with infrastructure variables such as distance to the nearest road and housing density or distance to the nearest settlement (Catry et al., 2009; Clarke et al., 2019; Miranda et al., 2012). Local weather variables, topography and average annual rainfall have also been shown to have an effect on human-caused ignitions (Catry et al., 2009; Clarke et al., 2019; Collins et al., 2015; Liu et al., 2012). Similarly, studies of lightning-caused fires have shown ignitions to be influenced by variables such as: aspect, slope and topographic position (Collins et al., 2015; Miranda et al., 2012); average annual rainfall (Clarke et al., 2019); fuel moisture indices (Dowdy and Mills, 2012; Liu et al., 2012; Miranda et al., 2012; Wotton and Martell, 2005); local weather (Clarke et al., 2019; Miranda et al., 2012); and soil moisture (Liu et al., 2012) are predicted to be influenced by changes to fire fuel loads with the time since last fire (Clarke et al., 2019).

2.3 GPATS lightning data

Lightning is an electrical discharge generated when positive and negative charges in clouds separate (Latham and

Williams, 2001). A flash of lightning from a cloud to the ground can contain a single or several return strokes. Each stroke can be described by its duration, strength (amplitude or current) and polarity (positive or negative). All strokes within a lightning flash can either follow the same channel to the ground or have several different terminals if the flash branches near to the ground (Larjavaara et al., 2005).

Data on lightning strokes were obtained from Global Position and Tracking System Pty. Ltd. (GPATS) Australia. GPATS uses triangulation of data from a network of radio receivers to determine the time and location of individual lightning strokes. This technique distinguishes between cloud-to-cloud strokes and cloud-to-ground strokes and detects the multiple strokes that can occur within a single lightning flash. The data contain information about the strength of each lightning stroke (amplitude), its polarity, its time and its location. There is some variation within the detection efficiency of the GPATS data owing to spatial and temporal variation in the systems used (Dowdy et al., 2017). The GPATS data obtained covered the state of Victoria from the period 2004 to 2019 (number of strokes = 3 977 126). The data were simplified for the analysis following the process outlined in Fig. 2.

The time and location of each stroke was used to extract information from raster layers representing natural and human built variables (Table 1). An ignition probability was calculated for each stroke following the method outlined in

Table 1. Environmental and human-mediated variables used as predictors in model development. The table provides a description of the variable and the source of the data.

Variable	Description	Time range	Source
<i>Topography</i>			
Elevation (m)	Calculated from 30 m Digital Elevation Model (DEM)		
Aspect (degrees)	Calculated from 30 m DEM		
Slope (degrees)	Calculated from 30 m DEM	n/a	https://www.data.vic.gov.au/
Topographic position index (TPI)	Calculated from 30 m DEM, combining slope position and landform category; positive TPI values indicate ridges, negative values indicate valleys and values near zero represent plains and areas of constant slope		(last access: 13 October 2020)
<i>Fire</i>			
Time since fire (years)	Derived from fire history maps; TSF was set to 100 for ignitions in areas with no mapped fire history	Annual 2000–2020	https://www.data.vic.gov.au/ (last access: 12 October 2020)
<i>Infrastructure</i>			
Housing density (houses per km ²)	Calculated from vector files of address locations following Clarke et al. (2019)	n/a	https://www.data.vic.gov.au/ (last access: 25 August 2020)
Distance to the nearest road (km)	Calculated from vector files of roads following Clarke et al. (2019)		
<i>Climate</i>			
Rainfall (mm)	Mean annual rainfall from Worldclim v2.1	Average from 1970 to 2000	https://www.worldclim.org/ (last access: 12 October 2020)
<i>Weather</i>			
Forest fire danger index (FFDI)	FFDI calculated from gridded hourly temperature, drought factor and relative humidity data within the VicClim database	Hourly	Country Fire Authority
Wind speed (km h ⁻¹)	Wind speed from gridded hourly data within the VicClim database	2000–2017	(CFA)
<i>Dryness</i>			
Soil moisture (sm_pct)	Extracted from the root zone soil moisture layer provided by the Australian Landscape Water Balance	Annual 2005–2020	http://www.bom.gov.au/ (last access: 1 October 2020)
<i>Fuel moisture</i>			
Live fuel moisture content Dead fuel moisture content	Fuel moisture within vegetation; calculated following Nolan et al. (2016b) (N.B. live fuel moisture was limited to areas of native vegetation)	Annual 2000–2019	
<i>Response variables</i>			
Historical fire ignitions data	Dataset containing latitude, longitude, timestamps and ignition cause for recorded fire ignitions across Victoria, Australia	2000–2019	Country Fire Authority (CFA); Department of Environment, Land, Water and Planning (DELWP)
GPATS lightning data	Dataset containing latitude, longitude and timestamps for individual lightning strokes across Victoria, Australia	2004–2019	Global position and tracking system Pty. Ltd. (GPATS) Australia

n/a stands for not applicable.




Figure 2. Lightning data manipulation process. Here, a lightning flash is composed of several strokes that can differ in their duration, strength (amplitude) and polarity. Strokes can follow the same channel to the ground or can branch and have several terminals near the ground. * Equations for the calculation of stroke ignition probability are described in text following the method outlines in Larjavaara et al. (2005).

Larjavaara et al. (2005) based on the temporal and spatial proximity of a stroke to an ignition caused by lightning. First, the proximity index (A) was calculated for each stroke within 10 km of the ignition and within 5 days preceding the ignition. Approximately 75 % of fire ignitions from lightning are detected within 3 d of the lightning occurring (Wotton and Martell, 2005). We allowed for 5 d to cover this period and ensure we captured the majority of ignitions. The proximity index was calculated following Eq. (1):

$$A = (1 - T/120)(1 - S/10), \tag{1}$$

where T is the delay in hours between the time of the stroke to the time of the ignition and S is the spatial distance in kilometres between the stroke and the fire.

The proximity index was then used to calculate ignition probability for each stroke (P_S) by dividing the proximity of each stroke (A_S) by the sum of all strokes (A_i) within 10 km and 5 d of the ignition following Eq. (2):

$$P_S = A_S / \sum_{i=1}^n A_i. \tag{2}$$

Lightning strokes were grouped into lightning flashes if they occurred within 0.5 s and 0.05° in both latitude and longitude. This reduced the size of the dataset ($n = 1\,994\,918$) and allowed inclusion of flash multiplicity (a potential indicator of ignition likelihood) (Flannigan and Wotton, 1991; Larjavaara et al., 2005) as a predictor in the statistical models. The ignition probability of a flash (P_F) was then calculated following the inclusion-exclusion principle described in Eq. (3):

$$P_F = 1 - \prod_{S=1}^n (1 - P_S). \tag{3}$$

The inclusion-exclusion principle was used rather than the sum of probabilities, as the latter could be greater than one when a lightning stroke was linked to more than one fire. More information on the inclusion-exclusion principle is included in Appendix A. By completing these calculations, all lightning strokes that were linked to a CFA or DELWP fire ignition using the proximity index were assigned an ignition probability greater than zero. Any lightning strokes that were not linked to a CFA or DELWP ignition by the proximity index were assigned an ignition probability of zero and were treated as absence data in the analysis. Finally, for each flash the environmental data (listed in Table 1) were calculated by averaging the data extracted for each stroke.

2.4 Random forest modelling

Random forests were used to determine the probability of an ignition occurring, with separate models built for lightning- and human-caused ignitions. Random forests are a non-parametric modelling technique with a higher classification accuracy and reduced risk of overfitting the data compared with other parametric modelling techniques (Breiman, 2001; Cutler et al., 2007). Both classification and regression trees are used in random forests, which are built using a random subset of the data (usually 70 %; termed out of bag (OOB) samples). Trees are then ensembled to calculate either the majority vote (classification) or average value (regression) of

predictions in the model (Breiman, 2001). Model accuracy is calculated by comparing the model built on the OOB samples with the data withheld during model development and averaging this across all observations (Cutler et al., 2007). For classification trees this results in an estimate of classification error, whereas for regression this gives a measure of variance explained and mean square error.

Variable importance is calculated following two different methods for the two types of random forest. For classification trees, variable importance is calculated by summing the decrease in Gini impurity that occurs every time a variable is chosen to split a node in the classification tree, giving the measure the Mean Gini Index (Cutler et al., 2007). For regression trees, variable importance is calculated by measuring the total decrease in the residual sum of squares that occurs every time a variable is chosen to split a node in a regression tree, giving a measure of Included Node Purity.

Random forests were used to model lightning- and human-caused ignitions separately. Each ignitions dataset was split again into those occurring within remnant native vegetation (hereafter “native forest”) and those on cleared or modified land (hereafter “matrix”) (based on the native vegetation layer from <https://www.data.vic.gov.au/>, last access: 13 November 2020). Models were therefore prepared for four different ignition datasets – human ignitions in native vegetation, human ignitions in matrix, lightning ignitions in native vegetation and lightning ignitions in matrix. Splitting the data into native forest and matrix vegetation was undertaken to allow the inclusion of live fuel moisture as a predictor in models of native forest and also resulted in much reduced computation times for the models. Live fuel moisture has only been modelled for native forest in the south-east of Australia, meaning that no data were available in matrix areas. As previous work has indicated that live fuel moisture thresholds can determine fire activity (e.g. Dennison et al., 2009) we wanted to determine its importance relative to other predictors in this area.

2.4.1 Human-caused ignitions model development

A classification random forest for human ignitions was built on the ignition data (presence) and a set of random points (absence). As the number of points in the presence and absence data was uneven, we used a down-sampling method to balance the two classes (Valavi et al., 2021). Down sampling is a method that takes subsamples of the data (with replacement) at each tree so that the classes are equal in sample size. The subsamples are replaced and resampled for every tree that is built. Models were built using all variables listed in Table 1. Soil moisture was removed from the models as it did not improve the overall accuracy of these models and it limited the dataset to ignitions from 2005 onwards. Similarly, of the weather variables, only the FFDI was retained in the model as including the other variables did not improve model accuracy. To ensure optimal model fit, model tuning

was conducted to determine the number of trees to grow and the number of variables to sample at each split.

Model validation was done using the OOB error described above and a 10-fold cross-validation procedure. The cross-validation procedure retained a 10 % subset of the data as a test set and built the model on the remaining data. By performing this cross-validation we were able to retain a complete section of the dataset to validate the model and account for any large variation present in the dataset. The model fit was assessed by comparing the training and testing errors produced by the two model validation procedures. Partial dependence plots (PDPs) were used to examine the relationship of each predictor variable with the probability of an ignition when the other predictors variables are held constant at their average (Friedman, 2001).

2.4.2 Lightning-caused ignitions model development

The lightning dataset was highly skewed towards lightning flashes with no probability of ignition (89 % of the dataset). Therefore, this analysis was conducted in two stages. The first stage reclassified the ignition probabilities to determine if a flash had no chance of starting a fire (ignition probability = 0) or if the flash had any chance of starting a fire (ignition probability > 0). The reclassified data were zero inflated; thus, a classification random forest was run using a down-sampling method for class imbalanced data (Valavi et al., 2021), as described above. The second stage of the analysis used a regression random forest on all the lightning flashes with any chance of starting a fire to predict what the probability of ignition would be. For consistency, we used the same variables in both stage one and stage two.

Models were built using all the variables listed in Table 1. Testing of variables was undertaken to determine which weather variables were important to include in the model. The best models were produced when both weather variables listed in Table 1 were used. Use of only the FFDI resulted in a drop in the accuracy of models. Additionally, the number of strokes within a flash and the average amplitude of these strokes were used as predictor variables. As with the human ignitions model, model tuning was conducted to determine the optimal number of trees and variables to use at each split. Model validation using OOB and cross-validation was conducted for both stages of the lightning model, and partial dependence plots were built to show the effect of the predictors on the probability of lightning causing an ignition.

All analyses were conducted in R v3.6.3 using the “randomForest” package to build random forest models (Liaw and Wiener, 2002). Partial dependence plots were produced using the “pdp” package (Greenwell, 2017) and all graphs were produced in the “ggplot2” package (Wickham, 2009).




Figure 3. Variable importance of predictors for human-caused ignitions in (a) matrix and (b) native forest, and lightning ignitions in (c) matrix and (d) native forest. Average and standard deviations of the mean decrease in the Gini Index for the models produced during the 10-fold cross-validation are shown.

3 Results

Models of ignition probability from human-caused sources used data from 59 984 ignitions in native forest and 94 034 ignitions in cleared or modified land (i.e. matrix areas). The classification model generated for each of these areas was very accurate, with both models predicting between 86.4 % and 90.3 % of ignitions and non-ignitions (i.e. random points) correctly (Table 2). Models of ignition probability from lightning sources used data from 888 604 flash events in native forest and 986 777 flash events in the matrix. Of these flashes, 11 % occurred within 10 km and 5 d of a recorded lightning fire event and so had an ignition probability calculated (average = 3 %). The first stage of the lightning modelling process (classification procedure) recorded all flashes with any probability of ignition as an ignition presence. These models performed well with low error classifying lightning flashes as either likely to start a fire, or with no chance of starting a fire (Table 2). The second stage of the lightning modelling process (regression procedure) used only flashes with a probability of starting a fire. This resulted in a reduced dataset containing only 77 193 lightning flash events in native forest and 82 762 flashes in matrix. The ability of these regression models to accurately predict the probability

that a fire was started from one of the lightning flashes was quite low (~ 15 %; Table 2).

3.1 Human-caused ignitions

The human ignitions models for both native forest and matrix were predominantly driven by variables measuring human infrastructure: distance to road and housing density (Fig. 3a and b). The probability of ignition in both models decreased rapidly as the distance from road increased to ~ 500 m and then levelled off (Fig. 4). This effect was not as large within the native forest model as it was in the matrix. Conversely, the probability of ignition increased rapidly in both models as housing density increased to ~ 100 and remained high above this threshold. The FFDI was the next most important variable, having a greater influence on the probability of ignition in native forests than in matrix (Fig. 3a and b). In both models, the probability of an ignition increased rapidly up to an FFDI of ~ 30 and remained high above this point (Fig. 4). Dead fuel moisture and rainfall had a weaker influence in both models (Fig. 3a and b). However, both showed thresholds of influence with a lower probability of ignition in parts of the state with low average annual rainfall (< 1000 mm)

Table 2. Random forest results for human ignitions and GPATS lightning data. Model parameters indicate where the random forest was a classification or a regression, how many trees were built (n_{tree}) and the number of variables tested at each split (m_{try}). Average and standard deviation from the 10-fold cross-validation of classification error for points classed as ignitions compared with points with no ignition are given for classification models. Average and standard deviations from the 10-fold cross-validation are provided for the variance explained (R^2) of regression models.

Model	Model parameters			Classification error		R^2
	Type	n_{tree}	m_{try}	Ignition	No ignition	
Human matrix	Classification	500	2	0.100 ± 0.004	0.101 ± 0.004	
Human native	Classification	500	2	0.136 ± 0.007	0.097 ± 0.006	
Lightning matrix	Classification	500	6	0.173 ± 0.005	0.127 ± 0.001	
Lightning native	Classification	500	6	0.226 ± 0.006	0.131 ± 0.002	
Lightning matrix	Regression	600	4			15.041 ± 1.269
Lightning native	Regression	600	4			15.358 ± 0.929




Figure 4. Partial dependence plot for the six top variables in the random forest model for human-caused fire ignitions in (a) matrix and (b) native forest. Variables are plotted in order of importance. Black lines are the average probability of ignition from the 10 models produced during the 10-fold cross-validation. Grey error bars represent the upper and lower estimates from the 10-fold cross-validation.

and higher probability of ignition in areas with dead fuel moisture content below 20 (Fig. 4).

3.2 Lightning-caused ignitions

In classification models predicting whether a lightning flash had any chance of starting a fire, fuel moisture, average annual rainfall, weather and soil moisture were the most important variables (Fig. 3). Dead fuel moisture was very important to both models in native forest and in the matrix, with a higher probability of ignition when dead fuel moisture was below 20 (Fig. 5). In matrix vegetation, rainfall was the

most important variable (Fig. 3c), with the probability of ignition in parts of the state with average annual rainfall above 1000 mm (Fig. 5a). Rainfall was less important in native forest (Fig. 3d) but followed a similar trend with the probability of an ignition increasing as average annual rainfall increased (Fig. 5b). FFDI was the most important variable in the native forest model (Fig. 3d), with the probability of an ignition increasing to an FFDI of ~ 30 and remaining stable at values above this threshold (Fig. 5b). For FFDI in matrix, the probability of ignitions followed the same trend (Fig. 5a). Soil moisture was of similar importance to both matrix and na-




Figure 5. Partial dependence plot for the six top predictor variables in the random forest models predicting whether a lightning flash has any probability of starting a fire in (a) matrix and (b) native forest. Variables are plotted in order of importance. Black lines represent the average probability of ignition from the 10 models used in the cross-validation. Grey error bars represent the upper and lower estimates from the 10-fold cross-validation.




Figure 6. Variable importance of predictors for the probability of a lightning flash causing an ignition in (a) matrix and (b) native forest. Average and standard deviations of the included node purity calculated by averaging the residual sum of squares over all trees in a random forest model for each model used in the 10-fold cross-validation.

tive vegetation models (Fig. 3). However, there was no strong trend in either vegetation type, with slightly higher ignition probability in areas with soil moisture below 0.25 (Fig. 5).

Fuel moisture, lightning stroke amplitude, weather and average annual rainfall were the most important predictors of ignition probability in the regression models for lightning ignition. Dead fuel moisture and average stroke amplitude had the strongest effect in both native and matrix vegetation, followed by FFDI and rainfall (Fig. 6). However, the overall

strength of these predictor variables to predict fire ignition probability from a lightning stroke was low (Table 2).

4 Discussion

Understanding spatial patterns of ignition probability is important for the assessment of risks from wildfires to human life, property and environmental values. We evaluated wild-

fire ignition probability across a broad range of environments in south-eastern Australia to determine key variables driving human-caused and lightning-caused ignitions. We found that human-caused ignitions were primarily influenced by human infrastructure and local weather conditions, whereas lightning-sourced ignitions were driven by fuel moisture, average annual rainfall and local weather conditions. In particular, dead fuel moisture influenced ignitions from both sources, with ignitions more likely to occur when dead fuel moisture dropped below 20 %.

Human-caused ignitions were strongly related to measures of infrastructure and local weather, with clear thresholds indicating the influence of these variables. These results are consistent with a number of other studies demonstrating the importance of infrastructure and weather for predicting spatial patterns in ignitions from human causes (Clarke et al., 2019; Collins et al., 2015; Faivre et al., 2014; Liu et al., 2012; Syphard et al., 2008). These studies also reported similar trends, with a higher probability of ignitions closer to roads and housing (Clarke et al., 2019; Faivre et al., 2011; Liu et al., 2012) and a steady increase in ignition probability as the FFDI reached 50 (Clarke et al., 2019). The accuracy of the models produced here (86.4 % and 90.3 %) are also comparable with that of models in other studies, although differences in the modelling approach may affect interpretation here. For example, Clarke et al. (2019) reported AUCs between 84.6 and 96.7, Collins et al. (2015) reported deviance explained of 83.8 % to 88.3 % and Liu et al. (2012) reported R^2 of 95 %. The similarity in predictive accuracies among these models suggests that the inclusion of fuel moisture might not improve overall model performance for human-caused ignitions, despite clear thresholds existing for ignition probability.

Lightning ignition probability was most strongly driven by rainfall, dead fuel moisture and local weather conditions (FFDI). Average annual rainfall is known to influence fuel load and accumulation (Thomas et al., 2014); thus, areas with higher average annual rainfall have greater fuel availability and consequently are likely to have an increased probability of ignition. The importance of average annual rainfall and FFDDI for lightning ignitions has previously been demonstrated in this region, with similar trends in ignitions for these variables reported (Clarke et al., 2019). Although studies have previously demonstrated the influence of fuel moisture on lightning ignitions (Dowdy and Mills, 2012; Miranda et al., 2012; Wotton and Martell, 2005), few have done so in conjunction with both local environmental and weather conditions (but see Liu et al., 2012). Incorporation of variables measuring fuel moisture, environmental factors and local weather may have contributed to a higher prediction accuracy for ignition following lightning flash events using a classification procedure in this study compared with others. Here, model classification accuracy was between 77.4 % and 87.3 % for models of lightning ignition. Previous studies using different methods and variables to measure classification

accuracy have found 67.2 % deviance explained (Collins et al., 2015), 73 % variability explained (Wotton and Martell, 2005) and an AUC of 79.3 (Clarke et al., 2019). Our estimates may be further improved by incorporating measures of weather in the days following a lightning event, as delays between lightning strike events and fire ignitions are known to occur (Wotton and Martell, 2005).

There is a clear dead fuel moisture threshold effect for both ignition types. Ignition probability was higher in areas where dead fuel moisture was below 20 % and dropped to almost zero at fuel moisture ratings above this level. Similar effects of fuel moisture or surface moisture have been found in previous studies, with decreases in ignition probability as moisture content increases for both lightning- and human-caused ignitions (Liu et al., 2012; Miranda et al., 2012; Wotton and Martell, 2005). The 20 % fuel moisture threshold in dead fuel moisture content mirrors the thresholds found in a study assessing the area burnt by fires in this region (Nolan et al., 2016a), indicating that dead fuel moisture levels above 20 % significantly reduce both the likelihood of an ignition and the size of a fire. Although dead fuel moisture influenced ignition probability in both lightning- and human-caused ignitions, live fuel moisture had only a limited effect in the models and is thus much less likely to determine ignition probability.

Despite the high accuracy achieved in the classification models, lower accuracies were recorded in the regression models of lightning ignition probability. This is likely due partly to discrepancies in the recording locations of both fire ignitions and GPATS lightning flashes. Within the fire ignitions dataset locations are sometimes recorded at the nearest road or intersection to the fire, rather than at the exact latitude and longitude of the ignition. There are also likely to be ignitions missing from the dataset, because many lightning ignitions occurring in remote areas may not be reported or may be reported days after the ignition has occurred. In the GPATS lightning dataset, there is potential recording errors of at least 1 km within key deployment areas (<http://www.gpats.com.au/>, last access: 25 January 2021) and potentially higher error in remote locations. Additionally, the use of different recording systems over the years has resulted in spatial and temporal variation in recorded lightning flash locations (Dowdy et al., 2017). These spatial and temporal discrepancies have likely contributed to the error in the regression lightning model, but may have also influenced classification accuracy in the lightning classification model.

Models of ignition probability were able to be produced with high predictive accuracy, despite the spatial error present within the predictor datasets. These models allow for spatial likelihood maps of ignition probability to be produced at a daily temporal resolution. High accuracy ignition probability models also have application in the estimation of areas where ignition probability may increase under different climate change scenarios. In turn, this would provide better information in models of future fire behaviour and risk in fire-prone landscapes. Further, the ignition probability mod-

els developed here use daily weather variables as inputs. This means that finer-scale, daily ignition probability maps could be produced on a regular basis. These could inform the placement of suppression resources for rapid attack when fires occur and also provide better information for the community, such as providing warnings on high fire danger days.

5 Conclusion

Globally consistent patterns were found in the drivers of both human- and lightning-caused ignitions. Human-caused ignitions are predominantly pre-determined by the proximity to human settlement and weather conditions, whereas lightning-caused ignitions are driven by fuel moisture, average annual rainfall and local weather. Relationships with remotely sensed values of fuel moisture provide a means of better understanding and predicting the likelihood of fires across the landscape on a daily time step. These high accuracy spatial and temporal likelihood maps of ignition probability will improve our models of wildfire risk and enable land managers to better allocate resources during the fire season.

Appendix A

Inclusion-exclusion formula for any two events:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B), \quad (\text{A1})$$

where

$$P(A \cap B) = P(A) \cdot P(B) \quad (\text{A2})$$

and

$$P(A) = 1 - P(\bar{A}) \text{ and conversely } P(\bar{A}) = 1 - P(A), \quad (\text{A3})$$

$$P(B) = 1 - P(\bar{B}) \text{ and conversely } P(\bar{B}) = 1 - P(B). \quad (\text{A4})$$

Therefore, the probability of an event not occurring,

$$P(\bar{A}) \cdot P(\bar{B}) = (1 - P(A)) \cdot (1 - P(B)), \quad (\text{A5})$$

becomes

$$P(\bar{A}) \cdot P(\bar{B}) = 1 - P(A) - P(B) + P(A \cap B). \quad (\text{A6})$$

And conversely, the probability of an event occurring:

$$P(A \cup B) = 1 - (1 - P(A)) \cdot (1 - P(B)). \quad (\text{A7})$$

Therefore, the inclusion-exclusion formula for the case when there are n number of events becomes:

$$1 - \prod_{i=1}^n (1 - P_i). \quad (\text{A8})$$

Data availability. All data processed could not be made publicly available. For access, the first author can be contacted by email: annalie.dorph@une.edu.au.

Author contributions. AD and TDP were responsible for conceptualisation. AD was responsible for data curation. AD and EM were responsible for formal analysis. TDP was responsible for funding acquisition. AD and KP were responsible for investigation. AD was responsible for methodology. AD, TDP and KAP were responsible for project administration. TDP and KP were responsible for supervision. AD was responsible for validation. AD was responsible for visualisation. AD, KAP, TDP and ED were responsible for writing the original draft, reviewing and editing.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. The fire ignitions data were made available for this study by the Victorian Department of Environment, Land, Water and Planning. Lightning strike data were made available from the Global Position and Tracking System Pty. Ltd. Australia. The authors would like to thank Stephen Deutsch, Frazer Wilson and Estrella Melero-Blanca for their thoughts and contributions to the development of the analytical approach. We would like to acknowledge the Traditional Owners of the land where this research took place, the Dja Dja Wurrung, Wadawurrung, Wurundjeri and Wida-geri peoples.

Financial support. This research has been supported by the Department of Environment, Land, Water and Planning, State Government of Victoria.

Review statement. This paper was edited by Ricardo Trigo and reviewed by Tomàs Artés Vivancos and one anonymous referee.

References

- Archibald, S., Roy, D. P., van Wilgen, B. W., and Scholes, R. J.: What limits fire? An examination of drivers of burnt area in Southern Africa, *Global Change Biol.*, 15, 613–630, <https://doi.org/10.1111/j.1365-2486.2008.01754.x>, 2009.
- Bar Massada, A., Syphard, A. D., Stewart, S. I., and Radeloff, V. C.: Wildfire ignition-distribution modelling: a comparative study in the Huron–Manistee National Forest, Michigan, USA, *Int. J. Wildl. Fire*, 22, 174–183, <https://doi.org/10.1071/WF11178>, 2013.

- Blanchi, R., Lucas, C., Leonard, J., Finkele, K., Blanchi, R., Lucas, C., Leonard, J., and Finkele, K.: Meteorological conditions and wildfire-related house loss in Australia, *Int. J. Wildl. Fire*, 19, 914–926, <https://doi.org/10.1071/WF08175>, 2010.
- Bowman, D. M. J. S., Balch, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A., D'Antonio, C. M., DeFries, R. S., Doyle, J. C., Harrison, S. P., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Marston, J. B., Moritz, M. A., Prentice, I. C., Roos, C. I., Scott, A. C., Swetnam, T. W., van der Werf, G. R., and Pyne, S. J.: Fire in the Earth System, *Science*, 324, 481–484, <https://doi.org/10.1126/science.1163886>, 2009.
- Bradstock, R., Penman, T., Boer, M., Price, O., and Clarke, H.: Divergent responses of fire to recent warming and drying across south-eastern Australia, *Global Change Biol.*, 20, 1412–1428, <https://doi.org/10.1111/gcb.12449>, 2014.
- Bradstock, R. A.: A biogeographic model of fire regimes in Australia: current and future implications, *Global Ecol. Biogeogr.*, 19, 145–158, <https://doi.org/10.1111/j.1466-8238.2009.00512.x>, 2010.
- Breiman, L.: Random Forests, *Mach. Learn.*, 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, 2001.
- Catry, F. X., Rego, F. C., Bação, F. L., and Moreira, F.: Modeling and mapping wildfire ignition risk in Portugal, *Int. J. Wildl. Fire*, 18, 921–931, <https://doi.org/10.1071/WF07123>, 2009.
- Chandler, C. H., Cheney, P., Thomas, P., Trabaud, L., and Williams, D.: Fire in forestry. Volume 1. Forest fire behavior and effects, in: Volume 2, Forest fire management and organization, Wiley-Interscience, ISBN 978-0471874423, 1983.
- Chuvieco, E., Aguado, I., and Dimitrakopoulos, A. P.: Conversion of fuel moisture content values to ignition potential for integrated fire danger asses, *Can. J. Forest Res.*, 34, 2284–2293, 2004.
- Chuvieco, E., González, I., Verdú, F., Aguado, I., and Yebra, M.: Prediction of fire occurrence from live fuel moisture content measurements in a Mediterranean ecosystem, *Int. J. Wildl. Fire*, 18, 430–441, <https://doi.org/10.1071/WF08020>, 2009.
- Clarke, H., Gibson, R., Cirulis, B., Bradstock, R. A., and Penman, T. D.: Developing and testing models of the drivers of anthropogenic and lightning-caused wildfire ignitions in south-eastern Australia, *J. Environ. Manage.*, 235, 34–41, <https://doi.org/10.1016/j.jenvman.2019.01.055>, 2019.
- Collins, K. M., Price, O. F., and Penman, T. D.: Spatial patterns of wildfire ignitions in south-eastern Australia, *Int. J. Wildl. Fire*, 24, 1098, <https://doi.org/10.1071/WF15054>, 2015.
- Cutler, D. R., Edwards, T. C., Beard, K. H., Cutler, A., Hess, K. T., Gibson, J., and Lawler, J. J.: Random forests for classification in ecology, *Ecology*, 88, 2783–2792, <https://doi.org/10.1890/07-0539.1>, 2007.
- Dennison, P. E., Moritz, M. A., Dennison, P. E., and Moritz, M. A.: Critical live fuel moisture in chaparral ecosystems: a threshold for fire activity and its relationship to antecedent precipitation, *Int. J. Wildl. Fire*, 18, 1021–1027, <https://doi.org/10.1071/WF08055>, 2009.
- Department of Environment, Land, Water and Planning: Native Vegetation Regulation Extent, Department of Environment, Land, Water and Planning [data set], <https://discover.data.vic.gov.au/dataset/native-vegetation-regulation-extent-20171> (last access: 13 November 2020), 2017.
- Dowdy, A. J. and Mills, G. A.: Atmospheric and fuel moisture characteristics associated with lightning-attributed fires, *J. Appl. Meteorol. Clim.*, 51, 2025–2037, <https://doi.org/10.1175/JAMC-D-11-0219.1>, 2012.
- Dowdy, A. J., Fromm, M. D., and McCarthy, N.: Pyrocumulonimbus lightning and fire ignition on Black Saturday in southeast Australia, *J. Geophys. Res.-Atmos.*, 122, 7342–7354, <https://doi.org/10.1002/2017JD026577>, 2017.
- Faivre, N., Roche, P., Boer, M. M., McCaw, L., and Grierson, P. F.: Characterization of landscape pyrodiversity in Mediterranean environments: contrasts and similarities between south-western Australia and south-eastern France, *Landsc. Ecol.*, 26, 557–571, <https://doi.org/10.1007/s10980-011-9582-6>, 2011.
- Faivre, N., Jin, Y., Goulden, M., and Randerson, J.: Controls on the spatial pattern of wildfire ignitions in Southern California, *Int. J. Wildl. Fire*, 23, 799–811, <https://doi.org/10.1071/WF13136>, 2014.
- Filkov, A. I., Ngo, T., Matthews, S., Telfer, S., and Penman, T. D.: Impact of Australia's catastrophic 2019/20 bushfire season on communities and environment. Retrospective analysis and current trends, *J. Saf. Sci. Resil.*, 1, 44–56, <https://doi.org/10.1016/J.JNLSSR.2020.06.009>, 2020.
- Flannigan, M. D. and Wotton, B. M.: Lightning-ignited forest fires in northwestern Ontario, *Can. J. Forest Res.*, 21, 277–287, <https://doi.org/10.1139/x91-035>, 1991.
- Flannigan, M. D., Stocks, B., Turetsky, M., and Wotton, B. M.: Impacts of climate change on fire activity and fire management in the circumboreal forest, *Global Change Biol.*, 15, 549–560, <https://doi.org/10.1111/j.1365-2486.2008.01660.x>, 2009.
- Friedman, J. H.: Greedy function approximation: a gradient boosting machine, *Ann. Stat.*, 29, 1189–1232, 2001.
- Greenwell, B. M.: pdp: an R package for constructing partial dependence plots, *R J.*, 9, 421–436, <https://doi.org/10.32614/RJ-2017-016>, 2017.
- Haynes, K., Handmer, J., McAneney, J., Tibbits, A., and Coates, L.: Australian bushfire fatalities 1900–2008: exploring trends in relation to the 'Prepare, stay and defend or leave early' policy, *Environ. Sci. Policy*, 13, 185–194, <https://doi.org/10.1016/J.ENVSCL.2010.03.002>, 2010.
- Keeley, J. E. and Syphard, A. D.: Historical patterns of wildfire ignition sources in California ecosystems, *Int. J. Wildl. Fire*, 27, 781–799, <https://doi.org/10.1071/WF18026>, 2018.
- Larjavaara, M., Pennanen, J., and Tuomi, T. J.: Lightning that ignites forest fires in Finland, *Agr. Forest Meteorol.*, 132, 171–180, <https://doi.org/10.1016/j.agrformet.2005.07.005>, 2005.
- Latham, D. and Williams, E.: Lightning and forest fires, in: Forest Fires: Behavior and Ecological Effects, edited by: Johnson, E. and Miyawaki, K., Academic Press, San Diego, California, ISBN 978-0123866608, 2001.
- Liaw, A. and Wiener, M.: Classification and regression by random Forest, *R News*, 2, 18–22, 2002.
- Liu, Z., Yang, J., Chang, Y., Weisberg, P. J., and He, H. S.: Spatial patterns and drivers of fire occurrence and its future trend under climate change in a boreal forest of Northeast China, *Global Change Biol.*, 18, 2041–2056, <https://doi.org/10.1111/j.1365-2486.2012.02649.x>, 2012.
- Miranda, B. R., Sturtevant, B. R., Stewart, S. I., and Hammer, R. B.: Spatial and temporal drivers of wildfire occurrence in the context of rural development in northern Wisconsin, USA, *Int. J. Wildl. Fire*, 21, 141–154, <https://doi.org/10.1071/WF10133>, 2012.

- Nieto, H., Aguado, I., Chuvieco, E., and Sandholt, I.: Dead fuel moisture estimation with MSG–SEVIRI data. Retrieval of meteorological data for the calculation of the equilibrium moisture content, *Agr. Forest Meteorol.*, 150, 861–870, <https://doi.org/10.1016/J.AGRFORMET.2010.02.007>, 2010.
- Nolan, R. H., Boer, M. M., Resco de Dios, V., Caccamo, G., and Bradstock, R. A.: Large-scale, dynamic transformations in fuel moisture drive wildfire activity across southeastern Australia, *Geophys. Res. Lett.*, 43, 4229–4238, <https://doi.org/10.1002/2016GL068614>, 2016a.
- Nolan, R. H., Resco de Dios, V., Boer, M. M., Caccamo, G., Goulden, M. L., and Bradstock, R. A.: Predicting dead fine fuel moisture at regional scales using vapour pressure deficit from MODIS and gridded weather data, *Remote Sens. Environ.*, 174, 100–108, <https://doi.org/10.1016/J.RSE.2015.12.010>, 2016b.
- Parisien, M. A., Miller, C., Ager, A. A., and Finney, M. A.: Use of artificial landscapes to isolate controls on burn probability, *Landsc. Ecol.*, 25, 79–93, <https://doi.org/10.1007/s10980-009-9398-9>, 2010.
- Pausas, J. G. and Paula, S.: Fuel shapes the fire-climate relationship: evidence from Mediterranean ecosystems, *Global Ecol. Biogeogr.*, 21, 1074–1082, <https://doi.org/10.1111/j.1466-8238.2012.00769.x>, 2012.
- Penman, T. D., Collins, L., Price, O. F., Bradstock, R. A., Metcalf, S., and Chong, D. M. O.: Examining the relative effects of fire weather, suppression and fuel treatment on fire behaviour – A simulation study, *J. Environ. Manage.*, 131, 325–333, <https://doi.org/10.1016/j.jenvman.2013.10.007>, 2013.
- Read, N., Duff, T. J., and Taylor, P. G.: A lightning-caused wildfire ignition forecasting model for operational use, *Agr. Forest Meteorol.*, 253–254, 233–246, <https://doi.org/10.1016/j.agrformet.2018.01.037>, 2018.
- Syphard, A. D., Radeloff, V. C., Keuler, N. S., Taylor, R. S., Hawbaker, T. J., Stewart, S. I., and Clayton, M. K.: Predicting spatial patterns of fire on a southern California landscape, *Int. J. Wildl. Fire*, 17, 602–613, <https://doi.org/10.1071/WF07087>, 2008.
- Syphard, A. D., Bar Massada, A., Butsic, V., and Keeley, J. E.: Land Use Planning and Wildfire: Development Policies Influence Future Probability of Housing Loss, *PLoS One*, 8, e71708, <https://doi.org/10.1371/journal.pone.0071708>, 2013.
- Thomas, P. B., Watson, P. J., Bradstock, R. A., Penman, T. D., and Price, O. F.: Modelling surface fine fuel dynamics across climate gradients in eucalypt forests of south-eastern Australia, *Ecography*, 37, 827–837, <https://doi.org/10.1111/ECOG.00445>, 2014.
- Valavi, R., Elith, J., Lahoz-Monfort, J. J., and Guillera-Aroita, G.: Modelling species presence-only data with random forests, *Ecography*, 44, 1731–1742, <https://doi.org/10.1111/ECOG.05615>, 2021.
- Wickham, H.: *ggplot2: elegant graphics for data analysis*, Springer-Verlag, New York, <https://doi.org/10.1007/978-0-387-98141-3>, 2009.
- Wotton, B. M. and Martell, D. L.: A lightning fire occurrence model for Ontario, *Can. J. Forest Res.*, 35, 1389–1401, <https://doi.org/10.1139/x05-071>, 2005.