



Dependence of seismic hazard assessment on the observation time interval: insights from a synthetic earthquake catalogue in southeastern Spain

Elena Pascual-Sánchez¹, José Antonio Álvarez-Gómez², Julián García-Mayordomo³, and Paula Herrero-Barbero³

¹Department of Topographical Engineering and Cartography, Universidad Politécnica de Madrid, Madrid, Spain

²Department of Geodynamics, Stratigraphy and Paleontology, Universidad Complutense de Madrid, Madrid, Spain

³Instituto Geológico y Minero de España – CSIC, Madrid, Spain

Correspondence: Elena Pascual-Sánchez (elena.pascual.sanchez@upm.es) and Julián García-Mayordomo (julian.garcia@igme.es)

Received: 2 October 2025 – Discussion started: 4 December 2025

Revised: 24 April 2026 – Accepted: 24 June 2026 – Published: 27 October 2026

Abstract. Traditionally Probabilistic Seismic Hazard Assessment (PSHA) is mainly based on historical and instrumental catalogues. The seismic catalogue in SE Spain covers a period of almost 1 kyr. Although this catalogue can be considered long, its time span is not enough to record the complete seismic cycle of the slow moving regional faults, for example, the ones which conform the Eastern Betics Shear Zone (EBSZ). To assess whether the seismic hazard at the EBSZ depends on the time interval in which the earthquake catalogue has been recorded, this study performs a PSHA using synthetic seismicity. The synthetic seismicity in the EBSZ consists of a 1 Myr catalogue generated in previous studies using the RSQSim earthquake simulator, applied to the fault system that forms the EBSZ. This catalogue has been divided into ten thousand sub-catalogues of the same duration as the historical and instrumental one and randomly distributed over time. The magnitude-frequency distributions of the synthetic sub-catalogues show significant variability in the maximum reached magnitude, in the slope of the Gutenberg-Richter relationship and in the annual rate of earthquakes. A hundred sub-catalogues have been selected to perform individual PSHA and have been compared with the results derived from historical and instrumental seismicity. Using R-CRISIS software, this study obtains seismic hazard curves for the main cities in the region, showing the estimated return period for different values of Peak Ground Acceleration (PGA). The hazard curves reveal that each sub-catalogue leads to different return period val-

ues. The obtained variability ranges from 11 % to 21 % for a $\text{PGA} = 0.04 \text{ g}$, and from 25 % to 58 % for a $\text{PGA} = 1 \text{ g}$. Our results show that there is a dependence between seismic hazard and the observation time interval in which an earthquake catalogue is recorded.

1 Introduction

Consequences of large earthquakes affect all aspects of human life (Daniell et al., 2012; Marano et al., 2010), both by direct and indirect effects (Gokhale et al., 2004). Characterization of seismic hazard allows us to prevent and mitigate the devastating effects of earthquakes. The development of a Probabilistic Seismic Hazard Assessment (PSHA) through a statistical analysis of the occurrence of earthquakes, leads to the determination of exceeding probabilities for different ground motion levels (Cornell, 1968; McGuire, 2008). The results derived from a PSHA are used in the definition of seismic actions in building codes (e.g. Bisch et al., 2012; Barbat et al., 2005), which makes them of crucial importance in the protection of human lives and the limitation of damages.

Recorded seismicity is the fundamental input data of PSHA, in which the so-called Gutenberg-Richter relationship (Gutenberg and Richter, 1944) and the maximum magnitude of the earthquake catalogue describe the seismic activity of a region. Gutenberg-Richter relationship is calculated fitting to a log-linear equation the cumulative magnitude dis-

tribution of an earthquake catalogue (Eq. 1).

$$\log_{10} N(M) = a - b \cdot M \quad (1)$$

where $N(M)$ represents the number of earthquakes with magnitude equal to, or higher than M , a is the annual activity rate of earthquakes and b is slope of the function, which depends on the relationship between the number of large and small events.

The probability density function associated with the Gutenberg-Richter relationship ($f_M(M)$, Eq. 2) describes a dependence with the minimum magnitude. This magnitude can be considered as the magnitude of completeness, M_c , from which all seismicity is assumed to have been recorded.

$$f_M(m) = 1 - e^{-\beta(M-M_c)} \quad M_c \leq M \leq \infty \quad (2)$$

In Eq. (2), β represents another way of expressing the slope of the Gutenberg-Richter relationship (Eq. 3).

$$\beta = b \cdot \ln(10) \quad (3)$$

Most of the implementations of the Gutenberg-Richter equation on PSHA are by means of the truncated, or modified, relationship (Cornell and Vanmarcke, 1969; Cosentino et al., 1977), in which probability density function ($f_M^t(M)$, Eq. 4) depends also on $M_{\max}$. In this case, $M_{\max}$ is considered to be the maximum possible value of the earthquake magnitude. This approximation avoids the overestimation of the mean return period of the largest earthquakes of the general Gutenberg-Richter exponential relationship.

$$f_M^t(m) = \frac{1 - e^{-\beta(M-M_c)}}{1 - e^{-\beta(M_{\max}-M_c)}} \quad (4)$$

At slow-moving fault systems, with tectonic deformation rates of less than 2 mm yr^{-1} and seismic cycles of thousands of years, recorded seismicity is usually moderate (Echeverría et al., 2013; Shi et al., 2014; Anderson and Luco, 1983; Perea et al., 2006; Alfaro et al., 2012; Gómez-Novell, 2021). An example is the case of SE Spain. In this region, seismicity is low to moderate, even though the EBSZ fault system has the potential to generate large earthquakes (Perea et al., 2012; Gómez-Novell et al., 2020; Herrero-Barbero et al., 2021).

Paleoseismological studies have shown that the time interval between major earthquakes at the EBSZ single faults is of the order of several thousands of years (Ortuño et al., 2012; Martínez-Díaz et al., 2012b, 2019; Masana et al., 2010, 2018; García-Mayordomo, 2005; Canora-Catalán et al., 2016; Ferrater et al., 2015; Moreno, 2011; Roquero et al., 2019; Insua-Arévalo et al., 2015; Martín-Banda et al., 2016). Meanwhile, the historical and instrumental record used in hazard studies at SE Spain, barely covers a thousand years: from 1048 to the present (IGN and UPM, 2013; González, 2017). This period is shorter than the seismic cycle of the faults. For example, in the case of the Carrascoy fault, the interval between

large earthquakes is estimated to be of three thousand years (Martín-Banda et al., 2016). Consequently, historical and instrumental record covers just one-third of the seismic cycle at this fault. Meaning a narrow part of the seismic cycle, earthquake catalogues neither may record the entire seismogenic potential of faults nor changes in their behaviour. An example of the latter is the study carried out by Martín-Banda et al. (2021) in which fault slip rate variability has been proposed during the last 210 kyr in the case of the Carrascoy fault at the EBSZ. This variability can be related to accelerations of the system and interactions between adjacent faults, which can lead to the coexistence of short seismic cycles with large ones (Sieh et al., 2008; Benedetti et al., 2013).

Characterization of slow-moving fault systems uses geological data to complement the scarce information about seismic activity. A better understanding of fault systems has turned into the inclusion of them as independent seismogenic sources in hazard studies around the world (Field et al., 2015; Stirling et al., 2012; Valentini et al., 2017; Peruzza et al., 2011; García-Mayordomo et al., 2012a). According to García-Mayordomo (2007), at cities located near active faults with recurrence intervals of 2 kyr, the inclusion of faults as seismogenic sources may represent a contribution of 10 %–30 % of hazard for return periods of 500 yr. Another approach has modeled the EBSZ as seismogenic source zone. To include the contribution of its main faults to a classical PSHA, a hybrid approach was used by defining an area source formed by the fault system (IGN and UPM, 2013).

Synthetic seismicity appears as another tool to address the limitations of seismic record (Robinson et al., 2011; Dieterich, 1994; Ward, 2000; Rundle et al., 2006). Computer models based on earthquake physics allow the creation of synthetic catalogues with the same characteristics as the instrumental ones but extended in time. Time expansion makes it possible to reproduce several seismic cycles (Robinson et al., 2011; Console et al., 2018) and to evaluate different models of seismogenic processes (Wilson et al., 2017). By using rate- and state- dependent friction numerical approach (Dieterich, 1979), it is also possible to simulate the physics of rupture processes of earthquakes (Richards-Dinger and Dieterich, 2012). All of these features make synthetic seismic models useful for gaining a better understanding of earthquake occurrence, as illustrated in this study.

In this study, a classical PSHA methodology (Esteva, 1967; Cornell, 1968) has been applied to 100 different 1 kyr long seismic catalogues extracted randomly from a 1 Myr synthetic catalogue of the EBSZ (Herrero-Barbero et al., 2021; Álvarez-Gómez et al., 2023). Depending on the time interval of analysis, each catalogue is supposed to cover a different part of the seismic cycle of faults. To compare the results obtained, a PSHA is also carried out with data from the historical and instrumental catalogue. The results derived from these analyses will show how the calculation of seismic hazard depends on the time interval in which the seismic catalogue is recorded.

2 Seismotectonic context

Southeastern Spain is characterized by weak to moderate seismicity (Stich et al., 2003; Rodríguez-Escudero et al., 2014). Nevertheless, it is considered the most seismically active region of the Iberian Peninsula (García-Mayordomo et al., 2007), due to the convergence between the African and Eurasian plates (DeMets et al., 1990; Argus et al., 1999). This study focuses on the Eastern Betic Shear Zone (EBSZ), located at the Betic Cordillera, specifically, at its Internal Zones. Betic Cordillera conforms the northern termination of the Rif-Betic Alpine orogenic belt (De-Larouziere et al., 1988). The EBSZ is an active deformation zone that conforms mainly left-lateral strike-slip faults oriented N-S to ENE-WSW (Ortuño et al., 2012). The main faults that constitute the system are (Fig. 1, Table 1), from south to north: the Carboneras Fault (CF), the Palomares Fault (PF), the Alhama de Murcia Fault (AMF), the Los Tollos Fault (LTF), the Carrascoy Fault (CAF) and the Bajo Segura Fault (BSF). The mean slip rate of the system is estimated to be $1.5 \pm 0.3 \text{ mm yr}^{-1}$ (Echeverría et al., 2013) unevenly distributed across different faults.

The EBSZ is characterized by a moderate seismic activity, although it has produced many damaging earthquakes ($M_w \geq 6.0$) in historical times: 1048 Orihuela (Silva et al., 2015), 1482 OHerrerorihuela (García-Mayordomo, 2005), 1487 Almería (IGN and UPM, 2013), 1518 Vera (García-Mayordomo et al., 2012a), 1522 Alhama de Almería (Reicherter and Hübscher, 2007), 1673 Orihuela (García-Mayordomo, 2005), 1674 Lorca (Martínez-Díaz et al., 2019), 1804 Dalias (Murphy, 2019), 1829 Torrevieja (García-Mayordomo and Martínez-Díaz, 2006) and 1910 Adra (Rodríguez-Escudero et al., 2014). The most recent damaging earthquake was the Lorca event of 2011, with $M_w = 5.1$, which claimed nine fatalities and huge economic losses (Martínez-Díaz et al., 2012b; López-Comino et al., 2012). All these major earthquakes have been associated with the faults of the area (Yazdi et al., 2023; Yazdi and García-Mayordomo, 2024). For producing a PSHA of the region, the EBSZ has been modeled previously as a seismogenic zone (García-Mayordomo et al., 2012a; IGME, 2015) which gathers the seismic features of the fault system.

3 Materials and methods

In order to gain insight into the influence of the observation time interval on seismic hazard results at the EBSZ, we have resorted to the use of synthetic seismic catalogues. From a 1 Myr-long computed catalogue (Herrero-Barbero et al., 2021; Álvarez-Gómez et al., 2023), we randomly selected ten thousand 1 kyr sub-catalogues to characterise the seismic hazard within a classical PSHA framework. These 1 kyr sub-catalogues were extracted to match the duration of the historical catalogue (spanning from 1048 AD and there-




Figure 1. Main faults which constitute the EBSZ (García-Mayordomo et al., 2012b, 2017): the Carboneras Fault (CF), the Palomares Fault (PF), the Alhama de Murcia Fault (AMF), the Los Tollos Fault (LTF), the Carrascoy Fault (CAF) and the Bajo Segura Fault (BSF). The representation of GNSS vectors (Echeverría et al., 2013; Borque et al., 2019) is made with fixed Eurasia. The earthquakes are sourced from the catalogue used in this study (IGN, 2022).

fore approximately 1 kyr long). Each sub-catalogue represents a possible realisation of the long-term seismic process, allowing us to compare the hazard estimated from multiple stochastic 1 kyr realisations with that derived from the observed historical and instrumental seismicity, which can likewise be interpreted as a single realisation.

Synthetic seismicity consists of a million-year catalogue generated using RSQSim earthquake simulator. RSQSim simulations require a fault system model that includes a representation of the geometry, kinematics and frictional parameters of the fault segments. The implementation of the EBSZ in RSQSim (Herrero-Barbero et al., 2021) is based on a fully-interacting 3D model built externally to the simulator. This model is composed of triangular cells of 1 km^2 resolution, allowing the simulation of events with magnitudes greater than $M_w = 3.0$ and approximating some geometric complexities. Rate- and state-dependent friction parameters (Dieterich, 1979) control the sliding resistance and are selected in this model following realistic relations between rupture area and earthquake magnitude. Seismic cycles are divided into several states (healing, nucleation and seismic rupture) that are passed by fault elements throughout the simulation. The evolution of stress, slip variation and state variable are controlled by analytical expressions that use event-driven steps. Mean slip rates of faults are used by the simulator to apply tectonic loading to the faults by the backslip method (Savage, 1983). The result of the simulations applied to this model is an earthquake catalogue characterized by a Magnitude Frequency Distribution (MFD) that tends

Table 1. Model input parameters for fault sections in the EBSZ obtained from Herrero-Barbero et al. (2021).

Fault name	Fault section ID	Rake	Deviation	Slip rate	Deviation	References
Alhama de Murcia Fault (AMF)	AMF-1	20	0–40	1.1	0.50–1.70	R from slickenlines measured by Martínez-Díaz (1998). SR based on trenches (Ferrater et al., 2016; Ortuño et al., 2012)
	AMF-2	39	19–59	0.9	0.80–1.00	R from 2011 Lorca eq. (Martínez-Díaz et al., 2012a) and structural analysis by Alonso-Henar et al. (2020). SR from trenching and morphotectonic analysis (Ferrater et al., 2016)
	AMF-3	42	25–58	0.2	0.07–0.32	Data inferred from AMF-4
	AMF-4	42	25–58	0.2	0.07–0.32	SR and R from cross-section restorations (Herrero-Barbero et al., 2020). Min. SR from Silva et al. (2003)
Carboneras Fault (CF)	CF-1	10	0–20	1.2	1.10–1.30	Data from deflected submarine channels in CF-1 (Moreno, 2011) and onshore fluvial channels in CF-2 (Moreno et al., 2015). Max. SR based on GPS data of Echeverría et al. (2015)
	CF-2	10	0–20	1.2	1.10–1.30	
Palomares Fault (PF)	PF-1	–5 (0)	(–25)–15	0.04	0.01–0.08	Data measured by Booth-Rea et al. (2004), comparing fluvial deposits and paleochannels in PF-1. R in PF-2 is consistent with the moment tensor of a recent M_w 3.6 event (IGN, 2019)
	PF-2	15	0–35	0.04	0.01–0.08	
	PF-3	15	0–35	0.1	0.04–0.16	R is inferred from PF-2 and SR is based on comparison between PF-3 and LTF-1
Los Tollos Fault (LTF)	LTF	15	0–35	0.16	0.06–0.25	R and SR obtained through trenching by Insua-Arévalo et al. (2015)
Carrascoy Fault (CAF)	CAF-1	90	75–90	0.37	0.29–0.45	R and SR estimated from trenches by Martín-Banda et al. (2016)
	CAF-2	15	5–25	0.85	0.50–1.20	Data from structural analysis (Martín-Banda, 2020; Sanz de Galdeano et al., 1998; Silva, 1994)
Bajo Segura Fault (BSF)	BSF-1	90	70–110	0.4	0.29–0.51 (0.6)	SR estimations from Alfaro et al. (2012) using stratigraphic markers and assuming pure reverse. Note that GPS data from Borque et al. (2019) show $0.6 \pm 0.2 \text{ mm yr}^{-1}$ of shortening for the entire BSF zone
	BSF-2	90	70–110	0.27	0.20–0.34	
	BSF-3	90	70–110	0.2	0.15–0.25	
	BSF-4	90	70–110	0.2	0.15–0.25	For R, see seismic profiles from Alfaro et al. (2012) and Perea et al. (2012). SR is inferred from BSF-3

to fit well with the historical and instrumental features. For more details on the physics-based model and the validation procedure of the best-fit catalogue, see Herrero-Barbero et al. (2021, 2023), Álvarez-Gómez et al. (2023), and Gómez-Novell et al. (2024).

The historical and instrumental earthquake catalogue in SE Spain is compiled by the *Instituto Geográfico Nacional* (IGN, National Geographic Institute). It can be downloaded freely online (IGN, 2022). The earthquake catalogue provides information on the date, the location, and the magnitude of earthquakes. To conduct this study, it has been necessary to process the catalogue collected until 2022, since the last hazard studies (IGN and UPM, 2013) were carried out in 2012. Originally, earthquakes were reported on heterogeneous magnitude scales. In order to obtain the size of earthquakes on the same magnitude scale, a homogenization process is needed. Using the equation proposed by Cabañas et al. (2015), from a Reduced Major Axis Regression (RMA), the magnitudes m_{bLg} and m_{b} are converted to moment magnitude, M_{w} . Then, earthquakes with magnitudes smaller than $M_{\text{w}} = 3.0$ are removed from the catalogue, since the equations of Cabañas et al. (2015) do not work in this range of values and the completeness of the catalogue cannot be guaranteed. In addition, earthquakes occurring at depths greater than 35 km are removed as they do not impact the seismic hazard (IGN and UPM, 2013).

According to the Poisson hypothesis (Cornell, 1968; Cornell and Winterstein, 1988), earthquakes must be considered as independent phenomena that occur randomly in time. For this purpose, a declustering process (Gardner and Knopoff, 1974) has been applied to the earthquake catalogue using a set of equations developed ad-hoc for the region (IGN and UPM, 2013) (Eqs. 5, 6, and 7).

$$\log(T) = 0.6 \cdot M_{\text{w}} - 0.7 \quad (5)$$

for $M_{\text{w}} < 5.9$

$$\log(T) = 0.04 \cdot M_{\text{w}} + 2.6 \quad (6)$$

for $M_{\text{w}} \geq 5.9$

$$\log(L) = 0.2 \cdot M_{\text{w}} + 0.4 \quad (7)$$

where T represents the time interval between earthquakes; M_{w} , the maximum magnitude of each cluster and L the linear distance that separates two earthquakes. M_{w} is determined by means of an iterative process (Wiemer, 2001) in which the largest earthquake is selected from a cluster of related events. The declustering process allows differentiation between mainshocks and fore- after-shocks. Mainshocks are selected as independent events with information that is representative of the cluster. This procedure has been applied to both historical-instrumental and synthetic catalogues as is the common procedure on classical PSHA analyses.

The characterisation of the EBSZ as a seismogenic source is represented by the Gutenberg-Richter relationship and the

maximum magnitude of each catalogue. Different scenarios for seismic hazard assessment have been generated using the Mersenne-twister random number generator (Matsumoto and Nishimura, 1998) and selecting intervals of 1 kyr from the general synthetic catalogue of 1 Myr. Each sub-catalogue is subjected to a PSHA, which results in an individual hazard curve for the selected sites.

The input data for the hazard calculation consist of the annual earthquake rate, the slope of the Gutenberg-Richter relationship and the maximum magnitude of each catalogue at the EBSZ. The seismic parameters of the seismogenic zones adjacent to the EBSZ zone are also required for the model. These data have been adopted from the ZESIS database (Table 2). These parameters were obtained within the framework of the National Seismic Hazard Map project, following a process of declustering and magnitude homogenization of the earthquake catalogue (IGN and UPM, 2013). In that work, different zoning models (García-Mayordomo et al., 2012b) and procedures for calculating the Gutenberg-Richter parameters were tested and eventually incorporated into the hazard calculations through a logic-tree approach. The maximum magnitude parameter was modelled using a normal probability distribution. The mean maximum magnitude and its standard deviation were derived for each zone based on criteria that balanced the available information on the presence and size of active faults and the magnitude of the largest historical events. Active fault parameters were obtained from the QAFI database (IGME, 2022) which at that time was version 2. Several procedures were evaluated to convert fault-size parameters into moment magnitude using empirical relationships from the literature (IGN and UPM, 2013). Table 2 shows the values referenced for the ZESIS model (IGME, 2015), which we adopt here.

For each scenario, a MFD has been created and adjusted to a Gutenberg-Richter power relationship using the maximum likelihood fitting technique (Eq. 8, Utsu, 1964; Aki, 1965; Sandri and Marzocchi, 2007).

$$b = \frac{\log_{10} e}{(\bar{M} - M_{\text{c}})} \quad (8)$$

where $\bar{M}$ describes the mean magnitude of earthquakes with a magnitude at least M_{c} , representing the magnitude of completeness of the catalogue.

The magnitude of completeness has been calculated using the Goodness-of-fit test (GMF, Wiemer and Wyss, 2000). Maximum magnitude has been calculated, following the procedure of IGN and UPM (2013), as an average between the geological maximum and the maximum magnitude of each catalogue.

In the development of seismic hazard studies, a set of Ground Motion Prediction Equations (GMPE) is often used in combination with a logic tree. In this case, in which we are interested in the variation of results linked to the observation time window, we use a single GMPE, to avoid including more parameters that might condition the variability of the

Table 2. Seismic parameters of the seismogenic zones adjacent to the EBSZ zone: β represents the slope of the Gutenberg-Richter relationship; $\tau(4.0)$ is the annual rate of earthquakes with magnitude $M_w \geq 4.0$; and “MM” is the mean maximum magnitude with its standard deviation (σ MM) (after IGN and UPM, 2013; IGME, 2015).

SEISMIC SOURCE	β	$\tau(4.0)$	MM	σ MM
Central Guadalquivir (CG)	2.35	0.191	6.6	0.4
Cazorla-Segura & Albacete pre-Betics (CS & ApB)	2.9	0.06	5	0.4
Valencian Platform & Alicante pre-Betics (VP & ApB)	2.05	0.259	6.6	0.4
Western Internal Betics (WIB)	2.3	0.198	6.6	0.3
Granada basin (Gb)	2.58	0.576	6.8	0.3
Sierra Nevada-Filábride & Guadix-Baza basin (SNF & GBb)	2.26	0.141	6.6	0.4
Murcian pre-Betics (MpB)	2.67	0.39	6.68	0.2
Central Internal Betics (CIB)	2.21	0.317	6.7	0.2
Eastern Internal Betics (EIB)	3.09	0.081	6.7	0.1
Western Alboran sea (WAs)	2.16	0.133	6.5	0.3
Argelian-Balear basin (ABb)	3.11	0.126	6.5	0.4
Alborán-Rif Central crest (ARCc)	2.46	0.889	7	0.2
Eastern Rif (ER)	2.07	0.3	6.4	0.4
Orán (O)	1.84	0.577	7.3	0.5

results. GMPE has been selected as the one derived by Campbell and Bozorgnia (2014) because it includes parameters that describe the behaviour of a fault system, such as the style of faulting and the depth of nucleation of earthquakes. Other studies of seismic hazard on the Iberian Peninsula had also chosen Campbell and Bozorgnia (2014) as GMPE (Rivas-Medina et al., 2018; Gómez-Novell et al., 2020).

Hazard curves reveal the estimated return period for different ground motion levels. Using three types of seismic parameters for the EBSZ: the ones that came from (1) synthetic seismicity, (2) historical and instrumental seismicity updated to the present and (3) ZESIS database, this study conducts a PSHA for six of the most populous cities in SE Spain (Fig. 2, Table 3). Hazard curves have been computed using R-CRISIS software (Ordaz and Salgado-Gálvez, 2020), which calculates seismic hazard using a fully probabilistic approach. Using seismic and geometric features of seismic sources, R-CRISIS performs a spatial integration by subdividing the original source. This procedure is based on the assumption that seismicity is evenly distributed over the area source (Ordaz et al., 2021). Considering the selected GMPE, R-CRISIS evaluates the exceedance probability at different values of ground motion at the selected sites. The probabilities of all the magnitudes and seismic sources are summed to obtain the total exceedance values. Exceedance values have been transformed into return periods. The results obtained consist of 100 curves from 100 synthetic catalogues, one curve for the historical and instrumental catalogue updated to the present, and one curve for ZESIS data. A hundred synthetic scenarios have been considered enough to show whether the variability within the time interval of analysis exists and to quantify it.




Figure 2. Seismogenic sources adjacent to EBSZ (IGME, 2015): Central Guadalquivir (CG), Cazorla-Segura & Albacete pre-Betics (CS-ApB), Valencian Platform & Alicante pre-Betics (VP-ApB), Western Internal Betics (WIB), Granada basin (Gb), Sierra Nevada-Filábride & Guadix-Baza basin (SNF-GBb), Murcian pre-Betics (MpB), Central Internal Betics (CIB), Eastern Internal Betics (EIB), Western Alboran sea (WAs), Argelian-Balear basin (ABb), Alborán-Rif Central crest (ARCc), Eastern Rif (ER) and Orán (O) and cities where seismic hazard analyses have been performed: Alicante (ALI), Torrevieja (TOR), Murcia (MUR), Lorca (LOR), Vera (VER) and Almería (ALM).


4 Results

4.1 Magnitude Frequency Distribution

MFD displays the rate of earthquakes over a range of magnitudes. The curve of accumulated events is described by the Gutenberg-Richter relationship (Fig. 3, in blue). In this study,


Table 3. Populous cities at SE Spain that have been selected to perform a PSHA based on synthetic seismicity.

CITY	GEOGRAPHIC COORDINATES	POPULATION (SSO, 2022)
Murcia	−1.130278, 37.986111	462 979
Lorca	−1.70, 37.683333	97 151
Alicante	−0.483056, 38.345278	338 577
Vera	−1.867778, 37.2475	18 224
Torrevecija	−0.683333, 37.977778	83 547
Almería	−2.4675, 36.84	199 237

**Figure 3.** MFD of the EBSZ obtained from the historical and instrumental record since 1048 to 2022. Discrete rate of annual earthquakes, in pink; cumulative annual rate, in black; Gutenberg-Richter relationship (Eq. 1) using maximum likelihood fitting technique (Eq. 8, Utsu, 1964; Aki, 1965), in blue; and the magnitude of completeness (M_c), in red.

the regression resulting from a maximum likelihood curve fitting technique (Eq. 8) for the historical and instrumental catalogue, has coefficient values of $a = 3.8$ and $b = 1.0$. On PSHA, the annual rate of earthquakes, τ , higher or equal to a given magnitude is a needed parameter. Following IGN and UPM (2013), the selected magnitude for this study is $M_w \geq 4.0$. In the case of the EBSZ, we obtained a value of $\tau(M_w \geq 4.0) = 0.63$, which means that, on average, less than one earthquake with a magnitude equal to or higher than four is recorded in this region per year.

In the case of synthetic sub-catalogues, a variety of frequency–magnitude distributions (Fig. 4) with their corresponding Gutenberg-Richter parameters (Fig. 5) can be observed. The distribution of a (Fig. 5a) shows two maxima between the values $a = 3.0$ – 3.5 and $a = 4.5$ – 5.5 . In the case of b (Fig. 5b), a similar pattern can be found. The slope of the Gutenberg-Richter relationship presents both maxima at $b = 0.8$ – 0.9 and $b = 1.2$ – 1.3 . If we evaluate the frequency distribution of a and b using ten thousand sub-catalogues as a Gaussian distribution, we obtained that it is centred at $a = 4.5$ and $b = 1.1$. These mean values are significantly

**Figure 4.** Frequency magnitude distribution for the 100 synthetic catalogues (grey lines) and the historical-instrumental catalogue (red line). The dashed line represents the magnitude of completeness for the full length of synthetic catalogues according to Herrero-Barbero et al. (2021). The magnitude of completeness for the historical-instrumental catalogue is M_w 3.5 (IGN and UPM, 2013).

close to the values which are obtained using the historical and instrumental seismic record, especially in the case of b .

Maximum magnitude values (Fig. 5c) shows a bimodal pattern with its main maximum at $M_{\max} = 7.3$ – 7.4 and a secondary maximum over $M_{\max} = 6.3$ – 6.4 . Given that we are considering the fault system as a whole, this bimodal pattern could be related to differences in the behaviour of faults. A segmentation of the synthetic sub-catalogues by faults has been carried out towards the evaluation of earthquake nucleation at each fault (Fig. 6). Obtained results reveal that, on the one hand, the Alhama de Murcia fault (AMF, Fig. 6) and the Carboneras fault (CF, Fig. 6) keep the same bimodal distribution as the whole system. On the other hand, the Carrascoy fault (CAF, Fig. 6) and the Bajo Segura fault (BSF, Fig. 6) only present a normal distribution centred in $M_{\max} = 5.6$ for both faults. The nucleation of large earthquakes in the model, with $M_{\max} > 7.1$ in the CAF and the BSF has not been notified in these sub-catalogues. According to previous research in the region (Herrero-Barbero et al., 2021), nucleation of




Figure 5. Statistical variability of (a) a – parameter (Gutenberg-Richter relationship), (b) b – parameter (Gutenberg-Richter relationship) and (c) $M_{\max}$ for 10 000 synthetic sub-catalogues and (d), (e) and (f) for 100 synthetic sub-catalogues.




Figure 6. Distribution of maximum magnitudes for 10 000 sub-catalogues divided by faults: the Alhama de Murcia Fault (AMF), the Carboneras Fault (CF), the Carrascoy Fault (CAF) and the Bajo Segura Fault (BSF). The “count” represents the number of times the 10 000 sub-catalogues show the corresponding magnitude as the maximum magnitude on the indicated fault.

large earthquakes is possible on the BSF, but less likely than in other faults of the system, such as the AMF and the CF.

4.2 Seismic Hazard Assessment

Hazard curves have been calculated for Murcia, Lorca, Alicante, Vera, Torrevieja and Almería (Fig. 7) using 100 synthetic subcatalogues. We consider a representative sample of the full set of 10 000 catalogues generated. The parameter distributions obtained from the 100 sub-catalogues reproduce the central tendency and dispersion of the reference

set of 10 000 sub-catalogues (Fig. 5d, e, f). No systematic bias is observed in the distributions of the a , b , and maximum magnitude values; therefore, the sample of 100 sub-catalogues is statistically representative of the variability of the dataset. The differences are thus attributable to expected fluctuations in finite samples. This number of catalogues offers a balance between statistical robustness and computational feasibility, ensuring that the variability of seismic behaviour is sufficiently captured while maintaining compatibility with the operational constraints of R-CRISIS. These hazard curves show the variation of return period for the same Peak Ground Acceleration (PGA). The hazard curves obtained using historical-instrumental and ZESIS data show less hazardous scenarios for almost all PGA values. Hazard curves are plotted on a logarithmic scale, which hinders the correct perception of variability. Evaluating specific return periods with seismic design interest (i.e. 100, 500, 1000, 2000, 5000 and 10 000 yr) for the cities of Murcia, Lorca, Alicante, Vera, Torrevieja and Almería (Fig. 8) it can be shown that variability in PGA is not uniform. From one side, for small return periods, the range of values is tiny and variability seems to be modest. On the other side, for return periods longer than 5000 yr, the range of PGA value increases, which translates into higher variability. For large return periods, there is an asymmetry in the dispersion of data with a predominance of higher PGA.

A comparison of the mean hazard curves between cities (Fig. 9a) reveals that shorter return periods are found for the city of Murcia for all values of PGA. In contrast, Alicante is the location where return periods are longer for all PGA with a significant difference. Meanwhile, the cities of Lorca, Vera, Torrevieja and Almería have quite similar values. Figure 9b represents the return period variability coefficient for each PGA at studied sites. What stands out in this




Figure 7. Seismic hazard curves obtained from 100 sub-catalogues of a thousand years build upon a million-year synthetic catalogue (Herrero-Barbero et al., 2021), in grey; upon updated to 2022 catalogue, in red; upon ZESIS database (IGME, 2015), in blue at cities: (a) Murcia, (b) Lorca, (c) Alicante, (d) Vera, (e) Torrevieja y (f) Almería.

figure is that variability presents a rapid decrease for small values of PGA, till a tipping point located at $\text{PGA} = 0.05\text{ g}$ for Murcia, $\text{PGA} = 0.05\text{ g}$ for Lorca, $\text{PGA} = 0.03\text{ g}$ for Alicante, $\text{PGA} = 0.05\text{ g}$ for Vera, $\text{PGA} = 0.04\text{ g}$ for Torrevieja and $\text{PGA} = 0.02\text{ g}$ for Almería. Henceforth, the variability coefficient presents a continual growth that reaches the following values for $\text{PGA} = 1\text{ g}$: 58 %, for Torrevieja; 57 %, for Murcia; 54 % for Vera; 53 %, for Lorca, 40 % for Alicante and 25 %, for Almería. It is notorious that, even though Murcia presents the larger values of PGA, the variability is slightly higher in the case of Torrevieja.

5 Discussion

The main objective of this work is to examine the variability between subcatalogues with analogous characteristics derived from a single synthetic catalogue. This approach ensures that the epistemic uncertainties associated with their generation are common to all subcatalogues, making the differences observed in their comparison independent of those uncertainties. Since our study focuses specifically on the variability introduced by the observation time window, we

have deliberately decided not to address other sources of epistemic uncertainty. These uncertainties have been previously explored in Herrero-Barbero et al. (2021), from which the synthetic seismic catalogue used here originates.

5.1 Implications for fault seismogenic behavior

A matter of particular attention in the discourse around physical earthquake simulators, such as RSQSim (Richards-Dinger and Dieterich, 2012), is the MFD obtained with them. These simulations depend on the frictional properties of fault systems, which are difficult to estimate. They control the generation of large earthquakes and, with that, the configuration of the MFD.

The model shows a decrease in the seismic rate between $M_w = 5.0\text{--}6.0$, suggesting a lack of moderate earthquakes compared to large ones. This scarcity may be related to the lack of secondary faults, such as the Torremendo (Alfaro et al., 2012) or Hinojar (Baena-Escudero, 1993) faults, in the model. It should be noted that this model only represents the main seismogenic sources in the region, which are responsible for the most dangerous earthquakes.




Figure 8. PGA variability for different return periods at the cities of (a) Murcia, (b) Lorca, (c) Alicante, (d) Vera, (e) Torrevieja y (f) Almería. The median is plotted in red; percentiles 25 % and 75 %, in blue; minimum and maximum values, in black and anomalous values as a “+” in red.




Figure 9. Comparison among the cities studied in this work: (a) mean hazard curves and (b) variability coefficient.

The description of the earthquake catalogue using a Gutenberg-Richter relationship does not contemplate the *characteristic earthquake* behaviour in the fault system (Youngs and Coppersmith, 1985; Wesnousky, 1994). The choice between Gutenberg-Richter relationship or the characteristic earthquake model depends on the knowledge of faults and data available (Valentini et al., 2017). The characteristic earthquake model suggests that earthquakes at large

magnitudes occur with a higher frequency than expected if a potential relationship is considered. According to Ben-Zion et al. (2003), the cyclic repetition of large earthquakes can be described as the dynamic regime of intermittent criticality caused by the spontaneous evolution of stress heterogeneities. The statistical distribution of earthquake magnitudes obtained by Ben-Zion et al. (2003) on a mature fault with positive dynamic weakening is consistent with the char-

acteristic earthquake model. The repeated presence of local maxima at high magnitudes in the synthetic MFDs compatible with the dimensions of the coherent brittle zones of faults (Ben-Zion and Rice, 1993) may suggest that EBSZ can also be described as an intermittent criticality dynamic system. Although these large events do not necessarily dominate hazard at engineering return periods in classical PSHA frameworks, their intermittent occurrence strongly influences the variability of hazard estimates when different observation time windows are considered. Large earthquakes in this region could be caused by multi-fault ruptures involving several fault segments which are kinematically compatible (Álvarez-Gómez et al., 2018; Herrero-Barbero et al., 2021; Herrero-Barbero, 2022). The results suggest that these multi-fault ruptures are more common than previously thought. The appearance of a bimodal pattern in the maximum magnitude frequency evaluation with a main maximum (Fig. 5c) at large magnitudes appears to support this assumption. The maximum magnitude count appears at $M_{\max} = 7.2$. According to the simulation, these magnitudes would be associated with large earthquakes nucleated mainly on the Alhama de Murcia Fault (AMF, Fig. 6) and on the Carboneras Fault (CF, Fig. 6).

5.2 Seismic Hazard Assessment

The hazard curves calculated in this study (Fig. 7) show different return periods for each catalogue. This variation results from the different values of b , $M_{\max}$ and $\tau(4.0)$ calculated for each catalogue, reflecting the variability of seismicity over finite time windows. Each 1 kyr catalogue represents a possible realisation of the long-term seismic process, and therefore produces a distinct hazard curve. Consequently, the observed differences highlight that seismic hazard calculation depends on the time interval in which an earthquake catalogue is recorded. The use of 100 catalogues for the calculation of the hazard curves is justified by the similarity in the mean parameters of the Gutenberg-Richter relation in both datasets (Table 4).

The notified variability of the hazard curves (Fig. 8) increases for long return periods and PGA values, both associated with large magnitude earthquakes. For a 500-year return period (Table 5), which is often used in seismic building codes for structures of normal importance, the values obtained for PGA show a range of scenarios that implies significant uncertainties that could affect the definition of seismic actions.

The variability of hazard among cities (Fig. 9a) shows the different influence that the EBSZ faults have on the calculation of seismic hazard in each location. The variability observed for the city of Torrevecija is higher, suggesting that there may be an important dependence between the hazard at Torrevecija and the seismic parameters of the Bajo Segura Fault at the NE extreme of the EBSZ. The return period variability coefficient (Fig. 9b) ranges from 11 % to 21 % for a

Table 4. Statistical parameters for earthquake catalogues using 100 and 10 000 sub-catalogues.

	100 cat.	10 000 cat.
$a_{\min}$	2.45	2.01
$\bar{a}$	4.40	4.55
$a_{\max}$	6.50	14.02
$b_{\min}$	0.66	0.62
$\bar{b}$	1.08	1.11
$b_{\max}$	1.52	2.95
$M_{w,\min}^{\max}$	5.80	5.30
$\overline{M_w^{\max}}$	6.75	6.79
$M_{\max}^{\max}$	7.60	7.60

Table 5. PGA associated to RP = 500 yrs for the cities considered in the PSHA calculations.

	RP = 500 yrs	
	PGA _{min} (g)	PGA _{max} (g)
Murcia	0.147	0.304
Lorca	0.141	0.281
Alicante	0.122	0.218
Vera	0.138	0.280
Torrevecija	0.139	0.301
Almería	0.155	0.234

PGA = 0.04 g, and from 25 % to 58 % for a PGA = 1 g. This range of variability means that the standard deviation is high in relation to the mean hazard curve. This fact illustrates the impact that the use of different seismic catalogues can have on the quantification of seismic hazard at specific locations.

5.2.1 Synthetic vs. historical-instrumental hazard curves

Hazard curves (Fig. 7) obtained using historical and instrumental records, as well as ZESIS database, represent one of the less conservative results for almost all the values of PGA. If return periods are compared (Fig. 10), it can be noticed that the values obtained through synthetic seismicity are, in general, lower than those derived from historical and instrumental data. The lower hazard obtained from historical and instrumental catalogues likely results from a limited sampling of large-magnitude earthquakes occurred within the observation window, making them representative of a less conservative seismic scenario. This would explain the longer return periods compared to those derived from synthetic catalogues. Still, the possibility that the synthetic model overestimates seismic activity rates cannot be entirely ruled out.




Figure 10. Comparison of the return periods obtained in the hazard curves for the synthetic scenarios and the historical-instrumental, HI, one (grey squares) in the cities of: **(a)** Murcia, **(b)** Lorca, **(c)** Alicante, **(d)** Vera, **(e)** Torrevieja and **(f)** Almería. The red line indicates the hypothetical scenario in which the results were identical for both datasets.




Figure 11. Frequency distribution of cumulative seismic moment for 100 sub-catalogues including the historical and instrumental result (in lilac).

The seismic moment released in the different sub-catalogues is calculated (Fig. 11) to assess the similarity between the synthetic scenarios and the real one. The total cumulative seismic moment for the historical and instrumental catalogue is $M_0 = 1.779 \times 10^{19}$ N m. Synthetic values range between $M_{0m} = 9.8 \times 10^{18}$ N m to $M_{0M} = 4.1 \times 10^{20}$ N m in the case of 100 catalogues. The mean cumulative seismic moment for 100 synthetic catalogues is $\overline{M_0} = 8.09 \times 10^{19}$ N m. The median $M_{0med} = 3.2 \times 10^{19}$ N m is slightly

lower than the mean and it is closer to the historical and instrumental value. The histogram of values of the cumulative seismic moment for the synthetic catalogues shows a maximum within the range that contains the cumulative moment of the historical and instrumental catalogue. However, this histogram spreads to values higher than the historical and instrumental one. This finding may suggest that the cumulative seismic moment released as earthquake occurrence could be overestimated. The assumption of total coupling likely inflates the cumulative seismic moment and, consequently, the estimated hazard. Introducing partial coupling or aseismic slip in future models could reduce this bias.

Our findings are consistent with the conclusions of Beauval et al. (2008), highlighting that short observation windows can significantly underestimate seismic hazard. By providing a broader sampling of earthquake variability, our synthetic catalogues underscore the importance of incorporating long-term fault behavior into hazard assessments, especially where the historical record is limited or incomplete.

5.3 Applicability of the model

The methodology applied in this study uses synthetic seismicity as data source. The comprehensive approach of this work, which requires evaluating a large amount of data spread out over time, hinders to replicate the experience with real data. Consequently, the model proposed is subordinated to the constraints associated with synthetic seismicity.

The main limitations are related to the absence of secondary faults, the lack of dynamic interactions and the uncertainties of the data input in the model. For a detailed revision see Herrero-Barbero et al. (2021).

The observed variability is likely not an artifact of the model but rather a reflection of the intrinsic complexity of the fault system's seismic cycle. The EBSZ has been proposed as the selected region for this study because of the wide knowledge that we have of this region. Importantly, the methodology is transferable and could be reproduced on other fault systems. It will be an exercise of great interest to develop the same kind of study in a region with faster slip rates and compare the differences with a region of slower ones as proposed in this study.

6 Conclusions

Seismic hazard estimates depend strongly on the characterization and parametrization of the seismogenic sources model used. When the source model characterization relies entirely on the historical earthquake catalogue, the incompleteness of the data becomes a major limitation, as fault seismic cycles are usually longer. On slow-moving fault systems, such as the EBSZ, the historical and instrumental record only represents a small part of the seismic cycle and the occurrence of large earthquakes could have not been recorded yet. In this study, we have shown how the variability of the time interval of analysis leads to different results in seismic hazard assessment. For the selected calculation sites, the return period for different PGA values showed a strong dependency on the time window in which the catalogue has been recorded. The variation coefficient ranges from 11 % to 21 % for a $\text{PGA} = 0.04 \text{ g}$, and from 25 % to 58 % for a $\text{PGA} = 1 \text{ g}$. Using synthetic seismicity, this study evidences that there is a dependence in the quantification of seismic hazard with the time interval in which the earthquake catalogue spans.

The evaluation of a statistically representative set of 10 000 sub-catalogues reveals the recurrent appearance of large-magnitude events ($M_w > 7.2$) associated with multi-fault ruptures. Although infrequent, these events reflect the long-term seismogenic behaviour emerging from the synthetic fault system, typically nucleating on the AMF and CF, which exhibit the highest slip rates within the EBSZ. Rather than implying a systematic dominance of large earthquakes in seismic hazard at engineering return periods, their presence highlights how rare events may significantly influence hazard variability when different observation time windows sample the seismic cycle unevenly.

At places where the geometry and behaviour of faults are well-known, synthetic models of seismicity can be built with confidence. This approach is especially useful in cases where systems have slow rates of deformation and seismicity is just low or moderate. In these cases, synthetic catalogues can reveal the inner variability of such systems and clarify the

seismogenic potential of faults. Nevertheless, they are constrained to the quality of the geological knowledge available for the active faults present in the region. To better understand the models and their impact on seismic hazard, further investigations should focus on issues such as fault segmentation, structural development, frictional properties, and degree of coupling (aseismic deformation and secondary faulting).

Data availability. The used data can be found at <https://doi.org/10.5281/zenodo.7994106> (Álvarez Gómez et al., 2023) and <https://doi.org/10.5281/zenodo.10286171> (Pascual-Sánchez et al., 2023).

Code availability. The ZMAP software has been developed by Wiemer (2001) and is available at <https://www.seismo.ethz.ch/en/knowledge/earthquake-data-and-analysis-tools/software-archive/> (last access: 11 September 2026). The R-Crisis software has been developed by Ordaz et al. (2021) and is available at <https://www.r-crisis.com/> (last access: 11 September 2026).

Author contributions. Elena Pascual-Sánchez has contributed to the design and implementation of the research, to the analysis of the results and to the writing of the manuscript; José Antonio Álvarez-Gómez has contributed to the design and implementation of the research, to the analysis of the results and to the writing of the manuscript; Julián García-Mayordomo has contributed to the design and implementation of the research, to the analysis of the results and to the writing of the manuscript; Paula Herrero-Barbero has contributed to the design and implementation of the research, to the analysis of the results and to the writing of the manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgements. The authors of this article would like to thank Chung-Han Chan and Francesco Visini for their valuable comments and thorough revision.

Financial support. This study has been supported by the projects Model_SHaKER (grant no. PID2021-124155NBC31) and NSources (grant no. PID2020-119772RB-I00) funded by MCIN/AEI/10.13039/501100011033. E. Pascual-Sánchez thanks the funding support from Programa Investigato of Madrid council

as part of the Recovery, Transformation and Resilience Plan for the development of the Next Generation EU recovery funds. The article processing charges for this open-access publication were covered by the CSIC Open Access Publication Support Initiative through its Unit of Information Resources for Research (URICI).

The article processing charges for this open-access publication were covered by the CSIC Open Access Publication Support Initiative through its Unit of Information Resources for Research (URICI).

Review statement. This paper was edited by Oded Katz and reviewed by Chung-Han Chan and Francesco Visini.

References

- Aki, K.: Maximum likelihood estimate of b in the formula $\log N = a - bM$ and its confidence limits, *Bull. Earthquake Res. Inst., Tokyo Univ.*, 43, 237–239, 1965.
- Alfaro, P., Bartolomé, R., Borque Arancón, M. J., Estévez, A., García Mayordomo, J., García-Tortosa, F. J., Francisco, J., Gil, A. J., Gracia, E., and Lacono, C.: The Bajo Segura fault zone: active blind thrusting in the eastern betic cordillera (SE Spain), *J. Iber. Geol.*, 38, 271–284, https://doi.org/10.5209/rev_jige.2012.v38.n1.39217, 2012.
- Alonso-Henar, J., Fernández, C., and Martínez-Díaz, J. J.: Application of the analytic model of general triclinic transpression with oblique extrusion to an active deformation zone: The Alhama de Murcia Fault (SE Iberian Peninsula), *J. Struct. Geol.*, 130, 103924, <https://doi.org/10.1016/j.jsg.2019.103924>, 2020.
- Álvarez Gómez, J. A., Insúa Arévalo, J. M., Herrero Barbero, P., Martínez Díaz, J. J., Canora Catalán, C., Alonso Henar, J., and García Mayordomo, J.: Potencial de encadenamiento de roturas sísmicas en la zona de cizalla de la Béticas orientales por transferencia de esfuerzos de Coulomb. Reunión Ibérica sobre Fallas Activas y Paleosismología (3^a. 2018. Alicante), <http://hdl.handle.net/10261/277616> (last access: 9 September 2026), 2018.
- Álvarez-Gómez, J. A., Herrero-Barbero, P., and Martínez-Díaz, J. J.: Seismogenic potential and tsunami threat of the strike-slip Carboneras fault in the western Mediterranean from physics-based earthquake simulations, *Nat. Hazards Earth Syst. Sci.*, 23, 2031–2052, <https://doi.org/10.5194/nhess-23-2031-2023>, 2023.
- Álvarez Gómez, J. A., Herrero Barbero, P., and Martínez Díaz, J. J.: Supplementary data for “Seismogenic potential and tsunami threat of the strike-slip Carboneras fault in the western Mediterranean from physics-based earthquake simulations”, Zenodo [data set], <https://doi.org/10.5281/zenodo.7994106>, 2023.
- Anagnos, T. and Kiremidjian, A. S.: A review of earthquake occurrence models for seismic hazard analysis, *Probabilist. Eng. Mech.*, 3, 3–11, 1988.
- Anderson, J. G. and Luco, J. E.: Consequences of slip rate constraints on earthquake occurrence relations, *B. Seismol. Soc. Am.*, 73, 471–496, 1983.
- Argus, D. F., Peltier, W. R., and Watkins, M. M.: Glacial isostatic adjustment observed using very long baseline interferometry & satellite laser ranging geodesy, *J. Geophys. Res.*, 104, 77–93, <https://doi.org/10.1029/1999JB000237>, 1999.
- Baena Escudero, R.: Evolución cuaternaria de la Depresión del Medio-Bajo Guadalquivir y sus márgenes (Córdoba y Sevilla), Tesis doctoral, Universidad de Sevilla (España), 1993.
- Barbat, H. A., Oller Martínez, S. H., and Vielma, J. C.: Cálculo y diseño sismorresistente de edificios: aplicación de la norma NCSE-02, Centre Internacional de Mètodes Numèrics en Enginyeria (CIMNE), <https://upcommons.upc.edu/server/api/core/bitstreams/07b43887-81be-4291-9069-61e5d0d6ae0a/content> (last access: 11 September 2026), 2005.
- Beauval, C., Bard, P. Y., Hainzl, S., and Gueguen, P.: Can strong-motion observations be used to constrain probabilistic seismic hazard estimates?, *B. Seismol. Soc. Am.*, 98, 509–520, <https://doi.org/10.1785/0120070006>, 2008.
- Benedetti, L., Manighetti, I., Gaudemer, Y., Finkel, R., Malavieille, J., Pou, K., Arnold, M., Aumaître, G., Bourlès, D., and Keddadouche, K.: Earthquake synchrony and clustering on Fucino faults (Central Italy) as revealed from in situ ^{36}Cl exposure dating, *J. Geophys. Res.-Solid Earth*, 118, 4948–4974, <https://doi.org/10.1002/jgrb.50299>, 2013.
- Ben-Zion, Y. and Rice, J. R.: Earthquake failure sequences along a cellular fault zone in a three-dimensional elastic solid containing asperity and nonasperity regions, *J. Geophys. Res.-Solid Earth*, 98, 14109–14131, 1993.
- Ben-Zion, Y., Eneva, M., and Liu, Y.: Large earthquake cycles and intermittent criticality on heterogeneous faults due to evolving stress and seismicity, *J. Geophys. Res.*, 108, 4, <https://doi.org/10.1029/2002JB002121>, 2003.
- Bisch, P., Carvalho, E., Degee, H., Fajfar, P., Fardis, M., Franchin, P., Kreslin, K., Pecker, A., Pinto, P., Plumier, A., Somja, H., and Tsionis, G.: Eurocode 8: seismic design of buildings worked examples, Luxembourg: Publications Office of the European Union, <https://doi.org/10.2788/91658>, 2012.
- Booth-Rea, G., Azañón, J.-M., Azor, A., and García-Dueñas, V.: Influence of strike-slip fault segmentation on drainage evolution and topography. A case study: the Palomares Fault Zone (southeastern Betics, Spain), *J. Struct. Geol.*, 26, 1615–1632, 2004.
- Borque, M., Sánchez-Alzola, A., Martín-Rojas, I., Alfaro, P., Molina, S., Rosa-Cintas, S., Rodríguez-Caderot, G., de Lacy, C., García-Armenteros, J. A., Avilés, M., Herrera-Olmo, A., García-Tortosa, F. J., Estévez, A., and Gil, J.: How Much Nubia-Eurasia Convergence Is Accommodated by the NE End of the Eastern Betic Shear Zone (SE Spain)? Constraints From GPS Velocities, *Tectonics*, 38, 1824–1839, <https://doi.org/10.1029/2018TC004970>, 2019.
- Cabañas, L., Rivas-Medina, A., Martínez-Solares, J. M., Gaspar-Escribano, J. M., Benito, B., Antón, R., and Ruiz-Barajas, S.: Relationships between M_w and other earthquake size parameters in the Spanish IGN seismic catalog, *Pure Appl. Geophys.*, 172, 2397–2410, <https://doi.org/10.1007/s00024-014-1025-2>, 2015.
- Campbell, K. W. and Bozorgnia, Y.: NGA-West2 ground motion model for the average horizontal components of PGA, PGV, and 5 % damped linear acceleration response spectra, *Earthquake Spectr.*, 30, 1087–1115, <https://doi.org/10.1193/062913EQS175M>, 2014.
- Canora Catalán, C., Roca, C., Martínez Díaz, J. J., Insúa Arévalo, J. M., Martín González, F., Alonso Henar, J., Gómez Ortiz, D., Martínez-Pagán, P., Masana, E., Ortuño, M., Ferrater Gómez, M., and Medialdea, A.: Nuevos datos de actividad paleosísmica de la falla de Alhama de Murcia en el abanico de La salud (segmento

- Lorca-Totana), Béticas orientales, *Geotemas* (Madrid), 16, 563–566, 2016.
- Cornell, C. A.: Engineering seismic risk analysis, *B. Seismol. Soc. Am.*, 58, 1583–1606, 1968.
- Cornell, C. A. and Vanmarcke, E. H.: The major influences on seismic risk. In *Proceedings of the fourth world conference on earthquake engineering*, Vol. 1, pp. 69–83, 1969.
- Cornell, C. A. and Winterstein, S. R.: Temporal and magnitude dependence in earthquake recurrence models, *B. Seismol. Soc. Am.*, 78, 1522–1537, 1988.
- Cosentino, P., Ficarra, V., and Luzio, D.: Truncated exponential frequency-magnitude relationship in earthquake statistics, *B. Seismol. Soc. Am.*, 67, 1615–1623, 1977.
- Console, R., Vannoli, P., and Carluccio, R.: The seismicity of the Central Apennines (Italy) studied by means of a physics-based earthquake simulator, *Geophys. J. Int.*, 212, 916–929, <https://doi.org/10.1093/gji/ggx451>, 2018.
- Daniell, J. E., Khazai, B., Wenzel, F., and Vervaeck, A.: The worldwide economic impact of historic earthquakes, in: *The 15 World Conference on Earthquake Engineering* (Vol. 15), Lisboa, https://www.researchgate.net/publication/258434390_The_Worldwide_Economic_Impact_of_Earthquakes_Paper_No_2038 (last access: 11 September 2026), 2012.
- De Larouzière, F., Bolze, J., Bordet, P., Hernandez, J., Montenat, C., and d’Estevou, P. O.: The Betic segment of the lithospheric Trans-Alboran shear zone during the Late Miocene, *Tectonophysics*, 152, 41–52, 1988.
- DeMets, C., Gordon, R. G., Argus, D. F., and Stein, S.: Current plate motions, *Geophys. J. Int.*, 101, 425–478, 1990.
- Dieterich, J.: Earthquake simulations with time-dependent nucleation and long range interactions, 9th Proc. Joint Meeting U.J.N.R. Panel Earthq. Predict. Technol., 9, 241–268, 1994.
- Dieterich, J. H.: Modeling of rock friction 1. Experimental results and constitutive equations, *J. Geophys. Res.*, 84, 2161–2168, 1979.
- Echeverría, A., Khazaradze, G., Asensio, E., Gárate, J., Dávila, J. M., and Suriñach, E.: Crustal deformation in eastern Betics from CuaTeNeo GPS network, *Tectonophysics*, 608, 600–612, <https://doi.org/10.1016/j.tecto.2013.08.020>, 2013.
- Echeverría, A., Khazaradze, G., Asensio, E., and Masana, E.: Geodetic evidence for continuing tectonic activity of the Carboneras fault (SE Spain), *Tectonophysics*, 663, 302–309, 2015.
- Esteva, L.: Criterios para la construcción de espectros de diseño sísmico. *Proceedings of the 3rd Pan-American Symposium of Structures*, Caracas, Venezuela, 1967.
- Ferrater, M., Ortuño, M., Masana, E., Pallàs, R., Perea, H., Baize, S., García-Meléndez, E., Martínez-Díaz, J. J., Echeverría, A., Rockwell, T. K., Sharp, W. D., Medialdea, A., and Rhodes, E. J.: Refining seismic parameters in low seismicity areas by 3D trenching: The Alhama de Murcia fault, SE Iberia. *Tectonophysics*, 680, 122–128, <https://doi.org/10.1016/j.tecto.2016.05.020>, 2016.
- Ferrater Gómez, M., Silva, P., Ortuño Candela, M., Rodríguez Pascua, M. Á., and Masana, E.: Archaeoseismologic Analysis of a Late Bronze Age Site on the Alhama de Murcia Fault, SE Spain, *Geoarchaeology, An Int. J.*, 30, 151–164, <https://doi.org/10.1002/gea.21505>, 2015.
- Field, E. H., Biasi, G. P., Bird, P., Dawson, T. E., Felzer, K. R., Jackson, D. D., and Zeng, Y.: Long-term time-dependent probabilities for the third Uniform California Earthquake Rupture Forecast (UCERF3), *B. Seismol. Soc. Am.*, 105, 511–543, <https://doi.org/10.1785/0120140093>, 2015.
- García-Mayordomo, J.: Caracterización y Análisis de la Peligrosidad Sísmica en el Sureste de España. Tesis Doctoral, Universidad Complutense de Madrid, España, <https://produccioncientifica.ucm.es/documentos/5d1df64729995204f7666fde> (last access: 11 September 2026), 2005.
- García-Mayordomo, J.: Incorporación de datos y criterios geológicos en el análisis de la peligrosidad sísmica en regiones de actividad moderadas: 1. Definición y caracterización de fuentes sismogénicas, *Geogaceta*, 41, 87–90, 2007.
- García-Mayordomo, J.: Creación de un modelo de zonas sismogénicas para el cálculo del mapa de peligrosidad sísmica de España. Publicaciones del Instituto Geológico y Minero de España, Serie Riesgos Geológicos/Geotecnia No 5, Madrid, 125 pp., free download: https://web.igme.es/publicaciones/Home/Publicaciones_Gratuitas (last access: 3 September 2026), 2015.
- García-Mayordomo, J. and Martínez-Díaz, J. J.: Caracterización sísmica del anticlinorio del Bajo Segura (Alicante): fallas del Bajo Segura, Torreveja y San Miguel de Salinas, *Geogaceta*, 40, 19–22, 2006.
- García-Mayordomo, J., Gaspar-Escribano, J. M., and Benito, B.: Seismic hazard assessment of the Province of Murcia (SE Spain): analysis of source contribution to hazard, *J. Seismol.*, 11, 453–471, <https://doi.org/10.1007/s10950-007-9064-0>, 2007.
- García-Mayordomo, J., Martín-Alfageme, R., Martín-Banda, R., Insua-Arévalo, J. M., Martínez-Díaz, J. J., Jiménez-Díaz, A., Rodríguez-Escudero, E., Rodríguez-Peces, M. A., Cabañas-Rodríguez, L., and Gaspar-Escribano, J. M.: Seismogenic fault-source characterization in SE Spain: Implications for probabilistic seismic hazard assessment. 15th World Conference of Earthquake Engineering, Lisboa, https://www.iitk.ac.in/nicee/wcee/article/WCEE2012_4044.pdf (last access: 11 September 2026), 2012a.
- García-Mayordomo, J., Insua-Arévalo, J. M., Martínez-Díaz, J. J., Jiménez-Díaz, A., Martín-Banda, R., Martín-Alfageme, S., Álvarez-Gómez, J. A., Rodríguez-Peces, M., Pérez-López, R., and Rodríguez-Pascua, M. A.: The Quaternary active faults database of Iberia (QAFI v. 2.0), *J. Iberian Geol.*, 38, 285–302, https://doi.org/10.5209/rev_JIGE.2012.v38.n1.39219, 2012b.
- García-Mayordomo, J., Martínez-Díaz, J. J., Capote, R., Martín-Banda, R., Insua-Arévalo, J. M., Álvarez-Gómez, J. A., Perea, H., González, Á., Lafuente, P., Martín-González, F., Pérez-López, R., Rodríguez-Pascua, M. A., Giner-Robles, J., Azañón, J. M., Masana, E., Moreno, X., Benito, B., Rivas, A., Gaspar-Escribano, J. M., Cabañas, L., Vilanova, S., Fonseca, J., Nemser, E., and Baize, S.: Modelo de zonas sismogénicas para el cálculo de la peligrosidad sísmica en España. VII Asamblea Hispano Portuguesa de Geodesia y Geofísica, San Sebastián (Spain), 23–28 June 2012, <https://doi.org/10.13140/RG.2.1.4536.5608>, 2012c.
- García-Mayordomo, J., Martín-Banda, R., Insua-Arévalo, J. M., Álvarez-Gómez, J. A., Martínez-Díaz, J. J., and Cabral, J.: Active fault databases: building a bridge between earthquake geologists and seismic hazard practitioners, the case of the QAFI v.3 database, *Nat. Hazards Earth Syst. Sci.*, 17, 1447–1459, <https://doi.org/10.5194/nhess-17-1447-2017>, 2017.

- Gardner, J. K. and Knopoff, L.: Is the sequence of earthquakes in Southern California, with aftershocks removed, Poissonian?, *B. Seismol. Soc. Am.*, 64, 1363–1367, 1974.
- Gokhale, V. A., Joshi, D. R., and Abhayankar, A.: The Psychological and Socio Economic Aspects of Earthquake Occurrence, Vancouver, 13th World Conference on Earthquake Engineering, 2004.
- Gómez Novell, O.: Paleoseismic transect across the Alhama de Murcia Fault and implications of a fault-based seismic hazard assessment for the Eastern Betics, Doctoral thesis, Universitat de Barcelona, España, <https://hdl.handle.net/2445/183755> (last access: 11 September 2026), 2021.
- Gómez-Novell, O., García-Mayordomo, J., Ortuño, M., Masana, E., and Chartier, T.: Fault System-Based Probabilistic Seismic Hazard Assessment of a Moderate Seismicity Region: The Eastern Betics Shear Zone (SE Spain), *Front. Earth Sci.*, 8, 579398, <https://doi.org/10.3389/feart.2020.579398>, 2020.
- Gómez-Novell, O., Visini, F., Pace, B., Álvarez-Gómez, J. A., and Herrero-Barbero, P.: A Benchmarking Method to Rank the Performance of Physics-Based Earthquake Simulations, *Seismol. Res. Lett.*, 96, 231–243, <https://doi.org/10.1785/0220240027>, 2024.
- González, Á.: The Spanish national earthquake catalogue: Evolution, precision and completeness, *J. Seismol.*, 21, 435–471, <https://doi.org/10.1007/s10950-016-9610-8>, 2017.
- Gutenberg, B. and Richter, C. F.: Frequency of earthquakes in California, *B. Seismol. Soc. Am.*, 34, 185–188, 1944.
- Herrero Barbero, P.: Modelización 3D de la estructura, la cinemática y el comportamiento sismogénico del sistema de fallas de las Béticas Orientales: aplicación a la amenaza sísmica. Doctoral dissertation. Universidad Complutense de Madrid, España, <https://hdl.handle.net/20.500.14352/3643> (last access: 11 September 2026), 2022.
- Herrero-Barbero, P., Álvarez-Gómez, J. A., Martínez-Díaz, J. J., and Klimowitz, J.: Neogene basin inversion and recent slip rate distribution of the northern termination of the Alhama de Murcia Fault (Eastern Betic Shear Zone, SE Spain), *J. Geophys. Res.-Tectonics*, 39, e2019TC005750, <https://doi.org/10.1029/2019TC005750>, 2020.
- Herrero-Barbero, P., Álvarez-Gómez, J. A., Williams, C., Villamor, P., Insua-Arévalo, J. M., Alonso-Henar, J., and Martínez-Díaz, J. J.: Physics-based earthquake simulations in slow-moving faults: a case study from the Eastern Betic Shear Zone (SE Iberian Peninsula), *J. Geophys. Res.-Solid Earth*, 126, e2020JB021133, <https://doi.org/10.1029/2020JB021133>, 2021.
- Herrero-Barbero, P., Álvarez-Gómez, J. A., Tsige, M., and Martínez-Díaz, J. J.: Deterministic seismic hazard analysis from physics-based earthquake simulations in the Eastern Betics (SE Iberia), *Eng. Geo.*, 327, 107364, <https://hdl.handle.net/20.500.14352/91475> (last access: 11 September 2026), 2023.
- Instituto Geológico y Minero de España (IGME): ZESIS: Base de Datos de Zonas Sismogénicas de la Península Ibérica y territorios de influencia para el cálculo de la peligrosidad sísmica en España, <https://info.igme.es/zesis> (last access: 3 September 2026), User's Guide: https://info.igme.es/zesis/docs/Guia-usuario-ZESIS_20150421.pdf (last access: 3 September 2026), 2015.
- Instituto Geológico y Minero de España (IGME): QAFI: Quaternary Active Faults Database of Iberia, <https://info.igme.es/QAFI> (last access: 3 September 2026), 2022.
- IGN: The Spanish seismic catalog, Catálogo Sísmico Nacional, IGN – Instituto Geográfico Nacional, <https://www.ign.es/web/ign/portal/sis-catalogo-terremotos> (last access: 3 September 2026), 2019.
- Instituto Geográfico Nacional (IGN): (04 de octubre de 2022) Catálogo de Terremotos, Instituto Geográfico Nacional, Centro Nacional de Información Geográfica (CNIG): Madrid, <https://doi.org/10.7419/162.03.2022>, 2022.
- Instituto Geográfico Nacional (IGN): Universidad Politécnica de Madrid (UPM): Actualización de Mapas de Peligrosidad Sísmica de España 2012. Centro Nacional de Información Geográfica, Madrid, 267 pp., <https://www.ign.es/web/libros-digitales/peligrosidad-sismica2012> (last access: 3 September 2026), 2013.
- Insua Arévalo, M., García Mayordomo, J., Salazar, Á., Rodríguez-Escudero, E., Martín Banda, R., Álvarez Gómez, J. A., Catalán, C., and Martínez-Díaz, J. J.: Paleoseismological evidence of Holocene activity of the los Tollos Fault (Murcia, SE Spain): a lately formed quaternary tectonic feature of the eastern betic shear zone, *J. Iberian Geol.*, 41, 333–350, https://doi.org/10.5209/rev_jige.2015.v41.n3.49948, 2015.
- Kotha, S. R., Bindi, D., and Cotton, F.: From ergodic to region-and site-specific probabilistic seismic hazard assessment: Method development and application at European and Middle Eastern sites, *Earthquake Spectra*, 33, 1433–1453, <https://doi.org/10.1193/081016eqs130m>, 2017.
- López-Comino, J. Á., Mancilla, F. D. L., Morales, J., and Stich, D.: Rupture directivity of the 2011, Mw 5.2 Lorca earthquake (Spain), *Geophys. Res. Lett.*, 39, <https://doi.org/10.1029/2011GL050498>, 2012.
- Matsumoto, M. and Nishimura, T.: Mersenne twister: a 623-dimensionally equidistributed uniform pseudo-random number generator, *ACM Transactions on Modeling and Computer Simulation (TOMACS)*, 8, 3–30, 1998.
- Marano, K. D., Wald, D. J., and Allen, T. I.: Global earthquake casualties due to secondary effects: a quantitative analysis for improving rapid loss analyses, *Nat. Hazards*, 52, 319–328, <https://doi.org/10.1007/s11069-009-9372-5>, 2010.
- Martín-Banda, R.: Segmentación y evolución reciente del Sistema de Fallas de la Sierra de Carrascoy: Implicaciones en su potencial sismogénico. Doctoral dissertation. Universidad Complutense, <https://hdl.handle.net/20.500.14352/11663> (last access: 11 September 2026), 2020.
- Martín-Banda, R., García-Mayordomo, J., Insua-Arévalo, J. M., Salazar, Á. E., Rodríguez-Escudero, E., Álvarez-Gómez, J. A., Medialdea, A., and Herrero, M. J.: New insights on the seismogenic potential of the Eastern Betic Shear Zone (SE Iberia): Quaternary activity and paleoseismicity of the SW segment of the Carrascoy Fault Zone, *Tectonics*, 35, 55–75, <https://doi.org/10.1002/2015TC003997>, 2016.
- Martín-Banda, R., Insua-Arévalo, J. M., and García-Mayordomo, J.: Slip rate variation during the last ≈ 210 ka on a slow fault in a transpressive regime: the Carrascoy Fault (Eastern Betic Shear Zone, SE Spain), *Front. Earth Sci.*, 8, <https://doi.org/10.3389/feart.2020.599608>, 2021.
- Martínez-Díaz, J. J.: Neotectónica y Tectónica Activa del oeste de Murcia y sur de Almería (Cordillera Bética). Doctoral dissertation.

- tion, Universidad Complutense de Madrid, <https://docta.ucm.es/entities/publication/ec6cc99d-a1aa-4eba-8893-3864d855ec50> (last access: 11 September 2026), 1998.
- Martínez-Díaz, J. J., Bejar-Pizarro, M., Álvarez-Gómez, J. A., Mancilla, F. d. L., Stich, D., Herrera, G., and Morales, J.: Tectonic and seismic implications of an intersegment rupture, *Tectonophysics*, 546–547, 28–37, 2012a.
- Martínez Díaz, J. J., Masana, E., and Ortuño, M.: Active tectonics of the Alhama de Murcia fault, Betic Cordillera, Spain, *J. Iberian Geol.*, 38, 253–270, https://doi.org/10.5209/rev_JIGE.2012.v38.n1.39218, 2012b.
- Martínez-Díaz, J. J., Alonso-Henar, J., Insua-Arévalo, J. M., Canora, C., García-Mayordomo, J., Rodríguez-Escudero, E., Álvarez-Gómez, J. A., Ferrater, M., Ortuño, M., and Masana, E.: Geological evidences of surface rupture related to a seventeenth century destructive earthquake in Betic Cordillera (SE Spain): constraining the seismic hazard of the Alhama de Murcia fault, *J. Iberian Geol.*, 45, 73–86, <https://doi.org/10.1007/s41513-018-0082-2>, 2019.
- Masana, E., Pallàs, R., Perea, H., Ortuño, M., Martínez-Díaz, J. J., García-Meléndez, E., and Santanach, P.: Large Holocene morphogenic earthquakes along the Albox Fault, Betic Cordillera, Spain, *J. Geodynam.*, 40, 119–133, <https://doi.org/10.1016/j.jog.2005.07.002>, 2005.
- Masana, E., Gràcia, E., Moreno, X., Bartolomé, R., and Dañobeitia, J. J.: Characterizing the seismic potential of the Eastern Betics Shear Zone (EBSZ), a major source of earthquakes in Southeastern Iberia. *Resúmenes de la 1^a Reunión Ibérica sobre Fallas Activas y Paleosismología*, Sigüenza, España, 101–104, <http://hdl.handle.net/10261/82453> (last access: 11 September 2026), 2010.
- Masana, E., Moreno, X., Gràcia, E., Pallàs, R., Ortuño, M., López, R., Gómez-Novell, O., Ruano, R., Perea, H., Stepancikova, P., and Khazaradze, G.: First evidence of paleoearthquakes along the Carboneras fault zone (SE Iberian Peninsula): Los trances site, *Geol. Acta*, 16, 461–476, <https://doi.org/10.1344/GeologicaActa2018.16.4.8>, 2018.
- McGuire, R. K.: Probabilistic seismic hazard analysis: Early history, *Earthq. Eng. Struct. D.*, 37, 329–338, <https://doi.org/10.1002/eqe.765>, 2008.
- Moreno, X.: Neotectonic and Paleoseismic Onshore-Offshore integrated study of the Carboneras Fault (Eastern Betics, SE Iberia). *Estudio integrado tierra-mar de la Neotectónica y Paleosismología de la Falla de Carboneras (Béticas Orientales, SE Península Ibérica)*. Tesis doctoral, Universitat de Barcelona (España), <https://hdl.handle.net/2445/34931> (last access: 11 September 2026), 2011.
- Moreno, X., Masana, E., Pallàs, R., Gràcia, E., Rodés, Á., and Bordonau, J.: Quaternary tectonic activity of the Carboneras Fault in the La Serrata range (SE Iberia): Geomorphological and chronological constraints, *Tectonophysics*, 663, 78–94, 2015.
- Murphy, P.: Los terremotos de Almería de 1804. En el archivo histórico nacional. Instituto Geográfico Nacional, <https://www.ign.es/web/libros-digitales/terremotos-almeria-1804> (last access: 11 September 2026), 2019.
- Ordaz, M. and Salgado-Gálvez, M. A.: R-CRISIS v20 Validation and Verification Document, ERN Technical Report, Mexico City, Mexico, 2020.
- Ordaz, M., Salgado-Gálvez, M. A., and Giraldo, S.: R-CRISIS: 35 years of continuous developments and improvements for probabilistic seismic hazard analysis, *B. Earthq. Eng.*, 19, 2797–2816, <https://doi.org/10.1007/s10518-021-01098-w>, 2021.
- Ortuño, M., Masana, E., García-Meléndez, E., Martínez-Díaz, J., Štěpančíková, P., Cunha, P. P., and Murray, A. S.: An exceptionally long paleoseismic record of a slow-moving fault: The Alhama de Murcia fault (Eastern Betic shear zone, Spain), *GSA Bull.*, 124, 1474–1494, <https://doi.org/10.1130/B30558.1>, 2012.
- Pascual-Sánchez, E., García-Mayordomo, J., and Álvarez-Gómez, J. A.: Catálogo sísmico actualizado para el cálculo de la peligrosidad sísmica en España, IGN/UPM/UCM/IGME, Zenodo [data set], <https://doi.org/10.5281/zenodo.10286171>, 2023.
- Perea, H., Masana, E., and Santanach, P.: A pragmatic approach to seismic parameters in a region with low seismicity: The case of eastern Iberia, *Nat. Hazards*, 39, 451–477, <https://doi.org/10.1007/s11069-006-0013-y>, 2006.
- Perea, H., Masana, E., and Santanach, P.: An active zone characterized by slow normal faults, the northwestern margin of the Valencia trough (NE Iberia): a review. *Journal of Iberian Geology*, Universidad Complutense de Madrid (UCM), https://doi.org/10.5209/rev_jige.2012.v38.n1.39204, 2012.
- Peruzza, L., Pace, B., and Visini, F.: Fault-based earthquake rupture forecast in Central Italy: Remarks after the L'Aquila M w 6.3 event, *B. Seismol. Soc. Am.*, 101, 404–412, <https://doi.org/10.1785/0120090276>, 2011.
- Reicherter, K. and Hübscher, C.: Evidence for a seafloor rupture of the Carboneras Fault Zone (southern Spain): Relation to the 1522 Almería earthquake?, *J. Seismol.*, 11, 15–26, <https://doi.org/10.1007/s10950-006-9024-0>, 2007.
- Richards-Dinger, K. and Dieterich, J. H.: RSQSim Earthquake Simulator, *Seismol. Res. Lett.*, 83, 983–990, <https://doi.org/10.1785/0220120105>, 2012.
- Rivas-Medina, A., Benito, B., and Gaspar-Escribano, J. M.: Approach for combining fault and area sources in seismic hazard assessment: application in south-eastern Spain, *Nat. Hazards Earth Syst. Sci.*, 18, 2809–2823, <https://doi.org/10.5194/nhess-18-2809-2018>, 2018.
- Robinson, R., Van Dissen, R., and Litchfield, N.: Using synthetic seismicity to evaluate seismic hazard in the Wellington region, New Zealand, *Geophys. J. Int.*, 187, 510–528, <https://doi.org/10.1111/j.1365-246X.2011.05161.x>, 2011.
- Rodríguez-Escudero, E., Martínez-Díaz, J. J., Álvarez-Gómez, J. A., Insua-Arévalo, J. M., and Capote del Villar, R.: Tectonic setting of the recent damaging seismic series in the Southeastern Betic Cordillera, Spain. *Bull. Earthq. Eng.*, 12, 1831–1854, <https://doi.org/10.1007/s10518-013-9551-3>, 2014.
- Roquero, E., Silva, P. G., Rodríguez-Pascua, M. A., Bardají, T., Elez, J., Carrasco-García, P., and Giner-Robles, J. L.: Analysis of faulted fan surfaces and paleosols in the Palomares Fault Zone (Betic Cordillera, SE Spain): Paleoclimatic and paleoseismic implications, *Geomorphology*, 342, 88–102, <https://doi.org/10.1016/j.geomorph.2019.06.003>, 2019.
- Rundle, P., Rundle, J., Tiampo, K., Donnellan, A., and Turcotte, D.: Virtual California: fault model, frictional parameters, applications, *Pure Appl. Geophys.*, 163, 1819–1846, https://doi.org/10.1007/978-3-7643-7992-6_7, 2006.

- Sandri, L. and Marzocchi, W.: A technical note on the bias in the estimation of the b -value and its uncertainty through the least squares technique, *Ann. Geophys.*, 50, 329–339, 2007.
- Sanz de Galdeano, C., López Garrido, Á. C., and García Tortosa, F. J.: Nuevos datos para la estimación de los valores de levantamiento desde el Tortonense Superior a la actualidad en la parte centro-occidental de la Sierra de Carrascoy (provincia de Murcia), *Geogaceta*, 23, 139–142, <http://hdl.handle.net/10261/29336> (last access: 11 September 2026), 1998.
- Savage, J. C.: A dislocation model of strain accumulation and release at a subduction zone, *J. Geophys. Res.-Solid Earth*, 88, 4984–4996, 1983.
- Serpelloni, E., Vannucci, G., Pondrelli, S., Argnani, A., Casula, G., Anzidei, M., Baldi, P., and Gasperini, P.: Kinematics of the Western Africa-Eurasia plate boundary from focal mechanisms and GPS data, *Geophys. J. Int.*, 169, 1180–1200, <https://doi.org/10.1111/j.1365-246X.2007.03367.x>, 2007.
- Shi, X., Kirby, E., Lu, H., Robinson, R., Furlong, K. P., and Wang, E.: Holocene slip rate along the Gyaring Co Fault, central Tibet, *Geophys. Res. Lett.*, 41, 5829–5837, <https://doi.org/10.1002/2014GL060782>, 2014.
- Sieh, K., Natawidjaja, D. H., Meltzner, A. J., Shen, C. C., Cheng, H., Li, K. S., Suwargadi, B. W., Galetzka, J., Philibosian, B., and Edwards, R. I.: Earthquake supercycles inferred from sea-level changes recorded in the corals of West Sumatra, *Science*, 322, 1674–1678, <https://doi.org/10.1126/science.1163589>, 2008.
- Silva, P. G.: Evolución geodinámica de la depresión del Guadalentín (Murcia) desde el mioceno superior hasta la actualidad: Neotectónica y geomorfología, Doctoral dissertation, Universidad Complutense de Madrid, <https://produccioncientifica.ucm.es/documentos/5d1ffb272999521e412dd4f3> (last access: 11 September 2026), 1994.
- Silva, P. G., Goy, J., Zazo, C., and Bardají, T.: Fault-generated mountain fronts in southeast Spain: geomorphologic assessment of tectonic and seismic activity, *Geomorphology*, 50, 203–225, 2003.
- Silva, P. G., Bardají, T., Roquero García-Casal, E., Martínez-Graña, A. M., Perucha, M. A., Huerta Hurtado, P., Lario, J., Giner-Robles, J. L., Rodríguez-Pascua, M. A., Pérez-López, R., Cabero, A., Goy, J. L., and Zazo, C.: Paleogeography and paleoseismicity. The ad 1048 Orihuela earthquake case study (lower segura depression, SE Spain). Una visión global del Cuaternario: El hombre como condicionante de procesos geológicos, 198–203 pp., JP Galve, <http://hdl.handle.net/10261/250060> (last access: 11 September 2026), 2015.
- Spanish Statistical Office (SSO) (14 de noviembre de 2022) Demografía y población, Padrón, INEbase, https://www.ine.es/dyngs/INEbase/es/categoria.htm?c=Estadistica_P&cid=1254734710990 (last access: 14 November 2022), 2022.
- Stich, D., Ammon, C. H., and Morales, J.: Moment tensor solutions for small and moderate earthquakes in the Ibero-Maghreb region, *J. Geophys. Res.*, 108, 2148, <https://doi.org/10.1029/2002JB002057>, 2003.
- Stirling, M. W., McVerry, G. H., and Gerstenberger, M.: New National Seismic Hazard Model for New Zealand: Changes to Estimated Long-Term Hazard, in: *Proc. 12th New Zealand society for earthquake engineering conference*, 13–15, 2012.
- Utsu, T.: read at the meeting of the Seismological Society of Japan, October, 1964.
- Utsu, T.: A statistical significance test of the difference in b -value between two earthquake groups, *J. Phys. Earth*, 14, 37–40, <https://doi.org/10.4294/jpe1952.14.37>, 1966.
- Valentini, A., Visini, F., and Pace, B.: Integrating faults and past earthquakes into a probabilistic seismic hazard model for peninsular Italy, *Nat. Hazards Earth Syst. Sci.*, 17, 2017–2039, <https://doi.org/10.5194/nhess-17-2017-2017>, 2017.
- Ward, S.: San Francisco Bay Area earthquake simulations: a step toward a standard physical earthquake model, *B. Seismol. Soc. Am.*, 90, 370–386, <https://doi.org/10.1785/0119990026>, 2000.
- Wesnowsky, S. G.: The Gutenberg-Richter or characteristic earthquake distribution, which is it?, *B. Seismol. Soc. Am.*, 84, 1940–1959, <https://doi.org/10.1785/BSSA0840061940>, 1994.
- Wiemer, S.: A software package to analyze seismicity: ZMAP, *Seismol. Res. Lett.*, 72, 373–382, <https://doi.org/10.1785/gssrl.72.3.373>, 2001.
- Wiemer, S. and Wyss, M.: Minimum magnitude of completeness in earthquake catalogs: Examples from Alaska, the western United States, and Japan, *B. Seismol. Soc. Am.*, 90, 859–869, 2000.
- Wilson, J. M., Yoder, M. R., Rundle, J. B., Turcotte, D. L., and Schultz, K. W.: Spatial evaluation and verification of earthquake simulators. *Earthquakes and Multi-hazards Around the Pacific Rim*, Vol. I. Pageoph Topical Volumes, 174, 2279–2293, https://doi.org/10.1007/978-3-319-71565-0_7, 2017.
- Yazdi, P. and García-Mayordomo, J.: Active fault interaction in the Eastern Betic Cordillera: A model of coseismic and postseismic stress transfer following historical earthquakes in SE Spain, *Tectonics*, 43, e2024TC008383, <https://doi.org/10.1029/2024TC008383>, 2024.
- Yazdi, P., García-Mayordomo, J., Álvarez-Gómez, J. A., Gaspar-Escribano, J. M., and Masana, E.: Exploring the connection of XVI-century major historical earthquakes in the Eastern Betic Cordillera, Spain: Insights from viscoelastic relaxation of the lithosphere, *Tectonics*, 42, e2023TC007917, <https://doi.org/10.1029/2023TC007917>, 2023.
- Youngs, R. R. and Coppersmith, K. J.: Implications of fault slip rates and earthquake recurrence models to probabilistic seismic hazard estimates, *B. Seismol. Soc. Am.*, 75, 939–964, 1985.