



**High streamflow  
forecasts 1–3 months  
in advance**

J. C. Bennett et al.

# The challenge of forecasting high streamflows in medium sized catchments 1–3 months in advance

J. C. Bennett<sup>1</sup>, Q. J. Wang<sup>1</sup>, P. Pokhrel<sup>1,\*</sup>, and D. E. Robertson<sup>1</sup>

<sup>1</sup>CSIRO Land and Water, Graham Road, Highett, Victoria 3190, Australia

\* now at: Entura, 89 Cambridge Park Drive, Cambridge, Tasmania 7170, Australia

Received: 19 June 2013 – Accepted: 25 June 2013 – Published: 8 July 2013

Correspondence to: J. C. Bennett (james.bennett@csiro.au)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Skilful forecasts of high streamflows a month or more in advance are likely to be of considerable benefit to emergency services and the broader community. This is particularly true for small-medium sized catchments ( $< 2000 \text{ km}^2$ ), where real-time warning systems are only able to give short notice of impending floods. In this study, we generate forecasts of high streamflows for the coming 1 month and coming 3 month periods using large-scale ocean/atmosphere climate indices and catchment wetness as predictors. Forecasts are generated with a combination of Bayesian joint probability modeling and Bayesian model averaging. High streamflows are defined as maximum single-day streamflows and maximum 5 day streamflows that occur during each 1 month or 3 month forecast period. Skill is clearly evident in the 1 month forecasts of high streamflows. Surprisingly, in several catchments positive skill is also evident in forecasts of large threshold events (exceedance probabilities of 25 %) over the next month. Little skill is evident in forecasts of high streamflows for the 3 month period. We show that including climate indices as predictors adds little skill to the forecasts, and thus catchment wetness is by far the most important predictor. Accordingly, we recommend that forecasts may be improved by using accurate estimates of catchment wetness.

## 1 Introduction

Skilful forecasts of high streamflows a month or more in advance have the potential to improve the management of floods. Flood warnings in Australia are presently derived from event-based forecast models that use real-time streamflow and rainfall observations to forecast floods with typical lead-times from hours to a few days, depending on flood travel time (Elliott et al., 2005). Real-time forecasts offer precise estimates of flood heights, but are only available around the time of the flood itself. This leaves emergency services a narrow window to prepare themselves and the community to mitigate flood impacts, particularly in small to medium sized catchments where flood

**NHESSD**

1, 3129–3168, 2013

### High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



warning systems can only give warning of floods from hours to one or two days in advance of an event. Ill-preparedness for floods can have serious implications. Pfister (2002) identified poor community preparedness to evacuate as the major cause of citizens' slow (and non-existent) responses to a flood evacuation order issued by emergency services. Australian emergency services rely heavily on volunteers for disaster response (Baxter-Tomkins and Wallace, 2009), and ensuring that sufficient volunteer-labour is available during emergencies is a challenge for flood-response agencies like the State Emergency Services (SES). Medium range forecasts (to forecast horizons of 3 months) of high streamflows are needed to enable both emergency services and the community to be better prepared for floods.

This study is a response to a request from the Australian Bureau of Meteorology to explore the skill of high streamflow forecasts at medium range forecast horizons. The Bureau of Meteorology is the lead agency for flood warnings in Australia, and emergency services are important users of these flood warnings. While medium range forecasts of high streamflows cannot hope to be as precise as real-time flood models, forewarning of conditions that could result in large or frequent flooding in the next month or more could allow emergency services to better plan and prepare for the impacts of floods, for example by informing volunteer emergency services personnel of heightened flood risk in the coming month(s).

Several studies have described teleconnections between Australian runoff variability and large-scale oceanic and atmospheric climate indices (hereafter, *climate indices*), particularly climate indices describing the El Niño Southern Oscillation (ENSO) (Chiew et al., 1998; Verdon et al., 2004; Schepen et al., 2012). These teleconnections have been used to produce forecasts or total seasonal streamflows that are skilful relative to forecasts derived from streamflow climatologies (Wang et al., 2009; Piechota et al., 1998; Sharma, 2000). Flood risk in south-eastern Australia has also been linked to ENSO (Kiem et al., 2003), but despite this no attempt has yet been made to use such a teleconnection to forecast high streamflows in Australia. Attempts to forecast high streamflows a month or more in advance are rarely reported for other continents, and

the examples that exist often focus on catchments where snowmelt makes a large contribution to seasonal floods (e.g. Kwon et al., 2009; Lindström and Olsson, 2011). Seasonal snow-melt is rarely an important feature of Australian rivers, and accordingly forecasts that rely on indicators of snow-melt have limited application in Australia.

5 The aim of this study is to apply a statistical technique, the Bayesian joint probability modelling approach (BJP), to the problem of forecasting high streamflows in medium sized catchments over the coming 1 month and 3 month periods. The BJP was developed to forecast seasonal total volumes of streamflows (Wang et al., 2009; Wang and Robertson, 2011; Robertson and Wang, 2012) and is now used operationally by  
10 the Bureau of Meteorology to issue forecasts for more than 70 sites across Australia (forecasts available at <http://www.bom.gov.au/water/ssf/>). The BJP produces probabilistic streamflow forecasts that are more accurate than climatology, and, importantly, it is able to reliably estimate uncertainty in the streamflow forecasts. Knowledge of the amount of water held in storage in a catchment (in the soil, as ground water, in surface stores, or as snow/ice – collectively, *catchment wetness*) often contributes more skill  
15 to next-month/next-season forecasts of streamflow than climate forecasts (Shukla and Lettenmaier, 2011; Li et al., 2009). The BJP is able to use multiple predictors to generate forecasts, meaning forecasts can be constructed from both catchment wetness and predictors of climate. For example, Wang et al. (2009) used the BJP to pair the  
20 initial catchment wetness with the southern oscillation index (SOI) to forecast seasonal streamflow totals.

A number of sets of predictors can be used to construct different forecast models, and forecasts can be improved by selecting models with the best predictive power (Robertson and Wang, 2012) or by weighting models according to predictive power (Wang et al., 2012a). Wang et al. (2012a) showed that Bayesian model averaging (BMA) out-  
25 performed predictor selection methods for merging rainfall forecast models generated with the BJP. In addition, predictor selection can lead to artificially inflated estimates of cross-validation skill if the predictor selection is not included in the cross-validation

---

**High streamflow forecasts 1–3 months in advance**J. C. Bennett et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(DelSole and Shukla, 2009; Robertson and Wang, 2013), a problem that is not present with the BMA method we use in this study.

Our study aims to test the ability of the BJP to forecast high streamflows up to three months in advance. To achieve this, we build a set of forecast models with the BJP by combining an estimate of initial catchment wetness with a suite of climate indices derived from oceanic and atmospheric variables. We combine the models with the BMA method described by Wang et al. (2012a) to maximise predictive power.

We next describe the study sites and give an overview of the forecast models. This is followed by descriptions of the verification measures we use to demonstrate the reliability and skill of the forecasts. We present the reliability and skill of these forecasts, and discuss the prospects for improving long lead forecasts of high streamflows. We conclude with a summary of the paper.

## 2 Data and methods

### 2.1 Study sites

Forecasts are generated for six catchments in south-eastern Australia shown in Fig. 1. Characteristics of the six catchments are summarised in Table 1 and Fig. 2. The catchments are selected as they have long (> 40 yr) streamflow records, are free of diversions or impoundments, and are minimally impacted by human activities. All the catchments are what we describe as “medium sized”, with drainage areas between 1000 km<sup>2</sup> and 2000 km<sup>2</sup>. The catchments are large enough to minimise the influence of highly localised storms (e.g. localised convective storms) on the streamflow records. Conversely, catchments are small enough so that flood travel times extend no more than two days, making it difficult to get advance warning of floods of more than two days with an event-based forecasting model. The catchments span a range of climate and hydrological conditions. Streamflows in the two north-eastern catchments, the Orara River (ORB) and the Nowendoc River (NOR), are only weakly seasonal, with the high-

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



est streamflows occurring in February and March (Fig. 2). The remaining catchments – Abercrombie River (ABH), Murray River (MUR), Mitta Mitta River (MMH) and Tarwin River (TAW) – have more strongly seasonal streamflow regimes, with high streamflows in the austral winter/spring, and low streamflows in the austral summer (Fig. 2). High-elevation areas in the MUR and MMH catchments often receive snowfalls in the Austral winter. However, even in these two catchments the contribution of seasonal snowmelt to streamflows is relatively small.

## 2.2 Forecast model

### 2.2.1 Overview

Forecasts are generated on the last day of each month for two periods: the coming month (January, February, . . . , December), and the coming three months (JFM, FMA, . . . , DJF). We refer to these as 1 month and 3 month forecast periods.

Figure 3 gives a schematic overview of how forecasts are generated. Thirteen forecast models are generated with the BJP method (Fig. 3a) for each forecast period and for each predictand. Forecasts from these individual models are then merged using BMA (Fig. 3b). We now describe the components shown in Fig. 3 in detail.

### 2.2.2 Predictands

We investigate two predictands to represent high streamflows:

1. The maximum 1 day streamflow for each forecast period (Max1D).
2. The maximum 5 day aggregated streamflow calculated for each forecast period (Max5D).

Neither Max5D nor Max1D is necessarily a large flood; for example, in the catchments with strongly seasonally delineated streamflows, Max5D streamflows in summer can

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



be very low compared to Max5D winter streamflows. In low streamflow months, medians of both Max1D and Max5D streamflows are sometimes not much larger than average monthly streamflows (Fig. 2). For this reason, we also evaluate the performance of the forecasts in terms of probabilities of events exceeding larger thresholds (see Sect. 2.3.3).

The BJP is able to generate forecasts jointly for multiple predictands. In addition to either Max1D or Max5D, we also include total rainfall for the forecast period as a predictand. Previous work has shown that using rainfall as a joint predictand with streamflow can elicit more useful information from the climate indices (Robertson and Wang, 2012). Rainfalls are aggregated to catchments from the Australian water availability project (AWAP) gridded dataset (Jones et al., 2009).

### 2.2.3 Predictors

We use catchment wetness and climate indices as predictors of high streamflows. We approximate catchment wetness with total streamflow in the previous month for both 1 month and 3 month forecast periods. Total streamflow can be a somewhat coarse measure of catchment wetness, and takes no account of differences in catchment wetness stores (e.g. snow). However, using total streamflow as an estimate of catchment wetness has the virtue of simplicity, and is adequate for this exploratory study.

Eleven lagged climate indices are evaluated as potential predictors in this study, and these are listed in Table 2. We select these climate indices as they have been linked to rainfall in south-east Australia. The teleconnection between south-east Australian rainfall and ENSO has been extensively described (e.g. Schepen et al., 2012; Chiew et al., 1998; Wang et al., 2009) including, as already noted, the link between flooding and ENSO (Kiem et al., 2003). We use five indices to describe ENSO: NINO3, NINO3.4, NINO4, the ENSO Modoki index (EMI) (Ashok et al., 2007) and the southern oscillation index (SOI) (Troup, 1965). The influence of Indian Ocean sea surface temperatures has also been linked to rainfall in south eastern Australia, with the teleconnection being most evident in winter months (Verdon and Franks, 2005; Schepen et al., 2012; Ashok

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2003). We use four Indian Ocean indices as predictors: the Indian Ocean west pole index (WPI), east pole index (EPI) and dipole mode index (DMI) (Saji et al., 1999), as well as the Indonesia index (II) (Verdon and Franks, 2005). Finally, extra-tropical sea surface temperatures and atmospheric features along Australia's east coast have been linked to south east Australian rainfall (Murphy and Timbal, 2008; Risbey et al., 2009; Pook et al., 2006). We use the Tasman Sea index (TSI) (Murphy and Timbal, 2008) and an index of atmospheric blocking (BI140) (Risbey et al., 2009) to represent extra-tropical climatic features. We note that Schepen et al. (2012) found no evidence of a relationship between lagged B140 and TSI with mean rainfall in any season. While it is unlikely that lagged TSI or B140 will contribute skill to high streamflow forecasts, we have included them in case they have a relationship with high rainfall events. Atmospheric blocking, for example, has been correlated with larger rain storms (Pook et al., 2006).

Mean monthly values of each climate index for the previous month are used for both 1 month and 3 month forecasts. For example, for a 1 month forecast for June we use catchment wetness and NINO3 calculated for May as predictors, while for a 3 month forecast for January-February-March we use predictors calculated for December.

Catchment wetness is combined with each of the 11 climate indices to create 11 forecast models for each predictand and for each forecast period. In addition, one forecast model is developed using only catchment wetness as a predictor, and one forecast model is developed based only on climatology (using no predictors). This gives a total of 13 forecast models for each predictand and for each forecast period.

While the effect of snow on the two alpine catchments (MUR and MMH) is expected to be small, we investigated the use of snow accumulation as a predictor for these two snow-affected catchments. Including snow accumulation as a predictor in these two catchments resulted in no increase in forecast skill and is not presented here.

# NHESSD

1, 3129–3168, 2013

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## 2.2.4 Bayesian joint probability modelling

The BJP is used to generate the 13 individual forecast models for each predictand and each forecast period (Fig. 3a), which we call *BJP forecast models*. Detailed mathematical formulations of the BJP are given by Wang et al. (2009), Wang and Robertson (2011) and Robertson and Wang (2012). In summary, the BJP is implemented as follows:

1. Predictands and predictors are transformed to normalise their distributions and stabilise their variances. Streamflow and rainfall are transformed with a log-sinh transform (Wang et al., 2012b), and climate indices are transformed with the Yeo–Johnson transform (Yeo and Johnson, 2000).
2. We assume that the set of transformed predictors and predictands can be described by a joint probability distribution – in this case a multivariate normal distribution.
3. The parameters of the log-sinh transform, the Yeo–Johnson transform, and the multivariate normal distribution are inferred jointly. Parameter inference is performed with Bayesian methods and Markov chain Monte Carlo (MCMC) sampling. Taken together, the parameters of the log-sinh transform, the Yeo–Johnson transform and the multivariate normal distribution define the statistical relationship between predictors and predictands, and allow us to generate forecasts.

Mathematically, if predictors are given by vector  $\mathbf{y}(1)$  and predictands by vector  $\mathbf{y}(2)$ , the probabilistic forecast is given by

$$f[\mathbf{y}(2)|\mathbf{y}(1)] = p[\mathbf{y}(2)|\mathbf{y}(1); Y_{\text{OBS}}, M] = \int p[\mathbf{y}(2)|\mathbf{y}(1); \theta] \cdot p[\theta|Y_{\text{OBS}}, M] \cdot d\theta \quad (1)$$

where  $M$  is the model used, and  $Y_{\text{OBS}}$  contains the historical data of both the predictors and the predictands used for model inference.  $\theta$  is the vector of parameters for the log-sinh transform, the Yeo–Johnson transform, and the multivariate normal distribution.

# NHESSD

1, 3129–3168, 2013

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.2.5 Bayesian model averaging

Forecasts from the thirteen BJP forecast models are merged with BMA to produce one *BJP-BMA forecast* for each predictand and for each forecast period (Fig. 3b). The BMA method we use is described in detail by Wang et al. (2012a). For a set of models

5  $M_k, k = 1, 2, \dots, K$ , each model is assigned a weight,  $w_k$ . The forecasts are then merged by:

$$f_{\text{BMA}}(\mathbf{y}(2)|\mathbf{y}(1)) = \sum_{k=1}^K w_k f_k(\mathbf{y}(2)|\mathbf{y}(1)). \quad (2)$$

We calculate  $w_k$  by maximizing the posterior distribution of the weights, which is proportional to:

$$10 \quad A = \prod_{k=1}^K (w_k)^{\alpha-1} \prod_{t=1}^T \sum_{k=1}^K w_k \cdot \rho(\mathbf{y}_{\text{OBS}}^t(2)|\mathbf{y}_{\text{OBS}}^t(1); Y_{\text{OBS}}^{(t)}, M_k) \quad (3)$$

where  $\alpha$  is the concentration parameter,  $\mathbf{y}_{\text{OBS}}^t(1)$  and  $\mathbf{y}_{\text{OBS}}^t(2)$  are the predictors and predictands for events  $t = 1, \dots, T$ , and  $Y_{\text{OBS}}^{(t)}$  is a matrix containing observed values

of predictors and predictands for all the events except event  $t$ .  $\prod_{k=1}^K (w_k)^{\alpha-1}$  is from the symmetric Dirichlet prior distribution used by Wang et al. (2012a). We use  $\alpha$  values

15 greater than 1 to distribute weights more evenly among models, which helps to stabilise the weights when there is significant sampling variability. Specifically,  $\alpha = 1 + a/K$  with  $a = 1$ . The remainder of the right side of Eq. (3) is the cross-validation likelihood function. By using the cross-validation likelihood function, we base each model weight on the predictive power of the model, rather than on the fitting ability of the model.  $A$  is

20 maximised with an iterative expectation-maximization (EM) algorithm, as described by Wang et al. (2012a).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.3 Forecast verification

Forecasts are verified using leave-one-out cross validation. Forecasts for events in year  $t = 1, 2, \dots, n$  are generated from all available historical data except those at year  $t$ . For each forecast variable  $y$ , this produces a series of forecast cumulative probability distributions  $y^t \sim F^t(y^t)$ . Forecasts are then verified against observations  $y_{\text{OBS}}^t$ .

Verifying the probabilistic forecasts is not straightforward, particularly when the aim is to forecast rare events. Here we evaluate forecast reliability to demonstrate that the probabilistic forecasts are neither too confident nor underconfident. We then assess forecast accuracy using three skill scores. We now describe each of the verification measures in detail.

### 2.3.1 Forecast reliability

For probabilistic forecasts to be meaningful, we must first demonstrate that the forecast probability distributions are reliable; that is, the uncertainty in the forecasts is reliably represented, and thus the forecast distributions are neither too wide (not confident enough) nor too narrow (overconfident). To achieve this, we present reliability diagrams. A reliability diagram plots the observed frequency against the forecast probability and shows how well the predicted probability of an event corresponds to its observed frequency (Wilks, 1995). We present reliability diagrams calculated from events that are larger than the 50% exceedance probability threshold of Max1D and Max5D streamflows.

### 2.3.2 Overall forecast accuracy: root mean square error in probability

The root mean square error in probability (RMSEP) works on the principle that if forecast and observed values are of similar exceedance probabilities then the forecast should be rewarded, even if the magnitudes of observed and forecast values are quite different (Wang and Robertson, 2011). RMSEP is calculated as follows:

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1. We represent the observed historical distribution (climatology),  $y$ , in the form of non-exceedance probability,  $F_{\text{CLI}}(y)$ .
2. For events  $t = 1, 2, \dots, n$ , we take the median of the forecast distribution,  $y_{\text{MED}}^t$ .
3. RMSEP is then calculated as

$$\text{RMSEP} = \left[ \frac{1}{n} \sum_{t=1}^n \left( F_{\text{CLI}}(y_{\text{MED}}^t) - F_{\text{CLI}}(y_{\text{OBS}}^t) \right)^2 \right]^{\frac{1}{2}}. \quad (4)$$

1. We calculate  $\text{RMSEP}_{\text{REF}}$  by substituting the forecast median,  $y_{\text{MED}}^t$ , in Eq. (4) with the climatology median. We then calculate the RMSEP skill score:

$$\text{SS}_{\text{RMSEP}} = \frac{\text{RMSEP}_{\text{REF}} - \text{RMSEP}}{\text{RMSEP}_{\text{REF}}}. \quad (5)$$

In addition, we calculate  $\text{SS}_{\text{RMSEP}}$  with  $\text{RMSEP}_{\text{REF}}$  represented by the BJP forecast generated with only catchment wetness as a predictor (i.e. no climate information is used to generate  $\text{RMSEP}_{\text{REF}}$ ). This allows us to show the relative contribution of catchment wetness and climate indices to forecast skill.

### 2.3.3 Accuracy of forecasts for large threshold events

For a given month, we consider a subset of larger “high” streamflows to assess forecast performance. These larger streamflows are defined as having exceedance probabilities of 50% ( $Q_{50}$ ), 25% ( $Q_{25}$ ) and 10% ( $Q_{10}$ ) for observed Max1D and Max5D. (These streamflows approximately correspond to annual exceedance probabilities (AEP) of 1 : 2 AEP, 1 : 4 AEP and 1 : 10 AEP. To keep the study as simple as possible, we have defined larger events on the basis of exceedance probabilities rather than fitting an extreme value distribution, so we continue to refer to large streamflows in terms of exceedance probabilities.) We treat these large streamflows as thresholds (we term them *large threshold events*), and measure forecast skill by comparing the forecast

probability of exceeding a large threshold event with the corresponding observation.  $Q_{50}$ ,  $Q_{25}$ , and  $Q_{10}$  thresholds are shown for 1-Month Max1D and Max5D streamflows are shown in Fig. 3.

Use of multiple skill scores is recommended to demonstrate robustness in the results (e.g. Cloke and Pappenberger, 2008). We use two measures of skill to verify forecasts at larger streamflow thresholds: the Brier Score and log-likelihood ratio.

### Brier score

The Brier score has been a staple for the verification of probabilistic forecasts since it was proposed by Brier (1950). We use the Brier score to verify forecasts of larger streamflows in order that our study can be compared to others.

Given forecast distributions  $y^t$  at events  $t = 1, 2, \dots, n$ , and streamflow thresholds  $Q_P$ , with exceedance probabilities  $P = 50\%$ ,  $25\%$ ,  $10\%$ , the forecast is presented as the probability of exceeding the streamflow threshold:

$$1 - F^t = p(y^t > Q_P). \quad (6)$$

We calculate the Brier score as:

$$BS = \frac{1}{n} \sum_{t=1}^n (1 - F^t - O^t) \quad (7)$$

where  $O^t$  takes the value of 1 if the threshold is exceeded, and 0 if it is not exceeded. We calculate  $BS_{REF}$  by substituting  $F^t$  with a forecast calculated from climatology,  $F_{REF}^t$ . We then calculate the Brier skill score:

$$SS_{BS} = \frac{BS_{REF} - BS}{BS_{REF}}. \quad (8)$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Log-likelihood ratio

The Brier score has been subject to criticism, particularly for producing unintuitive results for rare (and in our case, large) events when assessing very sharp forecasts (i.e. forecast probabilities of 100 % or 0 %) (Jewson, 2008; Benedetti, 2010). We adopt the recommendations of Benedetti (2010) and Jewson (2008), who both advocate variations on the likelihood to assess probabilistic forecasts. We term the measure we use the log-likelihood ratio (LLR).

The LLR is based on the likelihood ratio described by Jewson (2008). For all exceedance forecasts  $1 - F^t$ , let all the cases of  $t$  where  $1 - F^t$  exceeds a streamflow threshold  $Q$  be given by the set  $A$ , and all cases of  $t$  where the streamflow threshold is not exceeded be given by  $B$ . The log-likelihood for a forecast is calculated by:

$$LL = \log_e \left( \prod_A (1 - F^t) \prod_B F^t \right). \quad (9)$$

The log-likelihood of the reference forecast,  $LL_{REF}$ , is calculated by substituting  $F_{REF}^t$  (again, based on climatology) for  $F^t$  in Eq. (9). The LLR is then calculated by:

$$LLR = LL - LL_{REF}. \quad (10)$$

The LLR differs from skill scores like RMSEP or the Brier score in that it does not show proportional improvement over a reference forecast on a normalised scale (often from  $-\infty$  % to 100 %), making direct comparisons to other skill scores difficult. However, the LLR is essentially identical to the natural logarithm of the pseudo Bayes factor ( $\log_e(\text{PsBF})$ ) presented by Robertson and Wang (2012) and Schepen et al. (2012). Robertson and Wang (2012) showed that values of the  $\log_e(\text{PsBF})$  up to 2 are indistinguishable from statistical noise, while there is a 95 % chance that the relationship between a forecast and observations is true if the  $\log_e(\text{PsBF})$  is greater than 4. We adopt the qualitative categories for the LLR presented by Schepen et al. (2012) for our

## NHESSD

1, 3129–3168, 2013

### High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



study: little evidence of skill where  $LLR < 2$ ; positive evidence of skill where  $2 < LLR < 4$ ; strong evidence of skill where  $4 < LLR < 6$ ; very strong evidence of skill where  $LLR > 6$ .

### 3 Results

#### 3.1 Suitability of BJP for modelling high streamflows

5 The log-sinh transform used to normalise streamflows has been shown to be well-suited to hydrological data in general (Wang et al., 2012b; Del Giudice et al., 2013), but its applicability to high streamflows needs to be established. We show the log-sinh transformed normal distributions fitted to observed Max1D values for two example months, February and September (other months give very similar results). These two months represent low and high streamflow regimes: February is a month of low streamflows in MMH, MUR, ABH and TAW, and a month of high streamflows in ORB and NOR, while September is a month of high streamflows in MMH, MUR, ABH and TAW and a month of low streamflows in ORB and NOR. In general, the assumed log-sinh transformed normal distributions appear to adequately represent the marginal distribution of observations. Almost all observations fall within the confidence bounds of the fitted distributions, including large Max1D events. The log-sinh transformed normal distributions represent observed events well even in catchments with highly variable streamflows, such as ORB and ABH. In summary, the log-sinh transform is flexible enough to normalise the events we are attempting to forecast.

#### 3.2 Forecast reliability

In general, forecast uncertainty is reliably represented by the forecasts after cross-validation. Figure 5 shows reliability diagrams for the NOR and MUR catchments for Max1D 1-Month forecasts (other catchments produce similar results). In these diagrams, the range of forecast probabilities is divided into five bins (see inserts). The [0.05, 0.95] uncertainty interval of the observed relative frequency is calculated through

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



bootstrap resampling of the forecasts and observed streamflows. For the majority of forecast probability ranges, the uncertainty interval of the observed relative frequency intersects the theoretical 1 : 1 line, indicating that the forecasts of high streamflows are reliable. Similar results are obtained for the other catchments for all predictands and forecast periods (not shown). These results support the findings of Wang et al. (2009) and Wang and Robertson (2011), who showed the BJP produces reliable forecasts of seasonal streamflows.

### 3.3 Overall forecast skill

Figure 6 shows BJP-BMA cross-validated hindcasts of Max1D for an example 20 yr period for all catchments. Visual inspection of the hindcasts shows that the credible prediction intervals largely encompass the range of observations. In catchments with strongly seasonal streamflows (e.g. MUR, MMH), the mean of the ensemble forecast often gives realistic predictions of Max1D streamflows during seasons of high streamflows. Accuracy of forecasts in more variable catchments (e.g. NOR, ABH) is much more difficult to ascertain from these timeseries, and we now turn to formal measures of skill to assess these.

RMSEP skill scores are positive for Max5D forecasts for the 1 month forecast period for most months and catchments (Fig. 7b). Skill in Max5D 1 month forecasts is particularly strong in the winter-spring months (June–November). Skill in Max1D 1 month forecasts is generally lower than for Max-5D 1 month forecasts (Fig. 7a, b). Max1D streamflows are inherently more variable than Max5D streamflows, as Max5D streamflows are smoothed by the greater number of data included in their calculation. This makes forecasting Max1D streamflows more challenging. Nonetheless, RMSEP skill scores for Max1D 1 month forecasts are positive for most catchments and seasons (Fig. 7a). Max1D 1 month forecast skill is strongest in the winter-spring months. For the 3 month forecast period, RMSEP scores are generally lower for both Max1D and Max5D forecasts, although positive skill scores occur in winter-spring for the MUR,

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





MMH, and ABH catchments, and the NOR catchment shows skill intermittently through the year (Fig. 7c, d).

The reason for the reduced performance of the 3 month forecasts becomes evident when we review the contribution of climate indices to forecast skill. Figure 8 shows RM-SEP skill scores calculated relative to BJP forecasts generated using only streamflow as a predictor. The plot shows the skill gained by the inclusion of climate indices for Max1D 1 month forecasts. Figure 8 shows that almost no skill is gained in any month or catchment by including climate indices, meaning the forecasts depend heavily on catchment wetness for skill. Results are similar for Max5D (not shown). This finding is also supported by Robertson and Wang (2013), who found that climate indices made only weak contributions to the skill of forecasts of seasonal streamflow totals in the MMH and MUR catchments. The contribution of catchment wetness to forecast skill declines over longer forecast periods (Shukla and Lettenmaier, 2011; Li et al., 2009). Thus forecasts for longer periods are less accurate than for shorter forecast periods. This effect is also evident in individual catchments. The TAW catchment, for example, has the lowest autocorrelation of monthly streamflows of the six catchments (not shown), and forecasts for this catchment show poor skill in relation to streamflow climatology.

Nonetheless, 3 month forecasts can be skilful in certain catchments at times of the year when the influence of catchment wetness on high streamflows is strong. The influence of catchment wetness on streamflows is generally strongest on the receding limb of the annual hydrograph (Robertson and Wang, 2013). For the ORB and NOR catchments the annual hydrograph recedes in March–May, while in the ABH, MMH and MUR catchments the annual hydrograph recedes in August–November. This results in positive RMSEP skill scores for 3 month forecasts of these catchments during these months (Fig. 7c, d).

Overall, RMSEP generally shows positive skill scores for 1 month forecasts for both Max1D and Max5D streamflows, while 3 month forecasts are substantially less skilful. However, the positive RMSEP skill scores may be the result of good agreement of

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



forecasts with lower “high” streamflows, and not reflect forecasts at larger streamflows. We now turn to forecast skill at higher streamflows to determine the size of streamflows for which forecasts are skillful.

### 3.4 Forecast skill for large threshold events

In general, forecast skill declines as streamflows get larger (Figs. 9–12). Brier scores show more instances of positive skill than LLR scores, particularly for streamflows larger than  $Q_{10}$ . Because the Brier score has known problems with infrequent events (Benedetti, 2010), we focus on the LLR score to discuss forecast skill at larger streamflows.

Substantial skill is evident in forecasts where observed Max1D streamflows are larger than  $Q_{50}$  for 1 month forecasts, in both the Brier score (Fig. 9) and the LLR (Fig. 10). LLR scores are higher for Max5D streamflows than for Max1D streamflows, and the highest LLR scores generally occur in July–November. Skill is not related to seasonal changes in high or low Max1D/Max5D streamflows. The ARB, MUR, MMH and catchments show high skill during months of high streamflow (winter–spring, Figs. 10 and 2) while the ORB and NOR catchments only exhibit skill during months of low streamflow (July–November, Figs. 10 and 2). As with the RMSEP scores, the TAW catchment shows the lowest skill. Four of the six catchments show positive LLR scores in 6 or more months of the year for 1 month forecasts of Max5D streamflows above  $Q_{25}$  (Fig. 10). For Max1D streamflows greater than  $Q_{25}$ , three catchments show positive LLR scores in six or more months of the year (Fig. 10). Little skill is evident in any catchment or season for either Max1D or Max5D streamflows above  $Q_{10}$ .

Skill for 3 month forecasts of larger streamflows are generally low (Figs. 9 and 10). Except for one catchment (MUR), catchments show little forecast skill in the majority of months for any of the streamflow thresholds tested for either Max1D or Max5D streamflows. We find positive skill scores for 3 month forecasts in the MUR catchment of Max5D streamflows above  $Q_{50}$  and  $Q_{25}$  for six or more months, and also for Max1D streamflows above  $Q_{50}$  (Fig. 12). Indeed, forecasts for MUR performed best in most

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



measures and skill scores. It is not clear why this should be so. MUR receives reliable rainfall in the winter and spring, resulting in relatively low variability and strong autocorrelation in monthly streamflows. However these characteristics also apply to the nearby MMH catchment, for which forecasts perform no better than for ABH, ORB or NOR in a number of measures (e.g. Fig. 10).

Overall, forecast skill is positive to very strong for 1 month exceedance forecasts of streamflows exceeding  $Q_{50}$  for a majority of months in all but the TAW catchment. Skill is not related to seasonal cycles of high and low streamflows. Positive skill scores are also found in several catchments for 1 month exceedance forecasts of streamflows exceeding  $Q_{25}$ . The remaining large streamflow forecasts tested here showed little skill in most catchments.

## 4 Discussion

RMSEP skill scores reported here show the 1 month forecasts to be superior to climatology in forecasting high streamflows. Further, the skill in forecasts is not limited to the lowest of the “high” streamflows – forecasts of the probability of exceeding  $Q_{50}$  Max1D streamflows one month in advance show robust skill in a number of catchments. We note, however, that the  $Q_{50}$  Max1D streamflows are still not necessarily very large streamflows. Skill in forecasting large threshold events in two catchments, ORB and NOR, is restricted to months where “high” streamflows are small, and in which damaging floods are unlikely to occur. Conversely, skill in the MUR, ABH and MMH catchments is evident during periods of high streamflow. Accordingly, forecast skill in these catchments may be valuable to the Bureau of Meteorology when they are seeking to answer more general questions about the risks of high streamflows in a coming month. We note that the usefulness of the forecast is likely to vary with catchment in any case, both because forecast skill varies between catchments and because the prospect of flood damage varies greatly between catchments (i.e. in one

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



catchment a common high streamflow event may have deleterious consequences, in another catchment extremely large floods may be of little consequence).

The 1 month forecasts rely heavily on catchment wetness for skill. Improving estimates of catchment wetness are likely to be a simple way of improving forecasts.

Accumulated streamflow for a month can be a poor measure of catchment wetness. For example, a high value of total streamflow may be caused by a single intense rainfall event that causes infiltration-excess overland flow, resulting in a large streamflow but little infiltration. In this example the catchment wetness is overestimated by total streamflow. Catchment wetness can be modeled more effectively for forecasting with so-called “dynamical” approaches (Rosenberg et al., 2011; Robertson et al., 2013), and such dynamical catchment wetness models could improve forecasts.

The ability to forecast high streamflows a month or more in advance is limited by knowledge of climate during the forecast period. This problem is not likely to be easily surmountable. The much higher variability of larger rainfall events makes their prediction inherently difficult. In addition, climate indices that have the potential to forecast particular types of rain-bearing weather patterns may have little persistence from month to month. This is particularly so for climate indices calculated from atmospheric variables, which tend to be less persistent than oceanic variables. For example, we have used the atmospheric blocking index (B140, see Table 2) to attempt to account for atmospheric blocking and associated cutoff lows in our forecasts. Cutoff lows associated with atmospheric blocking bring a substantial proportion of rainfall to south-eastern Australia (Pook et al., 2006), and can counteract the drying associated with very strong El Niño years (Brown et al., 2009). However, we find that B140 adds little skill to forecasts of high streamflows, supporting Schepen et al. (2012) who showed that lagged B140 had no significant statistical relationship to mean rainfall anywhere in Australia.

While several studies have shown positive relationships between lagged climate indices and mean seasonal rainfall in Australia (e.g. Schepen et al., 2012), it appears from our work that the relationship between heavier rainfall events and lagged climate indices in general is weak for the purposes of forecasting high streamflows in south-

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



east Australia. Future improvements in dynamical climate models used for forecasting weeks to months advance (e.g. Marshall et al., 2011) may ultimately improve forecasts of high rainfalls, or may improve forecasts of sea-surface temperatures from which climate indices may be calculated. Using forecasts of climate indices from dynamical climate models shows promise in improving forecasts of monthly rainfall totals at lead-times of more than six months (Hawthorne et al., 2013), and avoids the use of lagged climate indices for forecasts. This method may also improve longer-term forecasts of high rainfalls.

The high streamflow forecasts we have developed here may be bolstered in future by the inclusion of Numerical Weather Prediction (NWP) models in hydrological forecasting. The Australian Bureau of Meteorology does not presently use NWP forecasts to quantify flood forecasts, although they are used qualitatively to inform flood warnings (Elliott et al., 2005). Very high resolution NWP forecasts have been shown to improve flood forecasts (Roberts et al., 2008). At present, however, NWP forecasts are skilful only for a few days (typically < 10 days); and even skilful NWP forecasts are often not accurate enough for use in hydrological forecasting systems, even in catchments substantially larger than those tested here (Cloke and Pappenberger, 2009; Shrestha et al., 2013; Cuo et al., 2011). As NWP models and post-processing of NWP forecasts improve, NWP forecasts may complement the simpler forecasts we have generated in this study.

## 5 Summary and conclusions

We have explored the ability of existing statistical forecasting methods to produce forecasts for high streamflows for the coming month and the coming three months. Forecast models are built from a combination of climate predictors and catchment wetness. Models are constructed with a Bayesian joint probability method, and the models are then weighted based on their predictive power using Bayesian model averaging.

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Skill is clearly evident in forecasts of high streamflows for the coming 1 month period. Forecasts of larger events, including maximum 1 day streamflows of exceedance probabilities as low as 25 %, are also skilful in comparison to long-term climatologies. Our 1 month high streamflow forecasts have the potential to complement existing real-time flood warnings currently used in Australia, to give emergency services and the community more warning of impending high streamflows.

Almost all forecast skill derives from the catchment wetness predictor. If the forecasts are to be extended to additional catchments, they are likely to be poor in catchments that have little month-to-month memory in streamflows. Forecasts in skilful catchments may be improved somewhat by using more refined estimates of catchment wetness.

We find substantially lower skill in forecasts of high streamflows for the coming 3 month period. The influence of catchment wetness on streamflows diminishes over longer periods, and climate predictors add little skill to the forecasts. Future improvements in forecasts of extreme rainfalls from dynamical climate models may be able to improve longer range forecasts of high streamflows.

*Acknowledgement.* This research has been supported by the Water Information Research and Development Alliance between the Australian Bureau of Meteorology and CSIRO Water for a Healthy Country Flagship. Thanks to Yong Song (CSIRO Land and Water), Christopher J. White (Bureau of Meteorology) and Senlin Zhou (Bureau of Meteorology) for their comments on earlier drafts.

## References

- Ashok, K., Guan, Z., and Yamagata, T.: Influence of the Indian Ocean dipole on the Australian winter rainfall, *Geophys. Res. Lett.*, 30, 1821, doi:10.1029/2003GL017926, 2003.
- Ashok, K., Nakamura, H., and Yamagata, T.: Impacts of ENSO and Indian Ocean dipole events on the Southern Hemisphere storm-track activity during austral winter, *J. Climate*, 20, 3147–3163, doi:10.1175/jcli4155.1, 2007.
- Baxter-Tomkins, T. and Wallace, M.: Recruitment and retention of volunteers in emergency services, *Australian Journal on Volunteering*, 14, 1–11, 2009.

---

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Benedetti, R.: Scoring rules for forecast verification, *Mon. Weather Rev.*, 138, 203–211, doi:10.1175/2009MWR2945.1, 2010.
- Brier, G. W.: Verification of forecasts expressed in terms of probability, *Mon. Weather Rev.*, 78, 1–3, doi:10.1126/science.27.693.594, 1950.
- 5 Brown, J. N., McIntosh, P. C., Pook, M. J., and Risbey, J. S.: An investigation of the links between ENSO flavors and rainfall processes in southeastern Australia, *Mon. Weather Rev.*, 137, 3786–3795, doi:10.1175/2009MWR3066.1, 2009.
- Chiew, F. H. S., Zhou, S. L., and McMahon, T. A.: Use of seasonal streamflow forecasts in water resources management, *J. Hydrol.*, 270, 135–144, doi:10.1016/S0022-1694(02)00292-5, 1998.
- 10 Cloke, H. L. and Pappenberger, F.: Evaluating forecasts of extreme events for hydrological applications: an approach for screening unfamiliar performance measures, *Meteorol. Appl.*, 15, 181–197, doi:10.1002/met.58, 2008.
- Cloke, H. L. and Pappenberger, F.: Ensemble flood forecasting: a review, *J. Hydrol.*, 375, 613–626, doi:10.1016/j.jhydrol.2009.06.005, 2009.
- 15 Cuo, L., Pagano, T. C., and Wang, Q. J.: A review of quantitative precipitation forecasts and their use in short-to medium range streamflow forecasting, *J. Hydrometeorol.*, 12, 713–728, doi:10.1175/2011JHM1347.1, 2011.
- Del Giudice, D., Honti, M., Scheidegger, A., Albert, C., Reichert, P., and Rieckermann, J.: Improving uncertainty estimation in urban hydrological modeling by statistically describing bias, *Hydrol. Earth Syst. Sci. Discuss.*, 10, 5121–5167, doi:10.5194/hessd-10-5121-2013, 2013.
- 20 DelSole, T. and Shukla, J.: Artificial skill due to predictor screening, *J. Climate*, 22, 331–345, doi:10.1175/2008JCLI2414.1, 2009.
- Elliott, J., Catchlove, R., Sooriyakumaran, S., and Thompson, R.: Recent advances in the development of flood forecasting and warning services in Australia, *International conference on innovation, advances and implementation of flood forecasting technology*, Tromsø, Norway, 17–19 October 2005, 1–10, 2005.
- Hawthorne, S., Wang, Q. J., Schepen, A., and Robertson, D. E.: Effective use of GCM outputs for forecasting monthly rainfalls to long lead times, *Water Resour. Res.*, submitted, 2013.
- 30 Jewson, S.: The problem with the Brier score, arXiv: physics/0401046v1 [physics.ao-ph], available at: <http://arxiv.org/abs/physics/0401046v1>, last access: June 2013, 2008.
- Jones, D. A., Wang, W., and Fawcett, R.: High-quality spatial climate data-sets for Australia, *Australian Meteorological and Oceanographic Journal*, 58, 233–248, 2009.

---

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kiem, A. S., Franks, S. W., and Kuczera, G.: Multi-decadal variability of flood risk, *Geophys. Res. Lett.*, 30, 1035, doi:10.1029/2002GL015992, 2003.
- Kwon, H.-H., Brown, C., Xu, K., and Lall, U.: Seasonal and annual maximum streamflow forecasting using climate information: application to the Three Gorges Dam in the Yangtze River basin, China, *Hydrolog. Sci. J.*, 54, 582–595, doi:10.1623/hysj.54.3.582, 2009.
- Li, H., Luo, L., Wood, E. F., and Schaake, J.: The role of initial conditions and forcing uncertainties in seasonal hydrologic forecasting, *J. Geophys. Res.*, 114, D04114, doi:10.1029/2008jd010969, 2009.
- Lindström, G. and Olsson, J.: A systematic review of sensitivities in the Swedish flood-forecasting system, *Atmos. Res.*, 100, 275–284, doi:10.1016/j.atmosres.2010.09.013, 2011.
- Marshall, A. G., Hudson, D., Wheeler, M. C., Hendon, H. H., and Alves, O.: Assessing the simulation and prediction of rainfall associated with the MJO in the POAMA seasonal forecast system, *Clim. Dynam.*, 37, 2129–2141, doi:10.1007/s00382-010-0948-2, 2011.
- Murphy, B. F. and Timbal, B.: A review of recent climate variability and climate change in south-eastern Australia, *Int. J. Climatol.*, 28, 859–879, doi:10.1002/joc.1627, 2008.
- Pfister, N.: The case of an evacuation from Grafton, *The Australian Journal of Emergency Management*, 17, 19–29, 2002.
- Piechota, T. C., Chiew, F. H. S., Dracup, J. A., and McMahon, T. A.: Seasonal streamflow forecasting in eastern Australia and the El Niño–Southern Oscillation, *Water Resour. Res.*, 34, 3035–3044, doi:10.1029/98WR02406, 1998.
- Pook, M. J., McIntosh, P. C., and Meyers, G. A.: The synoptic decomposition of cool-season rainfall in the southeastern Australian cropping region, *J. Appl. Meteorol. Clim.*, 45, 1156–1170, doi:10.1175/JAM2394.1, 2006.
- Risbey, J. S., Pook, M. J., McIntosh, P. C., Wheeler, M. C., and Hendon, H. H.: On the remote drivers of rainfall variability in Australia, *Mon. Weather Rev.*, 137, 3233–3253, doi:10.1175/2009MWR2861.1, 2009.
- Roberts, N. M., Cole, S. J., Forbes, R. M., Mooreb, R. J., and Boswell, D.: Use of high-resolution NWP rainfall and river flow forecasts for advance warning of the Carlisle flood, north-west England, *Meteorol. Appl.*, 16, 23–34, doi:10.1002/met.94, 2008.
- Robertson, D. E. and Wang, Q. J.: A Bayesian approach to predictor selection for seasonal streamflow forecasting, *J. Hydrometeorol.*, 13, 155–171, doi:10.1175/JHM-D-10-05009.1, 2012.



---

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Robertson, D. E. and Wang, Q. J.: Seasonal forecasts of unregulated inflows into the Murray River, Australia, *Water Resour. Manage.*, 27, 2747–2769, doi:10.1007/s11269-013-0313-4, 2013.

Robertson, D. E., Pokhrel, P., and Wang, Q. J.: Improving statistical forecasts of seasonal streamflows using hydrological model output, *Hydrol. Earth Syst. Sci.*, 17, 579–593, doi:10.5194/hess-17-579-2013, 2013.

Rosenberg, E. A., Wood, A. W., and Steinemann, A. C.: Statistical applications of physically based hydrologic models to seasonal streamflow forecasts, *Water Resour. Res.*, 47, W00H14, doi:10.1029/2010WR010101, 2011.

Saji, N. H., Goswami, B. N., Vinayachandran, P. N., and Yamagata, T.: A dipole mode in the tropical Indian Ocean, *Nature*, 401, 360–363, 1999.

Schepen, A., Wang, Q. J., and Robertson, D.: Evidence for using lagged climate indices to forecast Australian seasonal rainfall, *J. Climate*, 25, 1230–1246, doi:10.1175/JCLI-D-11-00156.1, 2012.

Sharma, A.: Seasonal to interannual rainfall probabilistic forecasts for improved water supply management: part 3 – a nonparametric probabilistic forecast model, *J. Hydrol.*, 239, 249–258, doi:10.1016/S0022-1694(00)00348-6, 2000.

Shrestha, D. L., Robertson, D. E., Wang, Q. J., Pagano, T. C., and Hapuarachchi, H. A. P.: Evaluation of numerical weather prediction model precipitation forecasts for short-term streamflow forecasting purpose, *Hydrol. Earth Syst. Sci.*, 17, 1913–1931, doi:10.5194/hess-17-1913-2013, 2013.

Shukla, S. and Lettenmaier, D. P.: Seasonal hydrologic prediction in the United States: understanding the role of initial hydrologic conditions and seasonal climate forecast skill, *Hydrol. Earth Syst. Sci.*, 15, 3529–3538, doi:10.5194/hess-15-3529-2011, 2011.

Troup, A. J.: The southern oscillation, *Q. J. Roy. Meteor. Soc.*, 91, 490–506, doi:10.1002/qj.49709139009, 1965.

Verdon, D. C. and Franks, S. W.: Indian Ocean sea surface temperature variability and winter rainfall: eastern Australia, *Water Resour. Res.*, 41, W09413, doi:10.1029/2004WR003845, 2005.

Verdon, D. C., Wyatt, A. M., Kiem, A. S., and Franks, S. W.: Multidecadal variability of rainfall and streamflow: eastern Australia, *Water Resour. Res.*, 40, W10201, doi:10.1029/2004WR003234, 2004.

# NHESSD

1, 3129–3168, 2013

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Wang, Q. J. and Robertson, D. E.: Multisite probabilistic forecasting of seasonal flows for streams with zero value occurrences, *Water Resour. Res.*, 47, W02546, doi:10.1029/2010WR009333, 2011.
- 5 Wang, Q. J., Robertson, D. E., and Chiew, F. H. S.: A Bayesian joint probability modeling approach for seasonal forecasting of streamflows at multiple sites, *Water Resour. Res.*, 45, W05407, doi:10.1029/2008WR007355, 2009.
- Wang, Q. J., Schepen, A., and Robertson, D. E.: Merging seasonal rainfall forecasts from multiple statistical models through Bayesian model averaging, *J. Climate*, 25, 5524–5537, doi:10.1175/JCLI-D-11-00386.1, 2012a.
- 10 Wang, Q. J., Shrestha, D. L., Robertson, D. E., and Pokhrel, P.: A log-sinh transformation for data normalization and variance stabilization, *Water Resour. Res.*, 48, W05514, doi:10.1029/2011WR010973, 2012b.
- Wilks, D. S.: *Statistical Methods in the Atmospheric Sciences*, Elsevier, New York, 648 pp., 1995.
- 15 Yeo, I. K. and Johnson, R. A.: A new family of power transformations to improve normality or symmetry, *Biometrika*, 87, 954–959, doi:10.1093/biomet/87.4.954, 2000.

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

**Table 1.** Characteristics of catchments used in this study.

Name	Short name	Streamflow record used	Area (km <sup>2</sup> )	Annual rainfall (mm)	Annual runoff (mm)
Orara River at Bawden Bridge	ORB	1956–2006	1823	1396	407
Nowendoc River at Rocks Crossing	NOR	1950–2006	1898	1155	258
Abercrombie River at Hadley No. 2	ABH	1960–2005	1626	842	117
Murray River at Biggara	MUR	1950–2005	1254	1178	446
Mitta Mitta River at Hinnomunjie	MMH	1950–2006	1528	1343	297
Tarwin River at Meeniyan	TAW	1955–2006	1066	1084	233

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Table 2.** List of oceanic and atmospheric climate indices used as predictors.

Index	Description
Southern Oscillation Index (SOI)	Troup (1965)
NINO3	Mean SST anomaly over 150–90° W and 5° N–5° S
NINO3.4	Mean SST anomaly over 170–120° W and 5° N–5° S
NINO4	Mean SST anomaly over 150–160° E and 5° N–5° S
ENSO Modoki Index (EMI)	Ashok et al. (2003)
Indian Ocean Dipole Mode Index (DMI)	Saji et al. (1999)
Indian Ocean West Pole Index (WPI)	Saji et al. (1999)
Indian Ocean East Pole Index (EPI)	Saji et al. (1999)
Indonesia Index (II)	Verdon and Franks (2005)
Tasman Sea Index (TSI)	Murphy and Timbal (2008)
140° E Blocking Index (B140)	Risbey et al. (2009)



**Fig. 1.** Catchments (shaded) and streamflow gauge sites (black dots) used in this study.

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## High streamflow forecasts 1–3 months in advance


J. C. Bennett et al.



**Fig. 2.** Catchment streamflow characteristics. Black dots show average monthly streamflows. Boxes show maximum five-day streamflow (Max5D – blue) and maximum 1 day streamflow (Max1D – red) occurring during each month for exceedance probabilities of 50 % ( $Q_{50}$ , bottom edge) to 10 % ( $Q_{10}$ , top edge), with box centreline showing Max5D/Max1D streamflows of exceedance probability of 25 % ( $Q_{25}$ ).

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.




**Fig. 3.** Schematic of forecast model. **(a)** Example of individual forecast model generated with the Bayesian joint probability method. In this example, catchment wetness (CW) and NINO3.4 predictors are used to predict Max1D streamflows. Rainfall is included as a joint predictand to elicit more information from the climate indices. Parameters for the transforms and joint probability distribution are inferred jointly. This process is repeated for thirteen different predictor-sets. **(b)** The forecasts from thirteen BJP models are weighted based on cross-validated predictive performance with Bayesian model averaging (BMA) to produce a merged BJP-BMA forecast. The use of a symmetric Dirichlet prior encourages even weights in instances of high sampling uncertainty. See text for details.

---

**High streamflow forecasts 1–3 months in advance**J. C. Bennett et al.

---




**Fig. 4.** Fit of log-sinh transformed normal distributions to Max1D values for two months. Red circles show actual values, black solid line shows fitted log-sinh transform, dashed lines show [0.1, 0.9] confidence intervals.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**High streamflow forecasts 1–3 months in advance**


J. C. Bennett et al.



**Fig. 5.** Forecast reliability diagrams at two catchments for Max1D streamflows of exceedance probability  $\leq 50\%$ . (Forecasts are divided into five bins. 1 : 1 dashed lines, perfectly reliable forecast; circles, observed relative frequency; vertical lines, [0.05, 0.95] uncertainty interval of observed relative frequency; inserts, number of events in the different forecast probability bins.)

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.



**Fig. 6.** Example forecast time series of cross-validated BJP-BMA for Max1D. Red circles show observed Max1D values, black points and lines show mean forecast and [0.1, 0.9] credible prediction intervals.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion




## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.



**Fig. 7.** RMSEP skill scores. Catchments are ordered by their location, from northernmost (top) to southernmost (bottom). **(a)** Max1D streamflows for 1 month forecasts, **(b)** Max5D streamflows for 1 month forecasts, **(c)** Max1D streamflows for 3 month forecasts, and **(d)** Max5D streamflows at 3 month forecasts. Scores show proportional improvement of forecasts over climatology forecasts.



**Fig. 8.** Skill added by climate indices to forecasts. Plot shows RMSEP skill scores for Max1D 1 month forecasts calculated with respect to BJP forecasts generated with only catchment wetness as a predictor. Scores show proportional improvement of BJP-BMA forecasts over BJP forecasts generated with only catchment wetness as a predictor.

## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.



**Fig. 9.** Brier skill scores calculated at three streamflow thresholds for 1 month forecasts. Scores show proportional improvement of BJP-BMA forecasts over climatology forecasts.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.



**Fig. 10.** Evidence of skill from the log-likelihood ratio (LLR) at three streamflow thresholds for 1 month forecasts. Scores show evidence of skill of BJP-BMA forecasts over climatology forecasts. Categories are taken from Schepen et al. (2012): little evidence of skill where  $LLR < 2$ ; positive evidence where  $2 < LLR < 4$ ; strong evidence where  $4 < LLR < 6$ ; very strong evidence where  $LLR > 6$ .

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.



**Fig. 11.** Brier skill scores calculated at three streamflow thresholds for 3 month forecasts. Scores show proportional improvement of BJP-BMA forecasts over climatology forecasts.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



High streamflow forecasts 1–3 months in advance

J. C. Bennett et al.



**Fig. 12.** Evidence of skill from the log-likelihood ratio at three streamflow thresholds for 3 month forecasts. Scores show evidence of skill of BJP-BMA forecasts over climatology forecasts. Categories are taken from Schepen et al. (2012): little evidence of skill where  $LLR < 2$ ; positive evidence where  $2 < LLR < 4$ ; strong evidence where  $4 < LLR < 6$ ; very strong evidence where  $LLR > 6$ .

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

