



This discussion paper is/has been under review for the journal Natural Hazards and Earth System Sciences (NHESS). Please refer to the corresponding final paper in NHESS if available.

Rogue waves, rogue events and extreme wave kinematics in spatio-temporal fields of simulated sea states

A. Sergeeva^{1,2} and A. Slunyaev^{1,2}

¹Institute of Applied Physics, Nizhny Novgorod, Russia

²Nizhny Novgorod State Technical University, Nizhny Novgorod, Russia

Received: 10 January 2013 – Accepted: 10 January 2013 – Published: 29 January 2013

Correspondence to: A. Sergeeva (a.sergeeva@hydro.appl.sci-nnov.ru)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

An approach to extensive studying rogue wave occurrence in numerical simulations is presented. As a result of numerical simulations of the unidirectional wave evolution, spatio-temporal fields of wave data of the size 20 min × 10 km are obtained with high resolution in time and space and are used for statistical analysis with the focus on extreme waves. Having the exhaustive information on the wave evolution enables us to capture the detailed picture of individual rogue waves; to detect intermittent rogue wave events, which last for significantly longer time, and hence, to draw the image of a rogue wave. Due to the benefit of owing full wave data, the question of relation between extreme wave kinematics and extremely high waves is discussed in the statistical sense.

1 Introduction

The rogue or freak wave topic is of a very high interest nowadays in marine engineering and geophysical communities. The huge oceanic waves seem to come from nowhere and to disappear with no trace, but at the transient moment of their occurrence they cause ultimate impact, damage and destroy boats and sea platforms, kill people. Rogue wave phenomenon challenges physical and mathematical researches, which are now numerous, see reviews in Kharif et al. (2009); Dysthe et al. (2008); Slunyaev et al. (2011). Though main physical mechanisms which can cause these waves seem to be formulated, realistic sea waves represent a quite complex object for investigation, so that the very vital problems: how rogue waves look like, how long they live and how probable they occur have not been answered with certainty so far.

Derivation of rogue wave probability from the primitive water wave equations is too complicate: the usual assumptions of the kinetic theory are violated; wave phase coherence and strong nonlinear effects must be taken into consideration. Direct measurements of sea waves have now resulted in an impressive amount of data. However, rogue waves seem to produce extreme effect on measuring devices, so that the number

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of reliable recorded time-series of rogue waves turns out to be insufficient to build the statistics of in-situ rogue waves (Christou and Ewans, 2011), what often leads to conflicting conclusions of different studies. Besides, the statistical ensemble must consist of homogeneous data; this request is difficult to fulfill, when processing in-situ records.

Laboratory and numerical simulations provide an efficient tool for studying stochastic wave evolution. Such experiments have been performed by many authors (Onorato et al., 2001, 2002, 2005; Dysthe et al., 2003; Janssen, 2003; Socquet-Juglard et al., 2005; Mori et al., 2007; Chalikov, 2005, 2009; Annenkov and Shrira, 2009; and many others), including the authors of the present study, see Pelinovsky and Sergeeva (2006); Shemer and Sergeeva (2009); Shemer et al. (2010a, b); Slunyaev and Sergeeva (2011). The simulations performed in the present research use a similar arrangement. A random superposition of sinusoidal waves with prescribed spectral shape is used as the initial condition at $t = 0$ and simulated in time. Then, the result of wave evolution is analyzed. We emphasize that the wave system may undergo essential statistical evolution, what yields violation of the stationarity assumption, see discussion in Shemer et al. (2010a, b); Slunyaev and Sergeeva (2011).

In most of the studies only a little amount of wave information is processed. For example, in the absolute majority of laboratory experiments the data consists of time series of surface elevation recorded by a few gauges. In vast majority of the numerical simulations only peak waves are analyzed at the moment when the maximum wave height is attained. The wave field data normally cannot be stored with too high frequency because this would slow down the simulation gradually, and would increase the amount of data. Meanwhile, the simulation of the initial problem employs integration in time, thus a fine time resolution of wave dynamics is naturally available.

If a rogue wave occurs (let us assume that the wave does not overturn, so that the simulation can be continued), it possesses a finite life-time, and it is a rogue wave *at every moment* of this time interval. We emphasize this obvious fact and analyze all the moments of the freak wave evolution. To produce one realization of a stochastic wave dynamics, we simulate quasi-random waves in time, and the simulated data is stored

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with a sufficiently high frequency for further analysis. Besides the surface elevation, velocity fields are also collected through the evolution, thus, the full wave data is available for the analysis.

Usually, 20-min time series represent the in-situ measurements; therefore the time interval of 20 min is considered in the present study. To ensure that all the waves, which pass one point in space during the 20 min period are indeed different, we take sufficiently long spatial domain of simulations: $C_{gr} \times 20 \text{ min} \approx 10 \text{ km}$, where C_{gr} is the typical group velocity. The data grid consists of 2048×2048 points, what corresponds to about $5 \text{ m} \times 0.6 \text{ s}$ mesh size. The moderate level of discretization is chosen with the purpose to analyze the data employing an ordinary PC; of course, a finer mesh in time and/or in space may be produced.

The spatial domain contains about 60 individual waves; their evolution is considered for about 120 wave periods (2048 instants). Twenty realizations of irregular waves for each initial condition, specified by the spectrum shape (the *sea condition*), build up the statistical ensemble, what results in about 2 500 000 individual waves in spatial domain (20 realizations \times 60 waves \times 2048 instants). All these waves are used for the analysis. It is important to beware of a direct comparison between the statistical results obtained in this paper and the statistics based on measurements in one point. In our case not all the waves are statistically independent; this issue will be discussed below in Sect. 3.1. The reverse side of the present approach is that if a rogue wave appears, then it is examined at all phases of its evolution. Possessing both the spatial and temporal data, the dynamics of a rogue wave is considered in Sect. 3.2, in particular, the life-time and prevailing shapes of rogue waves are obtained. We introduce the concept of a *rogue event*, which incorporates shorter-living extreme waves, which overpass the formal height criterion on a rogue wave with brief interruptions. With use of the wave velocity data, collected in our simulations, in Sect. 4 we discuss a very important issue of the wave-structure interaction: how extreme velocities in the simulated wave fields correlate with velocities, culminated during the rogue wave lifetime. The paper ends with conclusions.

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2 The approach for producing space-time wave data sheets

2.1 Numerical model

The primitive equations of potential hydrodynamics for the surface displacement $\eta(x, t)$ and the surface velocity potential $\Phi(x, t) = \phi(x, t, z = \eta)$ have the form

$$\frac{\partial \eta}{\partial t} = -\frac{\partial \Phi}{\partial x} \frac{\partial \eta}{\partial x} + \left(1 + \left(\frac{\partial \eta}{\partial x}\right)^2\right) \frac{\partial \phi}{\partial z}, \quad (1)$$

$$\frac{\partial \Phi}{\partial t} = -g\eta - \frac{1}{2} \left(\frac{\partial \Phi}{\partial x}\right)^2 + \frac{1}{2} \left(\frac{\partial \phi}{\partial z}\right)^2 \left[1 + \left(\frac{\partial \eta}{\partial x}\right)^2\right], \quad (2)$$

$$\frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial z^2} = 0, \quad z \leq \eta, \quad (3)$$

$$\frac{\partial \phi}{\partial z} \rightarrow 0, \quad z \rightarrow -\infty. \quad (4)$$

Here x is the spatial coordinate along the wave propagation, and t is time. The Oz axis is directed upward, and g is the acceleration due to gravity. Equation (3) is the Laplace equation for the velocity potential $\phi(x, z, t)$, Eqs. (1) and (2) constitute the surface boundary condition, and Eq. (4) is the bottom condition for it (an infinitely deep water case is considered).

In this research Eqs. (1)–(4) are integrated by means of the High-Order Spectral Method (HOSM), see Dommermuth and Yue (1987); West et al. (1987) with the 4th order Runge – Kutta integration in time. Strictly speaking, the HOSM is not a fully-nonlinear, but a strongly nonlinear algorithm, which resolves the wave-wave interactions up to the predetermined order M (for $M = 3$ the HOSM approach has the same accuracy as the Zakharov equation; Zakharov, 1968). In the present study parameters $M = 3$ and $M = 6$ are used for different simulations. The case $M = 6$ corresponds to nearly fully nonlinear case, see discussion in Clamond et al. (2006).

NHESSD

1, 39–72, 2013

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Initial conditions and preparation of the sea state

The initial condition for the simulations corresponds to a random superposition of sinusoidal waves with spectrum of the JONSWAP shape

$$S(\omega) = \alpha \left(\frac{\omega}{\omega_0}\right)^{-5} \exp\left[-\frac{5}{4}\left(\frac{\omega}{\omega_0}\right)^{-4}\right] \gamma \exp\left[-\frac{1}{2\Delta^2}\left(\frac{\omega-\omega_0}{\omega_0}\right)^2\right] \quad (5)$$

$$\Delta(\omega) = \begin{cases} 0.07, & \omega < \omega_0 \\ 0.09 & \omega > \omega_0 \end{cases}$$

Parameters of the sea states are specified in Table 1. There T_0 – is the wave peak period, H_s – is the significant wave height, and γ – is the peakedness. Parameter α in Eq. (5) is determined through the characteristic wave height, and the angular frequency relates to the wave period as $\omega_0 = 2\pi/T_0$. In this paper the majority of presented results concerns cases A and E. The first one is considered as a low sea state, while the second one corresponds to severe conditions.

Two functions at $t = 0$ are required to specify the initial condition for the simulation: the surface elevation, $\eta(x, t = 0)$, and the surface velocity potential, $\Phi(x, t = 0)$. Due to the nonlinearity these fields are in fact dependent in a non-trivial way, so that in the case of significant wave nonlinearity and broad spectrum it is very difficult to find the initial condition, which would be a realization of irregular waves with the proper relation between functions $\eta(x, t = 0)$ and $\Phi(x, t = 0)$.

Frequently, one of the fields is taken in the form of a linear superposition of Fourier harmonics with random phases, and the second one is obtained according to the linear relation between wave elevation and velocity. The result of such careless production of the initial condition is displayed in Fig. 1a, where the evolution of potential, E_p , and kinetic, E_k , energies of the wave system is given, where

$$E_k = \frac{1}{2} \int \frac{\partial \eta}{\partial t} \Phi dx, \quad E_p = \frac{1}{2} g \int \eta^2 dx. \quad (6)$$

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

According to the linear theory, the portions of kinetic and potential energy are equal at the initial moment. This balance is not retained by nonlinear waves. As a result, parasitic counter waves are naturally excited in course of evolution of the improper initial condition. They are well seen in the (ω, κ) Fourier diagrams of the wave evolution (see Slunyaev and Sergeeva, 2012), and also may be revealed due to the oscillations, which are readily seen in Fig. 1a. Similar oscillations with the period of half dominant wave period are typical for the case of standing waves (in other words: existence of two opposite wave systems).

To avoid this undesired effect in the simulations, the nonlinear parts of Eqs. (1)–(4) were turned on slowly to enable the waves to adjust to the nonlinearity normally. The dependences of potential and kinetic energies in this statement are shown in Fig. 1b. The time interval shown in Fig. 1 corresponds to about 20 wave periods, while the typical time scale of this adjustment is one wave period. A smooth departure of the kinetic and potential parts of energy is observed. Simultaneously, the amount of counter waves diminishes gradually.

Generation of the initial condition in the form of a superposition of sinusoids with random phases implicitly implies the assumption of a Gaussian wave statistics. Besides establishing correct proportions between the kinetic and potential energies due to generation of proper phase-locked bound waves, the initial stage of wave evolution may undergo a longer transition process, which is related to the forming of natural nonlinear wave groups. This effect is primarily due to the four-wave interactions, and has typical duration of order $\varepsilon^{-2}T_p$ (where ε is the wave steepness). This effect has been observed many times in laboratory and numerical experiments, see among others Shemer and Sergeeva (2009, 2011); Shemer et al. (2010a, b); Dysthe et al. (2003); Janssen (2003); Onorato et al. (2005); Annenkov and Shrira (2009).

For given wave intensity, this transition process occurs in case of sufficiently narrow spectrum and is characterized by a temporal spectrum widening with subsequent partial relaxation. During the transition stage the probability of extreme waves increases greatly. After this acute stage a quasistationary state is attained. Such a process is

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

most pronounced in the case of large values of the Benjamin–Feir Index what may happen when the JONSWAP spectrum is characterized by a large value of peakedness, γ . The transition process is not appreciable if the waves are weakly nonlinear or the spectrum is wide (BFI is small). The quasistationary condition is achieved within a few unites of the “nonlinear time” $\sim \varepsilon^{-2}T_p$.

In our numerical simulations first 200 s of the wave evolution are allocated for the transient processes and are not used for the analysis. The wave system is assumed to be the proper *sea state* for $t > 200$ s.

Figures 2 and 3 show the spectra which characterize the obtained sea states A and E correspondingly. These Fourier spectra are averaged; they are built on the basis of all simulated data using 20 realizations. Dashed curves in the figures indicate the corresponding JONSWAP spectrum, which is used for generation of the initial condition.

It may be noticed that the obtained spectrum is somewhat distorted due to the nonlinear processes and moderate number of realizations used for the averaging. However, the agreement between the modeled and the obtained spectra is pretty well. A significant variation of the wavenumber spectrum in time was not observed, what confirms the closeness of the simulated sea states to stationary conditions.

A peak frequency downshift may be seen for the steeper sea state in Fig. 3, which is a well-known effect. Besides, some faster decay of the simulated spectrum tail, in comparison with the JONSWAP curve, is noticeable in Fig. 3. This effect is due to the effective high-frequency wave dissipation, which was introduced to damp the processes of wave breaking in the way similar to Chalikov (2005). Otherwise, too steep waves, which appear during the wave evolution, cause numerical instabilities, and the simulation blows up shortly. Thus, the desired sea state E is not reproduced *in full*, the sea state corresponds to a slightly different condition of sea wave evolution.

The HOSM is unable to resolve very steep waves; it also suffers from instabilities what may normally reduce its capability of long-time simulations. High intensity of waves in the case E and slowly decaying spectrum tail of the JONSWAP function provide favorable conditions for occurrence of very steep waves. Sometimes the employed

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



high-frequency damping filter was not able to prevent occurrence of a small-scale instability. Then the simulation was interrupted, the steep wave was artificially removed from the field, and the simulation was continued. As a result of the introduced damping and removal of too much sharp waves, the root-mean-square surface elevation, σ , might decrease during the simulation of in total about 23 min up to about 10 %.

3 Analysis of the spatio-temporal surface elevations

The simulated spatio-temporal wave fields of the surface elevation with high resolution in time and space give the comprehensive information about wave shapes, and make possible a study of rogue wave's dynamics in detail. Since the considered spatial domain is as large as about 10 km, the probability of extreme waves in a certain time interval increases in comparison with the probability calculated for time records in a given spatial point. Thus, almost every simulated realization contains several abnormal waves, while the linear theory for the Gaussian sea predicts such a wave to occur in a single point once during about 8 h (Kharif et al., 2009).

The widely used criterion for rogue wave definition is based on the amplification factor, AI (abnormality index), formulated in terms of the wave height, H . The simple common definition of a rogue wave reads

$$AI > 2, \quad AI = \frac{H}{H_s}. \quad (7)$$

Here H_s – is the significant wave height determined as the average value of the one third part of highest waves in a record. For the Gaussian sea this parameter could be written with use of the root-mean-square of surface elevation as $H_s \approx 4\sigma$. We should mention that the statistics of simulated nonlinear waves deviates from the Gaussian one, and values of H_s , calculated according to the general definition, differ from the value of 4σ , what leads to a different criterion for the rogue wave selection. For simplicity, in this paper for selecting rogue waves we use criterion (7) with H_s taken equal to

4σ , i.e. for rogue waves $H > 8\sigma$. Totally about 800 rogue waves are found for the sea state A, and about 4500 waves for the more severe case E.

One of the aims of the present work is to consider the variety of shapes of rogue waves. To this end, individual waves must be singled out. Usually, this is performed by the zero-crossing approach, see e.g. Holthuijsen (2007). It is substantial that in the present case we deal with space series, but not time series. It is very well understood that in deep-water conditions waves have effectively narrower frequency spectrum in comparison to the wavenumber spectrum. As a consequence, the portion of short-scale waves which can be distinguished on water surface is greater in the spatial domain. They look like ripples on the background of wind waves; many of them are recognized as individual waves according to the naive application of the zero-crossing approach. The large amount of small-scale weak-amplitude waves distorts statistical properties (such as typical wave length, significant wave height) greatly. To avoid this spurious effect, wave filtering was introduced. First, individual waves were separated following the basic zero-crossing approach (hereafter this selection of individual waves in spatial series will be referred to as raw data). If an individual wave happens to be significantly shorter than the peak wave length (shorter than $L_0/5$, where $L_0 = T_0^2 g / (2\pi)$), it is merged by the neighbouring individual wave. Therefore, the number of individual waves after the filtering is smaller than the one in the raw data, and the significant wave height is, correspondingly, larger in the filtered data.

3.1 Probability distribution functions

We have already noted the difference between statistical data, used in this study, and composed by conventional time series retrieved at a single point: not all individual waves among about 2.5 millions are independent. Therefore, the direct comparison between the statistical results becomes impossible, and it is desirable to understand at least the main features of the difference.

First, we employ the advantage of fully-controllable numerical simulations, and conduct numerical simulations of 20 realizations for the initial condition A, when nonlinear

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



effects are switched off totally (this numerical experiment will be hereafter referred to as AL), and then produce the exceedence probability distribution function on the basis of the 2.5 millions of individual waves. The result of processing is shown in Fig. 4a. The probability distribution functions are produced for the raw data and for the filtered data as well and are denoted as H_s and H_s^f respectively; inequality $H_s^f \geq H_s$ takes place. The significant wave heights are defined as a mean of significant heights of zero down-crossing waves and up-crossing waves. The horizontal axis in Fig. 4 shows the normalized height, where the normalization factor, H_{ref} , is given in the figure legend. The Rayleigh probability function, which follows from the linear theory for narrow band waves, is given for the reference by a thick line.

The first important observation is that no rogue waves occur in simulation AL at all, since all the curves in Fig. 4a fall down at $H < 2H_{ref}$. This fact is obviously caused by insufficient volume of the statistical ensemble. The number of narrow-banded individual waves in the ensemble, which results in the Rayleigh distribution for wave heights, exceeding 4σ , was estimated in Pelinovsky and Kokorina (2002) to be about 10 000. Even a greater number of realizations should be simulated with the purpose to study the rogue wave statistics in conditions AL. There is practically no difference between curves for $H_{ref} = H_s^f$ and $H_{ref} = H_s$. The tail of the curve for $H_{ref} = 4\sigma$ is somewhat lower than the one for $H_{ref} = H_s$, what is because the value of 4σ exceeds significant wave height, and both the tails are lower than the Rayleigh function.

Figure 4b and c display the results of nonlinear wave evolution. Appearance of long tails is evident, waves exceeding $H > 2H_{ref}$ occur in all cases. The PDF's tails for the ensemble of non-filtered waves exceed theoretical curve when $H_{ref} = 2H_s$ both in Fig. 4b (moderately intense waves, A) and in Fig. 4 (severe wave conditions, E); for these cases $H_s < \sim 4\sigma$, similar to conditions AL. At the same time the calculated PDF with scaling $H_{ref} = 4\sigma$ is lower than the Rayleigh curve in case A (Fig. 4b) and seems to be a little bit higher than it in case E (Fig. 4c). Note that slightly different definition of the quantity H_s is able to change the positioning of the calculated curves with respect to the Rayleigh function.

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



The impact of the wave filtering, which sifts out short-scale waves, becomes very important in nonlinear simulations (Fig. 4b and c), in contrast to the linear case (Fig. 4a). Due to the filtering the amount of small waves decreases greatly, H_s^f grows respectively, and the number of waves with $H > 2H_s^f$ is much smaller than of waves with $H > 2H_s$ or $H > 8\sigma$. As a result, the probability distribution function for filtered data falls below the Rayleigh function. We emphasize that in the present study the wave processing is performed for the space series, but not conventional time-series. Because frequency wave spectrum of deep-water waves is about twice narrower than the wavenumber spectrum, it is natural to expect that the effect of small-scale waves on the distribution function in the case of time-series processing is less prominent.

The study of wave auto-correlation in time and space is illustrated in Fig. 5. The wave correlation is estimated with the help of the following function,

$$R(t, \Delta t) = \max_{\Delta x} \frac{\int \eta(t, x) \eta(t + \Delta t, x + \Delta x) dx}{\sqrt{\int \eta^2(t, x) dx} \sqrt{\int \eta^2(t + \Delta t, x) dx}}, \quad (8)$$

where the integration is made over the spatial domain of simulations. For simplicity, function $R(0, \Delta t)$ is examined and is plotted in Fig. 5. Three curves in the figure describe case A, when simulated within the linear framework (AL, dashed line), by means of the HOSM solver (thin solid line), and the case E (thick solid line). Figure 5 shows very different pictures for the linear and non-linear wave dynamics. It reports that the level of correlation decays much faster in the nonlinear case in comparison to the linear simulation. Sea states with relatively small waves and with steeper waves do not differ significantly.

According to the analysis, the effect of higher probability of high waves with respect to the linear theory, observed in the present simulations (see Fig. 4), is assumed by us to be twofold. First, nonlinearity is generally known to increase the likelihood of high waves; this effect was observed both, in deep and shallow water conditions (Onorato et al., 2001, 2002; Pelinovsky and Sergeeva, 2006; and many others). Qualitatively this

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



effect is in agreement with the formation of long-living nonlinear wave structures (coherent nonlinear wave groups in deep water, and solitary waves in shallow water). The second effect, which becomes vivid due to Fig. 5, is shortening of the wave correlation length, more waves among the 2.5 millions become effectively independent due to nonlinear effects, and hence the volume of the statistical ensemble effectively increases, and higher waves may be found.

3.2 Pictures of rogue waves and rogue events

An example of the simulated (x, t) sheet of surface elevations with detected rogue waves is given in Fig. 6. Filled circles mark extreme waves with $AI > 2$. As it is described above, the spatial domain is periodic, and space series are analyzed.

If to follow the evolution of a rogue wave, the wave height may get smaller than the threshold value (7), which specifies rogue waves, for some short time, and may exceed the threshold again shortly later. Thus, such extreme wave in the course of its evolution satisfies criterion (7) in a discontinuous manner. It is naturally to consider this sequence of extreme waves as a *rogue event*, even if the wave does not satisfy condition $AI > 2$ at all instants of its lifetime. Therefore, individual waves, which satisfy the rogue wave criterion (7), are assumed to belong to one *rogue event*, if they are located in space and time not further from each other than 2.5 characteristic wave lengths and 2.5 characteristic wave periods.

In Fig. 6 clustered rogue events are shown by frames, which assemble rogue waves (filled circles). Very short-living single rogue waves (of one wave period or less), continuous and intermittent long-living rogue events (lasting for many wave periods) may be observed in the figure. (Note periodic boundary conditions for the simulated spatial domain.) Even a longer rogue event might be distinguished in the figure (from 150 s to about 550 s), if a longer separation between the individual rogue waves is allowed.

Of course, the life time of a rogue event can be significantly longer than for continuous rogue waves. The spread of values of rogue event's life-times versus the abnormality index is pictured in Fig. 7 for series A, B, C and E. The figure claims that rogue events

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



may last for a few dozens of characteristic wave periods, up to about 10 min. This lifetime exceeds significantly some other estimations on the rogue wave lifetime including the results of our previous studies, where the lifetimes were estimated to be up to about 1.5 min. Figure 7 exhibits the trend that more intense waves live for longer times.

Typical shapes of rogue waves observed in the numerical simulations are shown in Fig. 8. The bold lines depict the rogue waves, how they are segregated by the present processing. Most of rogue waves are high crests (Fig. 8a) or skew-symmetric surfaces (Fig. 8b); besides, extreme waves with deep troughs (“holes in the sea”) are also found (Fig. 8c). Thin lines in Fig. 8 correspond to surface elevations at consecutive time instants close to the moment of the rogue wave occurrence.

Two approaches to selecting wave heights are usually applied to processing wave records: zero down-crossing or zero up-crossing (e.g. Holthuijsen, 2007). Note, that an up-crossing wave in a time series becomes a down-crossing wave, when its momentary shape is considered as a function of coordinate (and the coordinate axis is chosen along the wave propagation). In the present analysis, a rogue wave owns an extreme vertical excursion between the wave crest and the wave trough; these crest and trough, each localized according to the zero-crossing method, constitute the rogue wave, as shown in Fig. 8 by bold lines. A rogue wave, which we select, may be either up- or down-crossing depending on the mutual location of the wave crest and trough, which constitute the rogue wave. Waves in Fig. 8 represent space series of waves which propagate to the right. Extreme waves in Fig. 8a and c (given by the bold line) are zero down-crossing waves, and will be referred to as extreme rear slope waves. The spatial record of extreme wave given in Fig. 8b is an up-crossing wave and is represented by an extreme frontal slope.

Variety of geometries of rogue waves found in the numerical simulations is illustrated in Figs. 9 and 10. Figure 9 displays the relation between the ratio of crest amplitude to trough amplitude (vertical axis) versus the ratio of crest length to trough length (horizontal axis) for wave conditions A (Fig. 9a) and E (Fig. 9b). The distinction is made between the cases when the rogue wave has a huge frontal slope (shown with circles)

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



or an extreme rear slope (shown with crosses). In Fig. 9 all the rogue waves found in 20 realizations for cases A and E are collected. Figure 9 proves that extreme waves with high crests are prevalent in comparison with extremely deep troughs. Higher crests are shorter in length.

5 The proportion between rogue waves of different shapes is displayed in Fig. 10 for sea states A and E, respectively. Four kinds of rogue waves are distinguished regarding the vertical asymmetry and orientation of the wave (extreme frontal or rear slope). Both Fig. 10a and b claim prevalence of waves with high crests (“positive” waves, 85–95%), while waves with deep troughs (“negative” waves) are much less in number, what agrees with results in Fig. 9.

10 For less intense waves (Series A, Fig. 10a) the proportion between up-crossing and down-crossing rogue waves is about the same, whereas in the steeper wave case (Series E, Fig. 10b) the total number of recognized rogue waves is much greater, and the number of rogue waves with extreme rear slopes (61 %) is significantly larger than extreme fronts (34 %), thus in the majority of cases a ship should experience first a very high wave crest, and then a subsequent deep trough. This picture is opposite to the one known in relation to regions with strong opposite currents (Mallory, 1974; Lavrenov, 1998), when a deep trough precedes a high crest (a huge wave front), but agrees very well with the in-situ measurements by Pinho et al. (2004), where 276 cases of rogue waves result in the proportion between waves with large rear and frontal slopes 197 : 108.

4 Extreme kinematics of waves versus kinematics of extremely high waves

25 Establishing the relation between the extreme surface elevation and extreme wave kinematics is an important issue. Conventionally the rogue wave criterion is based on the amplitude characteristics (7). The reason for that is evident: although waves are in fact a complicate phenomenon represented by the movement of fluid particles, the surface elevation is much easier to observe and to register.

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Through the performed simulations a large amount of information on wave kinematics is available, what enables us to conduct a kind of statistical study of rogue wave kinematics characterizing particular sea states. The employed numerical model cannot simulate wave breaking; hence breaking waves are beyond the scopes of the present analysis.

In this section we consider the relation between extreme velocities in the simulated wave fields, and velocities of rogue waves (the latter are singled out on the basis of the wave height criterion (7)) and address the question: do rogue waves manifest faster velocities?

The Eulerian horizontal and vertical velocities of the water surface, V_x and V_z are defined by

$$V_x \equiv \left. \frac{\partial \phi}{\partial x} \right|_{z=\eta} = \frac{\Phi_x - \eta_t \eta_x}{1 + \eta_x^2}, \quad V_z \equiv \left. \frac{\partial \phi}{\partial z} \right|_{z=\eta} = \frac{\eta_t + \Phi_x \eta_x}{1 + \eta_x^2}. \quad (9)$$

The formula for V_z results from Eq. (1), and the formula for V_x comes from the identity $\Phi_x = \phi_x + \eta_x \phi_z$ at $z = \eta$.

A simulation of one irregular wave realization results in 2048×2048 values of velocities (9). They are plotted in diagrams in Fig. 11 by grey points (V_x versus V_z) for five realizations of sea states A (Fig. 11a) and E (Fig. 11b). The extreme values of horizontal (black points) and vertical (blue points) velocities, which are attained by the segregated rogue waves (see examples given by bold lines in Fig. 8), are shown for all 20 realizations, simulated for conditions A and E.

For the reference, velocities at the water surface for the 3rd order Stokes wave solution are shown in Fig. 11 (contours) according to formulas, see e.g. (Fenton, 1985; Slunyaev, 2005)

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$$V_x = \sqrt{k_0 g A} \left(1 - \frac{1}{8} k_0^2 A^2 \right) \exp(k_0 \eta) \cos \theta, \quad (10a)$$

$$V_z = \sqrt{k_0 g A} \left(1 - \frac{1}{8} k_0^2 A^2 \right) \exp(k_0 \eta) \sin \theta, \quad (10b)$$

$$\eta = A \cos \theta + \frac{k_0}{2} A^2 \cos 2\theta + \frac{3}{8} k_0^2 A^3 \cos 3\theta. \quad (10c)$$

Here A is the wave amplitude, and $0 < \theta < 2\pi$ is the wave phase; k_0 is the carrier wavenumber. The contours are plotted in Fig. 11 for two wave intensities: (i) when the wave height is equal to the significant wave height, H_s (then $A = H_s/2 - 3/64 k^2 H_s^3$, the smaller contour), and (ii) for the maximum wave height value registered in the simulated wave fields, H_{\max} (then $A = H_{\max}/2 - 3/64 k^2 H_{\max}^3$, the larger contour).

The larger contour corresponds to the highest possible velocity of surface waves if they would be described by the 3rd order Stokes wave solution (10). The area inside the contour corresponds well to the major amount of velocities in the both cases, Series A and E. The inner contour allocates the range of velocities of waves with heights smaller than H_s ; this area of velocities (V_x , V_z) is practically free of rogue waves. The 5th order Stokes wave solution for surface velocities does not change significantly the picture shown in Fig. 11a (Series A), it is quite similar to the 3rd order velocities. For severer Series E the value of H_{\max} seems to be so much that the asymptotic solution for wave velocities does not converge well.

Asymmetries of the vertical and horizontal velocity fields are well-pronounced, the velocities, which correspond to rogue waves, have qualitatively similar distribution asymmetry. The asymmetries strengthen for a steeper wave sea state.

Figure 11 proves inability of the high order Stokes wave theory to describe the rogue wave kinematics, although the major part of velocities attained by simulated rogue waves lies within the ranges, which correspond to the highest registered wave. Some “blind” areas in Fig. 11 are evident (where only grey points exist), which correspond to intense velocities (they may even exceed the limit set by the Stokes wave solution)

and at the same time to relatively small wave amplitudes (rogue waves with these fluid velocities are not found in the records). On the other hand, some rogue waves exhibit quite moderate values of velocities.

5 Conclusions

5 In the paper we present the approach to a more extensive use of data on rogue wave dynamics, statistics, and kinematics, which becomes available through stochastic simulations of irregular waves.

In this study waves characterized by the JONSWAP spectrum are simulated by means of the strongly nonlinear algorithm for solving the Euler equations, – the HOSM. 10 Two sea states are considered in more details – of moderately high waves and steeper waves. For each sea state 20 realizations are simulated, each results in 10 km × 20 min sheets of data with good resolution of the surface elevation, fluid velocity, etc.

The accumulated data may be used for generating time- or spatial wave records, and then used for verification and adjustment of processing techniques. The availability of 15 full spatio-temporal information about the wave field provides with opportunity to clarify the dispersion relation for nonlinear waves, where wave components (free, bound and counter-directional modes) are naturally separated (as discussed in Slunyaev and Sergeeva, 2012).

In the present study the main interest is focused on characteristics of rogue waves 20 which occur in the simulated fields. The simulated data contains a number of rogue waves, available for analysis. Space series are used, because of the peculiarity of periodic spatial boundary conditions. Spatial time series (one- and two-dimensional) of the surface elevation, which are recorded in laboratory and in-situ conditions become nowadays available due to developing of new measuring techniques. Hence, the direct 25 comparison between the theoretical simulation and measurements becomes potentially possible.

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The analysis of characteristics of individual rogue waves, found in our simulations, shows the variety of their shapes and appearances. Most of the simulated rogue waves have very high crests and shallow long troughs, or sign-variable shapes. “Holes in the sea”, characterized by deep troughs, are found in the simulation data as well. In severe conditions the predominance of rogue waves with extreme rear slopes is discovered compared to rogue waves with high fronts.

The employed approach allows observing a full picture of the rogue wave propagation. Some of extreme waves appear suddenly and live for a short time, less than a wave period. Large amount of rogue waves’ heights oscillate with time and exceed the formal criterion of a rogue wave, $H > 2H_s$, repeatedly, and thus may be considered as *rogue events* which can live significantly longer than the wave, which satisfies the rogue wave amplitude criterion continuously. As a result, the life-time of extremely high wave events varies from several seconds to tens of characteristic wave periods.

Water velocities at the surface are considered in relation to the rogue wave phenomenon. The velocities of extreme waves are compared versus extreme kinematics of all waves and with predictions of the 3-order and 5-order nonlinear theories for Stokes waves. The difference between the kinematics of extremely high waves and the extreme kinematics is emphasized, whereas the significant amount of data well correspond to each other. Rogue waves are typically characterized by large values of velocities, but high velocities do not necessarily correspond to rogue waves, and rogue waves may be characterized by moderate fluid velocities.

Acknowledgements. The research is supported by the EC’s Seventh Framework Programme FP7-SST-2008-RTD-1 project EXTREME SEAS – Design for Ship Safety in Extreme Seas under grant agreement No 234175, RFBR grants 11-02-00483, 11-05-92002, 12-05-33087 and 11-05-00216. ASe acknowledges support from the Volkswagen Foundation. ASI acknowledges support from the EC’s Programme FP7-PEOPLE-2009-IIF under grant agreements No 254389, 909389.

References

- Annenkov, S. Y. and Shrira, V. I.: Evolution of kurtosis for wind waves, *Geophys. Res. Lett.*, 36, L13603 doi:10.1029/2009GL038613, 2009.
- Chalikov, D.: Statistical properties of nonlinear one-dimensional wave fields, *Nonlin. Processes Geophys.*, 12, 671–689, doi:10.5194/npg-12-671-2005, 2005.
- Chalikov, D.: Freak waves: Their occurrence and probability, *Phys. Fluids*, 21, 076602 doi:10.1063/1.3175713, 2009.
- Christou, M. and Ewans, K.: Examining a comprehensive dataset containing thousands of freak wave events. Part 1 – description of the data and quality control procedure, *Proceedings of 30th International Conference on Ocean, Offshore and Arctic Engineering (OMAE-2011)*, OMAE2011-50168 doi:10.1115/OMAE2011-50168, 2011.
- Clamond, D., Francius, M., Grue, J., and Kharif, C.: Long time interaction of envelope solitons and freak wave formations, *Eur. J. Mech. B/Fluids*, 25, 536–553, 2006.
- Dommermuth, D. and Yue, D. K. P.: A high-order spectral method for the study of nonlinear gravity waves, *J. Fluid Mech.*, 184, 267–288, 1987.
- Dysthe, K. B., Trulsen, K., Krogstad, H. E., and Socquet-Juglard, H.: Evolution of a narrow-band spectrum of random surface gravity waves, *J. Fluid. Mech.*, 478, 1–10, 2003.
- Dysthe, K. B., Krogstad, H. E., and Müller, P.: Oceanic rogue waves, *Annu. Rev. Fluid. Mech.*, 40, 287–310, 2008.
- Fenton, J. D.: A fifth-order Stokes theory for steady waves, *J. Waterway, Port Coast. Ocean Eng.*, 111, 216–234, 1985.
- Holthuijsen, L. H.: *Waves in oceanic and coastal waters*, Cambridge Univ. Press, 2007.
- Janssen, P. A. E. M.: Nonlinear four wave interactions and freak waves, *J. Geophys. Oceanogr.*, 33, 863–884, 2003.
- Kharif, C., Pelinovsky, E., and Slunyaev, A.: *Rogue Waves in the Ocean*, Berlin Heidelberg, Springer-Verlag, 2009.
- Lavrenov, I.: The wave energy concentration at the Agulhas current of South Africa, *Nat. Hazards*, 17, 117–127, 1998.
- Mallory, J. K.: Abnormal waves on the south-east of South Africa, *Int. Hydrog. Rev.*, 51, 99–129, 1974.

NHESSD

1, 39–72, 2013

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Mori, N., Onorato, M., Janssen, P. A. E. M., Osborne, A. R., and Serio, M.: On the extreme statistics of long-crested deep water waves: Theory and experiments, *J. Geophys. Res.*, 112, C09011, doi:10.1029/2006JC004024, 2007.
- Onorato, M., Osborne, A. R., Serio, M., and Bertone, S.: Freak waves in random oceanic sea states, *Phys. Rev. Lett.*, 86, 5831–5834, 2001.
- Onorato, M., Osborne, A. R., and Serio, M.: Extreme wave events in directional, random oceanic sea states, *Phys. Fluids*, 14, L25–L28, 2002.
- Onorato, M., Osborne, A. R., Serio, M., and Cavaleri, L.: Modulational instability and non-Gaussian statistics in experimental random water-wave trains, *Phys. Fluids*, 17, 078101 doi:10.1063/1.1946769, 2005.
- Pelinovsky, E. and Kokorina, A.: The applicability of the Korteweg–de Vries equation for description of the statistics of the freak waves, *J. Korean Soc. Coastal Ocean Eng.*, 14, 308–318, 2002.
- Pelinovsky, E. and Sergeeva, A.: Numerical modeling of the KdV random wave field, *Eur. J. Mech. B/Fluid*, 25, 425–434, 2006.
- Pinho, U. F., Liu, P. C., and Ribeiro, C. E. P.: Freak Waves at Campos Basin, Brazil, *Geofizika*, 21, 53–67, 2004.
- Shemer, L. and Sergeeva, A.: An experimental study of spatial evolution of statistical parameters of unidirectional narrow-banded random wave field, *J. Geophys. Res.*, 114, C01015, doi:10.1029/2008JC005077, 2009.
- Shemer, L., Sergeeva, A., and Liberzon, D.: Effect of the initial spectral shape on spatial evolution of the statistics of unidirectional nonlinear random waves, *J. Geophys. Res.*, 115, C12039 doi:10.1029/2010JC006326, 2010a.
- Shemer, L., Sergeeva, A., and Slunyaev, A.: Applicability of envelope model equations for simulation of narrow-spectrum unidirectional random field evolution: experimental validation, *Phys. Fluids*, 22, 016601 doi:10.1063/1.3290240, 2010b.
- Slunyaev, A. V.: A high-order nonlinear envelope equation for gravity waves in finite-depth water, *J. Experiment. Theor. Phys.*, 101, 926–941, 2005.
- Slunyaev, A. V. and Sergeeva, A. V.: Stochastic simulation of unidirectional intense waves in deep water applied to rogue waves, *JETP Letters*, 94, 779–786, 2011.
- Slunyaev, A. V. and Sergeeva, A. V.: Numerical simulations and analysis of spatio-temporal fields of rogue waves, *Fundament. Appl. Hydrophys.*, 5, 24–36, 2012 (In Russian).

- Slunyaev, A. V., Didenkulova, I., and Pelinovsky, E.: Rogue Waters, Contemporary Physics, 52, 571–590, 2011.
- Socquet-Juglard, H., Dysthe, K. B., Trulsen, K., and Krogstad, H. E.: Probability distributions of surface gravity waves during spectral changes, J. Fluid Mech., 542, 195–216, 2005.
- 5 West, B. J., Brueckner, K. A., Janda, R. S., Milder, D. M., and Milton, R. L.: A new numerical method for surface hydrodynamics, J. Geophys. Res., 92, 11803–11824, 1987.
- Zakharov, V.: Stability of periodic waves of finite amplitude on a surface of deep fluid, J. Appl. Mech. Tech. Phys., 2, 190–194, 1968.

NHESSD

1, 39–72, 2013

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Table 1. Parameters of the sea states.

Series code	A	B	C	E
T_0 , s	10	10	10	10.5
H_s , m	3.5	7	9	7
Peakedness, γ	3	3	3	3.3

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion




Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev



a)



b)

Fig. 1. Potential and kinetic energy of waves in the numerical simulations. Case **(b)** corresponds to a slow switching on the nonlinearity in the governing equations. Sea state A.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	




Rogue waves, rogue events and extreme wave kinematicsA. Sergeeva and
A. Slunyaev

Fig. 2. Averaged frequency spectrum in linear **(a)** and semi-logarithmic **(b)** axis, and also the wavenumber spectrum in semi-logarithmic axis **(c)** for the sea state A. Dashes curves show the modeled JONSWAP spectrum.

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev




Fig. 3. Averaged frequency spectrum in linear **(a)** and semi-logarithmic **(b)** axis, and also the wavenumber spectrum in semi-logarithmic axis **(c)** for the sea state E. Dashes curves show the modeled JONSWAP spectrum.

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Wave height exceedence probability distribution functions for conditions A simulated within the linear framework (case AL) **(a)** and by means of HOSM ($M = 6$) **(b)**, and for condition E simulated by means of the HOSM ($M = 3$) **(c)**. The PDF are produced for the raw and filtered data; wave heights are normalized by different factors H_{ref} (see legend and discussion in the text). The theoretical Rayleigh distribution (thick line) is given for the reference.

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev




Fig. 5. Wave correlation $R(0, \Delta t)$ for conditions A (thick solid line) and E (thin solid line) and also for case A, when the evolution is simulated within the linear theory, AL (dashed line).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 6. Spatio-temporal sheet of the surface elevation. Markers indicate rogue waves, and rectangles assemble them into *rogue events*.

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion


Rogue waves, rogue events and extreme wave kinematicsA. Sergeeva and
A. Slunyaev

Fig. 7. Normalized life-time of rogue events versus the abnormality index, AI.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev




Fig. 8. Examples of rogue wave's profiles (bold lines), and surface elevations at consecutive time points: single crest (a), sign-variable wave (b), deep hole (c).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev




Fig. 9. Relation between vertical wave asymmetry (crest amplitude to trough amplitude) and horizontal asymmetry (crest length to trough length) for two types of rogue waves (circles show cases with extreme wave fronts, while crosses denote extreme rear slopes) for series A **(a)** and E **(b)**.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev




Fig. 10. The proportions of waves of different types in the total mass of rogue waves for series A (a) and E (b).

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Rogue waves, rogue events and extreme wave kinematics

A. Sergeeva and
A. Slunyaev




Fig. 11. Grey points are all the values of vertical versus horizontal surface velocities in the (x, t) fields (5 realizations are used). Black points are the maximum horizontal and blue points are the maximum vertical velocities attained by the detected rogue waves (all 20 realizations are used to plot these values): **(a)** Series A, **(b)** Series E. Contours correspond to the values for the 3rd order Stokes wave solution of the heights H_s (the smaller contour) and H_{\max} (the larger contour).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion