Nat. Hazards Earth Syst. Sci. Discuss., 1, C1494–C1495, 2013 www.nat-hazards-earth-syst-sci-discuss.net/1/C1494/2013/

© Author(s) 2013. This work is distributed under the Creative Commons Attribute 3.0 License.



NHESSD

1, C1494-C1495, 2013

Interactive Comment

Interactive comment on "Sensitivity of the WRF model to the lower boundary in an extreme precipitation event – Madeira Island case study" by J. C. Teixeira et al.

V. Levizzani

v.levizzani@isac.cnr.it

Received and published: 21 October 2013

This is indeed a very interesting paper on the sensitivity of the WRF model to such extreme conditions. I believe it will be of help when using the model to try account for similar cases, for example in the Mediterranean. The modeling progresses will certainly come from very high resolutions and thus need to take into account the effects described in the paper. Just for information, the authors may want to take a look at a recent paper depicting the orographic effect on precipitation as estimated from satellite passive microwave data during the same Madeira 2010 event:

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Levizzani, V., S. Laviola, E. Cattani, and M. J. Costa, 2013: Extreme precipitation on the Island of Madeira on 20 February 2010 as seen by satellite passive microwave sounders. European J. Remote Sensing, 46, 475-489, doi: 10.5721/EuJRS20134628.

Interactive comment on Nat. Hazards Earth Syst. Sci. Discuss., 1, 5603, 2013.

NHESSD

1, C1494-C1495, 2013

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

