1 Regional-scale analysis of high-mountain multi-hazard and risk in

2 the Pamir (Tajikistan) with GRASS GIS

3 Fabian E. Gruber, Martin Mergili*

4 Institute of Applied Geology, University of Natural Resources and Life Sciences (BOKU),

5 Vienna, Peter-Jordan-Straße 70, 1190 Vienna, Austria

6 *Corresponding author

7 Email: martin.mergili@boku.ac.at, Tel.: +43 47654 5412, Fax: +43 47654 5449

8 Abstract

9 We present a model framework for the regional-scale analysis of high-mountain multi-hazard and 10 -risk, implemented with the Open Source software package GRASS GIS. This framework is applied to a 98,300 km² study area centred in the Pamir (Tajikistan). It includes (i) rock slides, (ii) 11 12 ice avalanches, (iii) periglacial debris flows and (iv) lake outburst floods, First, a hazard indication score is assigned to each relevant object (steep rock face, glacier or periglacial slope, lake). This 13 14 score depends on the susceptibility and on the expected event magnitude. Second, the possible 15 travel distances, impact areas and, consequently, impact hazard indication scores for all types of 16 processes are computed using empirical relationships. These scores are finally superimposed with 17 an exposure score derived from the type of land use, resulting in a raster map of risk indication scores finally discretized at the community level. The analysis results are presented and discussed 18 19 at different spatial scales. The major outcome of the study, a set of comprehensive regional-scale hazard and risk indication maps, shall represent an objective basis for the prioritization of target 20 21 communities for further research and risk mitigation measures.

22 Keywords

23 Central Asia, GRASS GIS, Mass movements, Multi-hazard, Regional scale, Risk indication map

1 1 Introduction

High-mountain areas are commonly experiencing pronounced environmental changes such as
permafrost melting and the retreat of glaciers, caused by atmospheric temperature increase
(Beniston, 2003; Huber et al., 2005; IPCC, 2007; WGMS, 2008; Harris et al., 2009). Together
with earthquakes or volcanic eruptions, they disturb the dynamic equilibrium of the fragile highmountain geomorphic systems, leading to an increased occurrence of rapid mass movements
(Evans and Clague, 1994; Huggel et al., 2004a,b; Kääb et al., 2005; IPCC, 2007; Quincey et al.,
2007; Harris et al., 2009; Dussaillant, 2010; Haeberli et al., 2010a).

9 Whilst such mass movements often occur in remote areas and remain unrecognized, they may also evolve into long-distance flows affecting the communities in the valleys. Such processes are 10 11 referred to as remote geohazards. They are commonly related to the massive entrainment of 12 loose material or the interaction of two or more process types (process chain). Several cases are evident where slope failures including rock and/or ice have converted into long-distance 13 avalanches and consecutive processes. A striking example is the 1970 Huascarán event (Cordillera 14 Blanca, Peru) where several 1000 people lost their lives in the town of Yungay (Evans et al., 15 16 2009a). On September 20, 2002, a rock-ice avalanche in the Russian Caucasus entrained a glacier. 17 The resulting flow continued for 20 km as an avalanche of ice, rock and debris and for further 15 km as mud flow, resulting in approx, 140 fatalities (Kolka/Karmadon event; Huggel et al., 18 19 2005). On April 11, 2010, an ice avalanche from far upslope rushed into Laguna (Lake) 513 in the 20 Cordillera Blanca, causing a destructive outburst flood (Haeberli et al., 2010b).

Lakes are commonly involved in remote geohazard processes (Costa, 1985; Evans, 1986; Costa 21 22 and Schuster, 1988; Walder and Costa, 1996; Walder and O'Connor, 1997). Landslide-dammed lakes are of particular interest as most of them drain within the first year after their formation 23 24 (Costa and Schuster, 1988) whilst others persist for centuries. Glacial lakes, impounded by ice (Tweed and Russell, 1999) or (often ice-cored) moraines, are commonly coupled to retreating or 25 surging glaciers and therefore highly dynamic. Such lakes often occur in areas influenced by 26 27 permafrost. Some lakes are prone to sudden drainage (Glacial Lake Outburst Floods or GLOFs). 28 Studies of this phenomenon cover most glacierized mountain areas in the world such as the 29 Himalayas (Watanabe and Rothacher, 1996; Richardson and Reynolds, 2000; ICIMOD, 2011), the Karakorum (Hewitt, 1982; Hewitt and Liu, 2010), the Pamir (Mergili and Schneider, 2011), 30 the Tien Shan (Narama et al., 2010; Bolch et al., 2011), the Andes (Vilímek et al., 2005; 31 32 Harrison et al., 2006; Haeberli et al., 2010b), the North American mountains (Clarke, 1982), the Norwegian mountains (Breien et al., 2008) and the western Alps (Haeberli, 1983; Tinti et al., 33

1 1999; Huggel et al., 2002, 2003). GLOFs can evolve in different ways, for example by mass 2 movements into lakes, rising lake levels leading to overflow, progressive incision, mechanical 3 rupture or retrogressive erosion of a dam, hydrostatic failure or degradation of glacier dams or 4 ice-cores in moraine dams (Walder and Costa, 1996; Richardson and Reynolds, 2000). Peak 5 discharges are often some magnitudes higher than in the case of ordinary floods (Cenderelli and 6 Wohl, 2001). Entrainment may considerably increase the event magnitude and convert the flood 7 into a destructive debris flow.

8 A common feature of long-distance rock mass movements, ice avalanches, debris flows, lake 9 outburst floods and related process chains is their occurrence as rare (low frequency) or singular 10 events. The location, timing, magnitude and impact area of remote geohazard events are often hard or even impossible to predict, even though the governing processes are fairly well 11 understood and specific events were successfully back-calculated with deterministic computer 12 models (Evans et al., 2009a, b). This is particularly true where multiple hazards are evident over a 13 14 large area and/or where the resources for a broad-scale continuous monitoring of potentially hazardous situations are lacking, i.e. in developing countries. Here it is essential to identify 15 possible source and - particularly - impact areas of remote geohazard processes at the broad 16 (regional) scale in order to prioritize target areas for risk mitigation measures. Huggel et al. (2003, 17 2004a, b) and Mergili and Schneider (2011) have presented computer models suitable for the 18 19 regional-scale analysis of high-mountain hazards such as GLOFs, periglacial debris flows or ice 20 avalanches. Some of these models include process interactions. However, they neither attempt to account for the risk nor are they applied to very large areas. These gaps hamper a more focused 21 22 and comprehensive identification of possible target areas for risk mitigation.

Here we demonstrate a novel model framework for the regional-scale analysis of high-mountain 23 hazards and risks, including (i) rock slides, commonly converting into rock avalanches or, in 24 glacierized areas, into rock-ice avalanches, (ii) ice-avalanches, (iii) periglacial debris flows and (iv) 25 26 lake outburst floods, often evolving into flows of debris or mud. Examples of these results of processes, or of situations possibly leading to their occurrence, are illustrated in Fig. 2. Process 27 chains including more than one of the above process types are also considered. The study area in 28 the Pamir (Tajikistan, Central Asia) is introduced in Section 2. The data used for the study is 29 presented in Section 3 and the model framework is explained in detail in Section 4. Section 5 30 31 gives an overview of the model results which are discussed in Section 6. Section 7 summarizes the essence of the study. 32

1 2 Study area

A 98,300 km² study area in Central Asia is considered, extending from 1670 m a.s.l. near Khala-i-Khumb to 7495 m at the top of Ismoili Somoni Peak and largely corresponding to the headwaters of the Amu Darya River (Fig. 1). The northern and southern boundaries of the area are formed by the Alai and Hindukush ranges in Kyrgyzstan and Afghanistan. In between, the Pamir in the Gorno-Badakhshan Autonomous Oblast of Tajikistan represents the largest share of the study area.

8 The western Pamir is characterized by glacierized mountain ranges exceeding 6000 m a.s.l. and 9 deeply incised valleys. The eastern Pamir represents an arid highland above 3500 m a.s.l. with 10 glaciers covering only the highest peaks. The more humid northern Pamir with the Academy of 11 Sciences and Transalai ranges peaks above 7000 m a.s.l. and is extensively glacierized. The 12 Fedchenko Glacier extends over a length of >75 km and covers a surface area >700 km².

13 Intense tectonic uplift in combination with glacial and fluviatile erosion (Mahmood et al., 2008) 14 has resulted in a particularly pronounced relief. Consequently the geomorphic activity is high, including a large variety of mass wasting processes. They are commonly triggered by earthquakes 15 16 as the seismic activity and, therefore, the seismic hazard are significant (Giardini et al., 1999). Few 17 large historic events such as the 1911 Sarez rock slide (Schuster and Alford, 2004; Risley et al., 2006; see Fig. 2a) or the 1949 Khait rock avalanche (Evans et al., 2009b) are well documented. 18 19 The deposit of the 2 km² Sarez rock slide forms the 600 m high Usoi Dam, the highest dam worldwide. It retains the 60 km long Lake Sarez, the safety of which is still disputed (e.g., 20 Risley et al., 2006). 21

- The climate in the study area is temperate semi-arid to arid and continental with hot summers and cold winters. Most meteorological stations in the study area have recorded a positive trend of the mean annual air temperature (MAAT) in the period 1940–2000 (Makhmadaliev et al., 2008). The state of information suffers from a lack of up-to-date high-altitude meteorological data. According to the 4th IPCC report (IPCC, 2007), the median of the projected increase of the
- 27 MAAT from 1980–1999 to 2080–2099 for Tajikistan is 3.7 °C.
- Consequently, many glaciers are retreating (e.g. Khromova et al., 2006; Haritashiya et al., 2009; Mergili et al, 2012a), favouring the development of lakes in the glacier forefields or in subsiding areas on the glaciers. Mergili et al. (2013) detected a total number of 652 glacial lakes in the study area. A GLOF in 2002 caused dozens of fatalities, several more lakes are susceptible to sudden drainage (Mergili and Schneider, 2011; see Fig. 2d). Further, the retreat of glaciers over steep rock

cliffs (see Fig. 2b) may lead to the increased occurrence of ice avalanches. The shift of the
 permafrost boundary to higher areas results in the possible destabilization of rock and debris.
 Periglacial debris flows observed in the study area are most commonly associated with the termini
 of rock glaciers (see Fig. 2c).

5 The valleys in the study area are fairly densely populated, with Khorog as the only urban centre 6 (see Fig. 1). The local communities strongly depend on the natural resources and are therefore 7 affected by the consequences of the changing temperature regime in both positive and negative 8 ways (Kassam, 2009).

9 **3 Data**

The data the high-mountain multi-risk analysis builds on are summarized in Table 1. The ASTER GDEM V2, a product of METI and NASA, is used as input digital elevation model (DEM). It is provided at a cell size of approx, 30 x 30 m. Here persion resampled to 60 x 60 m is applied. Secondary data sets such as elevation with filled sinks, slope and flow direction are generated from the DEM which is further used to generate a gridded data set of the MAAT, making use of temperature data recorded at the stations of the Tajik HydroMet Agency and a vertical temperature gradient of 0.0062°C m⁻¹ (Müllebner, 2010; Fig. 3a).

The identification of areas with melting permafrost builds on the permafrost indication map for 17 Tajikistan presented by Mergili et al. (2012a): a set of rules-of-thumb for the lower boundaries of 18 19 sporadic and discontinuous permafrost in Switzerland (Haeberli, 1975) is adapted to the 20 conditions in Tajikistan. This set of rules is then combined with the DEM in order to produce a 21 gridded dataset indicating the possibility of permafrost occurrence for each raster cell. Applying the temperature gradient of Müllebner (2010), the effects of atmospheric temperature increase on 22 23 permafrost distribution are explored. Areas where the model predicts either sporadic or 24 discontinuous permafrost for the current state, but no permafrost of either of the two types for a temperature increase of +2°C or +4°C, represent separate classes. Such areas are of particular 25 26 interest for the permafrost susceptibility score S_{h} (Fig. 3b; see Section 4).

The seismic susceptibility of the area S_s is defined according to the peak ground acceleration with 10% chance of exceedance in 50 years (PGA), expressed in relation to gravity *g*. The Global

- 29 Seismic Hazard Map (Giardini et al., 1999), an outcome of the Global Seismic Hazard
- 30 Assessment Program (GSHAP), is employed (see Fig. 3c).

A raster map representing the glaciers in the study area is generated by a semi-automated classification of Landsat 7 satellite imagery of 2001. Three classes are distinguished: debriscovered glacier, glacier with exposed ice and no glacier (see Fig. 3c). The lakes in the study area are covered by the comprehensive lake inventory presented by Mergili et al. (2013), providing detailed information on 1640 lakes (see Table 1; see Fig. 3c). Besides the tabular information, a raster map with the unique ID of each lake is used.

7 The exposure of the communities in the study areas to high-mountain hazards (see Fig. 3d) is 8 generated from a raster map depicting the land use associated with each cell derived by the qualitative interpretation of ASTER, Landsat and Google Earth[®] imagery. Table 2 shows the key 9 used for deriving the exposure score E from the land use n, taking values in the range 0–4. 10 11 Linear structures such as roads or power lines are not considered. Each raster cell with E > 0 is associated to one of the 628 communities identified in the study area. The communities largely 12 correspond to the villages depicted in the Soviet Topographic Maps 1:50,000 and 1:100,000. 13 14 However, two or more villages are grouped to one community in cases where cells with E > 015 cannot clearly be assigned to one specific village.

16 **4 Model**

17 4.1 Concept of the multi-hazard and risk analysis

18 The high-mountain multi-hazard and -risk indication computer model is implemented with the Open Source software package GRASS GIS (Neteler and Mitasova, 2007; GRASS Development 19 20 Team, 2013). This software builds on a flexible modular design. Simple bash scripting can be 21 used to facilitate work flows by combining existing modules. Furthermore, new modules can be added by individual developers, so that the standard GIS functions are complemented by a large 22 23 array of more specialized applications. Such applications can be used individually or made publicly available. Examples of mountain hazard models implemented with GRASS GIS include 24 25 r.debrisflow (Mergili et al., 2012b), r.avalanche (Mergili et al., 2012c) and r.rotstab (Mergili et al., 26 submitted). The model presented here builds on a combination of newly developed or upgraded 27 modules and bash scripts. The logical framework of the model is illustrated in Fig. 4, the modules 28 dealing with the specific process types are detailed in Section 4.2 to Section 4.5. The model is 29 executed at a raster cell size of 60 x 60 m.

30 The high-mountain hazard analysis procedure applied at the regional scale aims at the 31 identification of possible (i) source areas and (ii) impact areas of hazardous processes. The risk analysis combines the hazard in the impact areas, the impact hazard *IH*, with the exposure *E*there in order to derive a risk indication score *R* in the range 0–6. Table 2 shows the matrix
employed for the combination of *IH* and *E*.

The following types of processes are considered: (i) rock slides, (ii) ice avalanches, (iii) periglacial debris flows and (iv) lake outburst floods. All of them show a potential for long travel distances and therefore represent a significant threat for the populated areas in the valleys. Even though each process type is considered separately, interactions are included in the model, such as triggering of a lake outburst flood by the impact of an upslope mass movement (see Fig. 4).

9 The scoring scheme employed for the hazard analysis follows the same basic principle for all 10 types of processes. It builds on susceptibility, hazard and risk indication scores to be understood 11 as ordinal numbers, not allowing for the use of arithmetic operations. Two-dimensional matrices 12 are therefore used, all scores can take values in the range 0–6 (Table 3 and Table 4).

The hazard indication score for the onset of a process H is computed by combining the score for the susceptibility S with the score for the expected magnitude M. The susceptibility is understood as the tendency of a lake, part of a glacier or slope to produce an event and acts as a surrogate for the frequency. The expected process magnitude is based on the possible onset volume (rock slides) or on the possible onset area (ice avalanches, lake outburst floods; see Table 3).

18 The impact susceptibility represents the tendency of a GIS raster cell to be affected by one of the 19 considered processes. It is derived by routing the mass movement from the onset area down through the DEM. At the regional scale, empirical relationships are suitable for relating the travel 20 distance L or the angle of reach ω_r of a flow to the involved volume V or the peak discharge Q_m 21 or for defining a global value of ω_r . The appropriate values or relationships are employed for each 22 process type, applying the lower envelope (ω_{rE} , maximum travel distance) and the average ω_{rA} 23 usually observed for the considered process (see Section 4.2 - Section 4.5). A random walk 24 25 procedure weighted for local slope and maintenance of flow direction is applied for routing. This, by applying a sufficiently large number of random walks, ensures a certain degree of lateral 26 27 spreading. Furthermore, the linear distance from the starting point has to increase with each step 28 of the routing procedure. For each passed cell, the average slope angle from the starting point, ω , 29 is updated. Each random walk terminates as soon as $\omega \leq \omega_{rE}$. The impact susceptibility score I of each cell builds on the maximum of the ratio 30

31
$$r_{\omega} = 1 - \frac{\tan \omega_{r,A} - \tan \omega}{\tan \omega_{r,A} - \tan \omega_{r,E}}$$
 Eq. 1

- 1 over all random walks. $r_{\omega} = 1$ at the average angle of reach and $r_{\omega} = 0$ at the per envelope (see 2 Table 4). *I* is determined separately for each hypothetic event. The impact hazard indication score
- 3 IH map, discretized on the basis of GIS raster cells, is derived by combining H and I.
- 4 As the final step of the hazard analysis, the impact hazard indication scores IH_i for all hypothetic
- 5 events *i* are combined in order to derive a raster map of the global hazard indication score *IH*.
- 6 The maximum score is used for each raster cell:

7 $IH = max(IH_1, IH_2, \dots, IH_n),$ Eq. 2

- 8 where the indices 1, 2, ..., *n* represent the hypothetic event IH_i is associated with, *n* is the total 9 number of possible onset areas for the considered process type.
- 10 Whilst the general concept outlined is applied to all types of hazards, the specific procedures for
- 11 each process type are detailed in Section 4.2 Section 4.5. Below, the subscript rs stands for rock
- 12 slides, ia for ice avalanches, pf for periglacial debris flows and lo for lake outburst floods. Maps
- 13 of *IH* and *R* are determined separately for each process.
- Given the uncertainties inherent to the regional-scale hazard and risk analysis, the discretization 14 of the results at a raster cell size of 60 x 60 m may pretend a level of detail not supported by the 15 16 methodology used. According to the purpose of the study, the prioritization of target 17 communities for risk mitigation measure, community-based risk indication scores for each 18 process type (CR_{rs} , $CR_{i\omega}$, $CR_{b\beta}$, CR_{b}) are derived. The maxima of the raster cell-based risk indication scores over all cells representing the considered village are applied. However, if the 19 20 highest risk indication score R assigned to a community applies to an area <10,000 m², CR is reduced by 1. In such cases a lower score of R, if it applies to a larger area, may determine the 21 22 score of CR for the community.

23 4.2 Rock slide hazard

- The GRASS raster module employed for the rock slide hazard analysis is named r.rockslide and, to some extent, builds on the approach of Hergarten (2012). The logical framework of r.rockslide is illustrated in Fig. 5.
- Loops over all raster cells within the study area are performed separately for four assumptions of sliding plane inclination $\beta_{s,i}$ (Table 5). If the local slope $\beta > \beta_{s,i}$ for a tested cell, the cell is considered as seed cell for a possible rock slide. In order to simulate a progressive failure, an inverse cone with a vertical axis and an inclination of $\beta_{s,i}$ is introduced. The apex of this cone coincides with the seed cell (see Fig. 5). All material above the cone surface (terrain elevation >

1 cone elevation) is considered as potential rock slide material, imitating a rock slide involving all 2 over-steepened terrain with respect to the base cell. For each seed cell, the volume V_{rs} removed 3 by the associated rock slide is recorded.

4 The susceptibility score S_{r} for each cell with terrain elevation > cone elevation is determined 5 according to Table 5, including the sliding plane inclination $\beta_{s,i}$ and the permafrost susceptibility 6 S_p as conditioning factors, and the seismic susceptibility S_s as possible triggering factor. S_{rs} can take values in the range 0–6. The rock slide hazard indication score H_n is computed according to 7 8 Table 3, with the possible event magnitude represented by the rock slide volume V_n . Each cell 9 may possibly be affected by rock slides from more than one seed cell. The final hazard indication 10 score for each raster cell is defined as the maximum of $H_{n,i}$ out of all relevant possible rock slides i: 11

- 12 $H_{rs} = max(H_{rs,1}, H_{rs,2}, ..., H_{rs,n}), Eq. 3$
- where the indices 1, 2, ... *n* denote the id of the considered possible rock slide, *n* is the number ofpossible rock slides.
- 15 The expected travel distance is estimated separately for each single possible slide (see Fig. 5), 16 using a relationship of the type
- 17 $\log_{10} \tan \omega_r = a \log_{10} V_r + b$, Eq. 4
- 18 where ω_r is the angle of reach and V_r is the rock slide volume. The curve has to be cut off at 19 tan $\omega = \tan \varphi$, where φ is the angle of repose. Eq. 4 is only valid as long as the slide starts from 20 rest. *a* and *b* depend on the process type, *b* can also be varied in order to account for uncertainties 21 of the relationship used. Two relationships are applied:
- (1) For rock slides in non-glacierized areas, the prediction curve suggested by Scheidegger (1973)
 is used. It was derived from a set of 33 historic and prehistoric events. The correlation coefficient
- is 0.82, the standard deviation is 0.14298. a = -0.15666, b = 0.62419 for the average and 0.36418 for the envelope.
- (2) It is well established that rock slides in glacierized areas often convert into rock-ice avalanches
 with longer travel distances (Evans and Clague, 1988; Bottino et al., 2002). If the rock slide starts
- 28 in a glacierized area, or as soon as it moves over a glacier, the relationship suggested by
- 29 Noetzli et al. (2006) is applied: a = -0.103, b = 0.165 for the average and -0.040 for the envelope.
- 30 The steeper regression line for non-glacierized areas results in the prediction of longer travel 31 distances by the Scheidegger (1973) model for very large volumes ($V_r > 361 \, 10^6 \, \text{m}^3$ for the

1 regression, $V_r > 34 \, 10^6$ m³ for the envelope). This phenomenon has no physical basis but can 2 most likely be attributed to a lack of very large events in the data set used by Noetzli et al. (2006). 3 In the r.rockslide model, the relationship yielding the longer travel distance is used for rock slides 4 in glacierized areas. Further, the runup height R at the opposite slope is limited by the envelope 5 of the regression derived from the dataset presented by Hewitt et al. (2008):

6 $\log_{10} R \le 0.375 \cdot \log_{10} V_r - 0.62077$. Eq. 5

100 random walks are performed for each rock slide or rock-ice avalanche, each of them starting at the highest raster cell of the hypothetic failure plane. The impact susceptibility score I_{rs} and the impact hazard indication score IH_{rs} are finally derived according to Eq. 1, Eq. 2 and Table 4.

10 Eq. 1 is here applied with the logarithms of tan $\omega_{r,E}$ and tan $\omega_{r,A}$.

11 4.3 Ice avalanche hazard

12 The logical framework of the ice avalanche hazard model r.iceaval is illustrated in Fig. 6. The slope beyond which glaciers or portions of glaciers are susceptible to produce ice avalanches 13 depends on the properties of the ice which are strongly determined by the ice temperature. As 14 15 data on ice temperature is not commonly available, mean annual air temperature is often used as 16 a surrogate. Huggel et al. (2004a) state that temperate glaciers produce ice avalanches at slopes 17 above 25°, cold glaciers at slopes above 45°. Here, a set of 11 cases (Alean et al., 1985; Huggel et al., 2004a) is taken as the basis for devising a scheme for ice avalanche susceptibility S_{ia} 18 (Fig. 7). A quadratic regression is fitted for this purpose, with 19

20 $\tan \beta = 3.2 \cdot 10^{-3} MAAT^2 - 2.03 \cdot 10^{-2} MAAT + \eta$, Eq. 6

21 where β is the slope and *MAAT* is the mean annual air temperature (°C). The intercept 22 $\eta = 0.5555$ for the regression and 0.357672 for the envelope. The thresholds applied to the ice 23 avalanche susceptibility classes $S_{ia} = 0-4$ are determined from Eq. 1 with η set in the way to split 24 the data set into quartiles (see Fig. 7). S_{ia} is increased according to the seismic susceptibility (see 25 Table 1) so that the possible score values cover a range from 0–6.

- Next, clusters of cells with $S_{ia}>0$ are identified. S_{ia} is increased by 1 for all clusters at glacier termini (no abutment). The ice avalanche hazard indication score H_{ia} is derived according to Table 3, combining S_{ia} and the area of each cluster.
- For each cluster, 100 random walks are applied for routing the possible ice avalanche down, starting at the highest point of the cluster. According to Huggel et al. (2004a), the travel path of

ice avalanches is constrained by an average slope of 17° , except for very large events (>5 10^{6} m³). 1 2 However, such events are most commonly rock-ice avalanches or complex process chains (e.g., 1962 and 1970 Huascarán events, 2002 Kolka/Karmadon event), which are covered separately 3 4 here, or related to volcanic processes (1980 Iliamna event, Alaska). Therefore, and since the ice 5 avalanche volume cannot be derived with the method applied, we constrain the impact area with an average slope of 17° (tan $\omega_{RE} = 0.31$). In the dataset used by Huggel et al. (2004a), the 6 7 minimum value of the average slope is tan $\omega = 0.44$, tan ω_{rE} is set to the average 0.375 and I_{ia} is 8 computed according to Eq. 1. Eq. 2 and Table 4 are applied in order to derive the ice avalanche 9 impact hazard indication score IHia.

10 4.4 Periglacial debris flow hazard

Melting permafrost on steep slopes leaves behind a certain amount of loose debris susceptible to 11 12 mobilization as debris flows. Such processes may occur in the active layer, but even more where permafrost is retreating. Here, we only consider areas where retreating permafrost is assumed (see 13 14 Table 1; Mergili et al., 2012a). Fig. 8 shows the logical frame work of the periglacial debris flow model r.periflow. Huggel et al. (2004b) noted that, in contrast to ordinary debris flows, 15 parameters such as slope curvature or the proximity to the stream network are hardly significant 16 17 for the onset of such processes. Further, they commonly occur at slope angles from 27-38°. 18 Table 6 shows the scheme applied here for deriving the susceptibility of each raster cell to periglacial debris flows S_{bf} in the range from 0–6. We follow the findings of Huggel et al. (2004b) 19 20 with regard to slope. Unfortunately, no means for the reliable distinction of bedrock and debris at 21 the relevant scale are known to the authors. Besides slope and the state of the permafrost, the seismic susceptibility is considered for deriving S_{bf} (see Table 1 and Table 6). 22

In contrast to the other processes considered in the present study, there are no means to approximate the onset volume and therefore the process magnitude. Clusters of susceptible cells are often large whilst the onset of debris flow processes is most commonly a rather localized process. We therefore use the approximation $H_{bf} = S_{bf}$ (see Fig. 8).

Consequently, the routing procedure (i) has to be started separately from each raster cell with $S_{pf} > 0$, and (ii) the average slope determining the impact area has to be independent from volume. Due to the commonly large clusters of starting cells, only 10 random walks are started from each cell. Huggel et al. (2004b) give an envelope average slope of the travel path of 11° (tan $\omega_{r,E} = 0.194$) which is also applied here. The maximum average slope is taken from Corominas et al. (2003) who provide a value of 26° (tan $\omega = 0.488$) for debris flows <800 m³ 1 propagating on undisturbed flow paths assumed for the study area. The average of the two 2 values, 0.341, is taken as tan $\omega_{r,A}$. I_{bf} is computed according to Eq. 1. For $I_{bf} < 4$, the runup on the 3 opposite slope is restricted. Eq. 2 and Table 4 are applied to derive the periglacial debris flow 4 impact hazard indication score IH_{bf} .

5 4.5 Lake outburst hazard

6 An improved version of the GRASS GIS raster module r.glof (Mergili and Schneider, 2011) is 7 used for the lake outburst hazard analysis. The logical framework of r.glof is illustrated in Fig. 9.

8 First, the susceptibility scores for (i) lake outburst caused by internal factors (dam failure) $S_{h,i}$ and

9 (ii) lake outburst triggered by external factors (impact of mass movements) $S_{lo,e}$ are considered

10 separately (see Fig. 9). $S_{lo,i}$ and $S_{lo,e}$ can take values in the range 0–6, negative values are set to 0.

The derivation of $S_{b,i}$ builds on the following key pointers: (i) lake type, indicating the dam 11 material; (ii) mode of lake drainage; (iii) lake evolution; (iv) dam geometry; (v) permafrost 12 susceptibility; (vi) seismic susceptibility (see Table 1). Table 7 shows the scoring scheme applied. 13 14 The lake type (Mergili et al., 2013) is taken as basis, with glacial lakes receiving the highest score. 15 Dams with seepage are considered more susceptible to failure than dams with surface runoff, and 16 growing lakes are considered more susceptible than stable or shrinking ones. The dam geometry 17 is expressed as an idealized average downstream slope of the dam: the dam width W is defined as 18 the Euclidean distance between the lake outlet and the closest raster cell along the downstream flow path with a lower elevation than the average lake bottom, using the average lake depth D_l 19 20 according to Huggel et al. (2002):

21
$$D_l = 1.04 \cdot 10^{-1} A_l^{0.42}$$
, Eq. 7

where A_l is the lake area (m²), D_l is given in m. The tangent of the average slope of the dam in outflow direction, $\tan \beta_d$, is derived as D_l/W . For very gentle downstream average slopes $\tan \beta_d < 0.02$, S_{lad} is decreased by 1 (see Table 7).

The event at Laguna 513 in the Cordillera Blanca (Haeberli et al., 2010b) has shown the need to include the entire catchment when analyzing lake outburst susceptibility. The topographic susceptibility *TS* is introduced in order to account for this need, employing the impact hazard indication scores for rock slides IH_{rs} , for ice avalanches IH_{ia} , for periglacial debris flows IH_{ps} and for outburst floods of lakes in the upper catchment IH_{lo} . The overall maximum score over the raster cells representing the considered lake ($IH_{ia,max}$, $IH_{rs,max}$, $IH_{pf,max}$ and $IH_{lo,max}$) applies, but the impact of periglacial debris flows and upstream lake outburst floods is down-weighted:

1
$$TS = max(IH_{rs,max}, IH_{ia,max}, IH_{pf,max} - 3, IH_{lo,max} - 3).$$
 Eq. 8

The topographic susceptibility is taken as the basis for the rating of the susceptibility to lake outburst triggered by external factors $S_{lo,e}$. If direct calving of ice into the lake is possible, the score for $S_{lo,e}$ is set to a minimum of 3.

5 The maximum of $S_{lo,i}$ and $S_{lo,e}$ is used as lake outburst susceptibility S_{lo} . S_{lo} is reduced for lakes with 6 a high freeboard F (defined as the difference between the DEM with filled sinks and the original 7 DEM for the lake centre): for lakes with F > 50 m the score is decreased by 3. For lakes with 8 F > 25 m it is decreased by 2, and for lakes with F > 10 m, the score is decreased by 1 in order to 9 derive the final value of S_{lo} .

10 The lake area is most likely the best surrogate for M_{b} since the lake volume is uncertain. Table 7

11 shows the matrix for the lake outburst hazard indication score H_{lo} which is discretized on th

12 basis of lakes.

Possible outburst floods are routed downwards through the DEM separately for each lake, the travel distance is determined according to the relationships listed in Table 8. After the deposition of the debris or mud, or if not much sediment is entrained at all, the flood may propagate much farther: Haeberli (1983) suggests an average angle of reach of 2–3°, but also travel distances exceeding 200 km are reported (e.g., Hewitt, 1982).

In order to achieve a robust estimate of the travel distance, the impact area of possible lake outburst floods and, consequently, the impact susceptibility I_{lo} , the approaches T1 – T4 shown in Table 8 are combined (Eq. 1 is not applied for lake outburst floods). The lake outburst flood is routed down starting from the outlet of the considered lake. 800 random walks are performed for each lake. A random walk is forced to terminate if it impacts a larger lake.

23 For T1, the debris flow volume V_d is set to five times the outburst volume (maximum sediment concentration in steep flow channels ~80% according to Iverson, 1997) in order to account for 24 25 sediment bulking. The outburst volume is set to the entire lake volume, lake area A_l multiplied with lake depth D_{i} . For T2, we use an angle of reach $\omega_r = 8^{\circ}$ based on the observation for the 26 2002 Dasht event (Mergili and Schneider, 2011) instead of $\omega_r = 11^{\circ}$ suggested by Haeberli (1983). 27 28 Several authors have introduced empirical relationships for relating the peak discharge Q_{b} (m³/s) 29 - required as input for the relationship T3 in Table 8 - to the outburst volume and the dam height (Costa, 1985; Costa and Schuster, 1988; Walder and O'Connor, 1997; Table 9). Q, is 30

31 determined from the maximum of the results computed with the relationships Q1 to Q6 shown

32 in Table 9. T1 and T3 are only applied to glacial lakes as there is no basis available for calculating

- 1 the depth or volume of lakes assigned to the other types. Instead, the angle of reach is set to 2 $\omega_r = 11^\circ$ (the value suggested by Haeberli, 1983) for T1 and to $\omega_r = 14^\circ$ for T3.
- The number of relationships T1 T4 (see Table 8) predicting an impact on a given raster cell determines the impact susceptibility: if all four relationships predict an impact, $I_{lo} = 6$, three relationships results in $I_{lo} = 5$ and so forth. For $I_{lo} < 6$, the runup on the opposite slope is restricted. If only an impact as flood is predicted, (T4; $I_{lo} \le 3$), the impact susceptibility is further differentiated according to ω : for $\omega \ge 6$, $I_{lo} = 3$, for $\omega \ge 4$, $I_{lo} = 2$ and for $\omega \ge 2$, $I_{lo} = 1$. T4 is only applied to lakes $\ge 50,000$ m². Furthermore, the criterion that the distance from the source has to
- 9 increase with each computing step is disabled for floods.
- 10 In an analogous way to the scores for the rock slides, ice avalanches and periglacial debris flows
- 11 (see Section 4.2 to Section 4.4), the impact hazard indication score is then derived by combining
- 12 $I_{l_{a}}$ with $H_{l_{a}}$ of the corresponding lake (see Table 4). The global impact hazard indication score $IH_{l_{a}}$
- 13 of a given raster cell is defined as the maximum of all lake-specific scores (see Eq. 2).

14 5 Results

The total area with significant periglacial debris flow (PF) susceptibility/hazard is much larger 15 16 than those for the other hazard types: 9.9% of the entire study area are designated as possible PF 17 source areas, based on the criteria defined in Table 6. 42.7% out of this area are assigned the three higher susceptibility scores 4-6 (Fig. 10a). This pattern indicates the ubiquity of hazardous 18 areas on the one hand, but also the limited means of a sharper delineation on the other hand. In 19 contrast, the ice avalanche (IA) susceptibility and the lake outburst (LO) susceptibility, due to 20 21 their confinement to glaciers and lakes, respectively, are constrained in a much sharper way. 1.6% of the total study area are identified as susceptible to IA, 64.5% out of this area are assigned the 22 23 susceptibility scores 4-6 (see Fig. 10a). The LO susceptibility is discretized on the basis of lakes. 70.9% of all lakes are assigned susceptibility scores >0, 50.0% of these lakes the susceptibility 24 scores 4-6 (see Fig. 10a). The rock slide (RS) susceptibility displays intermediate patterns in terms 25 26 of the total area identified as susceptible (4.7% of the total study area). However, only 16.2% of this area - a much lower value than those associated with the other process types - are assigned 27 the susceptibility scores 4-6 (see Fig. 10a). The reason for this phenomenon is the limited area 28 29 occupied by very steep slopes (see Table 5).

The distribution of the raster cells or lakes identified as susceptible among the six hazard indication score classes is illustrated in Fig. 10b, depending on the susceptibility and the possible process magnitude (see Table 3). 38.1% of the RS and 68.4% of the IA are assigned the hazard indication scores 4–6. In the case of PF, hazard and susceptibility are identical due to lacking means for an estimation of the magnitude (see Section 4.4). Comparatively few lakes (23.9%) are assigned the LO hazard indication scores 4–6. This phenomenon is explained by the large number of rather small but highly susceptible lakes.

Fig. 11 represents the hazard indication scores for each process type broken down to the level of small catchments identical to the output parameter basin of the GRASS GIS raster module r.watershed (GRASS Development Team, 2013) with a threshold parameter of 5000. The maximum out of all raster cell-based hazard indication scores is shown for each catchment, except for the LO hazard where the value assigned to each lake is illustrated.

As expected, the RS hazard (see Fig. 11a) is highest in areas with a particularly steep topography in the northern and central Pamir. More localized high-hazard areas are distributed throughout the study area. The IA hazard (see Fig. 11b) is high in most glacierized areas (see Fig. 3d), particularly in parts of the northern Pamir where large portions of steep glaciers are extremely abundant. Within these zones the inter-catchment differentiation of hazardous areas is rather poor.

17 The PF hazard is poorly differentiated at the catchment scale: steep slopes near the permafrost boundary are almost ubiquitous in the study area (see Fig. 3b), except for the elevated and 18 comparatively gently inclined south eastern portion. The most notable regional pattern is 19 therefore attributed to the seismic susceptibility. The patterns observed in Fig. 11c are further a 20 consequence of the limited input information that can be reasonably applied at the regional scale 21 22 (see Table 6). A detailed inventory of rock glaciers (comparable to the inventories prepared for 23 glaciers and lakes, see Fig. 3d) could help to sharpen the distinction between more and less 24 hazardous areas. However, as rock glaciers are extremely common throughout the study area, the 25 patterns at the inter-catchment level would most likely remain unchanged.

As the LO hazard is directly related to the well-known lake distribution it can be discretized at a 26 27 high level of detail (see Fig. 11d). Nine lakes are assigned the highest LO hazard indication score $H_{l_0} = 6$, the largest of them is Lake Sarez. Even though – or because – the safety of Lake Sarez is 28 29 highly disputed (e.g., Risley et al., 2006), this classification seems reasonable. The LO 30 susceptibility score $S_{la} = 5$ is a consequence of the high topographic susceptibility. The same is 31 true for Lake Zardiv, with 0.7 km² the second largest lakes with $H_{l_0} = 6$ (see Fig. 11d). Further 32 lakes of interest are, e.g., Lake Khavraz and Lake Shiva. Lake Khavraz is an 1.9 km² lake impounded behind a rock glacier at an elevation of 4000 m a.s.l., in the zone of possibly melting 33

permafrost. Here both the topographic susceptibility ($S_{loc} = 5$) and the LO susceptibility due to 1 2 internal factors ($S_{lo,i} = 5$) are at high levels. Lake Shiva is assigned a susceptibility in the medium 3 range of the scale ($S_{la} = 3$) and a sudden drainage is not likely but, due to the large size of 4 15.2 km², H_{lo} = 5. The lake is located close to several communities in the Panj Valley which could 5 be affected in the case of such an event. The largest lake in the study area, Kara Kul, is assigned 6 a hazard indication score of 4 (see Fig. 11d). The LO susceptibility of Kara Kul is rated with a 7 score of $S_{l_0} = 2$, only the very large lake area (405 km²) leads to the relatively high hazard 8 indication score. The fact that a closer look reveals no significant outburst hazard of Kara Kul 9 suggests that the approach used tends to overestimate the hazard for large lakes. One reason for this phenomenon is the topographic susceptibility: large lakes have the ability to alleviate the 10 11 impact of mass movements rather than small lakes. However, an objective basis to include the 12 dependence of the topographic susceptibility on lake size is missing. Table 10 summarizes the LO susceptibility and hazard by lake type. Whilst - as prescribed by the scheme shown in Table 7 13 14 - glacial lakes clearly display the higher scores of the LO susceptibility due to internal factors, this tendency is less pronounced - but still visible - for the LO susceptibility due to external factors. 15

16 The median and maximum travel distances computed for each process type are summarized in Table 11. The RS model commonly predicts travel distances of 3.0 km (average) - 5.6 km 17 (envelope), but for very large events (>800 10^6 m³) extending over vertical distances >4000 m the 18 model, when applied with the envelope, predicts a travel distance of almost 50 km (see Eq. 4). 19 20 The IA model ($\omega_{rE} = 17^{\circ}$) and the PF model ($\omega_{rE} = 11^{\circ}$) predict shorter travel distances. Note that, in both cases, the difference between the median and the maximum is only caused by 21 22 topography and not by the assumed process magnitude. The LO model predicts the possibility of 23 a significant debris flow for less than half of all lakes (no meaningful median value can therefore 24 be given in Table 11). The reason for this phenomenon is mainly the gentle slope observed 25 downstream from many lakes. However, lakes with steeper downstream slopes can produce debris flows with travel distances >15 km and floods >80 km, according to the model. 26

Fig. 12 shows the distribution of the impact hazard. For clarity, only the raster cell values along the main flow channels are shown. It is clear that the general patterns of the impact hazard at the broad scale resemble those of the hazard shown in Fig. 11: whilst a possible impact of rock slides and periglacial debris flows is shown for most valleys particularly in the western part of the study area (Fig. 12a and Fig. 12c), a more localized impact of ice avalanches and lake outburst floods is

32 suggested by the model (Fig. 12b and Fig. 12d).

The distribution of community risk over the study area reflects the patterns shown in Fig. 11 and 1 2 Fig. 12 on the one hand, and the distribution of the exposed communities on the other hand. Fig. 13 illustrates the relative frequency of the community risk indication score classes for 15 3 4 regions within the study area, each of them representing a catchment or section of a catchment. The eastern Pamir is considered as one single region due to the low number of communities 5 there. Except for the very western part of the study area, the eastern Pamir and the Kyrgyz part 6 7 of the study area (Chan-Alai Valley) in the north, all regions are dominated by communities with a significant rock slide risk, the highest scores are observed for the villages in the rugged Bartang 8 9 and middle Panj valleys as well as in the Gunt Valley (see Fig. 13a). The hot spots of ice 10 avalanche risk are identified in the Vanch and Bartang valleys, both deeply incised into glacierized 11 mountain ranges (see Fig. 13b). This type of risk plays a less prominent role in the other regions. 12 Also the risk of periglacial debris flows is highest in the deep gorges of the western Pamir, decreasing towards north where permafrost is less abundant (see Fig. 13c). However, the model 13 14 predicts a significant PF risk for most communities throughout the study area. This is not the case for the risk caused by lake outburst floods, which is significant mainly in the south western 15 Pamir and in part of the northern Pamir (see Fig. 13d). The LO community risk indication score 16 17 $CR_{in} = 6$ is not assigned to any village. Table 12 summarizes the relative frequency of villages 18 assigned to each class with respect to all four hazard types.

19 A composite hazard and risk indication map is prepared for the entire study area. It provides a 20 visual overlay of the hazard, impact hazard and community-based risk indication scores for each of the four process types considered in the study. Fig. 14 shows this map for a selected area 21 covering the Gunt Valley and its tributaries (see Fig. 1 for delineation). The area affected by the 22 prehistoric Charthem rock slide (see Fig. 14a) is very well reproduced by the model, therefore a 23 high RS risk is assigned to the nearby communities. However, also several other communities and 24 25 lakes are possibly impacted by rock slides. The patterns of IA hazard and risk illustrate the isolated appearance of this type of hazard in the area (see Fig. 14b). Even though the main effect 26 27 of the process is the possible impact on lakes, some communities in the main valley are at risk, but assigned rather low scores. Areas of PF hazard (see Fig. 14c) are strictly confined to steep 28 29 slopes near the permafrost boundary which are however very common along the slopes of most 30 valleys, confirming the broad-scale patterns shown in Fig. 11c. Therefore, most of the communities in the valleys are identified as at risk. The occurrence of periglacial debris flows in 31 this area, often starting from the termini of rock glaciers, is evident in the field and from remotely 32 sensed imagery (see Fig. 2c). The associated debris cones are located in zones of high impact 33

hazard identified by the model. In contrast, most lakes are located where permafrost is assumedstable.

Several lakes have developed near the termini of the glaciers of the tributary valleys (Mergili and 3 Schneider, 2011; Mergili et al., 2013). Three of them are assigned the hazard indication score 4 5 $H_{la} = 5$ (see Fig. 14d). The debris flow travel distances predicted by the LO model are relatively short, only debris flows from two lakes could reach the communities of the main valley: the 6 7 village of Varshedz is just located at the terminus of a possible debris flow starting from Lake 8 Varshedz (see Fig. 2d; lake area 0.16 km², $S_{lo} = 5$, $H_{lo} = 5$). Lake Nimats, an erosion lake with 9 S_{lb} = 4 and H_{lb} = 5, drains into a very steep channel heading directly down to the main valley. In 10 the case of a (not very likely) sudden drainage, the nearby villages would most likely suffer 11 substantial damage. The impact area of the distal floods resulting from the possible drainage of Lake Varshedz or Lake Nimats is characterized by lower to medium community risk indication 12 scores of the possibly affected villages. The largest lake shown in Fig. 14d is Rivakkul (1.2 km², 13 14 $S_{la} = 4$, $H_{la} = 5$). It is characterized by a very gently inclined downstream valley. As a result, the model predicts only a comparatively short travel distance of a possible outburst flood (4.6 km), 15 16 being alleviated far upslope from the villages in the main valley.

17 6 Discussion

The purpose of the model approach introduced in the previous sections and the resulting hazard and risk indication maps is to provide a reproducible basis for targeted hazard and risk assessment studies and mitigation measures at the community scale. The approach chosen is thought to be useful for the study area in the Pamir for two reasons.

First, the general difficulty of establishing frequencies for rare or singular events in combination with sparse historical data in the study area makes strictly quantitative approaches such as statistical methods inapplicable. Therefore a hazard and risk indication scoring scheme has to be applied, even though such a concept introduces a certain degree of subjectivity.

Second, the vulnerability of the local population to these types of hazards is high, even though NGOs have launched programs to improve the awareness of and the preparedness for geohazard events in the previous decade. This situation is comparable to other high-mountain areas in developing countries (e.g., Carey, 2005). The results of the present study shall highlight high-risk areas and serve as a baseline for in-detail studies and risk mitigation procedures. Consequently, the outcome of the study should not be seen as definite hazard and risk maps, but rather as conceptual hazard and risk indication maps. The hazard and risk indication score classes are therefore not given definite names such as Moderate hazard, Extremely high risk etc. Further, the interpretation of the model results on the basis of raster cells is appropriate for scientific discussion, but not for the design of risk mitigation measures. Here the scale of communities (see Fig. 14), catchments (see Fig. 12) or even regions (see Fig. 13) are much more suitable.

7 As far as a comparison with observed events is possible, it confirms the model results (e.g., 8 Charthem rock slide, see Fig. 14a). In the case of large rock slides such as the 1911 Sarez event 9 (see Fig. 2a) the comparison with the model results is of limited value due to the substantial change of the topography caused by such events. No records of ice avalance in the study area 10 11 are known to the authors whilst periglacial debris flows are very common. Their source and 12 impact areas are well recognized by the model, but the false positive rate is high. The two lakes with recorded sudden drainage are not characterized by exceptionally high susceptibility scores -13 14 the prediction of lake outburst floods is therefore particularly challenging.

15 The quality of the model results strongly depends on the input data used. The detail and accuracy of the ASTER GDEM is considered sufficient for the purpose of the present study, even though 16 17 the quality of the dam geometry estimates may suffer from artefacts and inaccuracies known for 18 this type of DEM. Also the quality of the carefully mapped lake, glacier and land use data sets is 19 largely considered sufficient. The potential permafrost areas were determined using a rule-ofthumb approach, adapting data obtained in the Alps (Haeberli, 1977) Mergili et al., 2012a). Even 20 21 though the predicted conditions and scenarios are likely to be realistic, uncertainties are hard to quantify. The seismic hazard map used (Giardini et al., 1999) is a highly generalized global 22 dataset. Other essential information such as the distinction bedrock - residual rock or the 23 24 orientation and dip of the bedding planes of geological layers are hardly manageable at the scale 25 relevant for a study of this type.

The scoring schemes used (see Table 2 to Table 7) are founded on expert knowledge. The interpretation of the model results have to consider the characteristics of the scheme used for each process. Necessarily, the schemes contain ±arbitrary thresholds such as those used for the event magnitude (see Table 3) or the 45° minimum slope for rock slides already used earlier by Hergarten (2012).

The modelling of the travel distance and the impact area of the considered processes is derived from the statistics of observed events. These statistics are reasonably robust for rock slides and rock ice avalanches (Scheidegger, 1973; Evans and Clague, 1988; Bottino et al., 2002;

Noetzli et al., 2006) and also for ice avalanches (Huggel et al., 2004a). However, they are based 1 2 on observations from other mountain areas such as the Alps. Their application is based on the hypothesis that (i) the patterns observed there are comparable to those in the Pamir and (ii) that 3 the - often rather small - datasets used for the derivation of the patterns and thresholds cover a 4 5 representative sample of the reality. This is equally true for the slope-temperature curve shown in 6 Fig. 7. The situation is even more difficult for periglacial debris flows (Huggel et al., 2004a) and 7 particularly for lake outburst floods. The threshold of $\omega_{r,E} = 11^{\circ}$ used by Huggel et al. (2004a, b) for debris flows from lake outburst events is not applicable to the Pamir as the 2002 Dasht event, 8 9 where $\omega_{rE} \sim 8.0^{\circ}$, has shown (Mergili and Schneider, 2011). Also the parameterization of floods 10 developing from lake outburst events is nothing more than a rough estimate so that the results 11 (such as the short travel distance predicted for a sudden drainage of Rivakkul, see Fig. 14d) have 12 to be interpreted with utmost care.

Further, the application of average slopes neglects the loss of energy due to changes of the flow direction. Strictly spoken, such concepts should only be used for straight flow paths. The criterion that the motion has to move away from the source with each step of the random walk partly accounts for this limitation.

17 Possible impact wave due to mass movements into lakes are explicitly accounted for by the 18 model. Other types of interactions are included indirectly: The conversion of rock slides into 19 rock-ice avalanches by the impact on glaciers is implicitly considered in the rock slide model, 20 even though there are no means to estimate the entrainment of snow or ice. In the case of rock 21 slides and rock-ice avalanches, the empirical relationships used implicitly include cascading effects such as the conversion into debris flows. Some process interactions are out of scope of the 22 present study, such as the damming of lakes by mass movements and possible subsequent 23 drainage. The same is true for the entrainment of debris, modelling of which remains a challenge 24 particularly at the scale of the present study. 25

The approach used does not allow for an analytical overlay of the susceptibility, hazard, impact hazard and risk indication scores associated to each process type. Even though attempted as far as possible, a homogenization of the scoring schemes for the different processes proves highly problematic due to the missing physical basis. The data the analysis is based on differs between the processes: e.g., the possible magnitude of rock slides is given in maximum volumes whilst only the maximum involved surface area allowed under the assumptions taken is used for possible ice avalanches and lake outburst floods (see Table 3). Also the schemes for susceptibility 1 can hardly be homogenized (see Table 5 to Table 7; Fig. 7), partly due to the varying level of

2 detail of the available input data.

3 7 Conclusions

A regional-scale multi-hazard and -risk indication model was introduced, including four selected 4 5 high-mountain processes: (i) rock slides and rock avalanches, (ii) ice-avalanches, (iii) periglacial debris flows and (iv) lake outburst floods. The model results for a very large area centred in the 6 7 Pamir (Tajikistan) were presented and discussed. The model shall help to distinguish areas with 8 higher from those with lower risk, even though the possibilities for comparison with observed 9 events are limited. The interpretation of the model results - preferably at the level of 10 communities, catchments or regions - has to take into account the characteristics of the scoring 11 schemes as well as the limitations of the input data and the methodology used.

12 Acknowledgements

The work presented is part of the project PAMIR supported by the European Commission (EC) and the Austrian Development Agency (ADA), as well as of the project TajHaz supported by FOCUS Humanitarian Assistance (an affiliate of the Aga Khan Development Network), the Swiss Agency for Development and Cooperation (SDC) and the UK Department for International Development (DFID). The Tajik Agency of Hydrometeorology has provided meteorological data. Special thanks for their support go to Matthias Benedikt, Johannes P. Müller and Jean F. Schneider, BOKU University, Vienna.

20 References

- Alean, J.: Ice avalanches: some empirical information about their formation and reach. J. Glac.,
 31(109), 324–333, 1985.
- Beniston, M.: Climatic Change in Mountain Regions: A Review of Possible Impacts. Clim.
 Change, 59(1–2), 5–31, 2003.
- Bolch, T., Peters, J., Yegorov, A., Prafhan, B., Buchroithner, M. and Blagoveshchensky, V.:
 Identification of potentially dangerous glacial lakes in the northern Tien Shan. Nat. Haz., 59,
- 27 1691–1714, 2011.

- 1 Bottino, G., Chiarle, M., Joly, A.and Mortara, G.: Modelling Rock Avalanches and Their Relation
- 2 To Permafrost Degradation in Glacial Environments. Permafr. Periglac. Proc., 13, 283-288,
- 3 2002.
- 4 Breien, H., DeBlasio, F. V., Elverhoi, A. and Hoeg, K.: Erosion and morphology of a debris
- 5 flow caused by a glacial lake outburst flood, Western Norway. Landslides, 5(3), 271–280, 2008.
- 6 Carey, M.: Living and dying with glaciers: people's historical vulnerability to avalanches and
- 7 outburst floods in Peru. Glob. Plan. Change, 47, 122–134, 2005.
- 8 Cenderelli, D. A. and Wohl, E. E.: Peak discharge estimates of glacial lake outburst floods and
- 9 "normal" climatic floods in Mount Everest region, Nepal. Geomorphology, 40, 57–90, 2001.
- 10 Clarke, G. K. C.: Glacier outburst floods from 'Hazard Lake', Yukon Territory, and the problem
- 11 of flood magnitude prediction. J. Glac, 28(98), 3–21, 1982.
- 12 Corominas, J., Copons, R., Vilaplana, J. M., Altamir, J. and Amigó, J.: Integrated Landslide
- 13 Susceptibility Analysis and Hazard Assessment in the Principality of Andorra. Nat. Haz., 30, 421–
- 14 435, 2003.
- 15 Costa, J. E.: Floods from Dam Failures. U.S. Geol. Surv. Open File Rep. 85-560, Denver,
 16 Colorado, 54 pp., 1985.
- 17 Costa, J. E. and Schuster, R.L.: The formation and failure of natural dams. Geol. Soc. Am. Bull.,
 18 100, 1054–1068, 1988.
- 19 Dussaillant A., Benito G., Buytaert W., Carling P., Meier C. and Espinoza F.: Repeated glacial-
- lake outburst floods in Patagonia: an increasing hazard? Nat. Haz., DOI 10.1007/s11069-0099479-8, 2009.
- 22 Evans, S. G.: The maximum discharge of outburst floods caused by the breaching of man-made
- 23 and natural dams. Can. Geotech.J., 23, 385–387, 1986.
- Evans S. G. and Clague, J. J.: Catastrophic rock avalanchesin glacial environments. In:
 Bonnard, C. (ed.): Landslides, A.A. Balkema, Rotterdam, 2, 1153–1158, 1988.
- Evans S. G. and Clague J. J.: Recent climatic change and catastrophic geomorphic processes in
 mountain environments. Geomorphology, 10(1-4), 107–128, 1994.
- 28 Evans, S. G., Bishop, N. F., Fidel Smoll, L., Valderrama Murillo, P., Delaney, K. B., Oliver-
- 29 Smith, A: A re-examination of the mechanism and human impact of catastrophic mass flows

- originating on Nevado Huascarán, Cordillera Blanca, Peru in 1962 and 1970. Engin. Geol., 108,
 96–118, 2009a.
- 3 Evans, S. G., Roberts, N. J., Ischuk, A., Delaney, K. B., Morozova, G. S., Tutubalina, O.:
- 4 Landslides triggered by the 1949 Khait earthquake, Tajikistan, and associated loss of life. Engin.
- 5 Geol., 109, 195–212, 2009b.
- 6 Giardini, D., Grünthal, G., Shedlock, K. and Zhang, P.: The GSHAP Global Seismic Hazard
- 7 Map. Ann. Geophys., 42(6), DOI 10.4401/ag-3784, 1999.
- 8 GRASS Development Team: Geographic Resources Analysis Support System (GRASS)
- 9 Software. Open Source Geospatial Foundation Project, http://grass.osgeo.org, 2013.
- 10 Haeberli, W.:Untersuchungen zur Verbreitung von Permafrost zwischen Flüelapass und Piz
- 11 Grialetsch (Graubünden). Dissertation, University of Basel, 1975.
- Haeberli, W.: Frequency and characteristics of glacier floods in the Swiss Alps. Ann. Glac., 4, 85–
 90, 1983.
- 14 Haeberli, W., Clague, J. J., Huggel, C. and Kääb, A.: Hazards from lakes in high-mountain glacier
- and permafrost regions: Climate change effects and process interactions. Avances de la
 Geomorphología en España, 2008-2010, XI Reunión Nacional de Geomorphología, Solsona,
- 17 439–446, 2010a.
- 18 Haeberli, W., Portocarrero, C. and Evans, S.: Nevado Hualcán, Laguna 513 y Carhuaz 2010 -
- 19 Observaciones, evaluación y recomendaciones (un corto informe técnico luego de las reuniones y
- 20 visita de campo en Julio 2010). Unpublished report on behalf of the Communidad Provincial de
- 21 Carhuaz, 2010b.
- 22 Haritashya, U. K., Bishop, M. P., Shroder, J. F., Bush, A. B. G., Bulley, H. N. N.: Space-based
- assessment of glacier fluctuations in the Wakhan Pamir, Afghanistan. Clim. Change, 94, 5–18,
 2009.
- 25 Harris, C., Arenson, L. U., Christiansen, H. H., Etzelmüller, B., Frauenfelder, R., Gruber, S.,
- 26 Haeberli, W., Hauck, C., Hölzle, M., Humlum, O., Isaksen, K., Kääb, A., Kern-Lütschg, M. A.,
- 27 Lehning, M., Matsuoka, N., Murton, J. B., Nötzli, J., Phillips, M., Ross, N., Seppälä, M.,
- 28 Springman, S. M., VonderMühll, D.: Permafrost and climate in Europe: Monitoring and
- 29 modelling thermal, geomorphological and geotechnical responses. Earth-Sci. Rev., 92(3–4), 117–
- 30 171, 2009.

- 1 Harrison, S., Glasser, N., Winchester, V., Haresign, E., Warren, C., Jansson, K.: A glacial lake
- 2 outburst flood associated with recent mountain glacier retreat, Patagonian Andes. Holocene, 16,
- 3 611–620, 2006.
- 4 Hergarten, S.: Topography-based modelling of large rockfalls and application to hazard
 5 assessment. Geophys. Res. Lett., 39, L13402, DOI 10.1029/2012GL052090, 2012.
- 6 Hewitt, K.: Natural dams and outburst floods in the Karakorum Himalaya. In: Glen J. W. (ed.):
- 7 Hydrological aspects of alpine and high-mountain areas. IAHS Publication 138, 259–269, 1982.
- 8 Hewitt, K., Clague, J.J. and Orwin, J.F.: Legacies of catastrophic rock slope failures in mountain
 9 landscapes. Earth-Sci. Rev., 87, 1–38, 2008.
- 10 Hewitt, K. and Liu, J.: Ice-dammed lakes and outburst floods, Karakoram Himalaya: historical
- 11 perspectives on emerging threats. Phys. Geog., 31(6), 528–551, 2010.
- 12 Huber, U., Bugmann, H. and Reasoner, M.: Global Change and Mountain Regions. An overview
- 13 of current knowledge. Advances in Global Change Research. Springer, Dordrecht, 2005.
- 14 Huggel, C.: Assessment of Glacial Hazards based on Remote Sensing and GIS Modeling.
- 15 Dissertation at the University of Zurich, Schriftenreihe Physische Geographie Glaziologie und
- 16 Geomorphodynamik, 2004.
- 17 Huggel, C., Kääb, A., Haeberli, W., Teysseire, P. and Paul, F.: Remote sensing based assessment
- of hazards from glacier lake outbursts: a case study in the Swiss Alps, Can. Geotech. J., 39, 316–
 330, 2002.
- 20 Huggel, C., Kääb, A., Haeberli, W. and Krummenacher, B.: Regional-scale GIS-models for
- 21 assessment of hazards from glacier lake outbursts: evaluation and application in the Swiss Alps.
- 22 Nat. Haz. Earth Syst. Sci., 3, 647–662, 2003.
- Huggel, C., Haeberli, W., Kääb, A., Bieri, D. and Richardson, S.: Assessment procedures for
 glacial hazards in the Swiss Alps. Can. Geotech.J., 41(6), 1068–1083, 2004a.
- Huggel, C., Kääb, A. and Salzmann, N.: GIS-based modeling of glacial hazards and their
 interactions using Landsat-TM and IKONOS imagery. Norwegian J. Geog., 58, 761–773, 2004b.
- 27 Huggel, C., Zgraggen-Oswald, S., Haeberli, W., Kääb, A., Polkvoj, A., Galushkin, I. and Evans,
- 28 S.G.: The 2002 rock/ice avalanche at Kolka/Karmadon, Russian Caucasus: assessment of
- 29 extraordinary avalanche formation and mobility, and application of QuickBird satellite imagery.
- 30 Nat. Haz. Earth Syst. Sci., 5, 173–187, 2005.

- 1 ICIMOD: Glacial lakes and glacial lake outburst floods in Nepal. ICIMOD, Kathmandu, 2011.
- 2 IPCC: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to
- 3 the Fourth Assessment Report of the Intergovernmental Panel on Climate Change [Solomon, S.,
- 4 Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H.L. (eds.)].
- 5 Cambridge University Press, Cambridge and New York, 2007.
- 6 Iverson, R. M.: The physics of debris flows. Rev. Geophys., 35, 245–296, 1997.
- 7 Jarvis, A., Reuter, H. I., Nelson, A. and Guevara, E.: Hole-filled seamless SRTM data V4.
- 8 International Centre for Tropical Agriculture (CIAT), available from http://srtm.csi.cgiar.org,
- 9 2008.
- 10 Kääb, A., Huggel, C., Fischer, L., Guex, S., Paul, F., Roer, I., Salzmann, N., Schlaefli, S., Schmutz,
- 11 K., Schneider, D., Strozzi, T. and Weidmann, Y.: Remote sensing of glacier- and permafrost-
- 12 related hazards in high mountains: an overview. Nat. Haz. Earth Syst. Sci., 5, 527–554, 2005.
- 13 Kassam, K. A.: Viewing Change Through the Prism of Indigenous Human Ecology: Findings
- 14 from the Afghan and Tajik Pamir. Hum. Ecol., 37, 677–690, 2009.
- 15 Khromova, T. E., Osipova, G. B., Tsvetkov, D. G., Dyurgerov, M. B. and Barry, R. G.: Changes
- 16 in glacier extent in the eastern Pamir, Central Asia, determined from historical data and ASTER
- 17 imagery. Rem. Sens. Environ., 102, 24–32, 2006.
- 18 Mahmood, S. A., Shahzad, F. and Gloaguen, R.: Remote Sensing Analysis of Recent Active
- 19 Tectonics in Pamir Using Digital Elevation Model: River Profile Approach. Proceedings of the
- 20 Geoscience and Remote Sensing Symposium IGARSS, 7 11 July 2008, II-1259–II-1262, 2008.
- 21 Makhmadaliev, B., Kayumov, A., Novikov, V., Mustaeva, N. and Rajabov, I. (eds.): The Second
- 22 National Communication of the Republic of Tajikistan under the United Nations Framework
- 23 Convention on Climate Change. State Agency for Hydrometeorology, Dushanbe, 2008.
- Mergili, M. and Schneider, J. F.: Regional-scale analysis of lake outburst hazard in the
 southwestern Pamir, Tajikistan, based on remote sensing and GIS. Nat. Haz. Earth Syst. Sci., 11,
 1447–1462, 2011.
- 27 Mergili, M., Kopf, C., Müllebner, B. and Schneider, J. F.: Changes of the cryosphere in the high-
- 28 mountain areas of Tajikistan and Austria: a comparison. Geog.Ann., A, 94(1), 79–96, 2012a.

- 1 Mergili, M., Fellin, W., Moreiras, S. M. and Stötter, J.: Simulation of debris flows in the Central
- 2 Andes based on Open Source GIS: Possibilities, limitations, and parameter sensitivity. Nat. Haz.,
- **3** 61(3), 1051–1081, 2012b.
- Mergili, M., Schratz, K., Ostermann, A. and Fellin, W.: Physically-based modelling of granular
 flows with Open Source GIS. Nat. Haz. Earth Syst. Sci., 12, 187–200, 2012c.
- 6 Mergili, M., Müller, J. P. and Schneider, J. F.: Spatio-temporal development of high-mountain
- 7 lakes in the headwaters of the Amu Darya river (Central Asia). Glob. Plan. Change, DOI
- 8 10.1016/j.gloplacha.2013.04.001, 2013.
- 9 Müllebner, B.: Modelling of potential permafrost areas in the Pamir and Alai mountains
- 10 (Tajikistan) using Remote Sensing and GIS techniques. Master thesis, BOKU University, Vienna,2010.
- 12 Narama, C., Duishonakunov, M., Kääb, A., Daiyrov M. and Abdrakhmatov, K.: The 24 July
- 13 2008 outburst flood at the western Zyndan glacier lake and recent regional changes in glacier
- 14 lakes of the Teskey Ala-Too range, Tien Shan, Kyrgyzstan. Nat. Haz. Earth Syst. Sci., 10, 647–
- 15 659, 2010.
- 16 Neteler, M. and Mitasova, H.: Open source GIS: a GRASS GIS approach. Springer, New York,17 2007.
- Noetzli, J., Huggel, C., Hoelzle, M. and Haeberli, W.: GIS-based modelling of rock-ice avalanches
 from Alpine permafrost areas. Computational Geosci., 10, 161–178, 2006.
- 20 Quincey D. J., Richardson S. D., Luckman A., Lucas R. M., Reynolds J. M., Hambrey M. J. and
- 21 Glasser N. F.: Early recognition of glacial lake hazards in the Himalaya using remote sensing
- 22 datasets. Glob. Plan. Change, 56(1-2), 137–152, 2007.
- 23 Risley, J. C., Walder, J. S. and Denlinger, R.: Usoi dam wave overtopping and flood routing in the
- 24 Bartang and Panj rivers, Tajikistan. Nat. Haz., 38(3), 375–390, 2006.
- 25 Richardson, S. D. and Reynolds, J. M.: An overview of glacial hazards in the Himalayas. Quatern.
- 26 Int., 65/66, 31–47, 2000.
- 27 Rickenmann, D.: Empirical Relationships for Debris Flows. Nat. Haz., 19, 47–77, 1999.
- 28 Scheidegger, A. E.: On the Prediction of the Reach and Velocity of Catastrophic Landslides.
- 29 Rock Mech., 5, 231–236, 1973.

- 1 Schuster, R. L. and Alford, D.: Usoi Landslide Dam and Lake Sarez, Pamir Mountains, Tajikistan.
- 2 Environ. & Engin. Geosci., 10(2), 151–168, 2004.
- 3 Tinti, S., Maramai, A. and Cerutti, A. V.: The Miage Glacier in the Valley of Aosta (Western Alps,
- 4 Italy) and the extraordinary detachment which occurred on August 9, 1996. Phys. Chem. Earth
- 5 (A), 24(2), 157–161, 1999.
- 6 Tweed, F. S. and Russell, A. J.: Controls on the formation and sudden drainage of glacier7 impounded lakes: implications for jökulhlaup characteristics. Prog. Phys. Geog., 23(1), 79–110,
 8 1999.
- 9 Vilímek, V., Zapata, M. L., Klimes, J., Patzelt, Z. and Santillán N.: Influence of glacial retreat on
- 10 natural hazards of the Palcacocha Lake area, Peru. Landslides, 2(2), 107–115, 2005.
- 11 Walder J. S. and Costa J. E.: Outburst floods from glacierdammed lakes: the effect of mode of
- 12 lake drainage on flood magnitude. Earth Surf. Proc. Landforms, 21, 701–723, 1996.
- 13 Walder, J. S. and O'Connor, J. E.: Methods for Predicting Peak Discharge of Floods Caused by
- 14 Failure of Natural and Constructed Earth Dams. Water Resour. Res., 33, 10, 2337–2348, 1997.
- Watanabe, T. and Rothacher, D.: The 1994 Lugge Tsho glacial lake outburst flood, Bhutan
 Himalaya. Mt. Res. Dev., 16, 77–81, 1996.
- 17 WGMS: Global Glacier Changes: Facts and Figures (Zemp, M., Roer, I., Kääb, A., Hoelzle, M.,
- 18 Paul, F., Haeberli, W., eds.), UNEP, World Glacier Monitoring Service, University of Zurich,
- 19 Switzerland, free download from WGMS and UNEP websites, 2008.

1 Tables

2 Table 1 Input data, rm = raster map, tc = table column.

Parameter	Data type	Source
Elevation	rm, m a.s.l.	ASTER GDEM V2, a product of METI and NASA
Glaciers	rm, boolean	Semi-automated classification of Landsat 7 imagery
Lake ID	rm, nominal	Manual mapping from ASTER and Landsat imagery (Mergili et al., 2013)
For each lake:		
Lake type	tc, nominal	Qualitative interpretation of ASTER, Landsat and
Lake drainage	tc, boolean	Google Earth [®] imagery (Mergili et al., 2013)
Calving of ice	tc, boolean	
Lake area A_l	t c, m ²	Derived from mapped lakes
Lake evolution	tc, boolean	Mapped, 75% confidence of growing trend in at least one of the periods 1968–2002 and 2002–2009 (Mergili et al., 2013)
Mean Annual Air Temperature MAAT	rm, °C	Temperature map of Müllebner (2010) based on regression of data recorded by the Tajik HydroMet Agency with elevation
Permafrost susceptibility S_p	rm, nominal	Permafrost indication map of Mergili et al. (2012a) for Tajikistan, based on the adaptation of the rules- of-thumb of Haeberli (1975)
Seismic susceptibility S_s	rm, g	GSHAP Global Seismic Hazard Map (Giardini et al., 1999): peak ground acceleration PGA with 10% chance of exceedance in 50 years
Exposure E	rm, nominal	Manual mapping of land use from ASTER, Landsat
Community ID	rm, nominal	and Google Earth [®] imagery

- 1 Table 2 Risk indication score R: combination of IH and E with scoring scheme for the exposure
- E as a function of land use.

$E\downarrow IH \rightarrow$	Land use	6	5	4	3	2	1	0
4 (Higher)	Built-up areas, often mixed with farmland or pastures	6	5	4	3	2	1	0
3	Farmland or pastures with some buildings	5	4	3	2	1	1	0
2	Farmland, pastures or forest with no or few buildings	4	3	2	1	1	1	0
1	Extensively used or temporarily unused land	3	2	1	1	1	1	0
0 (Lower)	No identifiable land use	0	0	0	0	0	0	0

1 Table 3 Hazard indication score H: combination of S and M, with thresholds of (a) rock slide

$M \downarrow S \rightarrow$	(a)	(b)	(c)	6	5	4	3	2	1	0
6 (Higher)	>24.3	>200.0	>200.0	6	6	5	5	4	3	0
5	8.1 - <24.3	100.0 - <200.0	100.0 - <200.0	6	5	5	4	3	2	0
4	2.7 - <8.1	50.0 - <100.0	50.0 - <100.0	5	5	4	4	3	2	0
3	0.9 - <2.7	25.0 - <50.0	25.0 - <50.0	5	4	4	3	3	2	0
2	0.3 – <0.9	12.5 - <25.0	12.5 - <25.0	4	3	3	3	2	2	0
1	0.1 - < 0.3	5.0 - <12.5	5.0 - <12.5	3	2	2	2	2	1	0
0 (Lower)	<0.1	<5.0	<5.0	0	0	0	0	0	0	0

2 volume V_{rs} (10⁶ m³), (b) area of hanging glacier A_a (10³ m²) and (c) lake area A_a (10³ m²).

$I \downarrow H \rightarrow$	r_{ω}	6	5	4	3	2	1	0
6 (Higher)	≥2.000	6	6	5	5	4	3	0
5	1.500-<2.000	6	5	5	4	3	2	0
4	1.000 - <1.500	5	5	4	4	3	2	0
3	0.667 - <1.000	5	4	4	3	3	2	0
2	0.333 -< 0.667	4	3	3	3	2	2	0
1	≥0.000 - <0.333	3	2	2	2	2	1	0
0 (Lower)	< 0.000	0	0	0	0	0	0	0

1 Table 4 Impact hazard indication score *IH*: combination of *H* and *I*, r_{ω} is computed according to 2 Eq. 1.

1 Table 5 Rock slide susceptibility score S_n . The initial values of S_n are determined from the sliding

2 plane inclination $\beta_{s,b}$ these values are then increased according to permafrost susceptibility and 3 seismic susceptibility, *g* is gravity (m s⁻²).

Criterion	Remarks	S_{rs}
Sliding plane inclination	$\tan(\beta_{s,i}) \ge 1.000$	1
$\beta_{s,i}$	$\tan\left(\beta_{s,i}\right) \geq 1.333$	2
	$\tan\left(\beta_{s,i}\right) \geq 1.667$	3
	$\tan\left(\beta_{s,i}\right) \geq 2.000$	4
Permafrost susceptibility	No permafrost or stable permafrost	± 0
	Susceptible to melting at Δ MAAT >0 – 4°	+1
Seismic susceptibility	PGA < 0.34 g	± 0
	$PGA \ge 0.34 - <0.65 g$	+1
	$PGA \ge 0.65 g$	+2

Criterion	Remarks	$S_{lo,i}$
Slope β	$\tan\beta < 0.5$	0
	$\tan \beta = 0.5 - <0.6$	1
	$\tan \beta = 0.6 - <0.7$	2
	$\tan \beta = 0.7 - 0.8$	3
	$\tan \beta > 0.8$	0
Permafrost	No permafrost or stable permafrost	0
susceptibility	Susceptible to melting at Δ MAAT >2 – 4°	±0
	Susceptible to melting at Δ MAAT >0 – 2°	+2
Seismic	PGA< 0.65 g	±0
susceptibility	$PGA \ge 0.65 g$	+1

1 Table 6 Scoring scheme for periglacial debris flow susceptibili	ty S _{pf} .
---	----------------------

1 Table 7 Scoring scheme for susceptibility to lake outburst triggered by internal factors S_{lo} . The 2 initial values of S_{lo} are determined from the dam material, these values are then increased or 3 decreased according to lake drainage, lake evolution, downstream slope of dam, permafrost 4 susceptibility and seismic susceptibility.

Criterion	Remarks	$S_{lo,i}$			
Lake type (dam	Erosion lake	0			
material)	Block- or debris-dammed lake	1			
	Glacial lake	3			
Lake drainage	Permanent or temporary superficial drainage	-1			
	No recognizable superficial drainage	±0			
Lake evolution	Stable or shrinking	±0			
	Growing trend in either the period 1968–2002 or 2002–2009	+1			
Downstream slope	$\tan\beta_d < 0.02$	-1			
of $dam\beta_d$	$\tan \beta_d \ge 0.02$	±0			
Permafrost	No permafrost or stable permafrost	±0			
susceptibility	Susceptible to melting at Δ MAAT >0 – 4°				
Seismic	PGA< 0.65 g	±0			
susceptibility	$PGA \ge 0.65 g$				

- 1 Table 8 Empirical relationships used for estimating the travel distance of lake outburst floods.
- 2 GLOF = glacial lake outburst flood, L = travel distance, V_d = debris flow volume, ΔZ = loss of
- 3 elevation, ω_r = average slope of reach, Q_p = peak discharge.

	Relationship	References	Remarks
T1	$L = 1.9 V_d^{0.16} \Delta Z^{0.83}$	Rickenmann (1999)	for debris flows in general
T2	$\omega_r = 11^{\circ}$	Haeberli (1983), Huggel et al. (2003), Huggel et al. (2004a)	for debris flows from GLOFs, applied with $\omega_r = 8^\circ$ in the present study (Mergili and Schneider, 2011)
Т3	$\omega_r = 18Q_p^{-0.07}$	Huggel (2004)	worst case for debris flows from GLOFs
Τ4	<i>ω</i> _{<i>r</i>} ≥2°	Haeberli (1983), Huggel et al. (2004a)	for floods from GLOFs

- 1 Table 9 Empirical regression equations relating peak discharge Q_p of glacial lakes to outburst
- 2 volume V_1 and lake depth (dam height) $D_r \rho_w =$ density of water (kg/m²), g = gravity (m/s²),
- 3 *envelope (worst case).

Reference		Q_p (m ³ /s) of glacial lakes
Costa (1985)	Q1	$113(10^{-6}V_l)^{0.61}$
	Q2	$3.8 (10^{-6} V_l D_l)^{0.61}$
Costa and	Q3	$1.3 \cdot 10^{-4} \big(\rho_{\scriptscriptstyle W} \cdot g \cdot V_{\scriptscriptstyle l} \cdot D_{\scriptscriptstyle l} \big)^{0.60}$
Schuster (1988)	Q4	$5.5 \cdot 10^{-6} \left(ho_W \cdot g \cdot V_l \cdot D_l ight)^{0.59}$
Walder and	and Q5*	$2.2 \cdot 10^{-1} V_l^{0.66}$
O'Connor (1997)	Q6*	$1.1(V_{l}D_{l})^{0.47}$

1 Table 10 Percentage of lakes assigned each class of lake outburst (LO) susceptibility (internal and

_
2
-

Lake type	LO sus	O susceptibility score (internal factors)							
	0	1	2	3	4	5	6	Sum	
Erosion lakes	33.1%	40.8%	24.7%	1.5%	0.0%	0.0%	0.0%	883	
Block- or debris-dammed									
lakes	12.4%	17.1%	44.8%	24.8%	1.0%	0.0%	0.0%	105	
Glacial lakes	38.2%	0.0%	0.0%	4.4%	31.1%	24.5%	1.7%	652	
	LO sus	ceptibilit	y score (external	factors)				
	0	1	2	3	4	5	6	Sum	
Erosion lakes	42.1%	5.3%	18.9%	16.5%	7.5%	7.7%	1.9%	883	
Block- or debris-dammed									
lakes	18.1%	1.0%	10.5%	14.3%	18.1%	32.4%	5.7%	105	
Glacial lakes	40.8%	0.2%	3.5%	11.3%	15.5%	21.3%	7.4%	652	
	LO haz	ard indic	cation sco	ore					
	0	1	2	3	4	5	6	Sum	
Erosion lakes	24.5%	5.8%	34.2%	19.0%	10.6%	5.3%	0.6%	883	
Block- or debris-dammed									
lakes	12.4%	0.0%	31.4%	23.8%	13.3%	16.2%	2.9%	105	
Glacial lakes	38.2%	0.0%	26.5%	20.2%	10.4%	4.4%	0.2%	652	

2	external	factors)	and	hazard	indica	ation	scores	accordi	ing to	lake ty	pe.
---	----------	----------	-----	--------	--------	-------	--------	---------	--------	---------	-----

1 Table 11 Maximum and median travel distances L (m) computed for each process. RS = rock

2 slides, IA = ice avalanches, PF = periglacial debris flows, LO = lake outburst floods,

	RS (A)	RS (E)	IA (E)	PF (E)	LO (dfl)	LO (fld)
Maximum	23,867	49,516	10,735	15,906	15,597	81,946
Median	2,964	5,628	2,013	2,618	_	12,589

3 A = average, E = envelope, dfl = debris flow, fld = flood.

1 Table 12 Per cent of communities assigned to each class of the community risk indication score.

Process type	Community risk indication score							
	0	1	2	3	4	5	6	
RS	12.1%	9.7%	8.6%	15.3%	27.5%	23.9%	2.9%	
IA	75.2%	8.9%	5.7%	4.8%	2.2%	2.4%	0.8%	
PF	11.5%	3.7%	6.1%	9.2%	24.4%	31.4%	13.9%	
LO	65.6%	12.3%	11.6%	6.2%	2.9%	1.4%	0.0%	

2 RS = rock slides, IA = ice avalanches, PF = periglacial debris flows, LO = lake outburst floods.





3 Fig. 1 Study area. The dashed red rectangle delimits the area shown in Fig. 14.



1

Fig. 2 Processes covered by the high-mountain multi-hazard and -risk indication model. (a) The km² rock slide deposit impounding lake Sarez, triggered by an earthquake in 2011, (b) hanging glacier in the Sauksay Valley prone to produce ice avalanches, (c) periglacial debris flow starting from a rock glacier terminus in the upper Gunt Valley, (d) Lake Varshedz in a southern tributary of the Gunt Valley, one of many glacial lakes possibly susceptible to sudden drainage. All photos taken by M. Mergili.



- 3 distribution, (c) peak ground acceleration, (d) mapped glaciers, lakes and land use.



Fig. 4 Logical framework of the high-mountain multi-hazard and -risk indication model.



2 Fig. 5 Logical framework of the rock slide model (r.rockslide).



2 Fig. 6 Logical framework of the ice avalanche model (r.iceaval).



Fig. 7 Scoring scheme applied for ice avalanche susceptibility according to data presented by
Alean (1985) and Huggel et al. (2004a).



- 2 Fig. 8 Logical framework of the periglacial debris flow model (r.periflow).



2 Fig. 9 Logical framework of the lake outburst model (r.glof).



Fig. 10 Relative abundance of (a) the susceptibility and (b) the hazard indication score values for the four considered processes. The values for rock slides (RS), ice avalanches (IA) and periglacial debris flows (PF) relate to raster cells, the values for lake outburst floods (LO) to lakes. Only those raster cells or lakes with a score of at least 1 are considered.



Fig. 11 Distribution of (a) rock slide, (b) ice avalanche, (c) periglacial debris flow and (d) lake
outburst hazard over the entire study area. The maximum score is shown for each catchment.



Fig. 12 Impact hazard for each process, (a) rock slides, (b) ice avalanches, (c) periglacial debris
flows, (d) lake outburst floods. For clarity, only the scores along the main flow lines are shown.



Fig. 13 Community risk, generalized to 15 regions. For each region, the pie chart illustrates the
relative abundance of the different community risk indication scores. The size of each chart is
proportional to the number of communities it represents.



1

Fig. 14 Hazard, impact hazard, topographic susceptibility of lakes and community risk associated with each process type. (a) Rock slide, (b) ice avalanche, (c) periglacial debris flow, (d) lake outburst flood. The extent of the map is shown in Fig. 1.