



This discussion paper is/has been under review for the journal Natural Hazards and Earth System Sciences (NHESS). Please refer to the corresponding final paper in NHESS if available.

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar^{1,2}, J. Brauch^{2,3}, A. Dobler⁴, K. Béranger⁵, and B. Ahrens^{1,2}

¹Institute for Atmospheric and Environmental Sciences, Goethe University Frankfurt am Main, Germany

²Biodiversity and Climate Research Center, Frankfurt am Main, Germany

³German Weather Service, Offenbach am Main, Germany

⁴Institute of Meteorology, Freie University Berlin, Germany

⁵Unité de Mécanique, ENSTA-ParisTech, Palaiseau, France

Received: 12 February 2014 – Accepted: 22 February 2014 – Published: 19 March 2014

Correspondence to: N. Akhtar (naveed.akhtar@iau.uni-frankfurt.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

So-called medicanes (Mediterranean hurricanes) are meso-scale, marine, and warm-core Mediterranean cyclones that exhibit some similarities to tropical cyclones. The strong cyclonic winds associated with medicanes threaten the highly populated coastal areas around the Mediterranean basin. To reduce the risk of casualties and overall negative impacts, it is important to improve the understanding of medicanes with the use of numerical models. In this study, we employ an atmospheric limited-area model (COSMO-CLM) coupled with a one-dimensional ocean model (1-D NEMO-MED12) to simulate medicanes. The aim of this study is to assess the robustness of the coupled model in simulating these extreme events. For this purpose, 11 historical medicane events are simulated using the atmosphere-only model, COSMO-CLM, and coupled model, with different setups (horizontal atmospheric grid-spacings of 0.44° , 0.22° , and 0.08° ; with/without spectral nudging, and an ocean grid-spacing of $1/12^\circ$). The results show that at high-resolution, the coupled model is able to not only simulate most of medicane events but also improve the track length, core temperature, and wind speed of simulated medicanes compared to the atmosphere-only simulations. The results suggest that the coupled model is more proficient for systemic and detailed studies of historical medicane events, and that this model can be an effective tool for future projections.

1 Introduction

The Mediterranean Sea is known to be one of the main cyclogenetic regions in the world (Pettersen, 1956; Hoskins and Hodges, 2002; Wernli and Schwierz, 2006). A certain type of cyclone in the Mediterranean Sea with physical and structural similarities to tropical cyclones is known as a medicane (Mediterranean hurricane). Medicanes are meso-scale cyclones (the diameter is usually less than 300 km), with a rounded structure and a cloudless area at the center. Other features include a warm-core and

NHESSD

2, 2117–2149, 2014

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

intense low sea level pressure, combined with strong cyclonic winds and heavy rainfall (Businger and Reed, 1989). In general, the intensity of medicanes is much weaker than tropical hurricanes (Moscatello et al., 2008); however, a few medicanes have reached tropical hurricane strengths (33 m s^{-1}). Strong surface heat fluxes and deep convection are important initial conditions in the formation of medicanes. The triggering mechanisms involved in this development are the presence of cold anomalies in the high troposphere, surface heat fluxes (latent and sensible), low wind shear, and high low-level vorticity (Cavicchia et al., 2013). The minimum value of sea surface temperature (SST) to develop a tropical hurricane is 26°C (Trenberth, 2005). However, studies show that the SST has to be higher than 15°C for medicanes (Tous and Romero, 2013). As with tropical hurricanes, the air–sea temperature difference plays an important role in the development of medicanes (Palmen, 1948; Miglietta et al., 2011). In the case of tropical hurricanes, SST creates the conditions of thermodynamical disequilibrium that trigger the development of a vortex, whereas in the case of medicanes, the interplay between the temperature at the surface and upper atmospheric layer plays a key role in their development (Cavicchia et al., 2013). The ocean also plays an important role in the intensity of tropical cyclones (Emanuel, 1986; Emanuel and Rotunno, 1987); similarly, the ocean feedback is also crucial in the development of medicanes. Table 1 shows a list of medicanes events from 1983 to 2003, their approximate times of mature phase, sizes and geographical positions (Tous and Romero, 2013). Spatially, the central and western parts of the Mediterranean Sea are the main genesis regions of medicanes. Most of the events occurred during autumn and winter, but some were observed during the spring season.

Medicanes have a high potential for destruction in the densely populated coastal areas around the Mediterranean Sea. To reduce the risk of casualties and overall negative impacts, better knowledge of medicanes is essential for potential prediction. Due to their meso-scale and marine characteristics, medicanes are not well represented in automatic detection methods. The lack of dense observations over sea, and the occasional occurrence of medicanes make it more difficult to recognize meteorological

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

features associated with them (Tous and Romero, 2013). Surface observations of such storms are limited to ships crossing nearby; in-situ weather reports are generally poor. One of the main sources of observations of medicanes with full coverage are satellite images available since 1980. Several studies have been carried out based on the observational evidence (e.g., Ernst and Matson, 1983; Rasmussen and Zick, 1987; Luque et al., 2007; Moscatello et al., 2008). Most of the contemporary modeling studies on medicanes are done by dynamical downscaling using regional atmosphere-only models (e.g., Homar et al., 2003; Fita et al., 2013; Miglietta et al., 2011; Cavicchia et al., 2013).

Coarse global climate models cannot fully resolve the complex orographic and other important local processes such as the Bora, Mistral, and Etesian winds, and the deep-water formations that characterize the Mediterranean region. Therefore, the air-sea fluxes over this region are not correctly represented in a coarse global climate model (Elguindi et al., 2009). Recent studies show that high-resolution coupled models over the Euro-Mediterranean region significantly improve the representation of air-sea fluxes (Gualdiet al., 2012; Dubois et al., 2012; Artale et al., 2010; Somot et al., 2008). In another study, Sanna et al. (2013) have shown that SST simulated through a high-resolution eddy-permitting ocean model have strong and beneficial effects on precipitation and cyclogenesis.

The SSTs in regional atmosphere-only runs are prescribed and derived from re-analysis data such as the European Center for Medium-Range Weather Forecasts (ECMWF) Reanalysis Interim (ERA-Interim), which also include satellite observations. The quality of SSTs in this reanalysis is in good agreement with the observations (Simmons et al., 2006). However, due to the small size and short lifetimes of medicanes, the quality of the reanalysis data is compromised by the coarse time and space resolution. Most reanalysis datasets are available in 6h intervals, whereas medicanes occur in short time scales (ranging from 12h to 90h). Fine scale feedback associated with air-sea interactions can influence the temporal and spatial structure of medicanes. A fully

NHESSD

2, 2117–2149, 2014

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

with simplified dynamics. In contrast, 3-D ocean models are computationally expensive, time consuming, but include full 3-D dynamics. The 1-D models are very useful for short time simulations, especially in coupled ocean–atmosphere modeling, to investigate extreme events. The validation of a 1-D compared to a fully 3-D configuration has been conducted in a previous study (Akhtar, 2013). The 1-D NEMO-MED12 is now coupled via the OASIS3 coupler (Valcke, 2013) to CCLM. The 1-D NEMO-MED12 is employed with a basin-wide (covering the whole Mediterranean Sea and part of the Atlantic Ocean), single column approach where each water column is isolated from the others. Each grid point in the 1-D NEMO-MED12 is treated as a single water column. The coupling process is executed every hour, where 1-D NEMO-MED12 passes SST to CCLM and, in exchange heat, momentum, and freshwater fluxes are obtained.

The different resolutions of CCLM, 0.44° (~ 50 km, 118×83 grid points, and 32 σ -levels), 0.22° (~ 25 km, 206×120 grid points, and 32 σ -levels), and 0.08° (~ 9 km, 536×295 grid points, and 40 σ -levels) that cover the Med-CORDEX domain (Ruti et al., 2012) are employed. For both coupled and atmosphere-only simulations, the horizontal diffusion parameters in CCLM are tuned for better realization of medicanes. The 1-D NEMO-MED12 has a resolution of $1/12^\circ$ (~ 9 km, 567×264 grid points, and 50 vertical levels), which remains the same for all the coupled runs. The atmospheric model uses ERA-Interim reanalysis data from the ECMWF for both the coupled and atmosphere-only runs. However, in the coupled runs, SST over the Mediterranean Sea is calculated by 1-D NEMO-MED12 and elsewhere prescribed and derived from the reanalysis data. The monthly mean seasonal climatology from MEDATLAS-II (Rixen, 2012) is used to initialize 1-D NEMO-MED12.

The coupled and atmosphere-only models are used to simulate the historically listed medicane events from 1983 to 1999 (Table 1), at three different resolutions (0.44° , 0.22° , and 0.08°). The same set of simulations are also performed with the spectral nudging technique (von Storch et al., 2000). In a study, Cavicchia and von Storch (2012) showed that the performance of CCLM to simulate the medicanes could be improved by applying the spectral nudging technique, particularly the spatial and tem-

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



poral locations of the simulated medicanes. The purpose of employing spectral nudging is to analyze its impact on the coupled simulations, as compared to the atmosphere-only simulations, and not to address the time and location of simulated medicanes. The main aim of this study is to examine the ability of the coupled model to simulate the medicanes compared to the atmosphere-only simulations with and without spectral nudging, and to validate it with observations.

For the remainder of this discussion, we used the abbreviations “CPLXXYY” for the coupled, and “CCLMXXYY” for the atmosphere-only simulations, where “XX” refers to the resolution (“44” for 0.44°, “22” for 0.22°, and “08” for 0.08°), and “YY” refers to spectral nudging (“sn”).

Based on the satellite (Infra-red Meteosat) imagery analysis, 12 cases are documented from 1983 to 2003 (Table 1). The selection criteria of these events were based on cloud structure, size, and lifetime of the cyclones (Tous and Romero, 2013). Because of the limited availability of the MEDATLAS-II climatology (1945–2002), only the first 11 medicane events from 1980 to 1999 are included in this study. Since the last event occurred in 2003 (Table 1), MEDATLAS-II climatology is not available to initialize the ocean model. For each medicane event, both the coupled and atmosphere-only simulations with and without spectral nudging are conducted over a period of one month, starting approximately three weeks prior to formation of the medicane.

The following datasets are used for validation:

- for wind speed, the NOAA “SeaWinds” dataset, available from 9 July 1987 to the present are used. The product contains global high-resolution ocean winds and wind stresses at 6h intervals on a 0.25° grid (Zhang et al., 2006). SeaWinds are generated by blending observations from multiple satellites.
- For sea level pressure and temperature, NASA’s MERRA reanalysis (Rienecker et al., 2011), presently available from 1 January 1979 to 30 September 2013 is used. The state-of-the-art MERRA reanalysis products are available globally for 6h intervals at 0.5° resolution, and 42 vertical levels.

3 Results and discussion

In this section, we discuss the ability of the coupled and atmosphere-only model to simulate the medicanes at different resolutions with and without applying the spectral nudging technique. The meteorological variables, sea level pressure, temperature at mid-troposphere (in our case, 700hPa level), and 10 m wind speeds, are analyzed. To simulate a medicane, one needs to find intense sea level pressure minima, a warm-core at mid-troposphere, and strong cyclonic winds (Tous et al., 2013). The medicane tracks are constructed by following the sea level pressure minima in hourly model outputs in a grid box with less than 40% of land friction. Due to the coarse temporal resolution of the MERRA reanalysis data (6h), it is not possible to track the full trajectories of medicanes. Therefore, only approximated tracks are shown here for MERRA reanalysis. Although the spatial resolution of the MERRA reanalysis data is coarse (0.5°), medicane features such as mean sea level pressure and warm-core structure are reasonably well represented due to the assimilation of observations such as satellite data. For the medicane's starting and ending times, we took the information available from satellite images on the website <http://www.uib.es/depart/dfs/meteorologia/METEOROLOGIA/MEDICANES>.

We selected 4 different cases from 11 simulated medicane events for detailed validation and discussion. Our choices are motivated by the locations and sizes of these particular medicane events. We chose two cases (ME08 and ME09) from the central region of the Mediterranean Sea, and two (ME06 and ME10) from the western region. Two cases, ME06 from the central region and ME08 from the west, had maximum lifetimes, and two cases, ME10 from the central region and ME09 from the west, had intermediate lifetimes (Table 1). Due to unavailability of the NOAA dataset before 9 July 1987, only cases that occurred after 1988 were selected.

NHESSD

2, 2117–2149, 2014

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.1 Case ME08

The ME08 case was recorded to have a lifetime of 90h, making it one of the longest listed medicane events (Table 1). The medicane started to develop north of Algeria and reached maximum strength while it was moving between the Balearic Islands and Sardinia. Strong winds and severe damage was reported in the Aeolian Islands. The medicane crossed Calabria and dissipated after making landfall in the coastal regions of Greece. The snapshots of coupled simulations at the three resolutions mentioned above, on 6 October at 03:00 UTC (30 min before the development of ME08), show a large-scale baroclinic disturbance that evolved over the western part of the Mediterranean Sea (Fig. 1). This intrusion of cold air into the upper and mid-troposphere can favor the development of medicanes. The synoptic scale analysis shows that medicanes are not fully isolated structures of atmospheric circulation (Emanuel, 2005; Homar et al., 2003; Pytharoulis et al., 2000). Two dynamical processes are involved in the development of medicanes: the first phase is dominated by the baroclinic development, and the second is driven by convective tropical-like activity and air–sea interactions. The large-scale disturbance is well represented in CPL44, CPL22, and CLP08 simulations and similar results are also observed in CCLM44, CCLM22, and CCLM08 (not shown here). According to the satellite observations, on the same day at 03:30 UTC, the disturbance turned into a medicane that reached its mature phase at 18:00 UTC (Table 1). Figure 2 shows the mean sea level pressure and temperature at mid-troposphere (700hPa pressure level) on 7 October, at 18:00 UTC, along with the medicane track (black dots). In this case the medicane eye developed twice (7 October at 06:30 UTC and 8 October at 12:00 UTC). In CPL44 and CCLM44 simulations, the criteria for medicanes are not met. However, the medicane track is reasonably well represented in CCLM22 and CPL22, and even finer and more intense in CPL08 and CCLM08 (Fig. 2). The length of the simulated medicane track in CPL22 is shorter (74 h) compared to CCLM22 (90 h). The lowest minimum sea level pressure values, lifetimes, and start and end dates of all the simulated medicanes are shown in Table 2.

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The track and warm-core structure of the medicane are better resolved and represented in 0.08° simulations, and are in good agreement with the MERRA reanalysis data (Fig. 2). This is due to the meso-scale characteristic of medicanes, which are not fully resolved at low resolutions (0.44° and 0.22°). The sea level pressure minimum appeared one hour earlier in CPL08 compared to the CCLM08 simulations, and the warm-core is more prominent in the CPL08 simulations. The medicane track length in CPL08 is 94h, 92h in CCLM08 and 72h in MERRA reanalysis (Table 2). The medicane track in CPL08 is improved compared to CCLM08, and according to the MERRA reanalysis, with a more accurate position near the Sicily channel.

Figure 3 shows the wind speed patterns in all the simulations (0.44° , 0.22° and 0.08°) of the ME08 event and the NOAA dataset on 7 October at 18:00 UTC. The high wind speeds associated with the medicane are not observed in the CPL44 and CCLM44 simulations. Compared to this, the wind speeds in the CPL22 and CCLM22 simulations are slightly improved, but still do not match the high values of the NOAA dataset. There is further improvement at 0.08° resolution; however, the maximum wind speed still does not match the maximum of 34ms^{-1} seen in the NOAA dataset. The wind speed patterns in the coupled simulations (CPL08) are more intense compared to the atmosphere-only (CCLM08) simulations, and the structures are more delicate. According to Cavicchia and von Storch (2012), the high wind speeds recorded in the NOAA dataset may be due to Mistral winds, which are not well resolved in simulations.

Most of the medicane features are well resolved at the higher resolution of 0.08° (Fig. 2). The correct times and locations of simulated medicane formations are not in good agreement with the satellite images and MERRA reanalysis data. This may be due to the small size of medicanes, boundary layer parameterizations, the role of moist microphysics, and deep convection (Tous et al., 2013).

The times and locations of simulated medicanes can be improved by using the spectral nudging technique. Thus, the same event is simulated again, this time with the spectral nudging technique applied to CCLM.

**Medicanes in an
ocean–atmosphere
coupled regional
climate model**

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Figure 4 shows the results of the CPL08sn and CCLM08sn simulations on 7 October at 18:00 UTC, along with the medicane track (black dots). The lowest minimum sea level pressure value during a fully developed medicane period is 995.94 hPa in CLP08sn, and 995.34 hPa in CCLM08sn. The lowest sea level pressure values with and without spectral nudging simulations (Table 2) are not very different. The first sea level pressure minimum appeared 3–4 h earlier in simulations with spectral nudging compared to those without. Initially, the location of the medicane in spectral nudging simulations is shifted more to the south, which compares better with the satellite observations and MERRA reanalysis data. Although the timing and location of the medicane is improved by applying spectral nudging as expected (Cavicchia and von Storch, 2012), compared to the satellite observations, the length of the track is reduced in this particular case. The length of the medicane track in CPL08sn (68 h) is longer than in CCLM08sn (51 h). In most cases, the length of a medicane's track in spectral nudging simulations is approximately the same as in cases without spectral nudging (Table 2). Figure 5 shows the wind speed patterns in CPL08sn and CCLM08sn simulations on 7 October at 18:00 UTC. The wind speed did not show any improvement in simulations with spectral nudging compared to simulations without spectral nudging (see Fig. 3).

The first 11 medicane events listed in Table 1 are also simulated by applying the spectral nudging technique to CCLM in both the coupled and atmosphere-only setups. In most cases, simulations with spectral nudging improved the precision in the time and location of simulated medicanes. A similar conclusion was reached by Cavicchia and von Storch (2012) for atmosphere-only, CCLM simulations. The results show that applying spectral nudging did not significantly improve the wind speed estimates in both the coupled and atmosphere-only simulations. In all spectral nudging simulations, the coupled model performance is improved to the same extent as the atmosphere-only model, showing more intense and finer structure of medicanes in coupled simulations. As mentioned before in this study, our focus is not to address the precision in times and locations of simulated medicanes. We are more interested in examining the ability and

added value of the coupled model to simulate medicanes. In subsequent examples, only the results of simulations without spectral nudging are described.

3.2 Case ME09

The ME09 case was a medicane event that was observed in the western part of the Mediterranean Sea. According to the satellite observations, the medicane started to develop on 8 December 1996 at 12:00 UTC and lasted for 48h (Table 1). The medicane started to develop northwest of Sardinia, and moved towards the Balearic Islands. After reaching the Balearic Islands, it started to move again towards Sardinia and dissipated after crossing it. Figure 6 shows the mean sea level pressure and temperature at 700hPa in CPL08, CCLM08, and the MERRA reanalysis data on 12 December at 18:00 UTC, along with the medicane track (black dots). The length of the medicane track in CPL08 is 54h, 50h in CCLM08, and 60h in the MERRA reanalysis data. Although the lowest minimum sea level pressure values are the same in both CCLM08 and CPL08, the length of the medicane track is longer in the latter (Table 2). Figure 7 shows the wind speed patterns in CPL08, CCLM08, and the NOAA dataset on 12 December at 18:00 UTC. The wind speed patterns in CPL08 are more intense and in good agreement with the NOAA dataset. Compared to the atmosphere-only simulations, the medicane tracks, warm-core structures, and wind speeds are improved in the coupled simulations.

3.3 Case ME06

The ME06 event is a well-known medicane that developed in the central Mediterranean Sea with striking similarities to a tropical cyclone. The satellite observations show that the medicane started to develop on 14 January 1995 at 12:00 UTC and ended on 18 January at 20:00 UTC. The eye of the medicane was clearly visible for the majority of the time (15 January at 07:00 UTC to 18 January at 06:30 UTC). It was one of the longest medicane events to ever occur in the central Mediterranean Sea (Table 1).

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



According to the satellite observations, a large-scale low pressure system moved from the central Mediterranean Sea and reached the coast of Greece in the late night of 14 January. It then started to move towards the coast of Libya. Strong winds, heavy rainfall, and positive temperature anomalies were reported during that time by ships cruising nearby (Cavicchia and von Storch, 2012). The storm dissipated after making landfall in the Gulf of Sirte on 18 January. Figure 8 shows the comparison of mean sea level and temperature at 700hPa on 16 January at 12:00 UTC in CPL08, CCLM08, and the MERRA reanalysis data along with the medicane track (black dots). The track does not appear in CPL08, CCLM08, and the MERRA reanalysis data during the first part of the medicane, when it was moving from south to north. This is due to the sea level pressure being less deep and steep in CLP08, CCLM08, and the MERRA reanalysis data. The length of the medicane track in CPL08 is 42h, 34h in CCLM08, and 48h in the MERRA reanalysis. The lowest minimum sea level pressure values and warm-core structures are similar in both the CPL08 and CCLM08 simulations (Table 2). However, the track length in CPL08 is longer in comparison to the CCLM08 simulations. The medicane tracks in CPL08 and CCLM08 are shifted toward the east compared to the MERRA reanalysis data. Compared to the atmosphere-only simulations, the medicane track and warm-core structure in the coupled simulations (CPL08) are in good agreement with the MERRA reanalysis data. Figure 9 shows the wind speed comparison in CPL08, CCLM08, and the NOAA dataset on 16 January 1995 at 12:00 UTC. The wind speed in CPL08 is more intense and in good agreement with the NOAA dataset, but the position of the maximum wind speed is shifted to the east in the simulations.

3.4 Case ME10

The ME10 medicane event occurred in the central part of the Mediterranean Sea. The satellite observations show a mature phase of the medicane on 26 January 1998 at 12:00 UTC (Table 1). The medicane started to develop in the southeast of the Sicily channel and dissipated after making landfall in the northeastern coast of Libya. The observed lifetime of this medicane was 30h (Table 1). Figure 10 shows the comparison of

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



the mean sea level and temperature at 700 hPa on 27 January at 00:00 UTC in CPL08, CCLM08, and the MERRA reanalysis data, along with the medicane track (black dots). The length of the simulated medicane track in CPL08 is 27h, 28h in CCLM08, and 36h in the MERRA reanalysis data. The first sea level pressure minimum appeared 1 h earlier in the CCLM08 simulations compared to the CPL08 simulations. The mean sea level pressure is slightly deeper in CPL08 compared to CCLM08 (Table 2) and the warm-core values are approximately the same in both simulations. However, the MERRA reanalysis data shows more intense values of mean sea level pressure (Table 2) and warm-core. The medicane tracks in CLP08 and CCLM08 are shifted towards the south compared to the MERRA reanalysis data. Figure 11 shows the wind speed comparison of CPL08, CCLM08, and the NOAA dataset on 27 January at 00:00 UTC. The wind speed patterns in CPL08 are stronger and finer compared to the CCLM08 simulations, and in good agreement with the NOAA dataset.

3.5 Cases ME01–ME05, ME07, and ME11

In the remaining ME02–ME05 and ME07 cases, finer structure are only observed in high-resolution (0.08°) simulations. Table 2 shows the lengths of medicane tracks and lowest minimum sea level pressure values of all simulated medicanes. The ME01 case was not well simulated at high-resolution (0.08°); the track length is very short in both the CPL08 (06h) and CCLM08 simulations (08h) compared to the MERRA reanalysis (66h) data (Table 2). For the ME11 case, the medicane tracks in both the 0.22° (CPL22; 13h and CCLM22; 13h) and 0.08° (CPL08; 12h and CCLM08; 06h) simulations are very short compared to the MERRA reanalysis data (24h). Tous et al. (2013) investigated the impact of surface heat fluxes on the intensity and trajectories of the 12 medicanes listed in Table 1. They found that surface heat fluxes do not play significant roles in the intensity and trajectories of these two medicanes (ME01 and ME11). This may be the reason that these two medicanes are not well simulated.

In the CPL22 simulations, the lengths of the medicane tracks in most cases are shorter than the CCLM22 simulations. The mean sea level pressure difference between

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

CPL22 and CCLM22 ranges from 0 to 4.7 hPa. However, the lengths of the medicane track in CPL08 are longer than CCLM08, except ME02, where both have the same track length, and ME10 where CPL08 shows a shorter (1 h) track length compared to CCLM08. The difference between the lowest minimum sea level pressure values in most of the 0.08° simulations ranges from 0 to 1.6 hPa, except ME03, where the difference is slightly higher (3.7 hPa). The results show that warm-core structures are also more intense in the CPL08 simulations compared to CCLM08.

The wind speed is strongly underestimated in all cases of the 0.22° simulations. However, the wind speed is significantly improved in the 0.08° simulations. Compared to CCLM08, the wind speed in CPL08 is more intense and in good agreement with the NOAA dataset. The results show that 0.08° is an appropriate atmospheric grid resolution to resolve most of the meso-scale characteristics associated with medicanes in coupled and atmosphere-only simulations. The coupled simulations at 0.08° also improved the results, particularly the medicane's track lengths, warm-core and wind speed structures compared to atmosphere-only simulations.

4 Conclusions

In this study we examined the ability of the coupled atmosphere–ocean model COSMO-CLM/1-D NEMO-MED12 with atmospheric grid-spacings of 0.44°, 0.22°, and 0.08° (about 50, 25, and 9 km, respectively) and an ocean grid-spacing of 1/12° to simulate medicanes. The results show that the model's performance depends strongly on the atmospheric grid resolution. The large-scale disturbance is well simulated in all three resolutions. Medicane signals are not observed in the 0.44° simulations. In the 0.22° simulations, the mean sea level pressure and warm-core are captured, and are more intense and finer in the 0.08° simulations. The wind speed is strongly underestimated in the 0.22° simulations compared to the NOAA dataset. Most of the medicane features are well resolved at high-resolution (0.08°) in both coupled and atmosphere-only simulations.

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Compared to the atmosphere-only simulations, the coupled model did not show any significant improvement at 0.44° and 0.22° resolutions. In most of the 0.22° simulations, the lengths of the medicane tracks in the atmosphere-only simulations are longer than in the coupled simulations. The lengths of the medicane tracks in the 0.22° atmosphere-only simulations are in good agreement with observations. The wind speed and warm-core structures in the 0.22° (both coupled and atmosphere-only) simulations are not well represented. However, the coupled simulations improved significantly compared to atmosphere-only simulations at higher atmospheric grid resolution (0.08°). The characteristic features of medicanes, such as warm-cores and high wind speeds, are more intense in coupled simulations compared to atmosphere-only simulations. In most cases, medicane tracks in the coupled simulations are longer compared to the atmosphere-only simulations, and therefore in good agreement with observations. These results suggest that a 0.08° grid resolution produces accurate detailed results in medicane simulations, particularly with the coupled model. An atmospheric grid resolution higher than 0.22° is vital to simulate the medicanes more realistically in both coupled and atmosphere-only models.

The coupled and atmosphere-only simulations with the spectral nudging technique increased the accuracy of times and locations of generated medicanes. However, the simulations with spectral nudging did not result in any significant improvement in the mean sea level pressure and wind speed estimates. In the spectral nudging simulations, the medicane tracks and wind speeds in the coupled simulations are better represented compared to the atmosphere-only simulations, similar to simulations performed without spectral nudging.

The present study shows that the coupled model is an effective tool for simulating extreme events such as medicanes. The presented coupled model can be a useful tool for studying tropical-like storms, particularly the ocean feedback effects. The impact of coupling on the vertical structures of medicanes and other important parameters such as precipitation and air–sea fluxes should be analyzed in detail. A full three-dimensional

ocean model can be used for long-term climate simulations and future projections of these extreme events.

Acknowledgements. The authors acknowledge funding from the Hessian Initiative for the Development of Scientific and Economic Excellence (LOEWE) through the Biodiversity and Climate Research Centre (BiK-F), Frankfurt am Main. The authors thank the Center for Scientific Computing (CSC) of the Goethe University Frankfurt am Main for providing computational facilities, especially LOEWE-CSC. We acknowledge Cindy Lebeauapin Brossier, Jonathan Beuvier, Thomas Arzouse, Samuel Somot and Philippe Drobinski for their help for the NEMO-MED12 model and the atmospheric coupling and Eric Maissonave for his support. We also acknowledge the French GIS and GMMC which have supported the NEMO-MED12 model development. N. Akhtar and B. Ahrens acknowledge the support from the German Federal Ministry of Education and Research (BMBF) under grant MiKliP: DECRET/01LP1118B. The authors also acknowledge the HyMex and Med-Codex community.

References

- Akhtar, N.: New approach with a basin-wide single column ocean model set-up to study the Mediterranean Sea dynamics, Master's Thesis, Goethe University, Frankfurt am Main, Germany, 2013. 2121, 2122
- Artale, V., Calmanti, S., Carillo, A., Dell'Aquila, A., Herrmann, M., Pisacane, G., Ruti, Sannino, G., Struglia, M. V., Giorgi, F., Bi, X., Pal, J. S., Rauscher, S., and The PROTHEUS Group: An atmosphere-ocean regional climate model for the Mediterranean area: assessment of a present climate simulation, *Clim. Dyn.*, 35, 721–740, 2010. 2120
- Businger, S. and Reed, R.: Cyclogenesis in cold air masses, *Weather Forecast.*, 20, 133–156, 1989. 2119
- Cavicchia, L. and von Storch, H.: The simulation of medicanes in a high-resolution regional climate model, *Clim. Dynam.*, 39, 2273–2290, 2012. 2122, 2126, 2127, 2129
- Cavicchia, L., von Storch, H. and Gualdi, S.: A long-term climatology of medicanes, *Clim. Dyn.*, 101, doi:10.1007/s00382-013-1893-7, 2013. 2119, 2120
- Dubois, C., Somot, S., Calmanti, S., Carillo, A., Déqué, M., Dell'Aquila, A., Elizalde, A., Gualdi, S., Jacob, D., L'Hévéder, B., Li, L., Oddo, P., Sannino, G., Scoccimarro, E., and Sevault, F.: Future projections of the surface heat and water budgets of the Mediterranean Sea

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

in an ensemble of coupled atmosphere–ocean regional climate models, *Clim. Dynam.*, 39, 1859–1884, 2012. 2120

Elguindi, N., Somot, S., Déqué, M., and Ludwig, W.: Climate change evolution of the hydrological balance of the Mediterranean, Black and Caspian Seas: impact of climate model resolution, *Clim. Dynam.*, 36, 205–228, 2009. 2120

Emanuel, K.: An air–sea interaction theory for tropical cyclones, Part I: Steady-state maintenance, *J. Atmos. Sci.*, 43, 585–605, 1986. 2119

Emanuel, K.: Genesis and maintenance of “Mediterranean hurricanes”, *Adv. Geosci.*, 2, 217–220, doi:10.5194/adgeo-2-217-2005, 2005. 2125

Emanuel, K. and Rotunno, R.: An air–sea interaction theory for tropical cyclones, Part II: Evolutionary study using a nonhydrostatic axisymmetric numerical model, *J. Atmos. Sci.*, 44, 542–561, 1987. 2119

Ernst, J. A. and Matson, M.: A Mediterranean tropical storm, *Weather*, 38, 332–337, 1983. 2120

Fita, L., Romero, R., Luque, A., Emanuel, K., and Ramis, C.: Analysis of the environments of seven Mediterranean tropical-like storms using an axisymmetric, nonhydrostatic, cloud resolving model, *Nat. Hazards Earth Syst. Sci.*, 7, 41–56, doi:10.5194/nhess-7-41-2007, 2007. 2120

Gualdi, S., Somot, S., Li, L., Artale, V., Adani, M., Bellucci, A., Braun, A., Calmanti, S., Carillo, A., Dell’Aquila, A., Déqué, M., Dubois, C., Elizalde, A., Harzallah, A., Jacob, D., L’Hévéder, B., May, W., Oddo, P., Ruti, P., Sanna, A., Sannino, G., Scoccimarro, E., Sevaut, F., and Navarra, A.: The CIRCE simulations: a new set of regional climate change projections performed with a realistic representation of the Mediterranean Sea, *B. Am. Meteorol. Soc.*, 94, 65–81, 2012. 2120

Homar, V., Romero, R., Stensrud, D., Ramis, C., and Alonso, S.: Numerical diagnosis of a small, quasi-tropical cyclone over the Western Mediterranean: dynamical vs. boundary factors, *Q. J. Roy. Meteor. Soc.*, 129, 1469–1490, 2003. 2120, 2125

Hoskins, B. and Hodges, K.: New perspectives on the Northern Hemisphere winter storm tracks, *J. Atmos. Sci.*, 59, 1041–1061, 2002. 2118

Lebeaupin, B. C., Béranger, K., Deltel, C., and Drobinski, P.: The Mediterranean response to different space-time resolution atmospheric forcings using perpetual mode sensitivity simulations, *Ocean Model.*, 36, 1–25, 2011. 2121

Liguang, W. U., Bin Wang and Braun, S. A.: Impacts of Air-Sea Interaction on Tropical Cyclone Track and Intensity, *J. Atmos. Sci.*, 133, 3299–3314, 2005.

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Luque, A., Fita, L., Romero, R., and Alonso, S.: Tropical-like Mediterranean storms: an analysis from satellite, EUMETSAT 07 proceedings, 2007. 2120
- Madec, G.: NEMO ocean engine, Note du Pole de modélisation, Institut Pierre-Simon Laplace (IPSL) France, 27, 1288–1619, 2008. 2121
- 5 Miglietta, M. M., Moscatello, A., Conte, D., Mannarini, G., Lacorata, G., and Rotunno, R.: Numerical analysis of a Mediterranean “hurricane” over south-eastern Italy: sensitivity experiments to sea surface temperature, *Atmos Res.*, 101, 412–426, 2011. 2119, 2120
- Moscatello, A., Miglietta, M. M., and Rotunno, R.: Observational analysis of a Mediterranean “hurricane” over south-eastern Italy, *Mon. Weather Rev.*, 136, 4373–4397, 2008. 2119, 2120
- 10 Niiler, P. P. and Kraus, E. B.: One-dimensional models of the upper ocean, in: *Modelling and Prediction of the Upper Layers of the Oceans*, 325, 1977. 2121
- Palmen, E.: On the formation and structure of tropical hurricanes, *Geophysica*, 3, 26–48, 1948. 2119
- Pettersen, S.: *Weather Analysis and Forecasting*, Mac Graw Hills Book Company, 1956. 2118
- 15 Pytharoulis, I., Craig, G., and Ballard, S.: The hurricane-like Mediterranean cyclone of January 1995, *Meteorol. Appl.*, 7, 261–279, 2000. 2125
- Rasmussen, E. and Zick, C.: A subsynoptic vortex over the Mediterranean with some resemblance to polar lows, *Tellus A*, 39, 408–425, 1987. 2120
- Rienecker, M. M., Suarez, J. M., Gelaro, R., Todling, R., Bacmeister, J, Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G. K., Bloom, S, Chen, J., Collins, D., Conaty, A., Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA-NASA’s Modern-Era Retrospective Analysis for Research and Applications, *J. Climate*, 24, 3624–3648, 2011. 2123
- 20 Rixen, M.: MEDAR/MEDATLAS-II, doi:10.6096/HyMeX.MEDAR/MEDATLAS-II.20120112, 2012. 2122
- Rockel, B., Will, A., and Hense, A.: A spectral nudging technique for dynamical down-scaling purposes, *The regional climate model COSMO-CLM (CCLM)*. *Meteorol Z*, 17, 347–348, 2008. 2121
- 30 Ruti, P. M., Somot, S., Dubois, C., Calmanti, S., Ahrens, B., Alias, A., Aznar, R., Bartholy, J., Bastin, S., Béranger, K., Brauch, J., Calvet, J. C., Carillo, A., Decharme, B., Dell’Aquila, A., Djurdjevic, V., Drobinski, P., Elizalde-Arellano, A., Gaertner, M., Galàn del Sastre, P., Gallardo, C., Giorgi, F., Gualdi, S., Bellucci, A., Harzallah, A., Herrmann, M., Jacob, D., Khodayar,

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

S., Krichak, S., Lebeau-pin, C., L'Heveder B., Li, L., Liguori, G., Lionello, P., Onol, B., Rajkovic. B., Sannino, G., and Sevault, F.: MED-CORDEX initiative for Mediterranean Climate studies, BAMS, in revision, 2012. 2122

5 Sanna, A., Lionello, P., and Gualdi, S.: Coupled atmosphere ocean climate model simulations in the Mediterranean region: effect of a high-resolution marine model on cyclones and precipitation, Nat. Hazards Earth Syst. Sci., 13, 1567–1577, doi:10.5194/nhess-13-1567-2013, 2013. 2120

Simmons, A., Uppala, S., Dee, D., Kobayashi, S.: New ECMWF reanalysis products from 1989 onwards, ECMWF Newsletter, 110, 26–35 2006. 2120

10 Somot S., Sevault F., Déqué M., and Crépon M.: 21st century climate change scenario for the Mediterranean using a coupled Atmosphere-Ocean Regional Climate Model, Glob. Planet. Change, 63, 112–126, doi:10.1016/j.gloplacha.2007.10.003, 2008.

Tous, M. and Remero, R.: Meteorological environments associated with medicane development, Int. J. Climatol., 33, 1–14, 2013. 2119, 2120, 2123, 2137

15 Tous, M., Romero, R., and Ramis, C.: Surface heat fluxes influence on medicane trajectories and intensification, Atmos Res., 123, 400–411, 2013. 2124, 2126, 2130

Trenberth, K.: Uncertainty in hurricanes and global warming, Science, 308, 41753–1754, 2005. 2119

20 Valcke, S.: The OASIS3 coupler: a European climate modelling community software, Geosci. Model Dev., 6, 373–388, doi:10.5194/gmd-6-373-2013, 2013. 2122

von Storch, H., Langenberg, H., and Feser, F.: A spectral nudging technique for dynamical down-scaling purposes, Mon. Weather Rev., 128, 3664–3673, 2000. 2122

Wernli, H. and Schwierz, C.: Surface cyclones in the ERA-40 dataset (1958–2001), Part I: Novel identification method and global climatology, J. Atmos. Sci., 2486–2507, 2006. 2118

25 Zhang, H. M., Bates, J. J., and Reynolds, R. W.: Assessment of composite global sampling: Sea surface wind speed, Geophys Res. Lett., 33, 714, doi:10.1029/2006GL027086, 2006. 2123

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Table 1. Code, date, approximate time of mature phase and geographical coordinates of medicane centers from 1983 to 2003 (Tous and Romero, 2013).

Code	Date	Time (UTC)	Lat (° N)	Lon (° E)	Maximum diameter (km)	Lifetime (h)
ME01	29 Sep 1983	12:00	41.1	6.8	220	90
ME02	7 Apr 1984	06:00	36.4	19.2	230	36
ME03	29 Dec 1984	06:00	35.4	11.6	220	60
ME04	14 Dec 985	12:00	35.5	17.6	290	54
ME05	5 Dec 1991	12:00	36.2	16.7	320	30
ME06	15 Jan 1995	18:00	36.4	19.1	300	78
ME07	12 Sep 1996	12:00	39.4	2.8	170	12
ME08	6 Oct 1996	18:00	37.2	3.9	240	90
ME09	10 Dec 1996	00:00	40.3	3.7	230	48
ME10	26 Jan 1998	12:00	36.7	17.9	250	30
ME11	19 Mar 1999	06:00	38.5	19.6	250	30
ME12	27 May 2003	00:00	40.1	2.8	280	42

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Code, start and end time, lifetime (h; in round brackets) and lowest minimum sea level pressure (hPa; in square brackets) of medicane track in CPL22, CCLM22, CPL08 and CCLM08, and the MERRA reanalysis data.

Code	CPL22	CCLM22	CPL08	CCLM08	MERRA
ME01	26 Sep 1983, 19:00– 28 Sep 1983, 14:00 (43 h) [1005.6]	26 Sep 1983, 19:00– 28 Sep 1983, 15:00 (44 h) [1006.0]	28 Sep 1983, 02:00– 28 Sep 1983, 08:00 (06 h) [1011.2]	28 Sep 1983, 01:00– 28 Sep 1983, 09:00 (08 h) [1010.4]	26 Sep 1983, 06:00– 28 Sep 1983, 23:00 (66 h) [1006.9]
ME02	05 Apr 1984, 19:00– 07 Apr 1984, 08:00 (37 h) [993.6]	05 Apr 1984, 20:00– 07 Apr 1984, 09:00 (37 h) [994.5]	05 Apr 1984, 23:00– 07 Apr 1984, 14:00 (39 h) [994.4]	05 Apr 1984, 00:00– 07 Apr 1984, 16:00 (39 h) [994.7]	06 Apr 1984, 18:00– 08 Apr 1984, 00:00 (30 h) [1000.7.7]
ME03	29 Dec 1984, 09:00– 31 Dec 1984, 18:00 (57 h) [1004.4]	29 Dec 1984, 08:00– 31 Dec 1984, 22:00 (62 h) [999.7]	29 Dec 1984, 03:00– 31 Dec 1984, 23:00 (68 h) [994.2]	29 Dec 1984, 04:00– 31 Dec 1984, 23:00 (67 h) [997.9]	29 Dec 1984, 06:00– 01 Jan 1985, 00:00 (66 h) [997.3]
ME04	13 Dec 1985, 16:00– 14 Dec 1985, 22:00 (28 h) [1008.3]	13 Dec 1985, 13:00– 15 Dec 1985, 02:00 (37 h) [1008.3]	13 Dec 1985, 14:00– 15 Dec 1985, 05:00 (37 h) [1004.8]	13 Dec 1985, 14:00– 15 Dec 1985, 03:00 (35 h) [1004.9]	13 Dec 1985, 12:00– 16 Dec 1985, 00:00 (60 h) [1010.6]
ME05	06 Dec 1991, 20:00– 08 Dec 1991, 11:00 (39 h) [985.1]	06 Dec 1991, 20:00– 08 Dec 1991, 09:00 (37 h) [988.4]	06 Dec 1991, 16:00– 08 Dec 1991, 17:00 (49 h) [982.6]	06 Dec 1991, 18:00– 08 Dec 1991, 15:00 (45 h) [984.2]	07 Dec 1991, 00:00– 08 Dec 1991, 06:00 (30 h) [987.4]
ME06	14 Jan 1995, 01:00– 15 Jan 1995, 15:00 (38 h) [995.5]	14 Jan 1995, 01:00– 15 Jan 1995, 14:00 (37 h) [995.2]	15 Jan 1995, 00:00– 16 Jan 1995, 18:00 (42 h) [991.7]	15 Jan 1995, 00:00– 16 Jan 1995, 10:00 (34 h) [991.1]	15 Jan 1995, 00:00– 17 Jan 1995, 12:00 (48 h) [993.1]
ME07	11 Sep 1996, 23:00– 13 Sep 1996, 04:00 (29 h) [990.3]	11 Sep 1996, 22:00– 12 Sep 1996, 07:00 (09 h) [991.1]	11 Sep 1996, 18:00– 12 Sep 1996, 17:00 (24 h) [992.0]	12 Sep 1996, 04:00– 12 Sep 1996, 17:00 (13 h) [992.6]	12 Sep 1996, 12:00– 12 Sep 1996, 18:00 (06 h) [998.1]
ME08	07 Oct 1996, 03:00– 10 Oct 1996, 05:00 (74 h) [999.8]	07 Oct 1996, 04:00– 10 Oct 1996, 14:00 (90 h) [995.6]	06 Oct 1996, 21:00– 10 Oct 1996, 19:00 (94 h) [995.6]	06 Oct 1996, 22:00– 10 Oct 1996, 18:00 (92 h) [994.1]	07 Oct 1996, 00:00– 11 Oct 1996, 00:00 (72 h) [996.0]
ME09	08 Dec 1996, 18:00– 11 Dec 1996, 02:00 (56 h) [984.0]	08 Dec 1996, 18:00– 10 Dec 1996, 10:00 (40 h) [984.0]	08 Dec 1996, 23:00– 11 Dec 1996, 05:00 (54 h) [988.8]	09 Dec 1996, 01:00– 11 Dec 1996, 03:00 (50 h) [988.8]	09 Dec 1996, 12:00– 12 Dec 1996, 00:00 (60 h) [1002.2]
ME10	26 Jan 1998, 01:00– 26 Jan 1998, 20:00 (07 h) [1000.6]	26 Jan 1998, 01:00– 27 Jan 1998, 07:00 (30 h) [1001.5]	26 Jan 1998, 04:00– 27 Jan 1998, 08:00 (27 h) [997.8]	26 Jan 1998, 03:00– 27 Jan 1998, 08:00 (28 h) [998.9]	26 Jan 1998, 06:00– 27 Jan 1998, 18:00 (36 h) [1001.5]
ME11	17 Mar 1999, 19:00– 18 Mar 1999, 08:00 (13 h) [995.01]	17 Mar 1999, 19:00– 18 Mar 1999, 08:00 (13 h) [995.01]	17 Mar 1999, 20:00– 18 Mar 1999, 08:00 (12 h) [994.55]	17 Mar 1999, 20:00– 18 Mar 1999, 02:00 (06 h) [994.4]	18 Mar 1999, 00:00– 19 Mar 1999, 00:00 (24 h) [992.8]

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.




Fig. 1. ME08; geo-potential height ($\text{m}^2 \text{s}^{-2}$; dotted contours lines at $300 \text{m}^2 \text{s}^{-2}$ intervals) and temperature ($^{\circ}\text{C}$; colored contours at 2°C intervals) at 500hPa in coupled (0.44° , 0.22° , and 0.08°) simulations on 6 October 1996 at 03:00 UTC.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.




Fig. 2. ME08; mean sea level pressure (hPa; dotted contours lines at 2hPa intervals) and temperature ($^{\circ}\text{C}$: colored contours at 2°C intervals) at 700hPa in coupled and atmosphere-only (0.44° , 0.22° , and 0.08°) simulations and the MERRA reanalysis data on 7 October 1996 at 18:00 UTC.

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.




Fig. 3. ME08; 10 m wind speed (ms^{-1}) in coupled and atmosphere-only (0.44° , 0.22° , and 0.08°) simulations and the NOAA “Blended Sea winds” on 7 October 1996 at 18:00 UTC.

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.




Fig. 4. ME08; mean sea level pressure (hPa; dotted contours lines at 2 hPa intervals) and temperature ($^{\circ}\text{C}$: colored contours at 2°C intervals) at 700 hPa in coupled and atmosphere-only (0.08°) with spectral nudging simulations, on 7 October 1996 at 18:00 UTC.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.




Fig. 5. ME08; 10 m wind speed (ms^{-1}) in coupled and atmosphere-only (0.08°) with spectral nudging simulations, on 7 October 1996 at 18:00 UTC.

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.




Fig. 6. ME09; mean sea level pressure (hPa; dotted contours lines at 2 hPa intervals) and temperature (°C: colored contours at 2°C intervals) at 700 hPa in coupled and atmosphere-only (0.08°) simulations and the MERRA reanalysis data on 10 December 1996 at 18:00 UTC.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.




Fig. 7. ME09; 10 m wind speed (ms^{-1}) in coupled and atmosphere-only (0.08°) simulations and the NOAA “Blended Sea winds” on 10 December 1996 at 18:00 UTC.

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.




Fig. 8. ME06; mean sea level pressure (hPa; dotted contours lines at 2 hPa intervals) and temperature (°C: colored contours at 2°C intervals) at 700 hPa in coupled and atmosphere-only (0.08°) simulations and the MERRA reanalysis data on 16 January 1995 at 12:00 UTC.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.




Fig. 9. ME06; 10 m wind speed (ms^{-1}) in coupled and atmosphere-only (0.08°) simulations and the NOAA “Blended Sea winds” on 16 January 1995, at 12:00 UTC.

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.




Fig. 10. ME10; mean sea level pressure (hPa; dotted contours lines at 2hPa intervals) and temperature (°C: colored contours at 2°C intervals) at 700 hPa in coupled and atmosphere-only (0.08°) simulations and the MERRA reanalysis data on 27 January 1998 at 00:00 UTC.

Medicanes in an ocean–atmosphere coupled regional climate model

N. Akhtar et al.




Fig. 11. ME10; 10 m wind speed (ms^{-1}) in coupled and atmosphere-only (0.08°) simulations and the NOAA “Blended Sea winds” on 27 January 1998 at 00:00 UTC.