



This discussion paper is/has been under review for the journal Natural Hazards and Earth System Sciences (NHESD). Please refer to the corresponding final paper in NHESD if available.

# Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa<sup>1</sup>, R. D. Field<sup>2,3</sup>, F. Pappenberger<sup>4,5,6</sup>, A. Langner<sup>7</sup>, S. Enghart<sup>8</sup>,  
U. Weber<sup>9</sup>, T. Stockdale<sup>4</sup>, F. Siegert<sup>8</sup>, J. W. Kaiser<sup>4,10,11</sup>, and J. Moore<sup>12</sup>

<sup>1</sup>Department Environment, Earth & Ecosystems, Open University, Milton Keynes, UK

<sup>2</sup>Department Applied Physics & Applied Mathematics, Columbia University, New York, USA

<sup>3</sup>NASA Goddard Institute for Space Studies, New York, USA

<sup>4</sup>European Centre for Medium-Range Weather Forecasts, Reading, UK

<sup>5</sup>School of Geographical Sciences, University of Bristol, Bristol, UK

<sup>6</sup>College of Hydrology and Water Resources, Hohai University, Nanjing, China

<sup>7</sup>Forest Resources & Climate Unit, Institute for Environment & Sustainability,  
EC-Joint Research Centre, Ispra, Italy

<sup>8</sup>Remote Sensing Solutions GmbH, Munich, Germany

<sup>9</sup>Department Biogeochemical Integration, Max Planck Institute for Biogeochemistry,  
Jena, Germany

<sup>10</sup>Department Atmospheric Chemistry, Max Planck Institute for Chemistry, Mainz, Germany

<sup>11</sup>Department of Geography, King's College London, London, UK

<sup>12</sup>College of Engineering, Mathematics & Physical Sciences, Exeter University, Exeter, UK

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Received: 22 July 2014 – Accepted: 23 July 2014 – Published: 6 August 2014

Correspondence to: A. C. Spessa (allan.spessa@open.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**NHESSD**

2, 5079–5111, 2014

---

**Seasonal forecasting  
of fire over  
Kalimantan,  
Indonesia**

A. C. Spessa et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Large-scale fires occur frequently across Indonesia, particularly in the southern region of Kalimantan and eastern Sumatra. They have considerable impacts on carbon emissions, haze production, biodiversity, health, and economic activities.

In this study, we demonstrate that severe fire and haze events in Indonesia can generally be predicted months in advance using predictions of seasonal rainfall from the ECMWF System 4 coupled ocean-atmosphere model. Based on analyses of up-to-date and long series observations on burnt area and rainfall, and tree cover, we demonstrate that fire activity is negatively correlated with rainfall, and is positively associated with deforestation in Indonesia. There is a contrast between the southern region of Kalimantan (high fire activity, high tree cover loss and strong non-linear correlation between observed rainfall and fire) and the central region of Kalimantan (low fire activity, low tree cover loss and weak non-linear correlation between observed rainfall and fire).

The ECMWF seasonal forecast provides skilled forecasts of burnt area with several months lead time explaining at least 70% of the variance between rainfall and with burnt area. Results are strongly influenced by El Niño years which show a consistent positive bias. Overall, our findings point to a high potential for using a more physical-based method for predicting fires with several months lead time in the tropics, rather than one based on indexes only. We argue that seasonal precipitation forecasts should be central to Indonesia's evolving fire management policy.

## 1 Introduction

The rainforests of equatorial Southeast (SE) Asia comprise some of the largest, oldest, most biodiverse forests on the planet (Page et al., 2011). Indonesian forests and peatlands are globally one of the largest reservoirs of terrestrial organic carbon, with an estimated 14 Gt of above-ground carbon in forests (Saatchi et al., 2011) and around 60

**NHESSD**

2, 5079–5111, 2014

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gt of carbon in the below-ground biomass of peatlands (Page et al., 2011). While fires in Indonesia have occurred throughout paleo-history, their frequency before the 1960s was comparatively rare; coinciding with exceptional but relatively infrequent droughts mostly associated with strong El Niño events (Field et al., 2009). The El Niño Southern Oscillation (ENSO) is the major driver of rainfall variability in the equatorial Pacific region, and occurs irregularly on a 2–7 year intervals lasting about one year but with varying strengths (Aldrian and Susanto, 2003; Dobles-Reyes et al., 2013). During the warm ENSO phase (El Niño) sea surface temperatures in the western Pacific tend to be cooler than normal leading to below normal dry season rainfall and extended dry season length, which increases the risk of mainly degraded forest areas becoming dry enough to burn (Siebert et al., 2001). Large-scale rainfall patterns in the region are also affected by other major weather systems such as the Indian Ocean Dipole (IOD) and the Madden Julian Oscillation (MJO), but their interaction with ENSO are highly complex (Field et al., 2009; Reid et al., 2012; Dobles-Reyes et al., 2013).

Since the early 1960s, however, large-scale fires and related widespread emissions episodes have occurred more frequently across Indonesia, particularly in the southern region of Kalimantan and eastern Sumatra (Field et al., 2008, 2009; Schultz et al., 2008). These episodes are nearly always associated with El Niño events, for example, emissions from biomass burning in Kalimantan were as much as 30 times greater during 2006, an El Niño year, than during 2000, a wet La Niña year (van der Werf et al., 2008). Furthermore, the destructive fires in Indonesia during the exceptionally strong El Niño in late 1997 and early 1998 rank as some of the largest peak emissions events in recorded history. Past studies estimate about 1Gt of carbon was released to the atmosphere from the Indonesian fires in 1997, which were mostly concentrated in carbon-rich forested peatlands of the southern region of Kalimantan. This amount was equivalent to over 10 % of the average global annual fossil fuel emissions released during the 1990s (Page et al., 2002; Schultz et al., 2008; van der Werf et al., 2010).

Over the past couple of decades, Indonesia has experienced some of the world's highest rates of deforestation and forest degradation, principally due to fire (Langner

**Seasonal forecasting  
of fire over  
Kalimantan,  
Indonesia**

A. C. Spessa et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

et al., 2007; Langner and Siegert, 2009; Hoscilo et al., 2011; Miettinen et al., 2011; Page et al., 2011; Hooijer et al., 2012). The general consensus is that the relatively low fire frequency prior to the 1960s in Indonesia, and the relatively higher fire frequency post-1960s are not due to any significant step-up in drought frequency per se, but rather an increase in human-caused ignitions associated with expansion of agriculture, palm and pulp paper plantations, industrial deforestation and peat forest reclamation. Land use activities such as these all make extensive use of fire to clear forest, especially during droughts where the impact of fire is maximized. The activities generally started in Indonesia during the 1960s, and intensified greatly in the early 1990s as exemplified by the *Transmigrasi* projects, including the ill-fated Mega Rice project in the southern region of Kalimantan and subsequent peat reclamation projects in that region (Langner et al., 2007; Langner and Siegert, 2009; Hoscilo et al., 2011; Miettinen et al., 2011; Page et al., 2011; Hooijer et al., 2012).

A clear negative correlation between fire activity and antecedent rainfall in Indonesia based EO-derived fire activity data (Field and Shen, 2008; van der Werf et al., 2008) or proxies of fire activity such as aerosols (Sudiana et al., 2003), haze (Field et al., 2004; Wang et al., 2004; Field et al., 2009), and mid-tropospheric CO (Nasser et al., 2009) has been established. Several studies have demonstrated a clear positive correlation between the fire activity and various indexes of El Niño strength (Fuller and Murphy, 2006; Reid et al., 2012; Wooster et al., 2012). Reid et al. (2012) also discussed the various possible contributions of ENSO, the Indian Ocean Dipole (IOD) and the Madden–Julian Oscillation (MJO) to fire occurrence over Western Indonesia during 2000–2010; concluding that while ENSO is the largest factor influencing fire occurrence, the ENSO Modoki (modified ENSO) and the IOD are also important. Due to the strong influence of ENSO on rainfall patterns across Indonesia, Wooster et al. (2012) demonstrated that statistical forecasting of the extent and magnitude of fire activity a few months in advance based on ENSO indexes is possible. Chen et al. (2011) developed an empirical model to forecast regional fire season severity

with lead times of 3–5 months in Amazonia based on a composite index derived from the Oceanic Niño Index and the Atlantic Multi-decadal Oscillation index.

Recent advances in seasonal climate forecasting based on the use of state-of-the-art dynamical models that couple atmosphere, ocean and land processes and assimilate a vast range of climate-related EO measurements (e.g. sea surface temperatures, SSTs) (Doblas-Reyes et al., 2013) opens up the possibility of forming a more physical-based method for predicting fires with several months lead time in the tropics, rather than one based on indexes only. The purpose of this study is to determine whether severe fire and haze events in Indonesia can be predicted in advance using one such system, the European Centre for Medium Weather Forecasting (ECMWF) Seasonal forecast system (System 4) (Molteni et al., 2011).

Ensemble seasonal climate forecasts issued with several months lead-time is a well-established field, and these forecasts have been applied broadly including managing and assessing risks in agricultural production (Hansen et al., 2011), malaria outbreaks (Jones and Morse, 2010), heatwaves (Lass et al., 2013), flooding and droughts (Pappenberger et al., 2013a, b). Comparatively less work has been done seasonal forecasting of fires, however. The USA has a long-standing seasonal fire danger prediction system (Roads et al., 2005, 2010), which is based on the National Center for Environmental Prediction's Coupled Forecast System (NCEP-CFS) (Saha et al., 2006, 2014). The latest version of the NCEP-CFS generates global and regional spectral model ensemble forecasts over a 3–7 month period, which in turn provides required input meteorological variables for calculating fire danger indices based on the National Fire Danger Rating System (Cohen and Deeming, 1985; Burgan, 1988). Roads et al. (2010) demonstrated the seasonal forecasts of fire danger indices driven by NCEP-CFS outputs had skill in predicting fire activity across western USA. The European Forest Fire Information System (EFFIS) (McInerney et al., 2013) currently provides temperature and rainfall anomalies that are expected to exist over European and Mediterranean areas during the next two months based on the multi-member ECMWF System 4 seasonal forecast system. Areas that are drier and

**Seasonal forecasting  
of fire over  
Kalimantan,  
Indonesia**

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



hotter than normal indicate higher forest fire danger (<http://forest.jrc.ec.europa.eu/effis/applications/long-term-forecast/>).

Since no equivalent seasonal fire forecasting system exists for Indonesia, we were motivated to develop and test one for the severely fire-affected region of southern Kalimantan as a case study, based on seven-month forecasts of monthly rainfall from the ECMWF System 4 for the period 1997–2010. The case study was identified from a comparison of observed burnt area and tree cover patterns across a broader region of southern-central Kalimantan, which generally exhibits a June–November dry season (Aldrian and Susanto, 2003). We evaluated the skill of the seasonal rainfall forecasts against observed rainfall, and in predicting observed burnt area. We further assessed forecast skill by analysing observed burnt area in relation to observed rainfall. Several previous studies have focused on the highly non-linear nature of precipitation and fire occurrence in the region; severe fire happens only below a threshold of seasonal precipitation (Field et al., 2004, 2008, 2009; van der Werf et al., 2008). It is only during years with a sufficiently strong precipitation deficit that disturbed peatlands can dry to their point of ignition and burn. In a practical sense, therefore, seasonal fire forecasting entails determining whether this threshold will be crossed during the upcoming dry season.

## 2 Methods

In order to understand and assess seasonal forecast skill, our analyses proceeded in two stages. We first determined the strength of precipitation-fire relationships over the region of southern-central Kalimantan for our 1997–2010 analysis period, knowing a priori that these relationships have been identified in other and earlier versions of the data used. We then evaluated the forecast skill of System 4, focusing on the degree to which the model could separate the precipitation associated with normal and severe fire years.

### Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.1 Study area and study periods

Borneo is divided among three countries: Indonesia, Malaysia and Brunei (Fig. 1). The broader region of southern-central Kalimantan (Indonesia) (Fig. 1) matches approximately the only region of Borneo identified by Aldrian and Susanto (2003) as having, on the average, a June–November dry season. Preliminary analyses of available burnt area data (see below) revealed that over 95 % of fires occurred during this period.

During our study period, 1997–2010, El Niño events were registered in 1997–1998, 2002, 2004, 2006, 2009 ([http://www.cpc.noaa.gov/products/analysis\\_monitoring/ensostuff/ensoyears.shtml](http://www.cpc.noaa.gov/products/analysis_monitoring/ensostuff/ensoyears.shtml)). However, drought and associated fires during 1998 phase of the 1997–1998 El Niño event were concentrated in East Kalimantan (Siegert et al., 2001; Field et al., 2004), which does not display as clearly defined wet and dry seasons (Aldrian and Susanto, 2003). Analyses of the GPCP rainfall data (see below) confirmed 1998 as a non-drought year for the southern-central region of Kalimantan. For the purposes of this paper, therefore, we regard 1997, 2002, 2004, 2006, 2009 as El Niño years, but not 1998.

## 2.2 Fire data

To increase the robustness of our analyses of burnt area in relation to rainfall and forest cover change, we used two independent monthly burnt area datasets spanning 1997–2010, aggregated to 0.5° gridcells. The first dataset is from the Global Fire and Emissions Database version 4 (GFED4) (Giglio et al., 2013), and the second is from the Remote Sensing Solutions (RSS) GmbH (after Langner and Siegert, 2009, with unpublished updates from RSS GmbH).

The Global Fire and Emissions Database version 4 (GFED4) burnt area is available at a monthly time step at 0.25° resolution, and is based on active fire detection from ERS (European Remote Sensing Satellite) ATSR (Along-Track Scanning Radiometer) World fire Atlas and TRMM (Tropical Rainfall Measuring Mission) VIRS

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





(Visible and Infrared Scanner) as well as the MODIS (Moderate Resolution Imaging Spectroradiometer) burnt area product (MCD64A1) (Giglio et al., 2013). For the MODIS era, the GFED4 monthly burnt area dataset was derived exclusively from the 500 m MCD64A1 burnt area product (Giglio et al., 2013). For the pre-MODIS era, burnt area was derived from calibrating monthly active fire counts from the VIRS and ATSR sensors to monthly burnt area supplied by the MCD64A1 product which was modified according to Giglio et al. (2010).

RSS burnt area data are only available for Borneo, and are based on active fire detections derived from NOAA (National Oceanic and Atmospheric Administration) AVHRR (Advanced Very High Resolution Radiometer), ATSR, and MODIS imagery (Langner and Siegert, 2009). Fires before 2000 were derived from hotspots recorded by NOAA 14 and ATSR by adding their datasets. ATSR was used in addition to NOAA because especially in 1997 and 1998 several large fires were not recorded by NOAA due to operation errors at the receiving station. Fires from 2000 onwards were derived from MODIS hotspots (MOD14/MYD14). One single hotspot pixel (sensor element) represents the area of the corresponding sensor resolution of 1 km. This pixel can be affected by a single fire or more than one fire (Langner and Siegert, 2009).

### 2.3 Forest cover data

Our analyses are based on three tree cover products: the global AVHRR 1993 continuous percentage tree cover product available at 1 km (de Fries et al., 2000), and two MODIS tree cover products available for SE Asia at 500 m for the years 2000 and 2010 (Miettinen et al., 2011).

### 2.4 Rainfall data

Version 1.2 of the Global Precipitation Climatology Project (GPCP) one-degree daily (1DD) rainfall data formed the basis for comparing burnt area in relation to observed rainfall and evaluating seasonal rainfall forecasts from System 4. The dataset is a global

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



product available at a 1° resolution, and combines precipitation estimates from several sources, including infrared (IR) and passive microwave (PM) rain estimates, and rain gauge observations (Adler, 2003; Huffman et al., 2009).

## 2.5 ECMWF seasonal forecast system (System 4)

5 Seasonal predictions of rainfall were derived from the ECMWF Seasonal forecast System (System 4), which provides operational seasonal predictions (Molteni et al., 2011). System 4 is the most recent ECMWF seasonal forecast system and includes 51 member ensembles and consists of 7 month forecasts initialized on the 1st day of every month. The seasonal forecast has a resolution of ~ 79 km (T255L91). It is  
10 coupled with an ocean model which has a horizontal resolution of 1°. Singular vectors and an ensemble of ocean analysis, including sea surface temperatures (SSTs) provide the initial perturbations. Atmospheric model uncertainties are included through a 3-time-level stochastically perturbed parameterized tendency scheme and the stochastic back-scatter scheme, which is similar to the ECMWF medium range forecasting  
15 system. The operational seasonal forecast are accompanied by a hindcast system which covers a 30 year period (from 1981 to today) where the land surface is initialized using an offline simulation driven by ERA Interim data (Dee et al., 2011). The hind cast has usually 15 ensemble members. Molteni et al. (2011) presented an overview of S4 model biases and forecast performance.

20 We restricted our use of the ECMWF System 4 rainfall forecasts to 1 May initializations, that is, seasonal forecasts spanning 1 May–30 November, each year 1997–2010. Mean daily rainfall for each month and each ensemble member was calculated.

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.6 Data preparation and statistical analyses

We aggregated both the burnt area data (GFED4, RSS) and the tree cover data (1993, 2002 and 2010) to 0.5° grid cells, and down-scaled the GPCP data to 0.5° grid cells by linear interpolation.

Total area burnt (GFED4) and total burnt area (RSS) were calculated for each 0.5° gridcell in the southern-central region of Kalimantan by summing the respective monthly burnt areas between June and November 1997–2010. Since a 1997 tree cover product does not exist for the study area, we derived a 1997 tree cover dataset based on simple linear interpolation between 1993 and 2000 tree covers. Spatial plots of total burnt area (RSS), total area burnt (GFED4) were then visually compared to spatial plots of tree cover (1993), tree cover (1997) and tree cover (2010) at 0.5° resolution.

For the central and southern regions of Kalimantan and each year, seasonal mean GFED4 burnt area per gridcell and seasonal mean RSS burnt area per gridcell were calculated by firstly summing the respective monthly burnt areas between June and November for each 0.5° gridcell, then summing these values over all gridcells, and finally dividing by the number of gridcells. The range of each seasonal mean burnt area per gridcell was calculated as  $\pm 1.96 \times SE_{\text{mean}}$ . This is similar to previous studies using seasonal (van der Werf et al., 2008; Wooster et al., 2012) or monthly mean (Wang et al., 2004; Field et al., 2008, 2009) estimates of fire activity or haze at an island-scale.

For the central and southern regions of Kalimantan and each year, seasonal mean GPCP rainfall per gridcell was calculated by firstly taking the average of the daily mean rainfall per month values between June and November for each 0.5° gridcell, then summing these values over all gridcells, and finally dividing by the number of gridcells. The range of each seasonal mean GPCP rainfall per gridcell was calculated as  $\pm 1.96 \times SE_{\text{mean}}$ .

For the southern region of Kalimantan only and each year, seasonal mean System 4 rainfall per gridcell was calculated by firstly restricting the System 4 outputs to the middle-ranked 17 members of the 51 member ensemble (middle tercile), then

**NHESSD**

2, 5079–5111, 2014

### Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



averaging over the mean daily rainfall per month values between June and November for each System 4 gridcell, then summing these values over all gridcells, and finally dividing by the number of gridcells. The mean of the highest-ranked 17 members of the 51 member System 4 ensemble (upper tercile) and the mean of the lowest-ranked 17 members of the 51 member System 4 ensemble (lower tercile) were calculated similarly. The upper range for each year was defined as  $\text{mean}_{\text{upper tercile}} - \text{mean}_{\text{middle tercile}}$ , and the lower range for each year was defined as  $\text{mean}_{\text{middle tercile}} - \text{mean}_{\text{lower tercile}}$ .

The following statistical analyses were undertaken:

- correlation between seasonal mean GFED4 burnt area per gridcell and seasonal mean RSS burnt area per gridcell;
- regression of seasonal mean GFED4 burnt area per gridcell on seasonal mean GPCP rainfall per gridcell, central and southern regions of Kalimantan;
- regression of seasonal mean RSS burnt area per gridcell on seasonal mean GPCP rainfall per gridcell, central and southern regions of Kalimantan;
- regression of seasonal mean GFED4 burnt area per gridcell on seasonal mean System 4 rainfall per gridcell, southern region of Kalimantan;
- regression of seasonal mean RSS burnt area per gridcell on seasonal mean System 4 rainfall per gridcell, southern region of Kalimantan; and
- correlation between seasonal mean GPCP rainfall per gridcell and seasonal mean System 4 rainfall per gridcell, southern region of Kalimantan.

Each regression analysis comprised two parts: linear regression of the form  $Y = a + b \times X$ , and non-linear regression of the form  $Y = a + b \times \ln(X)$ ; where  $Y$  is seasonal mean burnt area per gridcell (GFED4 or RSS),  $X$  is seasonal mean daily rainfall per gridcell (GPCP or System 4) and  $\ln$  is the natural log transformation.

### 3 Results

#### 3.1 Fire activity vs. tree cover change

Fire activity and tree cover loss were clearly both much higher across a distinct southern region of Kalimantan ( $N = 53$  gridcells at  $0.5^\circ$  resolution) compared with a central region of Kalimantan immediately above it ( $N = 40$  gridcells at  $0.5^\circ$  resolution) (Fig. 1). Seasonal mean GFED4 area burnt in the south was, on the average, 27.1 times higher than in the central region (Table 1); and seasonal mean RSS area burnt in southern region was, on the average, 4.2 times higher than in the central region (Table 1). In the southern region, the mean tree cover in 2010 was 17.9% lower than that for 1993, and 10.2% lower than that for 1997 (Table 1). In the central region, the mean tree cover in 2010 was 8.2% lower than that for 1993, and 4.4% lower than that for 1997 (Table 1).

#### 3.2 RSS vs. GFED4 burnt area

Seasonal mean RSS burnt area and seasonal mean GFED4 burnt area was highly correlated in the southern region ( $R^2 = 0.96$ ) and the central region ( $R^2 = 0.88$ ). Across the years in which El Niño affected the central-southern region (1997, 2002, 2003, 2004, 2006 and 2009), seasonal mean RSS burnt area was consistently much higher than seasonal mean GFED4 burnt area. In 1997 (pre-MODIS), seasonal mean RSS burnt area was 1.9 times higher than seasonal mean GFED4 burnt area in the south, and 10.2 times higher than the equivalent for GFED4 in the central region (Table 1). Averaged across the other El Niño affected years (all post-MODIS), RSS burnt area was 1.4 times higher than seasonal mean GFED4 burnt area in the south, and 9.6 times higher than the equivalent for GFED4 in the central region (Table 1).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.3 Burnt area vs. observed rainfall

Seasonal mean burnt area was non-linearly and highly correlated with seasonal mean GPCP rainfall for the GFED4 dataset (southern region:  $R^2 = 0.86$ ; central region:  $R^2 = 0.80$ ; Fig. 2a and b) and for the RSS dataset (southern region:  $R^2 = 0.90$ ; central region:  $R^2 = 0.70$ ; Fig. 3a and b). For the GFED4 dataset, the magnitude of the slopes and the intercepts determining the non-linear relationships were respectively 29.8 and 27.2 times greater for the southern region than the central region (Fig. 2a and b). For the RSS dataset, the magnitude of the slopes and the intercepts determining the non-linear relationships were respectively 5.2 and 4.6 times greater for the southern than in the central region (Fig. 3a and b). For the southern region, we diagnosed a clear rainfall threshold approximately equal to 6 mm, above which fire activity was almost negligible, and below which it increased exponentially (Figs. 2b and 3b).

### 3.4 Burnt area vs. System 4 rainfall

Focusing on the southern region, seasonal mean burnt area was non-linearly and significantly correlated with seasonal mean daily System 4 rainfall for the RSS dataset ( $R^2 = 0.70$ ) (Fig. 4a) and the GFED4 dataset ( $R^2 = 0.76$ ) (Fig. 4b). For each System 4-based relationship, the year 2006 was an obvious outlier. Both System 4-based analyses revealed a similar rainfall threshold approximately equal to 7 mm, but this is less clear-cut compared to the GPCP-based analyses (Fig. 4a and b).

### 3.5 System 4 rainfall vs. observed rainfall

Seasonal mean System 4 rainfall and seasonal mean GPCP rainfall were highly correlated ( $R^2 = 0.91$ ), and for each year, their respective values overlapped (Fig. 5). Seasonal mean System 4 rainfall was higher than seasonal mean GPCP rainfall during each El Niño affected year: 1997 (+0.19 mm), 2002 (+1.46 mm), 2004 (+0.52 mm), 2006 (+2.2 mm) and 2009 (+0.88 mm) (Fig. 5).

### 3.6 System 4 predicted vs. observed SSTs

Further analyses of the 15 member ensemble System4 hindcasts from 1997 to 2010 demonstrated that while most members correctly predicted observed SSTs, 2006 was particularly anomalous (Fig. 6). Only 2 out of 15 members predicted the SST cooling in this year. By contrast, in 1997, when the strongest cooling was observed, 13 out of 15 members predicted a strong cooling (Fig. 6).

## 4 Discussion

### 4.1 Observed fire, rainfall, tree cover patterns

Based on analyses of a comparatively up-to-date and long series observations on burnt area and rainfall (14 years from 1997 to 2010), and tree cover (1993, 1997, 2000 and 2010), our work supports a large body of work that demonstrates for different tropical regions, fire activity is negatively correlated with rainfall, and positively associated with deforestation in Indonesia (Langner et al., 2007; van der Werf et al., 2008; Langner and Siegert, 2009; Field et al., 2009), Amazonia (Aragao et al., 2008) and Columbia (Armenteras-Pascual et al., 2011). Our results highlight the stark contrast between the southern region of Kalimantan (high fire activity, high tree cover loss and strong non-linear correlation between rainfall and fire) and the central region of Kalimantan (low fire activity, low tree cover loss and weak non-linear correlation between rainfall and fire). The amount of forest cover loss between 1997 and 2010 we estimated is likely to be conservative. Forest cover immediately prior to the major El Niño induced fires of 1997 probably matches more closely forest cover in 1993, than that based on a simple linear interpolation between forest cover in 1993 and 2002. Although fire data covering the 1993 to 1997 period are not available, no major El Niño events occurred during this period, except for a relatively minor event in 1994–1995 ([http://www.cpc.noaa.gov/products/analysis\\_monitoring/ensostuff/ensoyears.shtml](http://www.cpc.noaa.gov/products/analysis_monitoring/ensostuff/ensoyears.shtml)). Our finding of strong

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Seasonal forecasting of fire over Kalimantan, Indonesia**

A. C. Spessa et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

non-linear relationships between fire and rainfall for southern Kalimantan confirms similar relationships reported for the region (van der Werf et al., 2008; Field and Shen, 2009). The exponential increase in fire activity in southern Kalimantan below a moisture threshold of around  $6 \text{ mm day}^{-1}$  supports the finding of a similar fire moisture threshold for the region (Field et al., 2008).

Our results support the general consensus that while fire activity is strongly linked to drought in Kalimantan, the strength of this relationship is mediated by land use and land cover change and the presence of disturbed peatlands. Numerous studies have reported the widespread deliberate use of fire in the southern region of Kalimantan to clear forests to establish plantations (mainly rice, oil palm and acacia), as well as the widespread incidence of escaped fires associated with these activities and logging (Langner et al., 2007; van der Werf et al., 2008; Langner and Siegert, 2009). Furthermore, extensive areas of peatland have been drained for plantation establishment, which has led to an increase in fire activity in these areas because drained peats lower the water table, exposing a greater depth of dry peat to burning (Page et al., 2011; Hooijer et al., 2012). By contrast, the central region of Kalimantan has undergone comparatively little development to date (Langner et al., 2007; Langner and Siegert, 2009; Margono et al., 2014). The central region has steep slope sections which reduce the ease of forest exploitation and clearing compared to the relatively more flat terrain of the southern region (Langner and Siegert, 2009), and due to its more varied topography, it contains relatively less peat (Page et al., 2011; Hooijer et al., 2012).

**4.2 Seasonal forecasting of fire**

Based on two independent burnt area datasets, we demonstrated that seasonal forecasts of rainfall from the ECMWF System 4 can predict fire activity during the June–November dry season in the southern region of Kalimantan. The empirical fits between System 4 rainfall and fire activity were highly non-linear, similar to those based on observed rainfall. The correlations between System 4 rainfall and fire activity while





## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

not as good as those between observed rainfall and fire activity were nonetheless significant (RSS burnt area dataset:  $R^2 = 0.70$ ; GFED4 burnt area dataset:  $R^2 = 0.76$ ). These significant System 4 rainfall–burnt area relationships demonstrate a prospective use of System 4 rainfall to predict seasonal fire activity as part of a region-wide fire management programme for the southern region of Kalimantan. Further development of any such programme would need to consider however three important factors potentially affecting the robustness of predicting burnt area as a simple function of modelled rainfall. These factors are: (i) a positive rainfall bias in the System 4 model, (ii) biases in the burnt area datasets used to construct these relationships, and (iii) the influence of land use and land cover change, as well as peat drainage, on rainfall vs. fire relationships for the southern region of Kalimantan.

### 4.3 Rainfall bias in seasonal forecast

Although ECMWF and observed rainfall were well-correlated ( $R^2 = 0.91$ ), ECMWF rainfall was consistently higher during the El Niño affected years of 1997, 2002, 2004, 2006 and 2009, that is, years with high fire activity. Furthermore, this bias was not consistent across these drought years. The smallest positive bias occurred during the strong 1997 El Niño year of 1997 (+0.19 mm), and the greatest positive bias occurred during the moderate El Niño year of 2006 (+2.2 mm). The later was clearly an outlier that reduced the goodness-of-fit of the empirical relationships between ECMWF rainfall and fire activity. Although the estimated moisture threshold diagnosed from the ECMWF rainfall vs. fire relationships (7 mm) was similar to that estimated from the GPCP rainfall vs. fire relationships (6 mm), it not as clear-cut because of the 2006 outlier.

Seasonal predictability in the maritime continent is to a large extent due to the El Niño–Southern Oscillation (ENSO); and the changes in SSTs associated with these oscillations (Oldenborgh et al., 2005; Lavers et al., 2009; Dobles-Reyes et al., 2013). In general, cooler SSTs indicate below-average rainfall, and warmer SSTs indicate above-average rainfall. Our analyses of SSTs from multi-member System 4 hindcasts



---

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



covering our study region clearly showed that while most members correctly predicted observed SSTs, the worst result was in 2006, where only 13% of members predicted the SST cooling in that year. The best was in 1997 where close to 90% predicted the SST cooling, corresponding with our observation that the smallest difference in System4 and GPCP rainfall during any drought year was in 1997. Whether the relatively poor prediction in 2006 is a consequence of model error, or is simply due to the event having low predictability, is not known. Some variations in SST and rainfall will always be unpredictable, but we do know that model error has the potential to play a role.

Molteni et al. (2011) reported a positive rainfall bias in Indonesia for System 4. They suggested this was consistent with too strong easterly trade winds simulated by the System 4 atmospheric model in the central and western Pacific, which act to reduce the movement of the western warm pool towards the east Pacific during El Niño onset, hence maintaining warmer than expected SSTs in the model for Indonesia. The impact of this bias on the specific 2006 forecast is not known, but it suggests that more work is needed to improve the feedback between SSTs and climate in System 4, with direct benefits for seasonal forecasting of fires in the region.

#### 4.4 Biases in the burnt area

Although the RSS and GFED4 burnt area data yielded similar patterns (southern region of Kalimantan burning much more than the central region, and non-linear relationships between observed rainfall and fire activity, and between System 4 rainfall and fire activity), RSS burnt area was consistently higher than GFED4 burnt area. During the MODIS era (2000 onwards), GFED4 is based on 500 m MODIS burnt area product (MDC64A1) (Giglio et al., 2013), and RSS is based on 1 km<sup>2</sup> MODIS active fire product (MOD14/MYD14) (Langner and Siegert, 2009). Previous work has shown that the MODIS burnt area product tends to underestimate fire activity because a 16 day cloud free mosaic is necessary to map the burnt areas (Roy et al., 2008). By contrast, the RSS data probably overestimates burnt area because these data are based on the assumption that each fire detection within 1 km<sup>2</sup> pixel means that the whole pixel is

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



burnt. Pre-MODIS RSS burnt area data are based on AVHRR 14 and ATSR hotspot data (Langner and Siegert, 2009) and pre-MODIS GFED4 data are based on ATSR and VIRS hotspots followed by a further correction (Giglio et al., 2010, 2013). For several reasons, AVHRR generally detects a much higher number of active fire events than ATSR (Langner and Siegert, 2009), which most likely explains the higher burnt area recorded by the RSS dataset compared with the GFED4 dataset in 1997. The AVHRR sensor records data in Borneo twice a day, but is saturated at low temperatures and sun glint, leading frequently to false alarms. While the spatial accuracy of the ATSR sensor is high and there are few false alarms due to night-time acquisition, it displays a high rate of omission because fire activity often peaks in the afternoon and the revisit cycle is only every 3 days.

### 4.5 Influence of land use and land cover change, and peat drainage

The southern region of Kalimantan has undergone extensive drainage of peatlands and widespread fire-induced loss of pristine forests due to logging activities and plantation establishment, as discussed above. This has created large tracts of forest that are highly degraded and fragmented (Siegert et al., 2001; Langer et al., 2007, 2009; Hoscilo et al., 2011). Future increases in the extent of degraded forests and drained peats will likely modify seasonal rainfall-burnt area relationships. Degraded forests respond more rapidly to rainfall deficits than undisturbed forests due to increased solar radiation reaching the surface and hence, higher evaporation rates; and provide a relatively greater source of flammable fuels for burning, such as invasive grasses and ferns, as well as debris from logging and land clearing operations (Siegert et al., 2001; Langer et al., 2007, 2009; Hoscilo et al., 2011). The moisture content of peats is controlled by the water table, and as previously discussed; the lower water table of drained peats means more dry peat available for burning than undisturbed peats (Field and Shen, 2008; Page et al., 2011; Hooijer et al., 2012). An increase in the availability of drier fuels from degraded forests and drained peats would therefore potentially increase the amount of burning beyond that forecasted by seasonal rainfall alone. This could be

manifested as an increase in the rate of change in burnt area with respect to a unit decrease in rainfall and/or a higher rainfall threshold controlling fire activity.

## 5 Conclusion

While operational seasonal fire prediction over Indonesia will have to incorporate the above factors, we have demonstrated, for the first time, that severe fire events (and potentially associated haze) are fundamentally predictable months in advance using state-of-the art seasonal rainfall forecasts. Seasonal forecasts are expected to continue to improve in the future and additional post-processing may increase the skill of the fire forecast (Peng et al., 2014). Given the considerable effort required in mobilising prevention and preparedness measures in Indonesia, we therefore argue that seasonal precipitation forecasts should be central to Indonesia's evolving fire management policies. Other potential applications of seasonal fire forecasting include improved risk assessments of biodiversity and carbon losses through fire; both important considerations for tropical forest protection programmes (e.g. REDD+; Barlow et al., 2012), and forest (re)insurance enterprises (Cottle, 2007).

*Acknowledgements.* A. Spessa appreciates funding support provided by the Open University Research Investment Fellowship scheme. The work by ECMWF was partially funded by the Copernicus program (<http://www.copernicus.eu/>). J. Kaiser was supported by the EU FP7 project MACC-II (grant agreement 283576).

## References

- Adler, R. F., Huffman, G. J., Bolvin, D. T., Curtis, S., and Nelkin, E. J.: Tropical rainfall distributions determined using TRMM combined with other satellite and rain gauge information, *J. Appl. Meteorol.*, 39, 2007–2023, 2000.
- Aldrian, E. and Susanto, R. D.: Identification of three dominant rainfall regions within Indonesia and their relationship to sea surface temperature, *Int. J. Climatol.*, 23, 1435–1452, 2003.

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Aragao, L. E. O., Malhi, Y., Barbier, N., Lima, A., Shimabukuro, Y., Anderson, L., and Saatchi, S.: Interactions between rainfall, deforestation and fires during recent years in the Brazilian Amazonia, *Philos. T. R. Soc. B*, 3631498, 1779–1785, 2008.
- Armenteras-Pascual, D., Retana-Alumbreros, J., Molowny-Horas, R., Roman-Cuesta, R. M., Gonzalez-Alonso, F., and Morales-Rivas, M.: Characterising fire spatial pattern interactions with climate and vegetation in Colombia, *Agr. Forest Meteorol.*, 151, 279–289, 2011.
- Barlow, J., Parry, L., Gardner, T. A., Ferreira, J., Aragão, L. E., Carmenta, R., and Cochrane, M. A.: The critical importance of considering fire in REDD+ programs, *Biol. Conserv.*, 154, 1–8, 2012.
- Burgan, R. E.: 1988 Revisions to the 1978 National Fire-Danger Rating System, Research Paper SE-273, USDA Forest Service, Southeastern Forest Experiment Station, Asheville, NC, 1988.
- Chen, Y., Randerson, J. T., Morton, D. C., DeFries, R. S., Collatz, G. J., Kasibhatla, P. S., Giglio, L., Jin, Y., and Malier, M. E.: Forecasting fire season severity in South America using sea surface temperature anomalies, *Science*, 334, 787–791, 2011.
- Cohen, J. D., Deeming, J. E.: The National Fire Danger Rating System: Basic Equations, General Technical Report PSW-82, USDA Forest Service Pacific Southwest Forest and Range Experimental Station, Berkeley, CA, 16 pp., 1985.
- Cottle, P.: Insuring Southeast Asian commercial forests: fire risk analysis and the potential for use of data in risk pricing and reduction of forest fire risk, *Mitig. Adapt. Strat. Glob. Change*, 12, 181–201 doi:10.1007/s11027-006-9042-9, 2007.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kallberg, P., Kohler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, 2011.
- DeFries, R. S., Hansen, M. C., and Townshend, J. R. G.: Global continuous fields of vegetation characteristics: a linear mixture model applied to multi-year 8 km AVHRR data, *Int. J. Remote Sens.*, 21, 1389–1414, 2000.

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Doblas-Reyes, F. J., García-Serrano, J., Lienert, F., Biescas, A. P., and Rodrigues, L. R.: Seasonal climate predictability and forecasting: status and prospects, *WIREs Clim. Change*, 4, 245–268, doi:10.1002/wcc.2174, 2013.
- Field, R. D., van der Werf, G. R., and Shen, S. S.: Human amplification of drought-induced biomass burning in Indonesia since 1960, *Nat. Geosci.*, 2, 185–188, 2009.
- Fuller, D. O. and Murphy, K.: The ENSO-fire dynamic in insular Southeast Asia, *Climatic Change*, 74, 435–455, doi:10.1007/s10584-006-0432-5, 2006.
- Giglio, L., Randerson, J. T., van der Werf, G. R., Kasibhatla, P. S., Collatz, G. J., Morton, D. C., and DeFries, R. S.: Assessing variability and long-term trends in burned area by merging multiple satellite fire products, *Biogeosciences*, 7, 1171–1186, doi:10.5194/bg-7-1171-2010, 2010.
- Giglio, L., Randerson, J. T., and Werf, G. R.: Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4), *J. Geophys. Res.-Biogeo.*, 118, 317–328, 2013.
- Hansen, J. W., Mason, S. J., Sun, L., and Tall, A.: Review of seasonal climate forecasting for agriculture in sub-Saharan Africa, *Exp. Agr.*, 47, 205–240, 2011.
- Hooijer, A., Page, S., Jauhainen, J., Lee, W. A., Lu, X. X., Idris, A., and Anshari, G.: Subsidence and carbon loss in drained tropical peatlands, *Biogeosciences*, 9, 1053–1071, doi:10.5194/bg-9-1053-2012, 2012.
- Hosillo, A., Page, S. E., Tansey, K. J., and Rieley, J. O.: Effect of repeated fires on land-cover change on peatland in southern Central Kalimantan, Indonesia, from 1973 to 2005, *Int. J. Wildland Fire*, 20, 578–588, 2011.
- Huffman, G. J., Adler, R. F., Bolvin, D. T., and Gu, G.: Improving the global precipitation record: GPCP version 2.1, *Geophys. Res. Lett.*, 36, L17808, doi:10.1029/2009GL040000, 2009.
- Jones, A. E. and Morse, A. P.: Skill of ENSEMBLES seasonal re-forecasts for malaria prediction in West Africa, *Geophys. Res. Lett.*, 39, L23707, doi:10.1029/2012GL054040, 2012.
- Langner, A. and Siegert, F.: Spatiotemporal fire occurrence in Borneo over a period of 10 years, *Glob. Change Biol.*, 15, 48–62, 2009.
- Langner, A., Miettinen, J., and Siegert, F.: Land cover change 2002–2005 in Borneo and the role of fire derived from MODIS imagery, *Glob. Change Biol.*, 13, 2329–2340, 2007.
- Lass, W., Haas, A., Hinkel, J., and Jaeger, C.: Avoiding the avoidable: towards a European heat waves risk governance, *International Journal of Disaster Risk Science*, 2, 1–14, 2011.

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Lavers, D., Luo, L., and Wood, E. F.: A multiple model assessment of seasonal climate forecast skill for applications, *Geophys. Res. Lett.*, 36, L23711, doi:10.1029/2009GL041365, 2009.
- Margono, B. A., Potapov, P. V., Turubanova, S., Stolle, F., and Hansen, M. C.: Primary forest cover loss in Indonesia over 2000–2012, *Nature Climate Change*, 4, 730–735, 2014.
- 5 McInerney, D., San-Miguel-Ayanz, J., Corti, P., Whitmore, C., Giovando, C., and Camia, A.: Design and function of the European forest fire information system, *Photogramm. Eng. Rem. S.*, 79, 965–973, 2013.
- Miettinen, J., Shi, C., and Liew, S. C.: Deforestation rates in insular Southeast Asia between 2000 and 2010, *Glob. Change Biol.*, 17, 2261–2270, 2011.
- 10 Molteni, F., Stockdale, T., Balmaseda, M., Balsamo, G., Buizza, R., Ferranti, L., Magnusson, L., Mogensen, K., Palmer, T., and Vitart, F.: The new ECMWF seasonal forecast system (System 4) (p. 49), European Centre for Medium-Range Weather Forecasts, Reading, UK, 2011.
- Nassar, R., Logan, J. A., Megretskaia, I. A., Murray, L. T., Zhang, L., and Jones, D.: Analysis of tropical tropospheric ozone, carbon monoxide, and water vapor during the 2006 El Niño using TES observations and the GEOS-Chem model, *J. Geophys. Res.*, 114, D17304, doi:10.1029/2009JD011760, 2009.
- 15 Page, S. E., Siegert, F., Rieley, J. O., Boehm, H. D. V., Jaya, A., and Limin, S.: The amount of carbon released from peat and forest fires in Indonesia during 1997, *Nature*, 420, 61–65, 2002.
- 20 Page, S. E., Rieley, J. O., and Banks, C. J.: Global and regional importance of the tropical peatland carbon pool, *Glob. Change Biol.*, 17, 798–818, 2011.
- Pappenberger, F., Stephens, E., Thielen, J., Salamon, P., Demeritt, D., Ansel, S. J., Wetterhall, F., and Alfieri, L.: Visualizing probabilistic flood forecast information: expert preferences and perceptions of best practice in uncertainty communication, *Hydrol. Process.*, 27, 132–146, 2013a.
- 25 Pappenberger, F., Wetterhall, F., Dutra, E., and C, H. L.: Seamless forecasting of extreme events on a global scale, in: *Climate and Land Surface Changes in Hydrology*, edited by: Boegh, E., Blyth, E., Hannah, D. M., Hisdal, H., Kunstmann, H., Su, B., and Yilmaz, K. K., IAHS Publication, Gothenburg, Sweden, 3–10, 2013b.
- 30



## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Peng, Z., Wang, Q. J., Bennett, J. C., Schepen, A., Pappenberger, F., Pokhrel, P., and Wang, Z.: Statistical calibration and bridging of ECMWF System4 outputs for forecasting seasonal precipitation over China, *J. Geophys. Res. Atmos.*, 119, 7116–7135, doi:10.1002/2013JD021162, 2014.

5 Roads, J., Fujioka, F., Chen, S., Burgan, R.: Seasonal fire danger forecasts for the USA, *Int. J. Wildland Fire*, 14, 1–18, doi:10.1071/WF03052, 2005.

Roads, J., Tripp, P., Juang, H., Wang, J., Fujioka, F., and Chen, S.: NCEP–ECPC monthly to seasonal US fire danger forecasts, *Int. J. Wildland Fire*, 19, 399–414, 2010.

10 Roy, D. P., Boschetti, L., Justice, C. O., and Ju, J.: The collection 5 MODIS burned area product – global evaluation by comparison with the MODIS active fire product, *Remote Sens. Environ.*, 112, 3690–3707, 2008.

15 Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T., Salas, W., Zutta, B. R., Buermann, W., Lewis, S. L., Hagen, S., Petrova, S., White, L., Silman, M., and Morel, A.: Benchmark map of forest carbon stocks in tropical regions across three continents, *P. Natl. Acad. Sci. USA*, 108, 9899–9904, 2011.

Saha, S., Nadiga, S., Thiaw, C., Wang, J., Wang, W., Zhang, Q., van den Dool, H. M., Pan, H.-L., Moorthi, S., Begringer, D., Stokes, D., Pena, M., Lord, S., White, G., Ebisuzaki, W., Peng, P., and Xie, P.: The NCEP climate forecast system, *J. Climate*, 19, 3483–3517, 2006.

20 Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y.-T., Chuang, H.-Y., Iredell, M., Ek, M., Meng, J., Yang, R., Peña Mendez, M., van den Dool, H., Zhang, Q., Wang, W., Chen, M., and Becker, E.: The NCEP climate forecast system version 2, *J. Climate*, 27, 2185–2208, 2014.

Schultz, M. G., Heil, A., Hoelzemann, J. J., Spessa, A., Thonicke, K., Goldammer, J. G., Held, A. C., Pereira, J. M. C., and van het Bolscher, M.: Global wildland fire emissions from 1960 to 2000, *Global Biogeochem. Cy.*, 22, GB2002, doi:10.1029/2007GB003031, 2008.

25 Sudiana, D., Kuze, H., Takeuchi, N., and Burgan, R. E.: Assessing forest fire potential in Kalimantan Island, Indonesia, using satellite and surface weather data, *Int. J. Wildland Fire*, 12, 175–184, 2003.

30 van der Werf, G. R., Dempewolf, J., Trigg, S. N., Randerson, J. T., Kasibhatla, P. S., Giglio, L., Murdiyarso, D., Peters, W., Morton, D. C., Collatz, G. J., Dollman, A. J., and Defries, R. S.: Climate regulation of fire emissions and deforestation in equatorial Asia, *P. Natl. Acad. Sci. USA*, 105, 20350–20355, 2008.



---

**Seasonal forecasting  
of fire over  
Kalimantan,  
Indonesia**A. C. Spessa et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S.,  
Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and  
the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009),  
Atmos. Chem. Phys., 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

5 van Oldenborgh, J. G., Balmaseda, M. A., Ferranti, L., Stockdale, T. N., and Anderson, D. L.:  
Did the ECMWF seasonal forecast model outperform statistical ENSO forecast models over  
the last 15 years?, J. Climate, 18, 3240–3249, 2005.

Wang, Y., Field, R. D., and Roswintiarti, O.: Trends in atmospheric haze induced by peat fires  
in Sumatra Island, Indonesia and El Niño phenomenon from 1973 to 2003, Geophys. Res.  
10 Lett., 31, L04103, doi:10.1029/2003GL018853, 2004.

Wooster, M. J., Perry, G. L. W., and Zoumas, A.: Fire, drought and El Niño relationships on  
Borneo (Southeast Asia) in the pre-MODIS era (1980–2000), Biogeosciences, 9, 317–340,  
doi:10.5194/bg-9-317-2012, 2012.

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

**Table 1.** Summary statistics for burnt area and tree cover in the southern and central regions of Kalimantan. Values shown are mean  $\pm 1.96 \times SE_{\text{mean}}$ . Number of gridcells in the southern region = 53. Number of gridcells in the central region = 40. Gridcell resolution =  $0.5^\circ$ .

	southern region of Kalimantan	central region of Kalimantan
Seasonal mean GFED4 burnt area per gridcell (averaged over 1997–2010) ( $\text{km}^2$ )	$81.3 \pm 15.0$	$3 \pm 1.3$
Seasonal mean RSS burnt area per gridcell (averaged over 1997–2010) ( $\text{km}^2$ )	$146.9 \pm 15.0$	$35.1 \pm 5.7$
Mean tree cover per gridcell 1993 (%)	$76.0 \pm 1.5$	$92.4 \pm 0.9$
Mean tree cover per gridcell 1997 (%)	$68.3 \pm 2.0$	$87.6 \pm 1.4$
Mean tree cover per gridcell 2010 (%)	$58.1 \pm 2.5$	$84.2 \pm 1.8$
Seasonal mean GFED4 burnt area per gridcell (1997) ( $\text{km}^2$ )	$363.1 \pm 54.5$	$9.5 \pm 4.4$
Seasonal mean RSS burnt area per gridcell (1997) ( $\text{km}^2$ )	$514.5 \pm 52.8$	$96.5 \pm 13.9$
Seasonal mean GFED4 burnt area per gridcell (averaged over El Nino years 2002, 2004, 2006, 2009) ( $\text{km}^2$ )	$155.6 \pm 34.7$	$5.7 \pm 2.8$
Seasonal mean RSS burnt area per gridcell (averaged over El Nino years 2002, 2004, 2006, 2009) ( $\text{km}^2$ )	$291.5 \pm 31.7$	$54.5 \pm 9.2$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.




**Figure 1.** The island of Borneo showing patterns in burnt area (top) and tree cover (bottom) across the southern-central region of Kalimantan: **(a)** total GFED4 burnt area, 1997–2010; **(b)** total RSS burnt area, 1997–2010; **(c)** tree cover 1993; **(d)** tree cover 1997; and **(e)** tree cover 2010. Data are plotted at  $0.5^\circ$  gridcell resolution. Number of  $0.5^\circ$  gridcells in the southern region of Kalimantan = 53. Number of  $0.5^\circ$  gridcells in the central region of Kalimantan = 40.


## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

a) southern region of Kalimantan



b) central region of Kalimantan



**Figure 2.** (a) The southern region of Kalimantan: seasonal mean GFED4 burnt area per gridcell as a function of seasonal mean GPCP rainfall per gridcell; and (b) the central region of Kalimantan: seasonal mean monthly GFED4 burnt area per gridcell as a function of seasonal mean GPCP rainfall per gridcell. The edges of the whiskers refer to the 5th and 95th percentiles, respectively ( $\pm 1.96 \times SE_{\text{mean}}$ ).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close


Full Screen / Esc

Printer-friendly Version


Interactive Discussion



a) southern region of Kalimantan



b) central region of Kalimantan



## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Figure 3. (a)** the southern region of Kalimantan: seasonal mean RSS burnt area per gridcell as a function of seasonal mean GPCP rainfall per gridcell; and **(b)** the central region of Kalimantan: seasonal mean RSS burnt area per gridcell as a function of seasonal mean GPCP rainfall per gridcell. The edges of the whiskers refer to the 5th and 95th percentiles, respectively ( $\pm 1.96 \times SE_{\text{mean}}$ ).

# NHESSD

2, 5079–5111, 2014

## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion




## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.

a) southern region of Kalimantan



b) southern region of Kalimantan



**Figure 4.** The southern region of Kalimantan: **(a)** seasonal mean GFED4 burnt area per gridcell as a function of seasonal mean System 4 rainfall per gridcell; and **(b)** seasonal mean RSS burnt area per gridcell as a function of seasonal mean System 4 mean rainfall per gridcell. Number of System 4 gridcells in the southern region of Kalimantan = 23. The edges of the burnt area whiskers refer to the 5th and 95th percentiles, respectively ( $\pm 1.96 \times SE_{\text{mean}}$ ). The upper edge of the System 4 whiskers =  $\text{mean}_{\text{upper tercile}} - \text{mean}_{\text{middle tercile}}$ , and the lower edge of the System 4 whiskers =  $\text{mean}_{\text{middle tercile}} - \text{mean}_{\text{lower tercile}}$  (refer data preparation in Sect. 3).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Seasonal forecasting of fire over Kalimantan, Indonesia

A. C. Spessa et al.




**Figure 5.** The southern region of Kalimantan: seasonal mean GPCP rainfall per gridcell versus seasonal mean System 4 rainfall per gridcell. The edges of the GPCP whiskers refer to the 5th and 95th percentiles, respectively ( $\pm 1.96 \times SE_{\text{mean}}$ ). The upper edge of the System 4 data point whiskers =  $\text{mean}_{\text{upper tercile}} - \text{mean}_{\text{middle tercile}}$ , and the lower edge of the System 4 whiskers =  $\text{mean}_{\text{middle tercile}} - \text{mean}_{\text{lower tercile}}$  (refer data preparation in Sect. 3).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




**Seasonal forecasting  
of fire over  
Kalimantan,  
Indonesia**

A. C. Spessa et al.



**Figure 6.** System 4 SST anomalies versus observed SST anomalies over the Indian ocean ( $0\text{--}10^\circ\text{ S}$ ,  $90\text{--}110^\circ\text{ E}$ ), 1997–2010. In this figure the variance has been corrected to match the variance of the observations. The observed anomalies are derived from ERA Interim and the forecasts shown are based on S4 hindcast system with 15 ensemble members.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)