



**Group
decision-making
approach for flood
vulnerability
identification**

G. Lee et al.

Group decision-making approach for flood vulnerability identification using the fuzzy VIKOR method

G. Lee¹, K. S. Jun¹, and E. S. Cung²

¹Department of Water Resources, Graduate School of Water Resources,
Sungkyunkwan University, Suwon, South Korea

²Department of Civil Engineering, Seoul National University of Science and Technology,
Seoul, Suwon, South Korea

Received: 1 July 2014 – Accepted: 28 August 2014 – Published: 30 September 2014

Correspondence to: E.-S. Cung (eschung@seoultech.ac.kr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

This study proposes an improved group decision making (GDM) framework that combines VIKOR method with fuzzified data to quantify the spatial flood vulnerability including multi-criteria evaluation indicators. In general, GDM method is an effective tool for formulating a compromise solution that involves various decision makers since various stakeholders may have different perspectives on their flood risk/vulnerability management responses. The GDM approach is designed to achieve consensus building that reflects the viewpoints of each participant. The fuzzy VIKOR method was developed to solve multi-criteria decision making (MCDM) problems with conflicting and noncommensurable criteria. This comprising method can be used to obtain a nearly ideal solution according to all established criteria. Triangular fuzzy numbers are used to consider the uncertainty of weights and the crisp data of proxy variables. This approach can effectively propose some compromising decisions by combining the GDM method and fuzzy VIKOR method. The spatial flood vulnerability of the south Han River using the GDM approach combined with the fuzzy VIKOR method was compared with the results from general MCDM methods, such as the fuzzy TOPSIS and classical GDM methods, such as those developed by Borda, Condorcet, and Copeland. The evaluated priorities were significantly dependent on the employed decision-making method. The proposed fuzzy GDM approach can reduce the uncertainty in the data confidence and weight derivation techniques. Thus, the combination of the GDM approach with the fuzzy VIKOR method can provide robust prioritization because it actively reflects the opinions of various groups and considers uncertainty in the input data.

Group
decision-making
approach for flood
vulnerability
identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Floods can cause major damages to social infrastructure, human lives, and property. They can harm both individuals and communities and produce negative social, economic, and hydrologic consequences. Flood risk management should consider complex circumstances in conjunction with stakeholders. Many studies have indicated that stakeholder involvement is crucial to vulnerability assessments. According to the Danube Flood Risk Project (2014), individual and institutional knowledge and the expertise of stakeholders are decisive factors in flood damage adaptation and risk assessment. However, the majority of stakeholders, including policymakers, communities, NGOs, engineers, and civil servants, have differing opinions, which are shaped by their personal experiences and areas of interest, such as social, economic, political, engineering, and environmental issues. Thus, flood vulnerability can be effectively identified if multi-criteria decision-making (MCDM) techniques are employed to aggregate various opinions and factors. MCDM can incorporate qualitative as well as quantitative data, which may be conflicting and uncertain, based on the aggregated opinions of all stakeholders.

However, flood vulnerability identification has become complicated by the increase in the number of decision criteria and involved groups, that is, flood vulnerability assessment is inevitably associated with a noticeable degree of uncertainty (Jun et al., 2013; Lee et al., 2013) that can be categorized as follows: (1) uncertainty due to the vagueness of data and (2) uncertainty caused by the subjective viewpoint of the decision maker. The first type of uncertainty is related to the precise data measurement and composition for the spatial and temporal values of social, economic, and hydrologic consequences, such as population, property value, and flood inundation area. The second type of uncertainty is caused by the differences among the personal characteristics of each decision maker. The degree of uncertainty may increase if the decision maker cannot impartially assess the decision process due to limited available knowledge or biased interest. Therefore, the main challenges in flood vulnerability

**Group
decision-making
approach for flood
vulnerability
identification**

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

identification are reducing the uncertainty of the both the crisp data of proxy variables and weights from different decision maker.

For the first uncertainty, some studies have combined MCDM methods with fuzzy set theory to address various uncertainties because fuzzy MCDM methods reduce the uncertainty of parametric approaches, which is inherent to weight determination and the derivation of crisp input data. These methods have been employed in various fields, such as groundwater vulnerability (Zhou et al., 1999), tourism (Tsaur, 2002), plant location selection (Chu, 2002), robot selection (Chu and Lin, 2003), water resources management (Simonovic and Nirupama, 2005), group decision making (Shih et al., 2007), reservoir operation (Fu, 2008; Afshar et al., 2011), knowledge management (Chang and Wang, 2009), the airline industry (Torlak et al., 2011), and environmental watershed planning (Chen et al., 2011), flood vulnerability (Lee et al., 2014). This study also utilized triangular fuzzy numbers (TFNs) to determine the criteria and their corresponding weights. Fuzzy numbers effectively reduce the first type of uncertainty.

The second type of uncertainty is caused by disagreements on the issue. In the case of decision making for flood management, stakeholders have different perceptions about flood hazards, which reflect their different socio-economic and socio-psychological backgrounds. Mutual frictions between the stakeholders, which are based on different goals and objectives, should be recognized and considered to determine the most preferred alternative using the preferences of individual decision makers. The group decision-making (GDM) method coupled with MCDM has also been proven to be an effective tool that enables a converging and compromising opinion for decision makers (Speller, 2005; Kim and Chung, 2013a). The GDM method is an approach in which an entire group of people can participate in consensus building. The method enables stakeholders to make better decisions using various information and knowledge of the decision makers. It has the advantage of sharing the responsibility for the consequences. However, many difficulties arise in the transition from a single-decision-maker process to a multiple-decision-maker-driven process. The latter becomes more complicated due to an increase in the number of

Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2 Methodology

2.1 Study procedure

This study consisted of four steps, as shown in Fig. 1. The first step is to identify the criteria of flood vulnerability and their weight sets using the Delphi survey because floods have a significant influence on society. In this study, flood vulnerability criteria were constructed using a hierarchy tree that consists of social, economic, and hydrologic components, which indicate the causal relationship between society and flood hazards. The survey advances through the modified three-round Delphi technique.

In the second step, the fuzzified data matrix was formulated using data collection and processing. The collected data from domestic public institutions served as the selected factor data. For example, population data on the number of households and asset values were obtained from the National Statistical Office, and rainfall data were obtained from the Korea Meteorological Administration. The majority of the remaining data were collected from the National Water Information System. The probabilities of the flood levels of river sections were calculated using the frequency analysis model developed by Jun and Kim (2011). The collected data on the criteria were fuzzified and standardized. Step 3 is to quantify flood vulnerability using each weight set from each decision-making group. The sites for each participant group were prioritized individually using data collection and processing in Step 3. In Step 4, the flood vulnerability priorities were derived using the VIKOR method.

2.2 Fuzzy VIKOR method

The fuzzy VIKOR method was developed by Opricovic (2011) to solve problems in an uncertain environment, in which both the criteria and weights may comprise fuzzy sets. To construct a formal fuzzy decision matrix, a typical fuzzy MCDM problem can

be expressed in the matrix form as

$$\tilde{D} = \begin{matrix} & C_1 & C_2 & \cdots & C_3 \\ \begin{matrix} A_1 \\ A_2 \\ \vdots \\ A_3 \end{matrix} & \begin{bmatrix} \tilde{x}_{11} & \tilde{x}_{12} & \cdots & \tilde{x}_{1n} \\ \tilde{x}_{21} & \tilde{x}_{22} & \cdots & \tilde{x}_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ \tilde{x}_{m1} & \tilde{x}_{m2} & \cdots & \tilde{x}_{mn} \end{bmatrix} \end{matrix}, \quad i = 1, 2, \dots, m; j = 1, 2, \dots, n \quad (1)$$

$$\tilde{W} = [\tilde{w}_1, \tilde{w}_2, \dots, \tilde{w}_n], \quad j = 1, 2, \dots, n$$

where \tilde{x}_{ij} is the rating of alternative A_i with respect to C_j , \tilde{w}_j is the importance weight of j assigned by the criterion, and \tilde{x}_{ij} and \tilde{w}_j are linguistic variables denoted by the TFNs.

The fuzzy VIKOR method includes the following steps:

1. Determine the fuzzy best value (FBV, \tilde{f}_j^*) and the fuzzy worst value (FWV, \tilde{f}_j^-) of all criterion functions

$$\tilde{f}_j^* = \tilde{x}_{ij}, j \in B; \quad \tilde{f}_j^- = \tilde{x}_{ij}, j \in C. \quad (2)$$

2. Compute the values \tilde{S}_i and \tilde{R}_i

$$\tilde{S}_i = \sum_{j=1}^n \tilde{w}_j (\tilde{f}_j^* - \tilde{x}_{ij}) / (\tilde{f}_j^* - \tilde{f}_j^-) \quad (3)$$

$$\tilde{R}_i = \tilde{w}_j (\tilde{f}_j^* - \tilde{x}_{ij}) / (\tilde{f}_j^* - \tilde{f}_j^-) \quad (4)$$

where \tilde{S}_i refers to the separation measure of A_i from the best fuzzy value and \tilde{R}_i is the separation measure of A_i from the worst fuzzy value.

3. Calculate the values \tilde{S}^* , \tilde{S}^- , \tilde{R}^* , \tilde{R}^- , and \tilde{Q}_i

$$\tilde{S}^* = \min_i \tilde{S}_i, \tilde{S}^- = \max_i \tilde{S}_i \quad (5)$$

$$\tilde{R}^* = \min_i \tilde{R}_i, \tilde{R}^- = \max_i \tilde{R}_i \quad (6)$$

$$\tilde{Q}_i = \frac{\nu (\tilde{S}_i - \tilde{S}^*)}{\tilde{S}^- - \tilde{S}^*} - \frac{(1 - \nu) (\tilde{R}_i - \tilde{R}^*)}{\tilde{R}^- - \tilde{R}^*} \quad (7)$$

where the index \tilde{S}_i is the maximum majority rule, the index \tilde{R}_i is the minimum individual regret of an opponent strategy, and ν is the weight of the strategy of the maximum group utility, with $\nu = 0.5$ being a typical situation.

4. Defuzzify TFN $\tilde{Q}_i = (Q_i^l, Q_i^m, Q_i^u)$ and rank the alternatives (\tilde{N}) by the index Q_i

$$\text{Crisp}(\tilde{N}) = (Q_i^l + 2Q_i^m + Q_i^u)/4.$$

3 Study area and decision-making groups

3.1 The south Han River

The south Han River is an upper tributary of the Han River, which is located in the middle of the Korean peninsula; it includes Yeosu, which is the satellite city of Seoul. The basin area and river length are approximately 5400 km² and 95.4 km, respectively; they extend from the Chungju regulating dam to the Paldang dam. The area includes six administrative districts: Gwangju, Yangpyeong, Yeosu, Icheon, Chungju, and Eumseong. Various development plans have been devised due to rapid population growth. However, the basin has experienced frequently repeated flood damages. During the last decade, the flood damage in the study area was estimated

to be more than USD 150 million; the average annual flood damage in Gwangju is USD 13 million.

This study employed a reach-based areal approach to include probabilistic flood risk values. In this study, the river reach was divided into constant lengths, and its corresponding watershed was developed based on its DEM, as shown in Fig. 2.

3.2 Participant decision-making groups

An extensive range of relevant stakeholders must get involved in sustainable flood management. For example, the government has the prime responsibility of ensuring the safety and well-being of its citizens. Therefore, all relevant departments at the national, state, district, and municipal levels are closely related to flood management strategies. Scientific institutions, universities, communities, NGOs, and engineering consultants can provide useful information for compromising solutions. Each of these categories also exhibits a significant diversity of opinions (Morss et al., 2005).

In South Korea, the Ministry of Land, Infrastructure, and Transport (MLIT), which plans and manages the country's major rivers, are responsible for the flood management of large river basins. MLIT operates a dedicated flood control office for the four large basins. Although no direct management body for flood disaster management exists, the Ministry of Security and Public Administration, which is responsible for the country's total safety management, and the Ministry of Environment, which is responsible for environment management, are associated with the management of river basins and environmental pollution caused by floods. Several research institutes are also concerned with water resources, including the K-water Research Institute, which is affiliated with the publicly owned water resource corporation, and government-funded research institutes, such as the Institute of Construction Technology, the National Disaster Management Institute, and the National Institute of Environmental Research. In addition, professional experts who specialize in water resource management and are affiliated with universities, engineering companies, and private research institutes, also address problems associated with floods.

**Group
decision-making
approach for flood
vulnerability
identification**

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Therefore, the opinions in this study were collected by conducting a questionnaire survey of people from these various institutions and organizations. Because it is difficult to obtain a consensus building for a weight set among various stakeholders and experts, this study assumed that each participator represents the group with which they are affiliated. That is, each interviewee assumed to represent each individual organization associated with important flood-related business organizations was assumed to be the opinion of the entire group.

We conducted focused interviews with 44 experts who were either senior managers or experienced group facilitators. The participator had experience in flood risk management. The majority (92 %) of the participants possessed master's degree-level or doctoral-level training. They worked in the following sectors: government-funded research institutes (34 %), private engineering companies (23 %), research centers (16 %), and universities (27 %). The age distribution of the respondents ranged from 20 to 60 years, with the majority (84 %) of participants ranging between 30 and 49 years of age.

4 Application

4.1 Identification of the criteria and their weights (Step 1)

Since identifying the appropriate criteria is very crucial in vulnerability study, this study approached the systematic hierarchy structure for sustainability. The criteria were divided into three characteristic groups: social, economic, and hydrologic components. The intent was to provide supporting results for appropriate policies by forming groups of similar properties.

The research staff established a draft criteria list based on the three components and distributed it to the decision-making groups during the first-round survey of the Delphi procedure. For the flood vulnerability assessment, the participants completed the survey in which they expressed their opinions regarding the importance of factors. Their

Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



responses were analyzed to develop 24 criteria, as shown in Table 1. The difference of opinions among the participating decision-making groups was not significant, which reveals similar opinions about the criteria that need to be reflected in the calculation of flood vulnerability. Detailed opinions about the necessity of factors were collected in the second Delphi round. These opinions immediately indicated the importance of factor evaluation for assigning the weight value. Each group determined the value of the weights for the importance of the criteria using linguistic variables, which can be expressed by a membership function of fuzzy sets. Table 2 presents the linguistic variables and TFNs for the weighting. Seven linguistic variables – very low, low, medium low, medium, medium high, high, and very high – were introduced using the fuzzy function. The research staff subsequently analyzed and distributed the summary of the third Delphi round, and the participants reconsidered the weights. As a result, relatively compromising weights were derived. This Delphi process was described in the previous study (Lee et al., 2013).

The Delphi process is known for its ability to effectively collect the opinions of decision makers; however, decision-making groups have a subjective tendency to assess importance based on the analysis of the collected weights. The results of the selected weight may exhibit significant differences due to the tendencies of the decision-making groups. For example, the histogram of the weight for a particular decision-making group is shown in Fig. 3. G6 determined that the weights only use two variables, MH and H (Fig. 3b), whereas G3 (Fig. 3a) presented a diversity of opinions. In Figs. 3c and 4d, G8 used M for the lowest value and weighted the high level as MH, H, and VH. Conversely, G12 selected primarily low-level values as VL, L, and ML. G30 typically selected middle-level weights, whereas G38 used various weight levels. Both groups used an extensive range of choices (Fig. 3e and f), which suggests that each group has distinct individuality.

Therefore, the collected weight sets of each decision maker were standardized as

$$W'_i = \frac{(W_i - W_{\min})}{(W_{\max} - W_{\min})} \quad (8)$$

where W_{\max} and W_{\min} are the maximum and minimum values of the set of weights ($W_i, i = 1, \dots, n$), respectively.

For example, a comparison of the distribution between the raw and standardized data for the weights for characteristic factors is presented in Fig. 4. Regarding the social factors, ML (medium low) received the most ballots for weights, followed by M (medium) and L (low). However, the majority of ballots were altered to L (low) by standardization.

Regarding the economic factor of the weights, M and ML have many more ballots compared with the remaining weights. The standardizing weights of the economic factor were distributed over an extensive range between L and H (high). The most influential weight value was the diverted L through standardization. VH (very high) received the most votes for hydrologic factors, followed by MH (medium high). Using standardization, the most weight was predominantly distributed as ML. The standardized results of the collected weights for the criteria were mainly distributed between L and ML (Fig. 5).

4.2 Construction of the data matrix (Step 2)

The performance value of all sections for all criteria were collected from reliable institutions, such as Statistics Korea, the Korea Meteorological Administration, the National Emergency Management Agency, the Water Management Information System, the local government, the National Geographic Information Institute, and the Han River flood control office. We assumed that if a province exhibits a value or extremely low variability, one value is used. The collected data were spatially analyzed, and the collected proxies were fuzzified using the TFN concept. Each TFN was statistically derived from the probability density function (PDF) using R-statistics software, as shown in Fig. 6.

**Group
decision-making
approach for flood
vulnerability
identification**

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.3 Vulnerability identification for individual decision group (Step 3)

In this step, the flood vulnerability for each decision group is evaluated using Eq. (9), and a performance matrix is constructed, as shown in Eq. (10). In this assessment, we employed the same sources of criteria to identify the TFN

$$\tilde{V} = \sum_{j=1}^m \tilde{w}_j \times \tilde{c}_j \quad (9)$$

$$\begin{matrix} & \text{Value} & & \text{Criteria value} & & \text{Criteria value} \\ & & & \mathbf{A}_1 \quad \dots \quad \mathbf{A}_k & & \mathbf{G}_1 \quad \dots \quad \mathbf{G}_m \\ \mathbf{V}_1 & \begin{bmatrix} \tilde{V}_{11} & \dots & \tilde{V}_{m1} \end{bmatrix} & = & \mathbf{C}_1 & \begin{bmatrix} \tilde{c}_{11} & \dots & \tilde{c}_{n1} \end{bmatrix} & \begin{bmatrix} \tilde{w}_{11} & \dots & \tilde{w}_{1m} \end{bmatrix} \\ \vdots & \begin{bmatrix} \vdots & \ddots & \vdots \end{bmatrix} & & \vdots & \begin{bmatrix} \vdots & \ddots & \vdots \end{bmatrix} & \begin{bmatrix} \vdots & \ddots & \vdots \end{bmatrix} \\ \mathbf{V}_k & \begin{bmatrix} \tilde{V}_{1k} & \dots & \tilde{V}_{mk} \end{bmatrix} & & \mathbf{C}_n & \begin{bmatrix} \tilde{c}_{1k} & \dots & \tilde{c}_{nk} \end{bmatrix} & \begin{bmatrix} \tilde{w}_{n1} & \dots & \tilde{w}_{nm} \end{bmatrix} \end{matrix} \quad (10)$$

where \tilde{w} denotes the weight sets of each group and \tilde{c} denotes the values of the criteria per unit area. They are expressed as TFNs (x_1, x_2, x_3). The values of \tilde{V} are calculated for each group by the WSM. For example, the results of area R4 are shown in Fig. 7. The figure revealed a substantial difference among the opinions of each group. Some groups (G2, G10, G16, G18, G27, G36, G39, and G41) exhibited a large range of values. Conversely, the results of other groups (G4, G7, G13, G21, G22, G26, G31, G40, and G42) were confined within a narrow range. Groups G5, G12, and G19 had a high upper bound (approximately 1) compared to the other groups, whereas G2 had an uncharacteristic lower bound on 0.

4.4 Determination of the priority ranking using the fuzzy VIKOR method (Step 4)

This step was intended to apply the fuzzy VIKOR GDM method to the preceding results of each group. Because the evaluated results are represented by TFNs, the priority was determined by a defuzzification value using Eq. (11). Defuzzification is another process that is used to transform a TFN into a crisp number. We used the following simple defuzzification method by Bojadziev and Bojadziev (1997):

G_A-tilde(x-tilde) = (x_1 + 2x_2 + x_3) / 4. (11)

S, R, and Q were calculated to derive the preference ranking of the alternatives, as shown in Table 3. In this study, the priority ranks of the alternatives were determined by the Q value to comprise two different types of regrets, i.e. S and R values. Based on the results of the fuzzy VIKOR GDM method, R11, R20, and R19 in the right bank and L18, L11, L20, and L7 in the left bank were adopted to the vulnerable area. S, R, and Q had similar ranking patterns for the alternatives, whereas R8, L6, and L10 exhibited significantly different results for S, R, and Q.

For comparison, the rank of the alternatives was estimated by the fuzzy TOPSIS method and fuzzy VIKOR method using the average weight value of the participating groups. Figure 8 presents the assessing ranks. L18 of the left bank was the most vulnerable area because it yielded the highest values of the three approaches, whereas the most vulnerable area of the right bank differed among the approaches. R11, which is the most vulnerable area based on the fuzzy VIKOR GDM method, ranked 5th for the fuzzy TOPSIS method and 9th for the fuzzy VIKOR method. In the case of the test area, the results of the fuzzy TOPSIS method were analogous to our GDM approach; however, the fuzzy VIKOR method yielded different results.

The deduced priorities shown in Table 4 facilitate the comparison of the evaluation results of the Borda, Condorcet, and Copeland methods, which are representative

Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



GDM methods that have been extensively used, with the fuzzy VIKOR GDM method proposed in this study. The evaluation rating results of the Borda, Condorcet, and Copeland methods were similar but differed significantly from the evaluation rating results of the fuzzy VIKOR GDM method.

In the case of the right bank, area R8 was ranked as the 5th most vulnerable area using the fuzzy VIKOR GDM method; however, it is ranked first for the remaining methods. R11 ranked first for the fuzzy VIKOR GDM method and 8th with the remaining methods. However, L7 and L11, which exhibited high vulnerability, ranked high for all methods, whereas L8 and L11, which exhibited low vulnerability, had significantly different rankings. Spearman's correlation coefficients were calculated to analyze the correlations among these evaluation methods, as shown in Table 5. The estimated coefficients exceeded approximately 75 %, demonstrating that the proposed fuzzy VIKOR GDM method (bold numbers in Table 5) is generally correlated with the other MCDM methods.

5 Conclusions

This study used the fuzzy VIKOR GDM method to spatially quantify flood vulnerability. The motivation behind this approach is the need for a method that considers the differences of opinions among groups, which becomes evident when collecting information about the weighted value of criteria in a survey. We applied the fuzzy VIKOR method to overcome the limitations of the classical vote-based GDM methods. We compared the evaluated rankings from the fuzzy VIKOR GDM method with those from fuzzy TOPSIS and fuzzy VIKOR. The classical GDM ranks were derived by averaging all ranks from the Borda, Condorcet, and Copeland methods.

In the case of the left bank, the GDM approach with the fuzzy VIKOR method typically yielded results similar to those obtained with the existing methods for the areas ranked as highly vulnerable, whereas significantly different results were obtained for the areas that are ranked low in vulnerability. Conversely, in the case of the right bank, all

**Group
decision-making
approach for flood
vulnerability
identification**

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

evaluation methods yielded different results for the high-vulnerability areas, whereas similar results were obtained for the low-vulnerability areas. These evaluation results were obtained because the collected criteria data respond sensitively to the weighted values, as a minimal difference exists between the areas with low numbers for the left bank and a minimal difference exists between the areas with high numbers for the right bank. The evaluated priorities can change significantly depending on the applied decision-making method.

Thus, the GDM approach with the fuzzy VIKOR method can provide useful results for MCDM because it is an evaluation method that actively reflects the opinions of various groups and considers uncertainty in input data.

Acknowledgements. This study was financially supported by the Basic Science Research Program of the National Research Foundation of Korea (NRF), which is funded by the Ministry of Education (NRF-2013R1A1A2060942).

References

- Afshar, A., Mariño, M. A., Saadatpour, M., and Afsahr, A.: Fuzzy TOPSIS multi-criteria decision analysis applied to Karun reservoirs system, *Water. Resour. Manage.*, 25, 545–563, 2011.
- Akter, T. and Simonovic, S. P.: Aggregation of fuzzy views of a large number of stakeholders for multi-objective flood management decision-making, *J. Environ. Manage.*, 77, 133–143, 2005.
- Bojadziev, G. and Bojadziev, M.: *Fuzzy Logic for Business, Finance, and Management*, World Scientific Publishing Co Pte Ltd, Singapore, 1997.
- Chang, T. H. and Wang, T. C.: Using the fuzzy multi-criteria decision making approach for measuring the possibility of successful knowledge management, *Inform. Sciences*, 179, 355–370, 2009.
- Chen, C. T.: Extension of the TOPSIS for group decision-making under fuzzy environment, *Fuzzy Set. Syst.*, 114, 1–9, 2000.
- Chen, V. Y. C., Lien, H.-P., Liu, C.-H., Liou, J. J. H., Tzeng, G.-H., and Yang, L.-S.: Fuzzy MCDM approach for selecting the best environment-watershed plan, *Appl. Soft Comput.*, 11, 265–275, 2011.

Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Chu, T. C.: Selecting plant location via a fuzzy TOPSIS approach, *Int. J. Adv. Manuf. Tech.*, 20, 859–864, 2002.
- Chu T. C. and Lin Y. C.: A fuzzy TOPSIS method for robot selection, *Int. J. Adv. Manuf. Technol.*, 21, 284–290, 2003.
- 5 Copeland, A. H.: A Reasonable Social Welfare Function. Seminar on Applications of Mathematics to Social Sciences, University of Michigan, Ann Arbor, 1951.
- Danube Flood Risk Project: Common stakeholders' involvement strategy for the Danube floodplain, available at: <http://www.danube-floodrisk.eu/> (last access: 16 September 2014), 2014
- 10 Fu, G.: A fuzzy optimization method for multi-criteria decision making: an application to reservoir flood control operation, *Expert Syst. Appl.*, 34, 145–149, 2008.
- Jun, K. S., Chung, E. S., Kim, Y. G., and Kim, Y.: A fuzzy multi-criteria decision approach to flood risk vulnerability in South Korea by considering climate change impacts, *Expert Syst. Appl.*, 40, 1003–1013, 2013.
- 15 Kim, Y. and Chung, E. S.: Assessing climate change vulnerability with group multi-criteria decision making approaches, *Climatic Change*, 121, 301–315, 2013a.
- Kim, Y. and Chung, E. S.: Fuzzy VIKOR approach for assessing the vulnerability of the water supply to climate change and variability in South Korea, *Appl. Math. Model.*, 37, 9419–9430, 2013b.
- 20 Kim, Y., Chung, E. S., Jun, S. M., and Kim, S. U.: Assessing climate change vulnerability with group multi-criteria decision making approaches, *Resour. Conserv. Recy.*, 73, 23–32, 2013.
- Lee, G., Jun, K. S., and Chung, E. S.: Integrated multi-criteria flood vulnerability approach using fuzzy TOPSIS and Delphi technique, *Nat. Hazards Earth Syst. Sci.*, 13, 1293–1312, doi:10.5194/nhess-13-1293-2013, 2013.
- 25 Lee, G., Jun, K. S., and Chung, E. S.: Robust spatial flood vulnerability assessment for Han River using fuzzy TOPSIS with a-cut level set, *Expert Syst. Appl.*, 41, 644–654, 2014.
- McLean, I.: The borda and condorcet principles: three medieval applications, *Soc. Choice. Welfare.*, 7, 99–108, 1990.
- Morss, R. E., Wilhelmi, O. V., Dwonton, M. W., and Grunfest, E.: flood risk, uncertainty, and scientific information for decision making: lessons from an interdisciplinary project, *B. Am. Meteorol. Soc.*, 86, 1593–1601, 2005.
- 30 Opricovic, S.: Multicriteria optimization of civil engineering systems (in Serbian, Visekriterijumska optimizacija sistema u gradjevinarstvu), Faculty of Civil Engineering, Belgrade, 1998.

**Group
decision-making
approach for flood
vulnerability
identification**

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Opricovic, S.: Fuzzy VIKOR with an application to water resources planning, *Expert Syst. Appl.*, 38, 12983–12990, 2011.
- Opricovic, S. and Tzeng, G. H.: Extended VIKOR method in comparison with outranking methods, *Eur. J. Oper. Res.*, 178, 514–529, 2007.
- 5 Sanayei, A., Farid Mousavi, S., and Yazdankhah, A.: Group decision making process for supplier selection with VIKOR under fuzzy environment, *Expert Syst. Appl.*, 37, 24–30, 2010.
- Shih, H. S., Shyur, H. J., and Lee, E. S.: An extension of TOPSIS for group decision making, *Math. Comput. Model.*, 45, 801–813, 2007.
- Simonovic, S. P. and Niruoama, N.: A spatial multi-objective decision-making under uncertainty for water resources management, *J. Hydroinform.*, 7, 117–133, 2005.
- 10 Speller, G.: Improving community and citizen engagement in FRM decision making, delivery and flood response, R & D Technical Report SC040033/SR3, product code SCHO1005BJTC-E-P, Environment Agency, Bristol, 2005.
- Torlak, G., Sevkli, M., Sanal, M., and Zaim, S.: Analyzing business competition by using fuzzy TOPSIS method: an example of Turkish domestic airline industry, *Expert Syst. Appl.*, 38, 3396–3406, 2011.
- 15 Tsaur, S., Chang T., and Yen C.: The evaluation of airline service quality by fuzzy MCDM, *Tourism. Manage.*, 23, 107–115, 2002.
- William, G. L.: Strategic voting and the borda method, *Public Choice*, 33, 85–90, 1978.
- 20 Yazdani-Chamzini, A. and Yakhchali, S. H.: Tunnel Boring Machine (TBM) selection using fuzzy multicriteria decision making methods, *Tunn. Undergr. Sp. Tech.*, 30, 194–204, 2012.
- Zhou, H. C., Wang, G. L., and Yang, Q.: A multi-objective fuzzy pattern recognition model for assessing groundwater vulnerability based on the DRASTIC system, *Hydrolog. Sci. J.*, 44, 611–618, 1999.

Table 1. Criteria for flood vulnerability asesement.

No.			Criteria (measure)
1	Social	Pressure	Population growth ratio (%)
2			Population (number)
3		State	Residential and industrial area ratio (%),
4			Population density (man/1 km ²)
5			Number of social overhead capital, cultural properties and natural monuments (number)
6		Impact	Annual casualties and sufferers due to floods and disasters (number/year)
7		Response	Number of flood and disaster prevention institutions (number/year)
8			Number of government officials for flood and disaster mitigation (number)
9	Economic	Pressure	Gross regional domestic product(KRW)
10		State	Urban area ratio (%)
11			Self-reliance ratio of finance (%)
12			Property value(KRW)
13		Impact	Annual flood damage (number/year)
14		Response	Annual recovery and preparation costs for floods and disasters (KRW/year)
15	Hydrologic	Pressure	Increased ratio of daily maximum precipitation (%)
16			Increased ratio of 1 h rainfall intensity (%)
17			Increased ratio of summer rainfall (%)
18			Watershed slope (deg)
19		State	Peak flow of the 200 yr floods
20			River stage of the 200 yr floods
21		Impact	Annual number of floods (number/year)
22		Response	Flood inundation area (km ²)
23			Number of flood mitigation infrastructures (number)
24			River improvement ratio (%)

**Group
decision-making
approach for flood
vulnerability
identification**

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I ◀

▶ I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Linguistic variables and fuzzy membership function for the importance of weight.

Importance	Membership function
Very low (VL)	(0.0, 0.0, 0.1)
Low (L)	(0.0, 0.1, 0.3)
Medium low (ML)	(0.1, 0.3, 0.5)
Medium (M)	(0.3, 0.5, 0.7)
Medium high (MH)	(0.5, 0.7, 0.9)
High (H)	(0.7, 0.9, 1.0)
Very high (VH)	(0.9, 1.0, 1.0)

Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 3.** Evaluation results.

Alternative	<i>S</i>	<i>R</i>	<i>Q</i>	Alternative	<i>S</i>	<i>R</i>	<i>Q</i>
R1	15	12	14	L1	7	5	6
R2	13	13	12	L2	17	18	18
R3	9	1	8	L3	9	10	10
R4	11	10	10	L4	12	15	16
R5	9	9	9	L5	19	19	19
R6	16	18	18	L6	6	13	11
R7	8	11	11	L7	3	4	4
R8	5	1	5	L8	18	20	20
R9	14	17	16	L9	14	14	14
R10	17	16	17	L10	20	11	13
R11	1	1	1	L11	2	3	2
R12	20	20	20	L12	16	17	17
R13	19	19	19	L13	13	16	15
R14	7	7	7	L14	8	8	8
R15	12	14	13	L15	15	12	12
R16	18	15	15	L16	11	9	9
R17	4	1	4	L17	10	7	7
R18	6	8	6	L18	1	1	1
R19	3	1	3	L19	5	6	5
R20	2	1	2	L20	4	2	3

Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 4.** Comparison of evaluated ranks by GDM approaches.

Alternative	Borda	Condorcet	Copeland	Group fuzzy VIKOR	Alternative	Borda	Condorcet	Copeland	Group fuzzy VIKOR
R1	17	15	17	14	L1	13	11	13	6
R2	14	13	14	12	L2	18	16	18	18
R3	10	10	10	8	L3	12	12	12	10
R4	13	14	13	10	L4	11	13	11	16
R5	9	9	9	9	L5	15	16	15	19
R6	15	17	15	18	L6	6	6	6	11
R7	5	1	5	11	L7	2	1	2	4
R8	1	1	1	5	L8	14	14	14	20
R9	11	11	11	16	L9	10	9	10	14
R10	16	17	16	17	L10	20	19	20	13
R11	8	8	8	1	L11	3	1	3	2
R12	20	19	20	20	L12	17	16	17	17
R13	19	19	19	19	L13	16	15	16	15
R14	7	7	7	7	L14	8	9	8	8
R15	12	11	12	13	L15	19	19	19	12
R16	18	16	18	15	L16	9	8	9	9
R17	3	1	3	4	L17	7	7	7	7
R18	6	6	6	6	L18	1	1	1	1
R19	4	1	4	3	L19	5	5	5	5
R20	2	1	2	2	L20	4	1	4	3

Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 5. Determined Spearman's correlation coefficients among the general MCDM methods and Group fuzzy VIKOR of this study (bold numbers).

Left-side bank						
	Borda	Condorcet	Copeland	Group fuzzy VIKOR	Fuzzy TOPSIS	Fuzzy VIKOR
Borda	–	97.3 %	100.0 %	77.0 %	83.0 %	54.0 %
Condorcet	–	–	97.3 %	80.1 %	85.6 %	53.9 %
Copeland	–	–	–	77.0 %	83.0 %	54.0 %
Group fuzzy VIKOR	–	–	–	–	87.7 %	74.8 %
Fuzzy TOPSIS	–	–	–	–	–	67.5 %
Fuzzy VIKOR	–	–	–	–	–	–
Right-side bank						
	Borda	Condorcet	Copeland	Group fuzzy VIKOR	Fuzzy TOPSIS	Fuzzy VIKOR
Borda	–	96.1 %	100.0 %	85.5 %	90.3 %	80.0 %
Condorcet	–	–	96.1 %	80.6 %	85.4 %	76.6 %
Copeland	–	–	–	85.5 %	90.3 %	80.0 %
Group fuzzy VIKOR	–	–	–	–	96.0 %	85.0 %
Fuzzy TOPSIS	–	–	–	–	–	88.2 %
Fuzzy VIKOR	–	–	–	–	–	–




Figure 1. Study procedure for group decision making.

**Group
decision-making
approach for flood
vulnerability
identification**

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Figure 2.** Study area and watershed section.

**Group
decision-making
approach for flood
vulnerability
identification**

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 3. Analysis of linguistic weight of each decision making group.

Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 4. Comparison of original and standardized data.

Group decision-making approach for flood vulnerability identification

G. Lee et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Figure 5. Standardized results of collected weights for criteria.

**Group
decision-making
approach for flood
vulnerability
identification**

G. Lee et al.

**Figure 6.** Examples for probability density function using R-statistics software.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Group decision-making approach for flood vulnerability identification

G. Lee et al.




Figure 7. Determined value for R4.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Group decision-making approach for flood vulnerability identification

G. Lee et al.




Figure 8. Comparison of assessed ranks by group fuzzy MCDM and typical MCDM approaches.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

