

Linking local wildfire dynamics to pyroCb development

Richard H.D. McRae¹, Jason J. Sharples² and Michael Fromm³.

^[1] Australian Capital Territory Emergency Services Agency, Canberra, Australia.

^[2] University of New South Wales, Canberra, Australia.

^[3] US Naval Research Laboratory, Washington, USA.

Correspondence to: R.H.D.McRae (rick.mcrae@act.gov.au)

Final rework. Marking up is as follows

[AR1]Changes arising from comments from Anonymous Reviewer 1.[/AR1]

[BP]Changes arising from comments from Reviewer 2, Brian Potter.[/BP]

[AR3]Changes arising from comments from Anonymous Reviewer 3.[/AR3]

[A]Changes made by the authors to improve final paper after all reviewers' comments included.[/A]

Deleted text is ~~crossed-out~~.

Additionally:

- Figure 4 has been amended to remove the A-Train overpass, in accord with BP comment.
- We have discussed with the Editor enlarging the sizes of the Figures. This will be done.
- We will discuss with the Editor the spelling of words such as "channelling" and "behaviour".

1 Abstract

2 Extreme wildfires are global phenomena that consistently result in loss of life and property,
3 and further impact the cultural, economic and political stability of communities. In their most
4 [A]severe[/A] form they cause widespread devastation of environmental assets and are
5 capable of impacting the upper troposphere – lower stratosphere through the formation of a
6 thunderstorm within the plume. Such fires are now often observed by a range of remote
7 sensing technologies, which together allow a greater understanding of a fire's complex
8 dynamics.

9 This paper considers one such fire that burnt in the Blue Mountains region of Australia in
10 late-November 2006, that is known to have generated significant pyrocumulonimbus clouds
11 in a series of blow-up events. Observations of this fire are analysed in detail to investigate the
12 localised processes contributing to extreme fire development. In particular, it has been
13 possible to demonstrate for the first time that the most [A]violent[/A] instances of
14 pyroconvection were driven by, and not just associated with, [A]atypical[/A]local fire
15 dynamics, especially the fire channelling phenomenon, which arises due to an interaction
16 between an active fire, local terrain attributes and critical fire weather, and causes the fire to
17 rapidly transition from a frontal to an areal burning pattern. The impacts of local variations in
18 fire weather and of the atmospheric profile are also discussed, and the ability to predict
19 extreme fire development with state-of-the-art tools is explored.

20

1 **1 Introduction**

2 Over the last decade southeastern Australia has experienced a number of extreme wildfire
3 events. These include the January 2003 Alpine Fires (Nairn, 2003), the 2006-2007 Great
4 Divide Fires (Smith, 2007) and the 2009 Victorian Fires (Teague et al. 2010), and also
5 multiple fires in more recent years that are still under review. These fires caused widespread
6 destruction of assets (both natural and anthropogenic) and multiple fatalities. While such fires
7 are undoubtedly tragic, they do offer rare opportunities to better understand the dynamics and
8 impacts of extreme wildfires. Of particular interest in this context is the development of
9 pyroconvective plumes that inject smoke and other wildfire emissions high into the
10 atmosphere.

11 [BP] The development of these vigorous plumes, which manifest as pyrocumulus or
12 pyrocumulonimbus (pyroCb) (American Meteorological Society, 2013), signals a transition in
13 fire development from what is primarily a surface phenomenon to a coupled fire[A]-
14 [/A]atmosphere phenomenon (Potter, 2002; McRae and Sharples, 2011), in which the fire is
15 able to interact with upper levels of the atmosphere (i.e. the mixed layer and above). Fires
16 which make such a transition may access stronger winds and very dry air, which if returned to
17 the surface (e.g. through convective mixing) can lead to [AR3]elevated[/AR3] levels of fire
18 behaviour and rates of spread (Potter et al., 2007; Mills, 2005, 2008). These fires would
19 therefore be difficult to suppress and would be very likely to do extensive damage to any
20 assets they impacted (McRae and Sharples, 2011).[/BP]

21 [BP] The injection height of smoke plumes from wildfires also has direct bearing on their
22 effects on air quality and climate,[/BP] as it is a major factor governing downwind dispersal
23 (Jost et al. 2004; Mazzoni et al. 2007; Kahn et al. 2008). In the most significant instances of
24 pyroconvection, emissions from wildfires can be injected into the lower stratosphere with
25 subsequent global-scale impacts on atmospheric chemistry (Fromm et al. 2004; Trentmann et
26 al. 2006; Rosenfeld et al. 2007). Stratospheric injection of pyrogenic aerosols could also play
27 an important part in the interaction between wildfire occurrence and climatic warming
28 (Fromm et al. 2004; Jost et al. 2004). Large fire plumes offer a direct transport path between
29 the planetary boundary layer and the stratosphere, which is not fully captured by large-scale
30 atmospheric transport models (Jost et al. 2004).

31 [AR1]In their analysis of the 2003 Canberra bushfires, Sharples et al. (2012) demonstrated
32 several cases where interactions between strong winds and rugged topography resulted in

1 rapid wildfire development. The resulting process, which they termed ‘fire channelling’,
2 resulted in a transition from the usual frontal burning pattern to an areal burning pattern
3 caused by a combination of rapid lateral fire propagation (transverse to the prevailing winds)
4 and downwind in-fill by short- to medium-range spotting. For the Flea Creek event (Sharples
5 et al., 2012, p288) they showed a direct link between the lateral spread and the pyroCb
6 development, which Fromm et al. 2006 has shown reached to 15 km a.g.l.[/AR1] [A]A series
7 of large fires merged on a rugged landscape on a day of elevated fire danger, eventually
8 covering over 90,000 ha in three hours. Linescan data showed that deep flaming in numerous
9 locations was primarily caused by the lateral spread associated with fire channelling. There
10 was a close association of this with pyroCb development, but in such a complex setting clear
11 linkages have not yet been demonstrated.[/A]

12 [AR3]PyroCb events such as those near Canberra in 2003 are at the most extreme end of the
13 pyroconvective spectrum.[/AR3] Indeed such high level smoke injection events hardly feature
14 at all in previous studies of fire plumes (e.g. Lavoue, et al. 2000; Val Martin, et al. 2010).
15 Fromm et al. (2010) discuss the global climatology of pyroCb events, and note in particular
16 the small number of published studies. Since 2001 there have been at least 30 instances of
17 significant pyroCb occurrence in Australia either demonstrated or inferred from their injection
18 of aerosols into the upper troposphere/lower stratosphere (Tables A1 and A2). [AR3]While
19 there is the possibility that additional pyroCbs have occurred in the past in association with
20 significant bushfires that predate the technology necessary to formally detect them, this
21 tabulation of pyroCb occurrence is informative both for the satellite and pre-satellite era. The
22 recent frequency of pyroCb occurrence in Australia necessitates on-going monitoring and
23 study.[/AR3]

24 Fromm et al. (2012) analysed pyroCb occurrence associated with two of the fires that
25 occurred in eastern Australia during the spring and summer of 2006/2007, at locations shown
26 in Fig. 1a. [AR3]The so-called Grose Valley and Wollemi fires were notable due to the fact
27 that their behaviour escalated quasi-simultaneously in the late morning, rather than in the late
28 afternoon when this typically occurs (Fromm et al., 2010).[/AR3] Based on remotely sensed
29 data from a range of satellites and weather radar, and surface and profile data from official
30 weather observation sites,

31 [AR1] Based on remotely sensed data from a range of satellites and weather radar, and
32 surface and profile data from official weather observation sites, Fromm et al. (2012)

concluded that the factors of significance for the development of the pyroconvection were significant departures from the norm in boundary layer temperature (as indicated by the elevated value of the Continuous Haines Index) and wind speed.[/AR1] [AR3]They also noted that neither factor alone was deterministic, reflecting a general need to further scrutinise the local drivers of the extreme fire behaviour.[/AR3]

The present paper therefore extends the study of Fromm et al. (2012) by exploiting data recorded for the Grose Valley Fire by an airborne multispectral linescanner (Cook et al. 2009), which provides critical insights into the wildfire dynamics. The combination of time-stamped, spatially-rectified fire data and radar data (vertically aggregated convective intensity and plume top height) permits detailed examination of the fire's development and the localised pyroconvective contributions. Moreover, the combined analyses permit further scrutiny of the role of dynamic fire processes, such as mass spotting and the fire channelling phenomenon, in extreme fire development. In particular, the link between fire channelling and pyroCb development, which hitherto was established in only a single instance and in only a very qualitative way, is examined in greater depth and in more quantitative detail.

2 Data

It should be noted that all times (and thus dates) used in this paper are in UTC. Local time, Australian Eastern Summer Time, is UTC plus 11 hours.

2.1 Satellite data

An initial examination of the areas of intense fire activity was conducted on LANDSAT archival imagery from USGS. One image that is both relatively cloud-free and free of scan-line correction problems showed the post-fire scars. An excerpt of the image is shown in Fig. 1d, and clearly indicates leafless eucalypt canopy left behind by intense fire activity.

~~[BP]As described in Fromm et al. (2012), much of the broad understanding of these fires derives from data acquired by the “A-Train” environment monitoring satellite flotilla.[/BP]~~

2.2 Weather data

Weather data were supplied by the Australian Bureau of Meteorology (BoM). This included automatic weather station (AWS) half-hourly observations and daily vertical profiles from radiosonde flights. Relevant weather data were recorded at Mt Boyce AWS (WMO Station

Number 94743), Nullo Mountain AWS (94754), Penrith Lakes AWS (94763) and Cessnock Airport AWS (95771), and radiosonde data were recorded at Williamtown (94776). The radar data were recorded by the Newcastle radar. The locations of all of the above recording stations are shown in Fig. 1a.

Fig. 2 shows time series plots of key weather measurements and fire indices for these stations. Fire behaviour estimates, using the equations of Noble et al. (1980), are based on the Forest Fire Danger Index (FFDI). As no direct measurements of fuel moisture content (FMC) were available, FMC was described using the dimensionless fuel moisture index (FMI), which has been shown to produce reliable estimates of fuel moisture content in a variety of fine fuel types (Sharples et al. 2009; Sharples and McRae, 2011; Sharples and Matthews, 2011). In particular, Sharples et al. (2009) showed that $FMI \leq 5$ consistently corresponded to eucalypt litter moisture contents below 3-4%, [AR3]a predictor of elevated fire intensity and energy release.[/AR3]

[AR1] Fromm et al. (2012, especially Figure 15c) discussed the synoptic situation, dominated by the passage, from the west, of a pressure trough linked to a cold front further south.[/AR1]

2.3 Radar data

The reflectivity data forming a full volume scan are available at 10-minute resolution over 22 November. Computations from scan azimuth, elevations and range permit estimation of the maximum height of radar returns (the “echotop”) (Fromm et al. 2012). [BP]We also employ vertical aggregated reflectivity, an indication of overall convective activity. The vertical aggregate of reflectivity is a simple summation of individual reflectivity values in the column at each reported range/azimuth location.[/BP] Note that echoes below the nominal cloud-base altitude (taken to be 4 km, based on the radiosonde profile at Williamtown) are attributable to non-hydrometeors such as debris lofted by the active fire (Lindley et al. 2011). Above the cloud-base stronger echoes are returned by hydrometeors, although these tend to be anomalously small (Rosenfeld, et al. 2007). [A]In Fig. 6 we use a rainrate product, which is derived from reflectivity.[/A]

2.4 Multispectral linescan data

The New South Wales Rural Fire Service aircraft used the Daedalus 1268 ATM multispectral linescanner supplied by Air Target Services Pty Ltd under contract (Cook et al. 2009, Sharples et al. 2012). For each run, flown at heights of 7km to 8km, data acquisition took generally 10 minutes to complete. In some cases a number of contiguous passes were merged to provide coverage of the whole fire area.

Indicative colour samples are provided [BP]in the legend for Fig. 3[/BP], however they relate to dynamically calibrated multispectral linescan products, taken in varying smoke scattering and absorption environments, which alter the pseudo-colour rendering.

3 Results

The available data for the Grose Valley Fire allow detailed analysis of its dynamics. Fig. 1b shows localities relevant to this. These are referred to using codes explained in the figure caption.

3.1 Fire spread prior to 21 November:

The Grose Valley fire began as two ignitions generally 10 km north of Mt Boyce, which were detected on 14 November. The exact date and time of the ignitions are unknown (Cronstadt, 2006).

Prior to 02:07 21 November (linescan 116), the fire was spreading with a single bounding fireline (as seen in Fig. 3b) in a manner that could be deduced from fuel, terrain and weather data. It had gone out or been extinguished in places.

3.2 [BP]Breakaway[/BP] on 21 November:

A [BP]breakaway (AFAC 2012)[/BP] can be seen in Fig. 3b, north of AR. As a result of this, the containment line for the fire in the north-east had to be extended to the south-east of ER. Fig. 3c shows this as cold edge (linescan 117M, at 00:20 22 November), but also shows active fire edge extending past BR to the south-east.

The fire has burnt parallel to the cliffline and bypassed containment and become established on the plateau around ER, as might be expected, given the terrain and the low FMI inferred from the weather observations (Fig. 2b).

1 The fire slowed significantly on the steep downslope runs off the range at about 08:30 21
2 November, when the fire weather conditions began to ameliorate. At approximately this time
3 the prevailing wind direction also changed from west-southwest to northwest (Mt Boyce, Fig.
4 2). The apparent lack of intense wind-driven fire activity at lower elevations in the Grose
5 Valley suggests that these winds did not penetrate significantly down into the gorge; however,
6 the fire on ER spread as a head fire down BR. By 09:00 21 November this part of the fire
7 complex had burnt approximately 350 ha, with some lesser runs later in the night indicated by
8 radar returns.

9 **3.3 Overnight foehn wind, 21 November:**

10 Observations at Penrith Lakes AWS indicated the presence of a pronounced foehn effect over
11 parts of the region, starting at about 17:00 21 November, the night before the pyroCb blow-
12 ups. Indeed, the synoptic weather pattern was consistent with that associated with the
13 development of foehn winds in southeast Australia (Sharples et al. 2010). As they noted, the
14 approach of a synoptic front causes winds to align nearly perpendicular to parts of the Great
15 Dividing Range. This primarily produces mechanical foehns and resultant elevated
16 temperatures and depressed humidity levels in the lee of the terrain. An abrupt increase in
17 temperature from 21.4°C to 30.3°C with a complementary drop in relative humidity from
18 46% to 19% was recorded over the period 16:30–17:00, 21 November at Penrith Lakes AWS
19 (Fig. 2). Coincident with these changes was a change in wind direction from southwest to
20 northwest and an increase in wind speed from 9 km h⁻¹ to about 30 km h⁻¹. This nocturnal
21 foehn event would have significantly inhibited the usual moisture absorption by fuels that
22 takes place overnight (Sharples, 2009, Sharples et al., 2010). It should be noted that while Mt
23 Boyce AWS is close to the Grose Valley Fire, it is on part of the terrain that would be
24 generating, not experiencing, the foehn effect. The easterly AWSs (Penrith Lakes and
25 Cessnock) are where the full effects of adiabatic processes would be expected. The fire
26 ground, being up to 800m lower than Mt Boyce AWS, might be expected to experience the
27 effects of such a foehn wind event.

28 The fire also spread to the south, from the end of the containment line near AR. At the white
29 arrow in Fig. 3c the fire crossed a creek-line at the cliff-top and moved uphill towards the
30 south. The fire also dropped below the cliff-line near CH and began to spread slowly
31 downslope to the east towards Govetts Creek.

3.4 Widespread Spotting in Grose Valley, 22 November:

Fig. 3c also shows the fire crossing the Grose River, approximately 1.5 km WSW of MB, after backing down the slopes below AR and PL. At 01:00 22 November the wind at Mt Boyce AWS shifted towards the west and increased in speed to above 35 km h^{-1} . The change also brought the winds into closer alignment with the valley axis. Up until 02:41 22 November (linescan 126, Fig. 3d), the fire produced dense medium-range spot fires that extended 3.5km downwind (in the area labelled “GV-C” in Fig. 1b), with a cross-wind extent of approximately 2 km. After merging, these spot fires covered an area of approximately 600 ha, as indicated in Fig. 3d. An extent of about 100 ha of this was on the plateau to the northwest of EDH, where the fire had merged with the right flank of the previous fire run over ER. Fig. 3d shows that the entire area burnt with high intensity, based on the relatively uniform spectral signature for active or decaying flame (Fig. 3 [BP]legend[/BP]).

3.5 Fire channelling, Banks Ridge, 22 November:

At a similar time, the left flank of the fire on the plateau (near KGB and BR) exhibited spread with a notable northerly component, despite the west-northwest direction of the prevailing winds. This part of the fire complex is labelled “GV-N” in Fig. 1b. [AR3]The relative intensity of the spectral signals indicates that this lateral spread produced higher rates of energy release compared to those inferred from Fig. 3c, two and a half hours earlier[/AR3]. The lateral fire propagation evident at GV-N is consistent with and best explained by the fire channelling phenomenon discussed by Sharples et al. (2012). Indeed, the steep lee slopes of the numerous side spurs leading north to CR satisfy the preconditions necessary for fire channelling to occur (Sharples et al. 2012). The downwind extension of the flaming zone extending towards MC is also consistent with that seen in other fire channelling events (Sharples et al. 2012).

There was a spatial alignment between the inferred fire channelling event at GV-N, which was captured at 02:41 22 November in Fig. 3d, and the strong radar returns recorded after 02:20 (Fig. 4b), the convective core of which has been outlined in Fig. 3d. Fig. 4b indicates that the pyroconvective activity in the vicinity of GV-N started at approximately 01:30, and Fig. 4a shows that the pyroconvective top exceeded an altitude of 9 km for approximately 40 mins between 02:30 and 03:10. This period covers that in which the intense lateral spread associated with fire channelling can be inferred to have been underway. [AR1][BP][AR3]This

establishes a compelling spatial and temporal connection between the intense, lateral spread associated with fire channelling and the violent pyroconvective activity detected by the radar. As stated by Finney and McAllister (2011), a large fire source will produce a plume which experiences less entrainment and is thus able to reach greater heights. This is consistent with the GV-N situation. The significance of this linkage between fire activity and convection is further demonstrated in the later event at CH, which will be discussed below.

3.6 Fire Channelling, Clarke Head, 22 November:

Fig. 3e (linescan 127M, 05:00 22 November) shows that, after traversing the heavily dissected cliff-line between PL and CH, the fire exhibited rapid lateral spread towards the south-southwest along a 1.3 km length of cliff-line. This part of the fire complex is labelled “GV-S” in Fig. 1b. The pattern of fire propagation depicted here is consistent with fire channelling and the terrain in this vicinity meets the criteria necessary for fire channelling occurrence (Sharples et al. 2012). In addition to the lateral spread the fire also spread downwind for 2.6 km across Govetts Creek, producing an active flaming area of approximately 400 ha. This expanse of active flame was constrained at the cliff-line below DFH with some spotting for a further 1.7 km onto the plateau above. The southern flank of this part of the fire complex may be seen to be a series of merging spot fires, which is again consistent with other instances of fire channelling (Sharples et al. 2012).

The cross-wind dimension of the flaming zone depicted to the south-southwest of CH in Fig. 3e was 1.3 km, and assuming that the lateral spread was initiated immediately after 02:41, the inferred minimum rate of lateral spread is 0.6 km h^{-1} . However, it is unlikely that a spectral signature corresponding to active flaming (i.e. bright yellow) would persist for over 2 hours without cooling and so it is far more likely that the fire covered the cross-wind distance over a time of the order of the burn-out time for the dominant vegetation type in the area, i.e. ~30 mins or less. Indeed, Fig. 4b indicates peak radar returns in the vicinity of GV-S sometime after 04:30, and if this is taken as an indicator of the onset of the lateral spread, then the estimate for the rate of lateral spread can be revised to 2.6 km h^{-1} . This latter estimate of the lateral rate of spread is of a similar order to that inferred in other cases of fire channelling using overlapping linescan data (Sharples et al. 2012).

Fig. 3e illustrates the spatial alignment between the fire channelling event at GV-S and the location of the strong radar returns used to populate Fig. 4. Fig. 4 indicates that this plume developed into a significant pyroCb that exceeded 9 km in height between 05:00 and 05:40. Moreover, Fig. 4a confirms that the radar detected an abrupt increase in the echotop altitude at approximately the same time as the fire channelling event at GV-S was underway. This [A]extreme[/A] pyroCb activity associated with the fire channelling event ultimately registered echotops near the tropopause at ~11.5 km.

[AR1]It is estimated that 650ha burnt within 2 hours at GV-S, yielding an estimated energy release of around 2×10^{11} kJ, broadly comparable to values estimated for the Canberra Fire in 2003 (Fromm et al. 2006). This and the entrainment concepts in Finney and McAllister 2011 support the connection between the fire and the pyroCb.[/AR1]

4 Discussion

[BP]“Blow-up” has been used as a term for unexpected fire escalation and has been discussed at a technical level by Arnold & Buck (1954), Byram (1954), Steiner (1976) and McRae & Sharples (2013). Byram suggested that the wind profile is a contributor to blow-up potential. The vertical profile recorded at Williamtown (the closest relevant observation site), shown in Fig. 5, indicates a wind velocity maximum at about 1km above the observation site’s elevation, cf. a jet point, Byram (1954).

Other work on blow-up fires has focussed on the role of vertical stability (Haines, 1988, Mills, 2008, Mills & McCaw, 2010). As discussed by Fromm et al. (2012), the C-Haines value was [A]elevated[/A] during these events. Of relevance to the stability component of the C-Haines, a band of small dry thunderstorms that passed over the region on the afternoon of November 22 ignited new fires in the region of the Wollemi Fire by means of lightning strikes (NSW Rural Fire Service, 2006). Radar data shows that this band passed over the Grose Valley Fire at about 03:20 and the Wollemi Fire at 04:20. Fig. 6 shows a radar image from [A]05:30[/A].

[AR1]The preceding analyses indicate that future forecasts of the occurrence of a blow-up event may need to consider some combinations of the fire, weather and terrain factors identified here (and their interactions), namely: the absence of a fine fuel moistening phase during the preceding night; the maximum fire danger at the surface, and its relation to the typical diurnal cycle; a jet point in the wind profile; the terrain on which the fire is burning (ruggedness) with respect to its orientation to the prevailing wind; the time of arrival of fire in

critical parts of the landscape; and the passage of pressure troughs, or other sources of atmospheric instability, that might exacerbate fire intensity. This is reflected in the model framework proposed by McRae and Sharples (2013).[/AR1]

Direct measurement of some of the important dynamics of a complex fire burning in a rugged landscape demonstrated that a variety of mechanisms may produce the required fire [A]characteristics[/A] for violent pyroconvection to commence, but that the most significant development of pyroCbs [BP]here[/BP] occurred in connection with the fire channelling phenomenon.

Recent numerical modelling by Simpson, et al. (2013 & 2014) has demonstrated that pyrogenic vertical vorticity can produce atypical lateral spread consistent with this. Future work may shed new light on the physical mechanisms underlying the Grose Valley Fire.[/BP]

[BP]5 Conclusions

[AR1]As noted by Fromm et al. (2010) there is a need for more case studies of blow-up events to improve our understanding, but, as we have shown here for this event, these must be based on sound fire and weather data.[/AR1]

In this study we have shown that atypical modes of fire propagation in a rugged landscape, primarily fire channelling exacerbated by antecedent anomalous fire weather arising from a foehn wind occurrence, were a major factor in [BP]these particular[/BP] pyroCb blow-ups.

We have shown that from time-to-time the evolution of a fire may produce localised surges in intensity which, in the right instability setting, [A]can[/A] produce pulses of violent pyroconvection. Fromm et al. (2012) showed there were only small differences between fire weather on this day and the preceding day. This makes it clear that for fire managers to ensure the safety of fire crews and the public, they need to think beyond established practices and assess the potential for atypical fire dynamics.[/BP]

~~[BP]As pointed out by Fromm et al. (2012), the fire development on 21 November would have been subject to similar surface weather, terrain and fire behaviour features to that of 22 November. Moreover, there is only a subtle difference in the C-Haines between the two days. Thus, in terms of the tools normally applied by fire managers to such events, there may have been no substantial reason to expect the fundamentally different fire behaviour that was observed on 22 November.~~

1 Fromm et al. (2012) demonstrated that two fires in the hinterland of Sydney generated pyroCb
2 clouds on 22 November 2006. Their broad analysis, which was based on satellite, radar and
3 weather data indicated that the pyroCb occurrence was best reflected in the extreme value of
4 the Continuous Haines Index (C-Haines), especially in its 850hPa temperature component,
5 and the surface winds, which reached high values earlier in the day than would typically be
6 the case. These observations suggested that there was more to the escalation of these fires
7 than the typical diurnal weather cycle.

8
9 While fire behaviour analyses based on fuel loads, surface weather and slope may be
10 reconciled with a fire's ultimate extent, they cannot resolve the key processes described here.
11 The demonstrated importance here of foehn winds, atypical fire propagation, wind profile and
12 their association with pyroCb development provides a significant enhancement to our
13 capability to mitigate the threats from extreme fires and blow-up events.

14 It should be noted that not all unexpected fire escalations are blow-up events as described
15 above, but rather are likely to be due to poorly forecast surface weather or underestimated fuel
16 loads. It is an increasingly important task of analysts investigating significant fires to
17 determine the true nature of any escalations. [/BP]

1 **Table A1**

- 2 List of events in Australia which, based on available evidence, are known to have been, or are
3 strongly suspected to have been, pyroCb occurrences.

YEAR	Date	State	Location / Fire	Longitude (°)	Latitude (°)
1995	25 Feb	Vic	Berringa	143.7	-37.8
1998	02 Jan	Vic	Caledonia River	146.8	-37.5
2001	18 Jan	WA	Splinter Rock	122.9	-33.1
2002	17 Dec	Victoria	Big Desert	141	-35.7
2003	17 Jan	ACT	Stockyard Spur	148.9	-35.5
	18 Jan	NSW	Broken Cart Fire	148.6	-35.5
	18 Jan	ACT	Stockyard Complex	148.9	-35.6
	18 Jan	ACT	Stockyard Complex	148.8	-35.5
	18 Jan	NSW	McIntyres Hut Fire	148.8	-35.3
	18 Jan	NSW	Alpine Complex	148.4	-36
	26 Jan	NSW/Vic	Alpine Complex	[147]	[-37]
	30 Jan	NSW/Vic	Alpine Complex	[148]	[-37]
2006	19 Jan	Victoria	Grampians	142.4	-37.3
	22 Nov	NSW	Wollemi	150.3	-32.8
	22 Nov	NSW	Grose Valley	150.3	-33.6
	29 Nov	NSW	Pilliga	149.2	-31
	06 Dec	Victoria	Alpine Complex	146.5	-36.8
	14 Dec	Victoria	Alpine Complex	146.2	-37.5
	16 Dec	Victoria	Alpine Complex	[147]	[-38]
2007	16 Jan	Victoria	Alpine Complex	146.4	-37.1
2009	07 Feb	Victoria	Kilmore Gap	145.1	-37.2
	07 Feb	Victoria	Murrindindi	145.5	-37.4
	07 Feb	Victoria	Bunyip	145.7	-37.9
	07 Feb	Victoria	Beechworth	146.7	-36.4
2010	18 Feb	WA	Esperance	122.2	-33.1
2012	18 Oct	WA	Deserts	125.1	-27.3
	18 Oct	WA	Deserts	124.8	-26.5
	18 Oct	WA	Deserts	126.7	-27.4
	18 Oct	WA	Deserts	126	-27.6
	18 Oct	WA	Deserts	126.7	-28.3
	29 Oct	WA	Deserts	122.8	-28.5
2013	04 Jan	Tasmania	Dunalley	147.7	-42.9
	13 Jan	NSW	Wambelong	149	-31.3
2014	16 Jan	Victoria	Grampians	142.4	-37
	17 Jan	Victoria	Grampians	142.3	-37.1
	09 Feb	Victoria	East Gippsland	145.5	-37.3

1 Note that geographic coordinates are accurate to 0.1° only, and those in brackets are
2 approximate.

3 **Table A2**

4 List of events in Australia which are currently under investigation as candidates for pyroCb
5 occurrences.

YEAR	Date	State	Location / Fire	Longitude (°)	Latitude (°)
1983	16 Feb	Vic/SA	Ash Wednesday	[144]	[-38]
1994	23 Jan	WA	WSW of Kalgoorlie	[120]	[-32]

6 Note that geographic coordinates are accurate to 0.1° only, and those in brackets are
7 approximate.

8

1 **Acknowledgements**

2 We thank: Tony Bannister, Bureau of Meteorology, for the level 2 radar data and AWS
3 hourly surface observations; Laurence McCoy, NSW Rural Fire Service, for arranging access
4 to the linescans (all linescans are courtesy of NSW Rural Fire Service GIS Section); Robert
5 Norman, Air Target Services Pty Ltd; and David Crust, Area Manager, Mudgee, NSW
6 National Parks and Wildlife Service

7

1 **References**

- 2 [BP] AFAC : Bushfire Glossary, AFAC Limited, Melbourne, January 2012.[/BP]
- 3 American Meteorological Society: Glossary of Meteorology, 2013. ([http://](http://http://glossary.ametsoc.org/wiki/Pyrocumulonimbus)
4 <http://glossary.ametsoc.org/wiki/Pyrocumulonimbus>. Accessed 24 April, 2014.)
- 5 Arnold, R.K. & Buck, C.C.: Blow-Up fires - Silviculture or Weather Problems? Journal of
6 Forestry, 52, 408-411, 1954.
- 7 Byram, G.M.: Atmospheric Conditions Related to Blowup Fires. Southeastern Forest
8 Experimental Station Paper, Asheville, NC, 35, 1954.
- 9 Cook, R., Walker, A. & Wilkes, S.: Airborne fire intelligence. In: Jones S, Reinke K (eds)
10 Innovations in remote sensing and photogrammetry. Springer, Heidelberg, 239-254, 2009.
- 11 Cronstedt, M.: Lawson's Long Alley Section 44 Incident Controller's Report. Prepared for
12 New South Wales Rural Fire Service, Sydney, New South Wales, 2006.
- 13 [AR1]Finney, M.A. and McAllister, S.S.: A Review of Fire Interactions and Mass Fires,
14 Journal of Combustion, Article ID 548328, 14 pp, doi:10.1155/2011/54832, 2011.[/AR1]
- 15 Fromm, M., Bevilacqua, R., Stocks, B. & Servranckx, R.: New Directions: Transport to the
16 Stratosphere: Add Fire-Convection to Volcanoes. Atmospheric Environment, 38, 163-165,
17 2004.
- 18 Fromm, M., Lindsey, D.T., Servranckx, R., Yue, G., Trickl, T., Sica, R., Doucet, P. & Godin-
19 Beekmann, S.: The Untold Story of Pyrocumulonimbus. Bulletin of the American
20 Meteorological Society. September 2010, 1193 – 1209, 2010.
- 21 Fromm, M. McRae, R., Sharples, J. & Kablick, G.: Pyrocumulonimbus Pair in Wollemi and
22 Blue Mountains National Parks, 22 November 2006. Australian Meteorological &
23 Oceanographic Journal, 62, 117-126, 2012.
- 24 Fromm. M., Tupper, A., Rosenfeld, D., Servranckx, R. & McRae, R.: Violent pyro-convective
25 storm devastates Australia's capital and pollutes the stratosphere. Geophysical Research
26 Letters, 33, L05815, DOI: 10.1029/2005GL025161, 2006.
- 27 Haines, D.A.: A lower atmosphere severity index for wildland fires. National Weather Digest,
28 13, 23-27, 1988.

1 Jost, H.-J., Drdla, K., Stohl, A., Pfister, L., Loewenstein, J.P., Hudson, P.K., Murphy, D.M.,
2 Cziczo, D.J., Fromm, M., Bui, T.P., Dean-Day, J., Gerbig, C., Mahoney, M.J., Richard, E.C.,
3 Spichtinger, N., Pittman, J.V., Weinstock, E.M., Wilson, J.C. & Xuereb, I.: In-situ
4 observations of mid-latitude forest fire plumes deep in the stratosphere. *Geophys. Res. Lett.*,
5 31, L11101, DOI: 10.1029/2003GL019253, 2004.

6 Kahn, R.A., Chen, Y., Nelson, D.L., Leung, F.-Y., Li, Q., Diner, D.J. & Logan, J.A.: Wildfire
7 smoke injection heights: Two perspectives from space. *Geophys. Res. Lett.*, 35, L04809,
8 DOI: 10.1029/2007GL032165, 2008.

9 Lavoué, D., Lioussé, C., Cachier, H., Stocks, B.J. & Goldammer, J.G.: Modeling of
10 carbonaceous particle emitted by boreal and temperate wildfires at northern latitudes. *Journal*
11 *of Geophysical Research: Atmospheres*, 105, 26871-26890, 2000.

12 Lindley, T., Vitale, J., Burgett, W. & Beierle, M.: Proximity meteorological observations for
13 wind-driven grassland wildfire starts on the southern high plains. *Electronic Journal of Severe*
14 *Storms Meteorology*, 6, 1-27, 2011.

15 Mazzoni, D., Logan, J.A., Diner, D., Kahn, R., Tong, L. & Li, Q.: A data-mining approach to
16 associating MISR smoke plume heights with MODIS fire measurements. *Remote Sensing and*
17 *Environment*, 107, 138-148, 2007.

18 McRae, R. & Sharples, J.: A conceptual model for assessing the risk posed by extreme
19 bushfires. *Australian Journal of Emergency Management*, 26, 47-53, 2011.

20 McRae, R. & Sharples, J.: A process model for forecasting conditions conducive to blow-up
21 fire events. *Proceedings, 2013 MODSIM Conference, Adelaide, SA*, 2013.

22 Mills, G.A.: On the sub-synoptic scale meteorology of two extreme fire weather days during
23 the Eastern Australian fires of January 2003. *Australian Meteorological Magazine*, 54, 265-
24 290, 2005.

25 Mills, G.A.: Abrupt surface drying and fire weather Part 2: a preliminary synoptic
26 climatology in the forested areas of southern Australia. *Australian Meteorological Magazine*,
27 57, 311-328, 2008.

28 Mills, G.A. & McCaw, L.: Atmospheric Stability Environments and Fire Weather in Australia
29 – extending the Haines Index. Report 20, Centre for Australian Weather and Climate
30 Research, Melbourne, Victoria, 2010.

- 1 Nairn, G.: A Nation Charred: Inquiry into the Recent Australian Bushfires. The Parliament of
2 the Commonwealth of Australia, Canberra, 2003.
- 3 New South Wales Rural Fire Service: Wollemi Western division Section 44 Report, New
4 South Wales Rural Fire Service, Sydney, New South Wales, 2006.
- 5 Noble, I.R., Bary, G.A.V., Gill, A.M.: McArthur's fire-danger meters expressed as equations.
6 Australian Journal of Ecology, 5, 201-203, 1980.
- 7 Potter, B.E.: A dynamics based view of atmosphere-fire interactions. International Journal of
8 Wildland Fire, 11, 247-255, 2002.
- 9 Potter, B.E., Charney, J.J., Heilmen, W.E., Bian, X.: Advances in fire convection dynamics.
10 In: M. Furniss, C. Clifton and K. Ronnenberg, (Eds.), Advancing the Fundamental Sciences:
11 Proceedings of the Forest Service National Earth Sciences Conference, San Diego, CA, 18-22
12 October 2004, PNWGTR-689, Portland, OR: U.S. Department of Agriculture, Forest Service,
13 Pacific Northwest Research Station, 2007.
- 14 Rosenfeld, D., Fromm, M., Trentmann, J. Luderer, G., Andreae, M.O., and Servranckx, R.:
15 The Chisholm firestorm: observed microstructure, precipitation and lightning activity of a
16 pyro-cumulonimbus. Atmos. Chem. Phys., 7, 645-659, 2007.
- 17 Sharples, J.J.: An overview of mountain meteorological effects relevant to fire behaviour and
18 bushfire risk. International Journal of Wildland Fire, 18, 737-754, 2009.
- 19 Sharples, J.J. & Matthews, S.: Evaluation of some simplified models for predicting the
20 moisture content of fine, dead fuels. In Chan, F., Marinova, D. and Anderssen, R.S. (eds)
21 MODSIM2011, 19th International Congress on Modelling and Simulation. Modelling and
22 Simulation Society of Australia and New Zealand, December 2011, Perth, Western Australia,
23 242-248, 2011.
- 24 Sharples, J.J. & McRae, R.H.D.: Evaluation of a very simple model for predicting the
25 moisture content of eucalypt litter. International Journal of Wildland Fire, 20, 1000-1005,
26 2011.
- 27 Sharples, J.J., McRae, R.H.D., Weber, R.O. & Gill, A.M.: A simple index for assessing fuel
28 moisture content. Environmental Modelling and Software, 24, 637-646, 2009.

1 Sharples, J.J., McRae, R.H.D. & Wilkes, S.R.: Wind-terrain effects on the propagation of
2 wildfires in rugged terrain: fire channelling. *International Journal of Wildland Fire*, 21, 282-
3 296, 2012.

4 Sharples, J.J., Mills, G.A., McRae, R.H.D. & Weber, R.O.: Foehn-like winds and elevated fire
5 danger conditions in south-eastern Australia. *Journal of Applied Meteorology and*
6 *Climatology*, 49, 1067-1095, 2010.

7 [BP]Simpson, C. C., Sharples, J. J., and Evans, J. P.: Resolving vorticity-driven lateral fire
8 spread using the WRF-Fire coupled atmosphere-fire numerical model, *Nat. Hazards Earth*
9 *Syst. Sci.*, 14, 2359-2371, doi:10.5194/nhess-14-2359-2014, 2014.

10 Simpson, C., Sharples, J., Evans, J., and McCabe, M.: Large eddy simulation of atypical
11 wildland fire spread on leeward slopes, *Int. J. Wildl. Fire*, 22, 282–296, 2013.[/BP]

12 Smith, R.: Report of the Key Issues Identified from Operational Reviews of Major Fires in
13 Victoria 2006/07. Melbourne: Department of Sustainability and Environment, 2007.

14 Steiner, J.T.: Blow-Up fires - The Byram Wind Profile. *Australian Meteorological Magazine*,
15 24, 139-142, 1976.

16 Teague, B., McLeod, R., Pascoe, S.: 2009 Victorian Bushfires Royal Commission Final
17 Report, State of Victoria, Melbourne, Victoria, 2010.

18 Trentmann, J., Luderer, G., Winterrath, T., Fromm, M.D., Servranckx, R., Textor. C.,
19 Herzog, M., Graf, H.-F., and Andreae, M.O.: Modeling of biomass smoke injection into the
20 lower stratosphere by a large forest fire (Part I): reference simulation. *Atmos. Chem. Phys.*
21 *Discuss.*, 6, 6041-6080, 2006.

22 Val Martin, M., Logan, J.A., Kahn, R.A., Leung, F-Y., Nelson, D.L. & Diner, D.J.: Smoke
23 injection heights from fires in North America: analysis of 5 years of satellite observations
24 *Atmos. Chem. Phys.*, 10, 1491-1510, 2010.

25

Figures

Figure 1. (a) Location map, showing: urban areas (grey), areas above 800m ASL (orange), burnt areas (black outline) and blow-up fire events (red). Also shown is the extent of the other figure panels. (b) Localities for the Grose Valley Fire. The outskirts of the town of Blackheath (lower left) are shown in magenta, while the final fire area is outlined in red. The contour interval is 100m, major clifflines are shown by heavy black lines. Elevation colours: 300 m = green; 600 m = white; 800 m = yellow; 1100 m = brown. The three fire areas are labelled GV-N, GV-C & GV-S. Locality codes are - AR: Anvil Rock; BR: Banks Ridge; CH: Clarke Head; CR: Carmarthen Ridge; DFH: Du Faur Head; EDH: Edgeworth David Head; ER: Explorers Range; KGB: King Georges Brook; MB: Mt Banks; MC: Mt Caley; PL: Perrys Lookdown. The white dashed lines are the easterly containment lines early on 22 November. (c) Landsat imagery from 2002. (d) Landsat imagery from 5 weeks after the Grose Valley Fire. The intensity of the red is a measure of fire intensity. (Landsat 5 Thematic Mapper, Path 9 Row 83, 27 December 2006.) Major clifflines are highlighted in (c) and in (d).

Figure 2. Weather graphs for four AWSs in the region, with surface weather (left-hand panels) and fire weather (right-hand panels). Dotted horizontal lines indicate FMI=5. For Cessnock and Penrith Lakes, the black dotted outlines highlight the nocturnal foehn event. The echotop time sequence (see Fig. 4 for detail) is included for cross-reference purposes.

Figure 3. (a) Summary of Grose Valley Fire spread dynamics. Arrows show, in a stylised manner, fire spread directions. Cross-hatched lines indicate the control lines that were in place from time-to-time.

(b, c, d & e) Linescan excerpts. Control lines are shown by white dashed lines.

(b) Run 116, flown 02:07, 21 November.

(c) Run 117M, flown at 00:20, 22 November.

The white arrow near PL is the critical point where fire was able to spread along the cliff-top into the GV-S area.

(d) Run 126, flown at 02:41, 22 November. The core of the contemporaneous radar [BP]reflectivity[/BP] pattern is outlined in white.

(e) Run 127M, flown at around 05:00, 22 November. The core of the contemporaneous radar reflectivity pattern is outlined in white.

[BP](Legend)[/BP] Linescan thematic legend – note the colours are indicative only, as there is considerable variation due to smoke density. Some colours arise from a composite of surface and atmospheric features.

Figure 4. Time series analysis of localised convective activity from radar data. The dataset spans 00:00 and 07:00 on 22 November only. (a) Echotop time series, from Newcastle radar data, for Grose Valley Fire. [BP]The horizontal dotted line indicates the tropopause altitude, determined from the Williamtown Radiosonde measurement at 00:00 on 22 November (12 km a.s.l., Fromm et al, 2012). [/BP] (b) Radar summaries. Black bars indicate pyro-tops over 9km ASL. Grey bars represent ten minute typical reflectivity, with the grey shade indicating reflectivity class values. Vertical black lines indicate key overhead remote sensing passes. The black dots indicate identified fire channelling events.

Figure 5. Radiosonde data from Williamtown, 00:00 22 November. The grey line indicates the elevation of the GVN fire area.

Figure 6. [A]Newcastle (Lemon Tree Passage) Radar rain rate product for 05:30 22 November. The black outlines are the burnt areas. The diagonal band of instability that is about to pass over the radar site from the west (situated between Scone and Lake Macquarie and extending to the SE into the Tasman Sea) is strongly interacting with the plume from the Wollemi Fire (between Putty and Nullo Mountain).[/A]