



Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Changes in the occurrence of rainfall-induced landslides in Calabria, Southern Italy, in the 20th century

S. L. Gariano^{1,2}, O. Petrucci³, and F. Guzzetti¹

¹CNR IRPI, via della Madonna Alta 126, 06128 Perugia, Italy

²University of Perugia, Department of Physics and Geology, via A. Pascoli, 06123 Perugia, Italy

³CNR IRPI, via Cavour 6, 87036 Rende (CS), Italy

Received: 5 May 2015 – Accepted: 8 May 2015 – Published: 4 June 2015

Correspondence to: S. L. Gariano (gariano@irpi.cnr.it)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We exploit a catalogue of 1466 rainfall events with landslides in the 90 year period 1921–2010 to study temporal and geographical variations in the occurrence of landslides in Calabria, Southern Italy. We use daily rainfall records obtained by a network of 318 rain gauges to reconstruct 448 493 rainfall events. Combining the rainfall and the landslide information, we obtain a catalogue of 1466 rainfall events with landslides (REL) in Calabria from 1921 to 2010, where a REL is the occurrence of one or more landslide during or immediately after a rainfall event. We find that the geographical and the temporal distributions of the rainfall-induced landslides have changed in the observation period. The average and the maximum values of the cumulated event rainfall that have resulted in landslides in the recent-most 30 year period 1981–2010 are lower than the values necessary to trigger landslides in previous periods, whereas the duration of the rainfall events that triggered landslides has remained the same. This can be considered evidence of variations in rainfall conditions, but also an increase in the vulnerability of the territory. We further find that the yearly distribution of rainfall-induced landslides has changed in the observation period, analysing the variations in the number of rainfall events with landslides occurred in each month in three 30 year periods. To investigate variations in the impact of REL on the population, we compared the number of REL in each of the 409 municipalities in Calabria, with the size of the population in the municipalities, measured by national Censuses conducted in 1951, 1981, and 2011. For the purpose, we adopted two strategies. The first strategy considered impact as $I_{REL} = \#REL/P$ and the second strategy measured impact as $R_{REL} = \#REL \times P$, where $\#REL$ is the total number of REL in a period, and P is the size of the population in the same period and geographical area. Considering the entire observation period, I_{REL} and R_{REL} have both increased in Calabria. However, considering the changes between the recent period 1981–2010 and the previous period 1951–1980, results are more variegated with a number of municipalities where I_{REL} and R_{REL} have increased,

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In this work, we exploit a catalogue of rainfall events with landslides in Calabria, Southern Italy, in the 90 year period 1921–2010 to study possible variations in the frequency, geographical distribution, and impact of rainfall-induced landslides on the population of Calabria.

2 Background

In the literature, the analysis of the effects of climatic and environmental changes on landslide activity is performed using modelling or empirical approaches (Crozier, 2010). The modelling approach investigates variations in the stability/instability conditions of single landslides driven by records rainfall and pore pressure measurements, and attempts to predict variations in the stability/instability conditions using synthetic future rainfall records obtained from downscaled global climate models (Buma and Dehn, 1998, 2000; Dehn and Buma, 1999; Dehn et al., 2000; Comegna et al., 2013; Rianna et al., 2014). The empirical approach exploits landslide records to determine variations in the activity or the frequency of the landslides, and can be separated in two groups, depending on the tools used to construct the landslide records. An approach exploits paleo-environmental evidences to construct and analyse landslide records, or periods of increased/decreased landslide activity. Borgatti and Soldati (2010) adopted this approach to establish relationships linking climate variations and landslides in the Italian Dolomites. Using stratigraphic methods and radiocarbon dates, these authors dated landslides of different types in the Late-glacial – Holocene transition period, and attributed the cluster of the dated landslides to increased permafrost melting. Another approach consists in the comparison of catalogues of historical landslide occurrences to records of rainfall measurements. Stoffel et al. (2014) used this approach to analyse changes in the frequency and seasonal distribution of shallow landslides in Piedmont, NW Italy, from 1960 to 2011. These authors identified two periods of increased landslide frequency attributed to an increase in the mean annual temperature. Polemio and Petrucci (2010) studied monthly rainfall (and temperature) records in Calabria from

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a number of regional and local catalogues of landslide events exists in Italy, including catalogues for e.g. the Emilia-Romagna (Emilia-Romagna SGSS, 2006), Umbria (Salvati et al., 2006), and Calabria (Petrucci and Versace, 2005, 2007; Petrucci et al., 2009; Palmieri et al., 2011) regions.

3 Method

We base our analysis of the possible variations in the rainfall-induced landslides in Calabria on two sources of information collected in the region, including: (i) a database of daily rainfall measurements, and (ii) a catalogue of rainfall-induced landslides. Both sources cover the 90 year period 1921–2010. Our approach relies on the construction and analysis of three catalogues i.e. (i) a catalogue of *landslide events*, (ii) a catalogue of *rainfall events*, and (iii) a catalogue of *rainfall events with landslides*.

We first define a *landslide event* (LE) as the occurrence of one or more landslides in a given municipality and in a given date (day, month, year). In the literature, no clear definition exists for a *rainfall event* (RE), and no common criteria exist to single out RE from rainfall records (Melillo et al., 2015). In this work, we define a *rainfall event* (RE) as a continuous sequence of rainy days (i.e. days with cumulated daily rainfall > 0 mm) preceded and followed by at least one dry day (i.e. a day with no measured rainfall, Gullà et al., 2012). We further define a *rainfall event with landslides* (REL) as the occurrence of a LE during or immediately after a RE. To single out the individual REL, we use two criteria. First, the geographical distance between the LE and the location of the rain gauge where the event is determined shall be < 5 km. Where two or more rain gauges meet this criterion, we select the rain gauge closest to the landslide. Second, the date of the LE must be between the start and the end dates of the RE, or no more than one day after the end of the RE.

Our analysis relies further on the definition of empirical rainfall thresholds for possible landslide occurrence in Calabria. To define the rainfall thresholds we adopt the method proposed by Brunetti et al. (2010) and modified by Peruccacci et al. (2012),

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where the threshold curve is a power law relationship linking the rainfall duration D to the cumulated event rainfall $E = (\alpha \pm \Delta\alpha) \times D^{(\gamma \pm \Delta\gamma)}$, α is a scale parameter that defines the intercept of the power law threshold model, γ is the shape parameter that defines the slope of the power law model, and $\Delta\alpha$ and $\Delta\gamma$ represent the uncertainties of α and γ , respectively. The method allows defining thresholds for different exceedance probabilities. For our analysis, we define 5% thresholds (T_5) i.e. threshold lines that leave 5% of the (D, E) empirical points below the threshold.

4 Study area

Our study area is Calabria, a region in Southern Italy that extends for 15 080 km² (Fig. 1a) and comprises 409 municipalities ranging in size from 2.4 to 292.0 km² (average 38.4 km²). Elevation in the region ranges from sea level to 2260 m a.s.l., and morphology is shaped by a tectonic uplift initiated in the Quaternary and that remains active. Allochthonous crystalline rocks, Palaeozoic to Jurassic in age, stacked over carbonate units in the middle Miocene represent the backbone of the region, with Neogene flysch filling tectonic depressions (Tortorici, 1982; Monaco and Tortorici, 2000). Mean annual precipitation averages 1150 mm, with the Ionian (E) side of the region less rainy than the Tyrrhenian (W) side. Annual rainfall depends on elevation, with the mountains significantly wetter (> 2000 mm) than the coastal plains (< 500 mm). About 70% of the annual rain falls from October to March, and 10% in the summer (Terranova, 2004). Rainfall events with large cumulated rainfall occur mainly between November and January, whereas high intensity events are most common in September and October (Terranova and Gariano, 2014).

A number of studies have investigated variations in the rainfall patterns and trends in Calabria. Ferrari and Terranova (2004) revealed a reduction in the annual and the winter amounts of rainfall for two overlapping periods (1920–2000 and 1960–2000), and Caloiero et al. (2011) recognized an augmented trend in the summer rainfall. Caloiero et al. (2008) showed that short-duration rainfall events were most frequent in November

of the cumulative curve, suggests a uniform reconstruction of the RE, in the considered period.

Figure 3a shows the number of severe RE per year in the 90 year observation period. The average value per year is 28, and was exceeded 43 times, of which 15 times in the 1921–1950 period, 18 times in the 1951–1980 period, and 10 times in the 1981–2010 period. The 1951–1980 period was characterised by the largest average number of severe RE per year (318). The maximum number of severe RE (609) was recorded in 1954, and the minimum (79) in 1922. In six years (1930, 1933, 1940, 1954, 1973, 1996) the number of reconstructed severe RE was larger than 500.

5.3 Catalogue of rainfall events with landslides

Using the method presented in Sect. 3 we reconstructed 1989 REL in Calabria between October 1921 and December 2010. The REL have an average duration $D = 6$ days, and an average cumulated event rainfall $E = 157.5$ mm. The reduced number of REL (1989) compared to the number of LE (7600) has many reasons. First, many landslides listed in the LE catalogue were not triggered by rainfall. In some cases they were triggered by earthquakes (e.g. in 1947), and in other cases by human activities. These LE were excluded from the analysis. Second, in some cases the rain gauges failed to measure or to record the rainfall, and the landslide information could not be used to reconstruct a REL. Third, for small shallow landslides triggered by intense, short-duration rainfall events lasting only a few hours, the daily rainfall measurements were not adequate to identify a triggering event, and the REL were not determined. Fourth, for some of the old events uncertainty in the date of the slope failure resulted in a mismatch between the landslide and the daily rainfall record, and the REL could not be determined. Finally we discarded from the analysis all the RE with mean rainfall intensity < 10 mm day⁻¹. The value was selected heuristically, following Terranova and Gariano (2014), to exclude from the analysis less significant events. The remaining 1466 RE with landslides have durations in the range $1 < D < 32$ days (average 7 days), and cumulated event rainfall in the range $10.0 < E < 1504.7$ mm (average 202.3 mm) (Table 1).

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 2e shows the number of REL per municipality. The average number of REL in a municipality is four, the minimum is zero (95 municipalities, located chiefly in the N part of the region and along the SW coast), and the maximum is 47, in the Catanzaro municipality (Fig. 2e). More than 30 municipalities experienced 10 or more REL in the investigated period. Figure 2f portrays the number of REL per year. On average, 16 REL occurred every year with a maximum of 139 REL in 1973. The three days with the largest number of REL were 18 February 2010 (39 REL), 29 February 1956 (33), and 27 November 1959 (30). The month with the largest number of REL was January (337), followed by November (255), and February (254). The decades with the largest number of REL were those between 1950 and 1959, between 1970 and 1979, and between 2000 and 2009. Inspection of the cumulated number of REL (black curve in Fig. 2f) reveals that in the early period 1910–1950, the rate of the REL is lower than in the later period 1950–2010. This is a result of the completeness of the information of the LE (Fig. 2b).

Figure 3b shows the number of REL per year in the 90 year observation period. In six years more than 60 REL occurred: 1973 (135 REL), 2009 (101), 1953 (92), 2010 (78), 1954 (76), and 1996 (72). The average value (16 REL per year) was exceeded 28 times: twice in the 1921–1950 period and 13 times in each of the following 30 year periods. The intermediate (1951–1980) and the recent (1981–2010) periods were characterised by average number of REL per year equal to 24 and 20, respectively. In both cases the values are larger than the 90 year average (16).

6 Analysis of rainfall events with landslides

Our catalogue lists 1466 severe REL in Calabria from October 1921 to December 2010, of which 534 (36.4%) have triggered “single” landslides and 932 (63.6%) “multiple” landslides. REL with “single” landslides have average and maximum cumulated event rainfall E lower than REL with “multiple” landslides (Table 1). For 924 REL (63.0%), qualitative information exists on the size of the landslides, with 610 REL (41.6%) that

Figure 6a–c shows the number of REL per municipality in three periods: 1921–1950 (old period), 1951–1980 (intermediate period), and 1981–2010 (recent period). Of the 1466 REL in the catalogue, 143 REL (9.8 %) occurred in the old period, 720 (49.1 %) in the intermediate period, and 603 (41.1 %) in the recent period (Table 1). The spatial distributions of the REL in the intermediate and the recent periods are similar, with a larger number of REL affecting the SE part of the region in the recent period. Figure 6d–f portrays the monthly distributions of the REL in the three periods. In the old period (Fig. 6d) REL occurred between October and March, with the majority of the REL (51, 35.7 %) in November. In the intermediate and the recent periods, the REL were also most abundant between October and March. In particular, in the intermediate period 1951–1980, REL were equally distributed in the autumn (322, 44.7 %) and the winter (329, 45.7 %), with peaks in October (145, 20.1 %) and November (170, 11.6 %, Fig. 6e). In the recent period, the majority of the REL (369, 61 %) occurred in winter, with a distinct peak in January (162 REL, 26.9 %) (Fig. 6f) and only 86 REL (14.3 %) in October and November. In the three considered periods, the REL exhibited similar ranges of rainfall duration D , and different ranges of cumulated event rainfall E (Table 1), with the REL in the recent period exhibiting larger maximum D , and lower average and maximum E than the corresponding values for the REL in the previous periods.

7 Discussion

Using the three catalogues of landslide events (LE), of rainfall events (RE), and of rainfall events with landslides (REL) in Calabria in the 90 year period 1921–2010, we first investigate the changes in the yearly distribution of rainfall events with landslides. This is followed by an analysis of the changes in the rainfall conditions that have resulted in landslides in Calabria, measured by empirical rainfall thresholds for landslide occurrence. Lastly, we compare the variations in the number and distribution in time of

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1.14 (+11.8 %) in the intermediate period, and increased further to 1.23 (+7.9 %) in the recent period. Table 2 lists the mean and maximum values of cumulated event rainfall E (in mm) for rainfall events reconstructed in the entire 90 year period and in the three considered 30 year periods. RE occurred in October and November in the intermediate period have larger (average, maximum) cumulated event rainfall E than RE occurred in the same two months in the recent period (Table 2). Conversely, REL occurred in December and January exhibit similar values of E (average, maximum) for all the three periods. The difference marks a dissimilarity in the distribution of the REL in the three 30 year periods.

7.2 Changes in the rainfall thresholds

To ascertain whether the rainfall conditions for possible landslide occurrence have changed in Calabria in the 90 year observation period, we defined cumulated event rainfall-rainfall duration (ED) thresholds for all the REL in the catalogue, and for different subsets. Table 3 lists the equations and the ranges of validity of the established 5 % ED thresholds. Except for the validity range, only minor differences exist in the parameters controlling the thresholds defined for the entire catalogue ($T_{5,CAL}$), for rainfall conditions that have resulted in “single” ($T_{5,SG}$) or “multiple” ($T_{5,ML}$) landslides, and for rainfall conditions that have caused “small” ($T_{5,SL}$) or “large” ($T_{5,LG}$) landslides.

Inspection of Table 3 reveals that the $T_{5,CAL}$, $T_{5,SG}$ and $T_{5,ML}$ thresholds have the same or very similar exponent (γ) and intercept (α) of the power law models. Although the average and the maximum cumulated event rainfall E responsible for “single” landslides are smaller than the E that have resulted in “multiple” landslides, the rainfall amount required to trigger “single” or “multiple” landslides has remained the same in Calabria in the 90 year observation period. The same is true for the rainfall conditions that have resulted in “small” or “large” landslides (Table 3). However, rainfall events that have resulted in “small” landslides have maximum and average rainfall duration D and cumulated event rainfall E lower than the rainfall events that have resulted in large landslides (Table 1).

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Considering the seasonal periods, the threshold defined for autumn ($T_{5,AT}$, $\gamma = 0.82 \pm 0.04$) is steeper than the threshold defined for winter ($T_{5,WT}$, $\gamma = 0.74 \pm 0.04$), and the threshold for the “wet” period ($T_{5,WET}$, $\gamma = 0.75 \pm 0.03$) is flatter than the threshold for the “dry” period ($T_{5,DRY}$, $\gamma = 0.99 \pm 0.04$). For the latter two thresholds the intercepts (α) are also different (Table 3). The number of the empirical data for the spring and the summer periods was not sufficient to determine reliable thresholds (Vennari et al., 2014), and a comparison for the two seasons is not possible.

Figure 7a compares the ED threshold defined in this work for Calabria ($T_{5,CAL}$) to the threshold defined by Vennari et al. (2014) ($T_{5,VEN}$) for shallow landslides using rainfall and landslide information in the period 1996–2011. The $T_{5,CAL}$ threshold is significantly steeper than the $T_{5,VEN}$ threshold ($\gamma = 0.74 \pm 0.03$ vs. $\gamma = 0.41 \pm 0.03$), and can be applied only for $D > 24$ h ($24 \leq D \leq 768$ h), whereas the $T_{5,VEN}$ is applicable for $1 \leq D \leq 451$ h. We maintain that the observed difference depends largely on the different resolution of the rainfall records i.e. daily measurements for our catalogue and hourly measurements for Vennari et al. (2014).

Figure 7b shows that the ED thresholds for landslide occurrence in Calabria are different for the three 30 year considered periods ($T_{5,21-50}$, $T_{5,51-80}$, $T_{5,81-10}$). Inspection of Table 1 confirms that the three thresholds are different, both for their intercepts (α) and the slopes (γ). However, considering the uncertainty associated to the thresholds (Brunetti et al., 2010; Peruccacci et al., 2012), we observe that the thresholds overlap for $40 < D < 400$ h. We further observe that $T_{5,21-50}$ is the steepest threshold ($\gamma = 0.93 \pm 0.05$) and $T_{5,81-10}$ is the less steep threshold ($\gamma = 0.66 \pm 0.04$). We infer that for rainfall events having $D \leq 72$ h (three days), less rain was required to trigger landslides in the old (1921–1950) period than in the intermediate and in the recent periods.

7.3 Changes in the landslide impact to the population

To investigate the variations in the impact of landslides on the population of Calabria, we compared the number of REL with the size of the population in each of

4242 km² of the region, 28.1 %), no variation in R_{REL} and I_{REL} between the intermediate and the recent period was observed (no REL in the catalogue in both periods).

Overall, I_{REL} has increased in 73 municipalities and decreased in 47 municipalities, covering 15.5 and 10.7 % of the region, respectively. In the 73 municipalities where I_{REL} has increased lives 18.5 % of the population of Calabria, and in the 47 municipalities where I_{REL} has decreased lives 20.8 % of the population. R_{REL} has increased in 30 municipalities, covering 6.7 % of the region and with 24.0 % of the total population. Conversely, R_{REL} has decreased in 147 municipalities, covering 39.0 % of the region and with 19.1 % of the population. Overall, 93 municipalities went from $R_{REL} > 0$ and $I_{REL} > 0$ in the intermediate period, to $R_{REL} = 0$ and $I_{REL} = 0$ in the recent period. In these municipalities the risk of REL to the population has decreased. Conversely, 44 municipalities changed the indexes from $R_{REL} = 0$ and $I_{REL} = 0$ in the intermediate period, to $R_{REL} > 0$ and $I_{REL} > 0$ in the recent period. In total, 42.5 % of the regional population suffered an increased impact, both in terms of I_{REL} and R_{REL} . Municipalities where I_{REL} has increased between the intermediate and the recent period are located mainly in the inner and mountainous areas of the region, and municipalities where I_{REL} has decreased between the two periods are mostly along the coasts and in the N part of the region. Municipalities where R_{REL} has increased in the recent period, compared to the intermediate period, are located mainly along the coasts, and the municipalities where R_{REL} are unevenly distributed in the entire regional.

8 Conclusions

We have investigated geographical and temporal changes in the occurrence of 1466 rainfall-induced landslide events in Calabria, in the 90 year period 1921–2010.

Our work revealed that the rainfall conditions that have resulted in rainfall-induced landslides in Calabria have changed in the observation period. We found that less cumulated event rainfall (E) was necessary to trigger landslides in the recent period (1981–2010) than in the preceding period (1951–1980). We consider this evidence of

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the increased propensity of the landscape to generate landslides in the recent period, and of the larger number of vulnerable elements exposed to landslides, in Calabria. We found that the monthly distribution of rainfall events with landslides (REL) has changed. In the earlier period (1951–1980) landslides were more frequent in autumn and winter, whereas in the recent period (1981–2010) landslides concentrated in the winter. We further observed significant variations in the geographical distribution of the REL. In the recent period (1981–2010) REL struck mainly the SE part of the region. We attribute the observed variations in the temporal and the geographical distributions of rainfall events with landslides in Calabria to variations in the frequency of the triggering rainfall events, and to an increased susceptibility to landslide of the territory.

We investigated the changes in the influence of rainfall events with landslides on the population of Calabria, adopting two complementary strategies. The analysis revealed a contrasting picture. The impact of REL on the population has increased in 37 municipalities covering 15.5% with 18.5% of the population, and has decreased in 47 municipalities covering 17.8% with 20.8% of the population. The risk posed by REL to the population has increased in 30 municipalities covering 6.7% and hosting 24.0% of the population, and has decreased in 47 municipalities covering 39.0% with 19.1% of the population. Overall, 42.5% (57.5) of the regional population has experienced an increased (decreased) level of landslide impact or risk.

Finally, we note that the observed changes in the impact of rainfall induced landslides on the population of Calabria in the investigated 90 year period are due to changes in the number of the events (a largely natural component) and to changes in the number of the exposed elements (a largely societal component). Several of the municipalities with an increased landslide risk between the intermediate and the recent periods have experienced a similar number of REL (hazard has remained about the same) in the two periods, but a large increase in the size of the population (exposure has increased) in the recent period.

We expect that the result of our study will be used for landslide hazard assessment and risk evaluation in Calabria, and in similar physiographic regions in the Mediterranean area.

**The Supplement related to this article is available online at
doi:10.5194/nhessd-3-3579-2015-supplement.**

Acknowledgements. The “Centro Funzionale Multirischi” of the Agenzia Regionale per l’Ambiente della Calabria (ARPACAL, <http://www.cfd.calabria.it>) made available the rainfall data. Paola Salvati and Cinzia Bianchi (CNR IRPI) provided updated information on landslides with human consequences in Italy and in Calabria. Massimo Melillo (CNR IRPI) calculated the rainfall thresholds.

References

- Borgatti, L. and Soldati, M.: Landslides as a geomorphological proxy for climate change: a record from the Dolomites (northern Italy), *Geomorphology*, 120, 56–64, 2010.
- Brunetti, M., Buffoni, L., Mangianti, F., Maugeri, M., and Nanni, T.: Temperature, precipitation and extreme events during the last century in Italy, *Global Planet. Change*, 40, 141–149, 2002.
- Brunetti, M., Caloiero, T., Coscarelli, R., Gullà, G., Nanni, T., and Simolo, C.: Precipitation variability and change in the Calabria region (Italy) from a high resolution daily dataset, *Int. J. Climatol.*, 32, 57–73, 2012.
- Brunetti, M. T., Peruccacci, S., Rossi, M., Luciani, S., Valigi, D., and Guzzetti, F.: Rainfall thresholds for the possible occurrence of landslides in Italy, *Nat. Hazards Earth Syst. Sci.*, 10, 447–458, doi:10.5194/nhess-10-447-2010, 2010.
- Brunetti, M. T., Peruccacci, S., Antronico, L., Bartolini, D., Deganutti, A. M., Gariano, S. L., Iovine, G., Luciani, S., Luino, F., Melillo, M., Palladino, M. R., Parise, M., Rossi, M., Turconi, L., Vennari, C., Vessia, G., Viero, A., and Guzzetti, F.: Catalogue of rainfall events with shallow landslides and new rainfall thresholds in Italy, in: *Engineering Geology for Society and Territory*, Vol. 2, edited by: Lollino, G., Giordan, D., Crosta, G. B., Corominas, J., Azzam, R.,

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Terranova, O. G. and Gariano, S. L.: Rainstorms able to induce flash floods in a Mediterranean-climate region (Calabria, southern Italy), *Nat. Hazards Earth Syst. Sci.*, 14, 2423–2434, doi:10.5194/nhess-14-2423-2014, 2014.

Tortorici, L.: Lineamenti geologico-strutturali dell'Arco Calabro-Peloritano, *Rendiconti della Società Italiana di Mineralogia e Petrologia*, 38, 940–972, 1982.

Trigila, A., Iadanza, C., and Spizzichino, D.: Quality assessment of the Italian landslide inventory using GIS processing, *Landslides*, 7, 455–470, 2010.

Varnes, D. J. and IAEG Commission on Landslides and other Mass Movements on Slopes: *Landslide Hazard Zonation: A Review of Principles and Practice*, UNESCO Press, Paris, 1–63, 1984.

Vennari, C., Gariano, S. L., Antronico, L., Brunetti, M. T., Iovine, G., Peruccacci, S., Terranova, O., and Guzzetti, F.: Rainfall thresholds for shallow landslide occurrence in Calabria, southern Italy, *Nat. Hazards Earth Syst. Sci.*, 14, 317–330, doi:10.5194/nhess-14-317-2014, 2014.

Wood, J. L., Harrison, S., and Reinhardt, L.: Landslide inventories for climate impacts research in the European Alps, *Geomorphology*, 228, 398–408, 2015.

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Table 2. Mean and maximum (max) values of cumulated event rainfall E (in mm) for rainfall events in the 90 year period 1921–2010, and in three 30 year periods. The largest monthly values in each period are shown in bold.

Month	1921–2010		1921–1950		1951–1980		1981–2010	
	mean	max	mean	max	mean	max	mean	max
Jan	170.7	999.9	169.3	985.8	168.1	758.9	174.7	999.9
Feb	160.8	959.5	155.5	753.9	163.2	959.5	163.6	771.8
Mar	164.5	1118.3	181.6	1118.3	149.3	806.1	162.7	656.2
Apr	145.2	1124.4	137.8	298.7	149.0	1124.4	148.9	355.7
May	148.6	811.8	138.5	351.2	142.6	452.6	164.6	811.8
Jun	163.9	813.0	136.5	331.0	143.2	741.2	212.1	813.0
Jul	163.7	403.4	134.2	314.8	133.8	266.7	223.0	403.4
Aug	130.3	257.4	133.6	255.9	121.8	239.7	135.4	257.4
Sep	165.4	748.4	159.8	470.5	143.0	425.0	193.3	748.4
Oct	170.2	1504.7	158.3	533.5	198.6	1504.7	153.7	528.6
Nov	174.6	872.0	169.8	734.0	194.2	872.0	159.7	477.4
Dec	175.0	1250.7	181.6	1216.2	174.8	1250.7	168.6	696.0

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.




Figure 1. (a) Map of Calabria, Southern Italy, showing terrain elevation (shades of green to brown), main cities (yellow squares), and location of rain gauges used in the study (triangles). Shades of blue show number of years with measurements for each rain gauge, in five classes. (b) Number of rain gauges per year in Calabria between 1920 and 2010.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 2. (a) Number of landslide events (LE) in each municipality in Calabria in the 90 year period 1921–2010. (b) Temporal distribution of LE (orange bars), and cumulated number of LE (black line) in the period 1921–2010. The three single days with the largest number of LE are shown. (c) Number of rainfall events (RE) reconstructed for each rain gauge in the period 1921–2010. Reference area for each rain gauge is shown using Thiessen polygons. (d) Temporal distribution of reconstructed RE (blue bars), and cumulated number of RE (black line) in the period 1921–2010. The three single days with the largest number of RE are shown. (e) Number of rainfall events (REL) in each municipality in the period 1921–2010. (f) Temporal distribution of REL (red bars), and cumulated number of REL (black line). The three single days with the largest number of REL are shown.

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.




Figure 3. (a) Number of rainfall events (RE) and (b) number of rainfall events with landslides (REL) per year in Calabria in the 90 year period 1921–2010. Solid lines show average values for the 90 year period. Dashed lines show the average values for the three 30 year periods 1921–1950, 1951–1980, and 1981–2010.

Title Page

Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.




Figure 4. Maps show the number of rainfall events with landslides (#REL) per municipality in Calabria in the (a) Autumn (22 September–20 December), (b) Winter (21 December–20 March), (c) Spring (21 March–20 June), and (d) Summer (21 June–21 September). Pie charts show number and percentage of municipalities in each seasonal class.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 5. Top: maps show the number of rainfall events with landslides (#REL) per municipality in Calabria in **(a)** the dry period (April to October) and **(b)** in the wet period (November to March). Pie charts show number and percentage of municipalities in each class. Bottom: bar charts show temporal distribution of REL in the **(c)** dry and in the **(d)** wet periods.

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.




Figure 6. Maps show the number of rainfall events with landslides (#REL) in each municipality in Calabria in the three 30 year periods, **(a)** 1921–1950, **(b)** 1951–1980, and **(c)** 1981–2010. Pie charts show number and percentage of municipalities in each class. Middle: red bars in **(d, e)** and **(f)** show number of REL per month in the three 30 year periods. Bottom: blue bars in **(g, h)** and **(i)** show number of RE per month in the three 30 year periods.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 7. (a) Cumulated event rainfall-rainfall duration (ED) 5% thresholds for rainfall-induced landslides in Calabria defined in this study ($T_{5,Cal}$, 1), compared to the 5% threshold for shallow landslides in Calabria defined in Vennari et al. (2014) ($T_{5,Ven}$, 2). (b) Cumulated event rainfall – rainfall duration (ED) 5% thresholds defined for the three 30 year periods 1921–1950 ($T_{5,21-50}$, 3), 1951–1980 ($T_{5,51-80}$, 4), and 1951–2010 ($T_{5,81-10}$, 5). Grey areas show uncertainties associated to the thresholds. Data shown in log-log coordinates.

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.




Figure 8. Top: maps show density of population in the 409 municipalities in Calabria for three 30 year periods, **(a)** 1921–1950, **(b)** 1951–1980, and **(c)** 1981–2010. Middle: maps show landslide impact I_{REL} given by the number of rainfall events with landslides (#REL) per 1000 people in each municipality in Calabria, for the **(d)** 1921–1950, **(e)** 1951–1980, and **(f)** 1981–2010 periods. Bottom: maps show risk of rainfall events with landslides R_{REL} , given by the product of the number of rainfall events with landslides (#REL) and of inhabitants in each municipality in Calabria, for the **(g)** 1921–1950, **(h)** 1951–1980, and **(i)** 1981–2010 periods. Key: null, $R_{REL} = 0$; Low, $0 < R_{REL} \leq 10\,000$; Medium, $10\,000 < R_{REL} \leq 100\,000$; High, $R_{REL} > 100\,000$. Pie charts show number and percentage of municipalities in each class.




Figure 9. Map showing variations in I_{REL} and R_{REL} in the municipalities of Calabria, between the intermediate (1951–1980) and the recent (1981–2010) periods. Legend: red, municipalities with an increased I_{REL} (increased number of REL and a smaller population). Orange, municipalities with an increased R_{REL} (increased number of REL and a larger population). Yellow, municipalities with a decreased R_{REL} (decreased number of REL and a smaller population). Green, municipalities with a decreased I_{REL} (decreased number of REL and a larger population). White, municipalities with I_{REL} and R_{REL} null (no REL listed in the catalogue in the intermediate and in the recent periods).

Changes in the occurrence of rainfall-induced landslides in Calabria

S. L. Gariano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

