

## ***Interactive comment on “The 21st Century Decline in Damaging European Windstorms” by L. C. Dawkins et al.***

**L. C. Dawkins et al.**

[l.c.dawkins@exeter.ac.uk](mailto:l.c.dawkins@exeter.ac.uk)

Received and published: 14 July 2016

General comments

Referee comment:

The main comment is that there is insufficient evidence that the A20 metric is valid for its purpose of documenting a recent decline in damaging European windstorms. The validation in the article consists of finding a value of X% containing 23 significant windstorms, and a lower value of X indicates a better metric. This metric analyses a subjective subset of individual severe windstorms, and is not robust to outliers. Further, the validation concerned an A25 rather than the A20 metric used in this article.

Response:

[Printer-friendly version](#)

[Discussion paper](#)



This is a very good point which should be addressed within the text. This validation method was used in the Roberts et al (2012) paper and remains the best method here, because the insured loss value for the whole of Europe is only available to us for a very limited number of events, all of which are extreme storms (see Table 1 in Roberts et al. (2012)), not enough to validate the SSI. Even the Barredo (2010) paper only includes losses from 54 extreme storms between 1970 and 2008, still thought to be too few to validate these SSIs. Since the A25 measure was validated in the Roberts et al (2012) paper and has a strong positive relationship with A20 (see attached scatter plot), we feel this is sufficient evidence that the A20 SSI is valid for its purpose in this paper, having no further way to validate it.

This issue will be addressed by adding a paragraph on page 5, line 13:

‘The SSI validation method used by Roberts et al. (2012) is based on a subjective subset of extreme windstorms and is not robust to outliers. However, since the value of insured loss for the whole of Europe is only available for a very limited number of windstorm events, all of which are extreme (Roberts et al., 2012 (Table 1); Barredo, 2010), this is thought to be the most appropriate available method for validation here. If further loss data were made available for validation the exploration of the most successful European-wide SSI would be a hugely beneficial area of future research.’

We will amend Figure 2 to include the scatter plots of A25 and A20 and add text on page 5, line 24:

‘Figure 2 (a) shows a scatter plot of the logarithm of the damage area SSIs A25 and A20, and Figure 2 (b), a scatter plot of the logarithm of the SSI developed by Klawa and Ulbrich (2003), labelled L98 and the damage area SSI A20, for the top 50% of the 6103 windstorm events in the data set in each SSI.’

Define A25 on page 5, line 8/9:

‘They found that the SSI characterising the area of the footprint exceeding 25 ms–1

[Printer-friendly version](#)[Discussion paper](#)

over land, A25, outperformed . . .’

And page 5, line 29:

‘Figure 2 (a) shows that there is a strong positive relationship between A20 and A25, indicating that the A20 SSI is appropriate for representing insured loss, as shown by Roberts et al. (2012) for the A25 SSI.’

Referee comment:

However, there is a much bigger validation problem: this article analyses annual integrated A20 rather than individual severe windstorms, hence the proposed validation has little relevance. The distinction between individual event damage and annual integrated values is very substantial here: we know that a single storm such as Daria or Lothar produced more damage than the long-term annual average, whereas the A20 estimate for Daria is 1 or 2% of long-term annual average and points to far too much weight on weak storms in A20. The authors have to choose a metric which has been validated as an annual integrated measure of damage. Articles such as Barredo (2010) can provide some data on annual integrated damage to help with such a validation. This is not viewed as a major change in direction since the authors use metrics such as Klawa and Ulbrich (2003) in this article - instead, it is a change in emphasis on the best metric.

Response:

We acknowledge the difference between A20 for individual severe storms and annual integrated A20, however we are of the view that an SSI that is able to rank extreme storms highly and therefore weaker storms lower, can represent the insured loss in general. In addition, since we have the footprints for all windstorm events in each year, the sum of A20 over all of these events in one year, should be equivalent to the annual aggregate loss.

We are unsure of where these figures (1 or 2%) have come from. For windstorm Daria,

[Printer-friendly version](#)

[Discussion paper](#)



A20 is approximately 3000 (Figure 2 is on a logarithmic scale  $\exp(8) \sim 3000$ ), which is about 500% of the long-term annual average A20 ( $\sim 600$ ). Has the Figure been misread or can this comment be explained further please?

As mentioned above, the Barredo (2010) paper contains losses for 54 extreme events between 1970 and 2008, therefore we are of the view that the annual aggregate losses used in this paper will not be comparable to the annual aggregate A20 values.

Additional mention of the caveats of the A20 SSI will be included in the conclusion-Page 8, line 17:

‘This conclusion is based on a subjective set of extreme windstorms and could therefore benefit from further validation based on insured loss data, if more were available. In addition, other SSIs could have been included in the validation investigation, using different damage thresholds and exposure variables. Many such SSIs are compared by Dawkins (2016), where the damage area SSI A25 is still shown to be most successful at representing insured loss. Further, this validation is based on relatively low resolution footprints and could give different conclusions if higher resolution footprints were available for exploration.’

Referee comment:

In general, the article follows the paradigm ‘large-scale climate patterns force weather events’ with statements similar to ‘NAO explains changes in A20’. However, the aggregate of the individual weather events contribute significantly to the large-scale pattern, see text and references within “Contrasting interannual and multidecadal NAO variability” by Woollings et al. (2014). Would the authors consider changing their description of climate-weather link to something like ‘changes in NAO are consistent with annual damage metric variations’, if they find such behaviour?

Response:

This is a very good point. The language is too causal in some places and will be

[Printer-friendly version](#)

[Discussion paper](#)



changed.

Page 8, line 32:

‘This suggests that windstorms with large values of A20 may be associated with positive NAO and hence that much of the variation in A20, and therefore windstorm losses, \*is consistent with variation in NAO\*. ‘

Page 8, line 27:

‘A strong positive association was also found between winter total A20 and winter averaged NAO, showing that changes in NAO are consistent with the variation in annual A20 and is therefore related to annual loss.’

Specific Comments

Referee comment:

Page 1, lines 18-22: Roberts et al. do not provide loss estimates for the four named storms, could the source of these losses be given?

Response:

Table 1 in Roberts et al. (2012) gives the losses estimates for 16 extreme windstorms including these four named storms. These loss estimates are sourced from the Sigma technical reports between 2004 and 2013.

Referee comment:

Page 2, line 11: the loss function used five stations, rather than four.

Response:

Thank you, this will be amended

Referee comment:

Page 2, lines 17-19: the published work by Wallace would be a better reference for the

[Printer-friendly version](#)

[Discussion paper](#)



NAO?

Response:

This is a good point. On reconsideration of this reference, we think the most appropriate reference is the 2003 book by James W. Hurrell et al:

Hurrell, J. W., Y. Kushnir, G. Ottersen and M. Visbeck, 2003: The North Atlantic Oscillation, Climate Significance and Environmental Impact. American Geophysical Union, Washington, DC.

Referee comment:

Page 6, lines 3-7: The statement “..number of very damaging windstorms has decreased in recent decades” is not supported by the evidence in Figure 3. Instead, the decline in total A20 is caused by reduced A20 per event (Fig 3b) and there is no evidence specific to the subset of very damaging storms in Figure 3.

Response:

Thank you, this is a very good point. This statement will be removed from the text since this conclusion cannot be taken from Figure 3.

Referee comment:

Page 7, lines 8-9: this sentence is tautological: A20 counts number of occurrences of wind > 20 m/s, or put another way, is a measure of frequency of occurrence of winds > 20 m/s.

Response:

This sentence will be reworded to bring out the emphasis on north-west Europe: ‘The largest contribution to the decline in A20 comes from north-west Europe.’

Referee comment:

Page 7, lines 11-16: Figure 5 helps to explain A20 changes in Paris, but it also raises

some major issues. First, the peak gusts never exceed 30 m/s, yet storm Lothar was measured above 40 m/s by multiple weather stations in and around Paris. Second, the top 10 or 20 storms in the period are responsible for the vast majority of damage at Paris, and the top 10 points in this plot show the recent period to have consistently higher gusts. This indicates the extreme gusts from XWS footprints are very different from observed behaviour. Do the authors if XWS wind values have been compared to actual weather station gusts, and if so, has a trend been found such that modelled hazard for older storms have more negative bias with respect to observed, compared to newer storms?

Response:

These are very interesting points. Weather stations, which are uncertain themselves, provide point observations of wind speeds, whereas what we present is a gridded analysis so we would not expect them to be the same. However, it is true that this bias can be large. The XWS gridded footprints used here have been shown to underestimate wind gust speeds above 25m/s – see Figure 8 f) in Roberts at al. (2012), where an observed wind gust of  $\sim 60$ m/s is modelled as being  $\sim 20$ m/s. This bias is likely due to the low resolution of the gridded analysis.

When developing the XWS catalogue, only the 50 storms included in the catalogue were recalibrated, and therefore compared to weather station data. No temporal trend in the bias was noticed, however, this could be interesting to explore in more detail in future work.

We will address these comments by including extra text in data section – page 4, line 3:

‘In addition, the gridded analysis is shown to underestimate extreme wind gust speeds greater than  $\hat{\text{Lij}} 25\text{ms}^{-1}$  to varying degrees. Figure 8 c) and f) in Roberts et al (2012) show how this underestimation is small for some locations but in others this bias is much larger, with observed wind gust speeds greater than  $40\text{ms}^{-1}$ , modelled as being

[Printer-friendly version](#)

[Discussion paper](#)



below 30ms<sup>-1</sup>. This bias was found to be due to several mechanisms, for example, the underestimation of convective effects and strong pressure gradients, likely due to the limitations of the model horizontal resolution.'

By adding text after the q-q plot – page 7, line 15:

'The most extreme wind gust speeds are higher for the 21st century, however, the bias in modelled wind gust speeds at this high level, shown by Roberts et al (2012) and discussed in Section 2, suggest that these high quantiles should be treated with great uncertainty. For example, it is known that during windstorm Lothar in December 1999 wind gust speeds near 50ms<sup>-1</sup> were recorded in Paris (Ulbrich et al 2001).'

Additional reference: Ulbrich, U., A. H. Fink, M. Klawns and J. G. Pinto, 2001: Three extreme storms over Europe in December 1999. *Weather*, 56: 70-80. doi:10.1002/j.1477-8696.2001.tb06540.x

And lastly, adding a couple of sentences to the conclusion – page 9, line 32:

'This investigation is based on model generated gridded analysis windstorm footprints. While the large number of footprints within the data set is beneficial in validating conclusions, the modelled wind gust speeds are known to be biased, particularly in areas of high altitude and when wind gust speeds exceed 25ms<sup>-1</sup>. Further exploration of this bias could improve the validity of the investigation and carrying out the same analysis using both observations and gridded analysis could be an interesting extension of this work.'

Referee comment:

Page 8, lines 4-10: could the Scandinavia Pattern be included in analysis? The eastwards extent of the spatial pattern in Figure 6 suggests the SP.

Response:

We are of the view that the Scandinavian Pattern will not relate to correlation pattern

[Printer-friendly version](#)

[Discussion paper](#)





over both north and south Europe. In addition, we think that in adding more climate indices, a longer time series is required to identify relationships and we are restricted to 35 years of footprint data.

Referee comment:

Page 8, lines 20-21: the conclusion to be drawn from lower total A20 and higher number of events is the mean A20 per event is lower. No conclusion can be drawn about very damaging windstorms, since they are a very small part of this particular A20 metric (see comments about Paris above, where the q-q plot indicates more severe storms in past 15 years).

Response:

Again, this is a very good point. This statement will be removed from the text since this conclusion cannot be taken from Figure 3

However, this does not relate to the q-q plot because, here, extreme is defined as exceeding the 20m/s damage threshold, not the actual wind intensity.

Referee comment:

Page 8, lines 22-24: these two sentences are redundant.

Response:

Again, this statement will be reworded: 'The largest contribution to the decline in A20 between 20th and 21st centuries comes from north-west Europe.'


---

Interactive comment on Nat. Hazards Earth Syst. Sci. Discuss., doi:10.5194/nhess-2016-121, 2016.

[Printer-friendly version](#)

[Discussion paper](#)





**Fig. 1.**

[Printer-friendly version](#)

[Discussion paper](#)

