



# Review article: Potential application of surface methods for the monitoring of organic matter dynamics in marine systems

Galja Pletikapić<sup>1,2</sup>, Nadica Ivošević DeNardis<sup>1</sup>

<sup>1</sup>Division for Marine and Environmental Research, Ruđer Bošković Institute, POB 180, 10000 Zagreb, Croatia

5 <sup>2</sup>FOM Institute AMOLF, POB 41883, 1009 DB Amsterdam, Netherlands (*current address*)

*Correspondence to:* Nadica Ivošević DeNardis (ivoševic@irb.hr)

## Abstract

Marine organic matter has a highly reactive nature; it continuously undergoes biotic and abiotic transformations and consequently exhibits high variation in both time and space, with a tendency to accumulate at the interfaces. To study its role  
10 in natural systems, it is of crucial importance to preserve its original state. We present the potential of non-invasive surface methods, namely, the electrochemical method of chronoamperometry at mercury electrode applied together with atomic force microscopy imaging (AFM), and their suitability on the basis of field studies undertaken in the northern and southern Adriatic Sea (Boka Kotorka Bay, Montenegro). Electrochemical approach meets the requirements for monitoring purposes due to the simple and fast analysis of a large number of raw seawater samples enabling simultaneous differentiation of  
15 organic constituents. On the other hand, atomic force microscopy allows direct visualization of biotic and abiotic particles and provides an insight into structural organization of marine organic matter at micro- and nanometer scales. Application of these methods can contribute to the prediction of mucilage events and to the monitoring of organic pollution in aquatic systems. Moreover, understanding the dynamics of organic matter in seawater is of crucial importance for improvement of national legislation on environmental protection, sustainable development and for management of marine systems.

## 20 1 Introduction

Based on the traditional oceanographic routines, organic matter in the marine environment is classified as dissolved (DOM), consisting of truly dissolved and colloidal fractions, and particulate (POM). DOM is mainly produced by primary production in surface waters (Carlson 2002), and 10 % of it can self-assemble to form nano-, micro- and even macrogels (Verdugo et al., 2004; 2012). Because of the great complexity and heterogeneous composition of marine organic matter, its  
25 characterization remains a challenge. Characterization of DOM in terms of its composition and molecular structure has advanced substantially in the last decade, mainly owing to the development of advanced instrumental analytical methods and



corresponding procedures, for example, Fourier-transform ion cyclotron, mass spectrometry, homo- and hetero-correlated multidimensional nuclear magnetic resonance spectroscopy, and excitation emission matrix fluorimetry with parallel factor analysis for UV-fluorescence spectroscopy, along with advances made in sample preparation (i.e., desalting/extraction techniques, Nebbioso and Piccolo 2013; Roth et al., 2015; Kujawinski et al., 2016; Mopper et al., 2007). In addition, development of particle counters (electronical, electrochemical), histological staining and sophisticated microscopical methods in the last decades helped establish new particle classes, predominantly non-living, which had remained undetected due to their small size, transparency and susceptibility to degradation (Žutić and Svetličić 2000). These particle classes have shown large abundance and reactivity, consequently having a crucial role in aquatic environments, particularly in the cycling of nutrients, aggregation of organic matter and food web (Koike et al., 1990; Wells and Goldberg 1991; 1993; 1994; Alldredge et al., 1993; Long and Azam 1996). These particles are usually classified as: (i) colloidal particles (5-200 nm) collected by ultracentrifugation and observed under transmission electron microscopy, reaching concentrations up to  $10^{12}/L$  at the thermocline of the North Atlantic and Northwest Pacific (Wells and Goldberg 1991; 1993); (ii) submicrometer particles (0.4-1 $\mu$ m) detected by Coulter particle counters reaching concentrations up to  $10^{10}/L$  in the top 40 m of the North Pacific of Japan, Northwest Atlantic Schlep (Koike et al., 1990; Longhurst et al., 1992); (iii) surface-active particles (1-500  $\mu$ m), mostly formed from decomposed and excreted phytoplankton products, can reach concentrations up to  $10^8/L$  in the estuarine mixing zones, the sea surface or halocline of the Adriatic Sea (Žutić et al., 1984; Žutić and Legović 1987; Marty et al., 1988; Svetličić et al., 2005; 2006; Žutić et al., 2004), detected electrochemically. These particles play an important role as precursors of massive aggregate formation; (iv) transparent exopolymeric particles (TEP, 3-100  $\mu$ m) are flexible, polysaccharide microgels serving as surfaces for the bacterial colonization and transfer by adsorption, are detected by Alcian blue staining or spectrophotometrically and may reach concentrations up to  $5 \times 10^6/L$  at the Californian coast (Alldredge et al., 1993; Passow 2002; Villacorte et al., 2015); (v) protein containing particles (CSP, 2-500  $\mu$ m) are visualized by Coomassie blue staining, may reach concentrations up to  $10^8/L$  off the Californian coast (Long and Azam 1996), providing a rich nitrogen source for bacteria and other organisms, and (v) giant aggregates (>1m) observed by a scuba diver in the Northern Adriatic Sea (Stachowitch et al., 1990; Leppard et al., 1997; Svetličić et al., 2005; Gianni et al., 2005a). Recently, more attention has been paid to marine gels ranging from nano to macro dimensions (Verdugo et al., 2004; 2012; Mišić Radić et al., 2011). Marine gels are defined as three-dimensional networks of solvated biopolymers embedded in seawater. Their importance in marine ecosystems has been widely recognized among scientists from different fields due to their role in the microbial loop and sedimentation processes, biogeochemical carbon cycling, marine carbohydrate chemistry, and particle dynamics in the ocean (Žutić and Svetličić 2000; Azam and Long 2000; Passow 2002; Verdugo et al., 2004; 2012). These gels are also bioreactive, since they compartmentalize nutrients and can thus enhance microbial activity (Kepkay, 1994; Azam 1998). Marine gels can exhibit even macroscopic dimensions when massive gelatinous macroaggregates cover a few square kilometers of the sea surface and form clouds in the water column (Fig.1). This can have harmful consequences for the marine systems and cause substantial damage to the economy of the region. Even though evidence accumulated from satellite observations to micro-scale studies and a large number of hypotheses on the mechanism of the event have emerged



over the last decades (Žutić et al., 2004), what triggers such a large-scale phase transformation of organic matter is still a subject of debate.

While relatively broad scientific literature is available on the application of different techniques for characterization of marine organic matter, most of the methods include relatively expensive equipment and/or complex sample preparation procedures that effects the state of the fragile fractions of organic matter. It is now generally agreed that organic matter of a given chemical composition can occur in a large range of forms under the influence of the dynamic aquatic conditions: monomer molecules, polymer chains, globules, vesicles and complex networks such as gels (Žutić and Svetličić 2000). All of these will undoubtedly affect organic matter's persistence, stability, aggregation behaviour and bioavailability. Little is known about the distribution of environmental particles due to the difficulties in sampling, sample handling, insufficient sensitivity and selectivity of analytical methods but also due to the heterogeneity of natural samples (Buffle and Leeuwen 1992). For example, the seawater sample needs to be unperturbed in order to preserve the heterogeneous distribution and primary structure of organic particles. High heterogeneity makes both the analysis and data interpretation challenging. In this review, we present the potentials of non-invasive surface methods, namely, the electrochemical method of chronoamperometry when applied together with atomic force microscopy imaging, and their suitability on the basis of field studies undertaken in the northern and southern Adriatic Sea (Boka Kotorka Bay, Montenegro). We emphasize the importance of no need of sample pretreatment in order to capture the physical structure and organization of organic matter as closely as it is in the natural system. Every sample pretreatment such as filtration, centrifugation, staining, fixation, reverse osmosis, dialysis introduces artefacts. We particularly focus on marine organic particles in the size range of only a few nanometers to several hundreds of micrometers. This fraction mainly involves soft, hydrophobic, hydrophilic and reactive particles, susceptible to degradation. Micrometer sized surface-active particles can be measured electrochemically on the single particle level based on their interfacial properties, structural organization and hydrophobic/hydrophilic character. Electrochemical approach is fast compared to other methods, with sample characterization taking only a couple of minutes, making it suitable for analyses of a large number of raw seawater samples applicable for monitoring. On the other hand, structural organization of organic matter on the molecular level and three dimensional images of fine, fragile and soft structures can be revealed using AFM and correlated to the electrochemical data in order to better understand the biotic as well as abiotic transformation of organic matter at micro and nanoscales as the root of macroscopic giant gel formation in the sea (Ciglenčki and Svetličić, 2015). The advent of AFM opened a possibility to directly explore these processes at the scale that determines the fate of organic matter and its interactions at the interfaces.

## 2 Surface methods for characterization of marine organic matter

### 2.1 Electrochemical method

The electrochemical method used here is chronoamperometry of dissolved oxygen at a charged dropping mercury electrode (Barradas and Kimmerle 1966; Zvonarić et al., 1973). Chronoamperometry at a mercury electrode is an important and



convenient tool for the *in situ* single particle analysis in an aqueous electrolyte solution. Mercury as a substrate has several very unique advantages: it is atomically smooth, fluid and chemically inert with a large set of interfacial data in various aqueous electrolyte solutions. Fluidity of mercury perfectly mimics interactions with other fluid interfaces in contact (e.g. biological membranes, vesicles, bubbles). Electrochemical measurement is performed by immersing the charged dropping mercury electrode directly into the electrochemical vessel containing a fresh seawater aliquot of 15 mL. The mercury electrode has a surface area of only 4 mm<sup>2</sup> with the drop life time of 2 seconds. Each 2 seconds, a new and reproducible mercury surface is formed at the end of the glass capillary and the analysis can be repeated many times. This is an important aspect of the method, since the arrival of particles to the interface is a stochastic process and the representative behaviour can be determined only by analysing a larger set of data collected under identical experimental conditions (Kovač et al., 2000).

The electrochemical method is based on the sensing of the interfacial properties, hydrophobicity and supramolecular organization of particles, rather than on the chemical composition (Žutić et al., 2004). The first advantage of the electrochemical approach is simultaneous and direct detection of different organic constituents in seawater based on their different electrochemical responses (Fig. 2) while inorganic particles remain undetected (Ivošević and Žutić 1997; Baldi et al., 1999; Kovač et al., 1999; Svetličić et al., 2006; Ivošević DeNardis et al., 2007a). Organic constituents of seawater can be classified according to their electrochemical responses at the dropping mercury electrode based on the phenomena of molecular adsorption, adhesion and spreading, and particle collision (Žutić et al., 1990; 1993; Tsekov et al., 1999). On the chronoamperometric curve, adsorption of dissolved organic matter and submicron particles is manifested cumulatively as a gradual decrease of the oxygen reduction current (Žutić et al., 1999). On the other hand, adhesion of fluid surface-active particles with hydrophobic properties is detected on the single particle level through spike-shaped signals (Žutić et al. 1993; Svetličić et al., 2001, Ivošević et al., 2007b). Collision of gel microparticles with the flexible three-dimensional network and thus possessing hydrophilic character is detected also on the single particle level through specific dip-shaped signals (Svetličić et al., 2006; Ivošević DeNardis et al., 2007a). The dropping mercury electrode (DME), as an *in situ* sensor, enables characterization of fluid surface-active particles in the aqueous environment (Žutić et al., 1984; Žutić and Legović 1987; Žutić et al., 2004) in terms of concentration, size, reactivity and hydrophobicity (Žutić et al., 1993; Kovač et al., 2000; Ivošević et al., 2007b). Fig. 3 shows the attractive interaction between an oil droplet and the charged electrode/seawater interface. Oil droplet, as a subclass of surface-active particles, adheres and spreads at the charged interface causing double layer charge displacement and establishment of the flow of the compensating current known as adhesion signal. Signal amplitude reflects the particle size, signal shape reflects the physicochemical properties of the particle and signal frequency indicates particle concentration in the seawater sample. Signal frequency is translated into particle concentration using the corresponding calibration curve with *Dunaliella tertiolecta* cells as standard particles (Kovač et al., 2000). Chronoamperometric measurements meet the requirements for monitoring based on the following advantages (i) simple and fast analysis (100 seconds per sample) of soft micrometer particles in the size and concentration ranges of 1-500 μm and 10<sup>5</sup>-10<sup>8</sup> particles/L, respectively, and (ii) no sample pre-treatment. Recently, an effort has been made to develop a mathematical model that would provide interpretation of the adhesion signal of lipid vesicles and algal cells on the kinetic level, helping



explain the mechanism of adhesion and spreading of soft particles at the interface (Ivošević DeNardis et al., 2012; 2015). On the other hand, determination of adsorbable organic matter (dissolved organic matter and submicron particles) in seawater is based on a smooth and gradual decrease of reduction current proportional to surfactant concentration in the sample, known as surfactant activity (Zvonarić et al., 1973; Žutić et al., 1977). Surfactant activity of seawater is expressed as the equivalent amount of the nonionic synthetic surfactant Triton-X-100 (polyethylene glycol tert-octyl phenyl ether, MW=600) in milligrams per litre. The polarographic maximum of Hg(II) ions offers an alternative approach to measuring dissolved organic carbon of seawater (Hunter and Liss 1981).

## 2.2 Atomic force microscopy (AFM)

Atomic force microscopy has been recently introduced as a tool in marine ecology (Svetličić et al., 2005; Malfatti et al., 2009; 2010; Urbani et al., 2012; Bosak et al., 2012; Svetličić et al., 2013 and references therein). The advantages of AFM over conventional light or electron microscopy are: (i) 3D high resolution imaging of different seawater constituents (biotic and abiotic) connecting micro-meter and nano-metric dimensions, (ii) samples can be non-destructively imaged in both the air and under near natural aqueous conditions, (iii) nanomechanical measurements (Young's modulus (as a measure of elasticity), deformation, hydrophobicity), and (iv) determination of intra- and inter-molecular forces in heterogeneous molecular assemblies (Pletikapić et al., 2014). All these unique advantages make AFM a powerful tool in marine environmental studies, well suited either for investigating the mechanism of marine mucilage formation or for closer morphological description of marine cells (Pletikapić et al., 2012a; Francius et al., 2008). AFM is based on a relatively simple principle; it involves raster scanning of a sharp and hard tip (probe) located at the free end of a flexible cantilever. The tip is scanned over the surface of a sample, sensing the interaction forces between the tip and sample (Fig. 5). The sample is mounted on a piezoelectric scanner, which allows three-dimensional positioning with subnanometer accuracy. Interaction between the tip and the surface of the sample leads to cantilever bending, which is measured by laser light reflected from the cantilever to a position sensitive photodetector. As changes in cantilever deflection result in variation of the distance between the tip and the sample, a constant distance is re-established with a feedback loop between the sample/tip positioning system and a computer controlled piezoelectric scanner. Registered values of cantilever deflection are electronically converted into a pseudo 3D image of the sample. As a result, AFM gives real 3D images of the sample with a vertical resolution of 0.1 nm and lateral resolution of 1 nm.

## 3 Field study in the Northern Adriatic Sea

### 3.1 Monitoring of marine organic matter distribution by electrochemical based detection

A field-oriented study in the Northern Adriatic was carried out in the framework of the long-term National Monitoring Programme (Systematic Study of the Adriatic Sea as a Base for Sustainable Development of the Republic of Croatia, 1998-



2012). The study refers to the monitoring of organic microparticle distribution as an early warning sign for mucilage formation. We introduced a direct electrochemical method to probe the state of marine organic matter without perturbing its original heterogeneous distribution. Data were collected during a period of 10 years and those referring to station SJ 105 are shown in Fig. 4. This particular sampling station was selected because it is distant from the coast line and thus the influence of direct input of inland freshwater is avoided. SAP concentrations ranged from  $5 \times 10^5$  to  $6 \times 10^6$ /L showing spatiotemporal variability with special patterns depending on the sampling depth in the water column and/or season. For example, significantly higher SAP concentrations (about two times) occurred in the region under the direct influence of nutrient-rich freshwater inputs of the Po River (SJ 101, 103, 108). In the first three year period (2002, 2003, 2004), a mucilage event appeared. In the period just before the mucilage event, SAP concentration rise to  $1 \times 10^7$ /L in the subsurface layer (0-10 m, SJ 101, 103, 108), indicating the forthcoming event. In contrast, low concentration of SAP in the subsurface layer at station SJ105 in 2007 is in line with the absence of mucilage events. Hence, SAP was identified as a reactive precursor for mucilage formation in seawater. Concentrations of GeP were usually around  $10^5$ /L in the years without mucilage events ( $0.2 \times 10^5$ /L is lower electrochemical detection limit). Elevated concentration of GeP, by about one order of magnitude ( $15 \times 10^6$ /L), was determined in all sampling stations at the time of mucilage events (2002-2004). Abundance of GeP was at average 10% of SAP values (Project Jadran). Surfactant activity reflecting the concentration of submicron and dissolved organic fractions varied in the range from 0.5 to 2 mg/L in the years without mucilage events and decreased with the depth profile. Elevated surfactant activity up to 3.4 mg/L was determined during the mucilage event. In addition, a net increase of surfactant activity in the whole water column was observed at the end of years without mucilage events as a consequence of prolonged accumulation of organic matter in the water column. It was therefore concluded that the likelihood of mucilage formation depends mainly on new primary production, specific structural organization of the newly produced organic matter and, above all, hydrographic and meteorological conditions in the seawater (Project Jadran).

### 3.2 Structural organization of marine organic matter at the nanometric level visualized by AFM

25 In one of the first AFM studies of marine organic matter, Santschi and coworkers (1998) used AFM to image individual fibrillar polysaccharides in marine macromolecular organic matter. From there on, potential AFM has been recognized to study both the networks secreted by marine bacteria (Malfatti et al., 2009) and marine fibrils forming giant gel macroaggregates formed in the Northern Adriatic (Volleinder et al., 1999; Giani et al., 2005b), their self-assembly and molecular structure at different levels of association (Mišić Radić et al., 2011). However, application of AFM imaging for monitoring by itself is often not suitable. This is due to the large number of samples required in the monitoring studies, making AFM analysis, if used alone, a time consuming process. It is therefore recommended to analyse seawater samples first with some more rapid and “bulk” technique (i.e. chronoamperometry) in order to identify samples of interest. For AFM imaging, the drop deposition method modified for marine samples and imaging in air were found to be most adequate for visualization of macromolecular organic matter organization (Pletikapić et al., 2011; Mišić Radić et al., 2011 for more



detail). Just a drop of seawater (5 $\mu$ L) deposited on freshly cleaved mica substrate is enough to reveal the structural organization of organic matter after rinsing of sea salts and evaporation of water excess. Representative images of marine fibril forming networks with different levels of biopolymer entanglements and with different pore sizes are shown in Fig. 6. It is interesting to note that the appearance of marine gel networks at the nanometric scale, as shown in the corresponding Figure coincided with the occurrence of gel microparticles detected electrochemically, as shown in Fig. 4, at the same sampling time. Image analysis revealed that biopolymer heights were in the 0.5 nm - 2.5 nm range, while pore sizes ranged from  $\sim$  40 nm to more than 1 $\mu$ m along the longest axis. Svetličić et al., 2011, found that these parameters (pore size, fibril height) can be related to the aging of gel aggregates in the water column, i.e. older aggregates contained smaller pores and thicker fibrils, while “young” aggregates had a looser structure with smaller fibrils and bigger pore sizes. In addition, gel network formation was visualized at station SJ 107 in August 2010, when concentrations of micrometer sized GeP and SAP were low (0.2x10<sup>6</sup>/L and 2.3x10<sup>6</sup>/L, respectively). Significantly higher concentration of submicron organic content (surfactant activity, 2.88 mg/L) was detected 2 months prior to the mucilage event at the depth of 10 m, in 2010, which could act as a pool for biopolymer self-organization towards gel network formation. One of the most comprehensive chemical characterization studies of Northern Adriatic gel macroaggregates showed that carbohydrates are the main organic fraction and that the average ratio of polysaccharides to total organic carbon is 41.5 % (Gianni et al., 2005b). Exopolysaccharides released by the diatom *Cylindrotheca closterium* under the conditions of increased production comprised a simpler system, which had the same chemical composition of polysaccharides as marine gels, indicating that the diatom exopolymeric substance was the most likely origin of marine gels (Urbani et al., 2005). In addition, striking similarity was found in both marine gels and reconstituted polysaccharide networks (obtained from the same diatom) in terms of their mechanical properties as well as in the morphological features (Pletikapić et al., 2014). Marine gels were characterized in the non-imaging mode of AFM, the so called “fishing”, where single polymers and/or their more complex associations are “picked up from the surface (fished) by the AFM probe and extended and/or unfolded/unzipped by increasing the distance of the AFM tip from the surface. This approach enabled relating mechanical properties to different biopolymer morphologies (from single fibril, helical structures to more complex junction zones in networks with a higher degree of crosslinking), revealing new information about network architecture and connectivity. Networks were characterized as physical gels in line with DCS measurements (Mišić et al., 2011), where the level of association can be controlled by dilution, adjusting the ionic strength and stirring (Pletikapić et al., 2014). Due to their dense 3D network structure and “sticky” nature, marine gels show a potential to scavenge nanoparticles that accumulate in the junction zones of the network (Svetličić et al., 2011; Pletikapić et al., 2012b). In this way, marine gels might stabilize nanoparticles and prevent aggregation and export from the water column. As native marine gels are constantly exposed to mechanical perturbations caused by local hydrodynamics (turbulence and shear) as well as large-scale physical processes (strong winds and currents), the application of hundreds of piconewtons of force is highly biologically relevant. It is expected that AFM will reveal new classes of submicron particles in the future, and contribute to the understanding of the dynamic process of transformation and biogeochemical cycling of organic matter in marine systems. Vesicular structures are another structural form of organic matter in seawater (Fig 6).



Existence of vesicle-like assemblies has already been inferred by Nagata and Kirchman (1997) but vesicles were for the first time electrochemically detected in seawater samples (Žutić et al., 1984). Recently, it was reported that membrane vesicles of 100 nm in diameter can be released by marine cyanobacteria and are frequently found in marine ecosystems. Such vesicles may entrap nutrients, toxins, concentrate and preserve them in the inner cavity and in the hydrophobic membrane variety of organic molecular species present in seawater (Biller et al., 2014). This ability of vesicles to deliver diverse compounds in discrete packages adds another layer of complexity to the flow of information, energy and biomolecules in marine microbial communities. For identification of vesicle like structures, AFM can be a very suitable tool due to the high resolution imaging and easy sample preparation. Such vesicles (Fig. 7) were also periodically encountered during the diatom bloom experiment and during the Northern Adriatic field studies by AFM (Svetličić et al., 2008). It is likely that the abundance of such structures is highly underestimated. Due to the small size and relatively soft and fragile character of these structures, during AFM imaging with relatively higher forces, particularly when contact mode is used, vesicles are swept away by the AFM tip and remain undetermined. Proper care while setting imaging parameters, scanning at low force (i.e. Peak Force Tapping mode) and coupling AFM with other bulk methods such as electrochemistry to identify the sample of interest with higher concentrations of vesicles may resolve the issue. Such an approach was used to study vesicle like particles in the Boka Kotorska Bay (Ivošević DeNardis et al., 2013; 2014). The most important findings show that both surface methods, electrochemical and AFM, provide comparable results on different scales. To summarize, application of AFM opens a possibility to directly explore the main organic assemblies found in seawater, their molecular organization and stability, providing new data and enhancing our understanding of the gel formation mechanism and the role of marine gels in ocean biogeochemical cycles. This information is difficult to access by conventional analytical methods. However, despite all, AFM is not yet exploited up to its full potential. Recent progress has been made by the construction of a new generation of soft probes using the advantage of mercury as an atomically smooth, fluid and conductive surface in order to properly characterize soft surfaces like biomembranes under physiological conditions (Schon et al., 2013).

### 3.4 A field case study in the Southern Adriatic Sea (Boka Kotorska Bay, Montenegro): Oil spill incident

The field case study was undertaken due to the recent accidental sinking of a ship and spilling of diesel fuel in the bay of Boka Kotorska (Fig. 8, Ivošević DeNardis et al., 2014). The ship had an overall length of 32.7 meters and weighed 325 tons, carrying oil and diesel oil (D2). The ship sank on October 11, 2013 at Rt Murva (15 m depth). One month after the mechanical removal of oil spill, dispersed oil droplets were still present in the whole water column. Characterized oil droplets were in the continuum size range from micro to nano. Smaller oil droplets tend to accumulate at the halocline, while larger rise to the surface where they may coalesce. It was electrochemically detected that the highest concentration of micrometer droplets was  $2.0 \times 10^5/L$  at the 5 m depth (i.e., at the density gradient), while droplet concentrations were below detection limit at other depths. Dispersed droplets were in the size range from 1  $\mu\text{m}$  up to 30  $\mu\text{m}$  in measured samples (determined by microscopy and Nile red staining). Bacterial colonization of micrometer sized fuel droplets indicated that a





biodegradation process occurred, as revealed by epifluorescence microscopy and DAPI staining (Fig. 9a). Hydrocarbon-degrading bacteria could account for up to 10 % of the total bacterial population (Hassanshahian and Cappello 2013), thus the biodegradation process is rather slow. Hydrocarbon-degrading bacteria utilize oil droplets as a carbon and energy source. Bacterial efficiency in the degradation of diesel fuel droplets can be measured as a change in droplet concentration, distribution, size, bacterial film formation, bacterial oxygen consumption and through chemical changes of oil composition (Baldi et al., 1997; 1999). On the other hand, atomic force microscopy enables visualization of oil droplets in the seawater sample at the nanometer level. Larger and smaller nanometer sized oil droplets spread over the mica substrate forming spherical caps, as shown in Figs. 9b and c. The imaged droplet diameter in seawater would correspond to 520 nm and 80 nm, respectively (Ivošević DeNardis et al., 2014). Nanometer sized droplets were detected in the whole water column as reactive and stable fractions that persist longer and can therefore cause a significant decrease of phytoplankton primary production. Recovery of phytoplankton variability and abundance after a heavy fuel oil spill in the Marmara sea, Turkey, required a time period of three years (Tas and Erdogan, 2007). The consequences of anthropogenically caused oil spills are known to provoke serious ecosystem disturbances influencing the variability in planktonic dynamics, survival of fish eggs, larvae, shellfish, hatching, swimming activity, fertility in copepods, and overall trophic interactions. For example, dispersed diesel oil had a harmful effect on the cardiac activity of Mediterranean mussels (Bakhmet et al., 2009; Martinović et al., 2015) important for adequate protection of shellfish farms, breeding control and preventing food poisoning of humans. Monitoring of water quality should be recommended due to the intensive boat trafficking and shipyard activity in the Boka Kotorska Bay in order to decrease the risk to the sensitive marine ecosystem. Boka Kotorska Bay is a semiclosed basin of Montenegro characterized by karst rivers and underground springs, which affect the temperature, density and salinity of seawater (Campanelli et al., 2009). To understand the fate and behaviour of dispersed diesel fuel under the complex hydrodynamic conditions requires a comprehensive and interdisciplinary approach. Supplementary modeling data of the corresponding geographical area would be of benefit in order to assess the scale of oil spreading and trajectory, important for emergency responses and decision matters. Such approach can help understand the scale of the problem and efficient coordination between local authorities, research and industry in order to enable environmental protection as well as sustainable development.

#### 4 Conclusions

Organic matter in the water column has a highly reactive nature; it continuously undergoes biotic and abiotic transformations and consequently exhibits high variation in time and space, with a tendency to accumulate at the interfaces. To study the role that organic matter plays in marine systems, it is of crucial importance to characterize it in such a way as to preserve its original state. Hence, the electrochemical method of amperometry at mercury electrode meets the requirements of non-invasive, simple and fast analysis of raw aquatic samples, enabling simultaneous differentiation of organic constituents. On the other hand, atomic force microscopy allows direct visualization of biotic and abiotic particles in seawater at micro- and



nanometer scales without the need of vacuum and complex sample preparation. A combination of complementary methods enables us to set a continuous scale description of marine organic matter from the micrometer to nanometer scale. The first field oriented study showed how distribution of organic microparticles in the Northern Adriatic can be related to mucilage formation. A few months prior to the mucilage event, the concentration of accumulated surface-active particles in the halocline was  $1 \times 10^7/L$ , supported by certain meteorological and hydrodynamics conditions, and a large scale sol to gel organic matter phase transformation occurred. Such microparticles were therefore identified as reactive precursors of macroaggregate formation in seawater. Formation of a gel network at the nanometric scale coincided with elevated concentrations of flexible gel microparticles or surfactant activity prior or after the macroaggregation event. The second field oriented study showed the potential of surface methods for testing the cleanup performance after oils spill events during accidental sinking of the ship in the Boka Kotorska Bay (Montenegro). In spite of the mechanical removal of the major oil spill content, the remaining micro and nanodroplets detected in the collected sample represent a threat for the nearby mussel farms, fisheries and tourism. Possible applications of these surface methods include prediction of mucilage events and monitoring of organic pollution. Interdisciplinary approach and long-term measurements of relevant parameters are required in order to better understand the spatiotemporal distribution of organic matter in aquatic systems and consequently its role in the biogeochemical cycles. In the future, it is expected that integration of experimental data (on nanometer and micrometer scales) with mathematical models (on kilometer scale), (taking into account meteorological data, oceanographic conditions, circulation patten, satellite data) could contribute to a better understanding of organic matter dynamics in aquatic systems as well as to evaluation of the ecological status of the corresponding system. Potential application of surface methods to the monitoring of organic matter dynamics in marine systems would be of great benefit to environmental protection, development and management of the coastal area, especially for tourism, fishery and cruiser trafficking.

### *Acknowledgments*

This work was supported by the (i) Croatian Ministry of Science, Education and Sports through the projects Surface Forces on Atomic Scale Applied in Marine Science and Nanotechnology, (ii) National Monitoring Programme (Project Jadran), Systematic Research of the Adriatic Sea as a Base for Sustainable Development of the Republic of Croatia, (iii) Croatia-Montenegro bilateral scientific cooperation Impact assessment and determination of organic pollutants in the waters (project leaders: Zoran Kljajić and NID). The authors thank V. Žutić and V. Svetličić for coordinating the organic microparticle study in the framework of the Project Jadran. We thank many coworkers who contributed to the activities described in this review. This review is dedicated to the memory of the late Dr. Zoran Kljajić of the Institute of Marine Biology, Kotor, Montenegro.



## References

- Allredge, A.L., Passow, U., and Logan, B. E.: The abundance and significance of a class of large, transparent organic particles in the ocean, *Deep-Sea Res.*, I 40, 1131-1140, doi: 10.1016/0967-0637(93)90129-Q, 1993.
- Azam, F.: Microbial control of oceanic carbon flux: The plot thickens, *Science*, 280, 694-696, doi: 10.1126/science.280.5364.694, 1998.
- Azam, F. and Long, A.R.: Sea snow microcosmos, *Nature*, 414, 495-498, doi:10.1038/35107174, 2000.
- Bakhmet, I. N., Fokina, N. N., Nefedova, Z. A., and Nemova, N. N.: Physiological-biochemical properties of blue mussel *Mytilus edulis* adaptation to oil contamination, *Environ. Mon. Assess.*, 155, 581–591, doi:10.1007/s10661-008-0457-5, 2009.
- Baldi, F., Pepi, M., Fani, R., Di Cello, F., Da Ros, L., and Fossato, V. U.: Complementary degradation of fuel oil in superficial waters and in axenic cultures of aerobic Gram-negative bacteria isolated from Venice Lagoon, *Croat. Chem. Acta*, 70, 333–346, 1997.
- Baldi, F., Ivošević, N., Minacci, A., Pepi, M., Fani, R., Svetličić, V., and Žutić, V.: Adhesion of *Acinetobacter venetianus* to diesel fuel droplets studied by in situ electrochemical and molecular probes, *Appl. Environ. Microbiol.*, 65, 2041-2048, 1999.
- Barradas R. G. and Kimmerle, F. J.: Effect of highly surface- active compounds on polarographic electrode processes, *J. Electroanal. Chem.*, 11, 163–170, doi:10.1016/0022-0728(66)80079-7, 1966.
- Biller, S. J., Schubotz, F., Roggensack, S. E., Thompson, A.W., Summons, R. E., and Chisholm, S. W.: Bacterial vesicles in marine ecosystems, *Science*, 10, 183-186, doi: 10.1126/science.1243457, 2014.
- Bosak, S., Pletikapić, G., Hozić, A., Svetličić, V., Sarno, D., and Viličić, D.: A novel type of colony formation in marine planktonic diatoms revealed by atomic force microscopy, *PLoS One*, 7, e44851, doi:10.1371/journal.pone.0044851, 2012.
- Buffle, J. and van Leeuwen, H. P.: *Environmental particles*, Lewis publishers, London, 1992.
- Campanelli, A., Bulatović, A., Cabrini, M., Grilli, F., Kljajić, Z., Mosetti, R., Paschini, E., Pierluigi, P., and Marini, M.: Spatial distribution of physical, chemical and biological oceanographic properties, Phytoplankton, nutrients and Coloured Dissolved Organic Matter (CDOM) in the Boka Kotorska Bay (Adriatic Sea), *Geofizika*, 26, 1–14, 2009.
- Carlson, C.A.: Production and removal processes, in: *Biogeochemistry of marine dissolved organic matter* Hansell, D. A. and Carlson, C. A. (eds.), Academic Press, San Diego, 91–151, 2002.
- Ciglencčki, I. and Svetličić, V.: Nanoparticles and Marine Environment: An Overview, in: Camesano, T. A., (ed.), *Nanotechnology to Aid Chemical and Biological Defense*, Springer International Publishing, Netherlands, 95-112, 2015.
- Croatian National Monitoring Programme (Project Jadran), *Systematic Research of the Adriatic Sea as a Base for Sustainable Development of the Republic of Croatia (1998-2012)*, Ministry of science, education and sport, Republic of Croatia.
- Francius, G., Tesson, B., Dague, E., Martin-Jezequel, V., and Dufrene, Y. F.: Nanostructure and nanomechanics of live *Phaeodactylum tricornutum* morphotypes, *Environ. Microbiol.*, 10,1344– 1356, doi: 10.1111/j.1462-2920.2007.2008.



- Giani, M., Degobbi, D., and Rinaldi, A.: Mucilages in the Adriatic and Tyrrhenian seas, *Sci. Total. Environ.*, 353,1-380, doi:10.1016/j.scitotenv.2005.09.006, 2005a.
- Giani, M., Berto, D., Zangrando, V., Castelli, S., Sist, P., and Urbani, R.: Chemical characterization of different typologies of mucilaginous aggregates in the Northern Adriatic Sea, *Sci. Total. Environ.*, 353, 232–246, doi:10.1016/j.scitotenv.2005.09.027, 2005b.
- Hassanshahian, M., and Cappello, S.: Crude oil Biodegradation in the marine environments in biodegradation-engineering and technology, Chamy, R. and Rosenkranz, F. (eds) InTech, Rijeka, Croatia, 101-135, doi.org/10.5772/55554, 2013.
- Hunter, K. A. and Liss, P. S.: Polarographic measurement of surface-active material in natural waters, *Water Res.*, 15, 203-215, doi: 10.1016/0043-1354(81)90113-5, 1981.
- 10 Ivošević, N. and Žutić, V.: Polarography of marine particles: A model study, *Croat. Chem. Acta*, 70, 167-178, 1997.
- Ivošević DeNardis, N., Žutić, V., Svetličić, V., and Urbani, R.: In situ amperometric detection of vesicles and microgel phases in an aggregating system: Calcium Alginate, *Electroanalysis*, 19, 473-478, doi: 10.1002/elan.200603750, 2007a.
- Ivošević DeNardis, N., Žutić, V., Svetličić, V., Frkanec, R., and Tomašić, J.: *In situ* amperometric characterization of liposome suspensions with concomitant oxygen reduction, *Electroanalysis*, 19, 2444-2450, doi: 10.1002/elan.200703999, 15 2007b
- Ivošević DeNardis, N., Ružić, I., Pečar Ilić, J., El Shawish, S., and Zihel, P.: Reaction kinetics and mechanical models of liposome adhesion on charged interface, *Bioelectrochemistry*, 88, 48-56, doi: 10.1016/j.bioelechem.2012.05.003, 2012.
- Ivošević DeNardis, N., Šegota, S., Svetličić, V., Castelli, A., and Kljajić, Z.: Characterization of marine organic matter in the bay of the Boka Kotorska by electrochemical and atomic force microscopy imaging, *Stud. Mar.*, 26, 5-22, 2013.
- 20 Ivošević DeNardis, N., Šegota, S., Svetličić, V., Pletikapić, G., and Kljajić, Z.: Presence of dispersed diesel fuel in water column in the Boka Kotorska bay: A case study, *Stud. Mar.*, 27, 43-64, 2014.
- Ivošević DeNardis, N., Pečar Ilić, J., Ružić, I., and Pletikapić, G.: Cell adhesion and spreading at a charged interface: Insight into the mechanism using surface techniques and mathematical modelling, *Electrochim. Acta*, 176, 743-754, doi: 10.1016/j.electacta.2015.07.068, 2015.
- 25 Kepkay, P.: Particle aggregation and the biological reactivity of colloids, *Mar. Ecol. Prog. Ser.*, 109, 293–304, doi: 10.3354/meps111293, 1994.
- Koike, I., Hara, S., Terauchi, K., and Kogure, K.: Role of sub-micrometre particles in the ocean, *Nature*, 345, 242-244, doi: doi: 10.1038/345242a0, 1990.
- Kovač, S., Svetličić, V., and Žutić V.: Molecular adsorption vs. cell adhesion at an electrified aqueous interface, *Colloids Surf. A*, 149, 481–489, doi: 10.1016/S0927-7757(98)00519-6, 1999.
- 30 Kovač, S., Kraus, R., Geček, S., and Žutić, V.: Cell suspension as a model system for electrochemical analysis, *Croat. Chem. Acta*, 73, 279-291, 2000.



- Kujawinski, E. B., Longnecker, K., Barott, K. L., Weber R. J. M., and Kido Soule M.C.: Microbial community structure affects marine dissolved organic matter composition, *Front. Mar. Sci.*, 3, 1-15, doi:10.3389/fmars.2016.00045, 2016. |
- Leppard, G. G., West, M. M., Flannigan, D. T., Carson, J., and Lott, J. N. A.: A classification scheme for marine organic colloids in the Adriatic Sea: colloid speciation by transmission electron microscopy, *Can. J. Fish. Sci.*, 54, 2334-2349, 10.1139/f97-140, 1997.
- 5 Long, R. A. and Azam, F.: Abundant protein-containing particles in the sea, *Aquat. Microb. Ecol.*, 10, 213-221, doi: 10/a010p213, 1996.
- Longhurst, A. R., Koike, L., Li, W., Rodriguez, J., Dickie, P., Kepkay, P., Partensky, F., Bautista, B., Ruiz, J., Wells, M. L. and Bird, D.: Sub-micron particles in north-west Atlantic shelf water, *Deep-Sea Res., Rapid Response Paper*, 39, 1-7, 1992.
- 10 Malfatti, F. and Azam, F.: Atomic force microscopy reveals microscale networks and possible symbioses among pelagic marine bacteria, *Aquat. Microb. Ecol.*, 58, 1-14, doi: 10.3354/ame01355, 2009.
- Malfatti, F., Samo, T. J., and Azam, F.: High-resolution imaging of pelagic bacteria by atomic force microscopy and implications for carbon cycling, *ISME J.*, 4, 427-439, doi:10.1038/ismej.2009.116, 2010.
- Martinović, R., Gačić, Z., and Kljajić, Z.: The Influence of oil, dispersed oil and the oil dispersant SD-25 on the heart rate of the Mediterranean mussel (*M. galloprovincialis* L.), in: Sustainable Development of Sea-Corridors and Coastal Waters, Stylios, C., Floqi, T., Martinski, J., and Damiani L. (eds.), Springer International Publishing, Switzerland, 21-27, doi: 10.1007/978-3-319-11385-2\_2, 2015.
- 15 Marty J-C., Žutić V., Precali R., Saliot A., Čosović B., Smoldaka N., and Cauwet G.: Organic matter characterization in the Northern Adriatic Sea with special reference to the sea surface microlayer, *Mar.Chem.*, 26: 313-330, doi:10.1016/0304-4203(89)90037-6, 1988.
- 20 Mišić Radić, T., Svetličić, V., Žutić, V., and Boulgaropoulos, B.: Seawater at the nanoscale: marine gels imaged by atomic force microscopy, *J. Mol. Recognit.*, 24, 397-405, doi: 10.1002/jmr.1072, 2011.
- Mopper, K., Stubbins, A., Ritchie, J. D., Bialk, H. M., and Hatcher, P. G.: Advanced instrumental approaches for characterization of marine dissolved organic matter: extraction techniques, mass spectrometry, and nuclear magnetic resonance spectroscopy, *Chem. Rev.*, 107, 419-442, 2007.
- 25 Nagata, T. and Kirchman, D. L.: Roles of submicron particles and colloids in microbial food webs and biogeochemical cycles within marine environments, *Adv. Mar. Biol.*, 15, 81-103, 1997.
- Nebbioso, A. and Piccolo, A.: Molecular characterization of dissolved organic matter (DOM): a critical review, *Anal. Bioanal. Chem.*, 405, 109-124, doi: 10.1007/s00216-012-6363-2, 2013.
- 30 Passow, U.: Transparent exopolymer particles (TEP) in aquatic environments, *Prog. Oceanogr.*, 55, 287-333, doi:10.1016/S0079-6611(02)00138-6, 2002.
- Pletikapić, G., Mišić Radić, T., Hozić Zimmermann, A., Svetličić, V., Pfannkuchen, M., Marić, D., Godrijan, J., and Žutić, V.: Extracellular polymer release AFM imaging of extracellular polymer release by marine diatom *Cylindrotheca closterium* (Ehrenberg) Reiman & J.C. Lewin, *J. Mol. Recognit.*, 24, 436-445, doi: 10.1002/jmr.1114, 2011.



- Pletikapić, G., Berquand, A., Mišić Radić, T., and Svetličić, V.: Quantitative nanomechanical mapping of marine diatom, *J. Phycol.*, 48, 174–185, doi: 10.1111/j.1529-8817.2011.01093, 2012a.
- Pletikapić, G., Vinković Vrček, I., Žutić, V., and Svetličić, V.: Atomic force microscopy characterization of silver nanoparticles interactions with marine diatom cells and extracellular polymeric substance, *J. Mol. Recognit.*, 25, 309–317, doi: 10.1002/jmr.2177, 2012b.
- Pletikapić, G., Svetličić, V., Lannon, H., Murvai, U., Kellermayer, M., and Brujić, J.: Self-assembly of polysaccharides gives rise to distinct mechanical signatures in marine gels, *Biophys. J.*, 107, 355–364, doi:10.1016/j.bpj.2014.04.065, 2014.
- Roth, V-N., Dittmar, T., Gaupp, R., and Gleixner, G.: The Molecular composition of dissolved organic matter in forest soils as a function of pH and temperature, *Plos One*, 1-23, doi:10.1371/journal.pone.0119188, 2015.
- Stachowitsch M., Fanuko N. and Richter M.: Mucus aggregates in the Adriatic Sea: An overview of stages and occurrences. *Mar.Ecol.* 11, 327-350, doi: 10.1111/j.1439-0485.1990.tb00387, 1990.
- Santschi, P. H., Balnois, E., Wilkinson, K. J., Zhang, J., and Buffle, J.: Fibrillar polysaccharides in marine macromolecular organic matter as imaged by atomic force microscopy and transmission electron microscopy, *Limnol. Oceanogr.*, 43, 896–908, doi: 10.4319/lo.1998.43.5.089, 1998.
- Schön, P., Geerlings, J.; Tas, N., and Sarajlic, E.: AFM Cantilever with in situ renewable mercury microelectrode, *Anal. Chem.*, 85, 8937-8942, doi: 10.1021/ac400521p, 2013.
- Svetličić, V., Ivošević, N., Kovač, S., and Žutić, V.: Charge displacement by adhesion and spreading of a cell, *Bioelectrochemistry*, 53, 79-86, doi:10.1016/S0302-4598(00)00115-X, 2001.
- Svetličić, V., Žutić, V., and Hozić Zimmermann, A.: Biophysical scenario of giant gel formation in the northern Adriatic sea, *Ann. N.Y. Acad. Sci.*, 1048, 524–527, doi: 10.1196/annals.1342.082, 2005.
- Svetličić, V., Balnois, E., Žutić, V., Chevalet, J., Hozić Zimmermann, A., Kovač, S., and Vdović, N: Electrochemical detection of gel microparticles in seawater, *Croat. Chem. Acta*, 79, 107-113, 2006.
- Svetličić, V., Žutić, V., Hozić Zimmermann, A., and Mišić, T.: Marine vesicles: Adhesion based detection of vesicles-like microparticles in seawater, *Biophys. J.*, 94 (S) 655a-655a, 2008.
- Svetličić, V., Žutić, V., Mišić Radić, T., Pletikapić, G., Hozić Zimmermann, A., and Urbani R.: Polymer networks produced by marine diatoms in the northern Adriatic sea, *Mar. Drugs*, 9, 666–679, doi:10.3390/md9040666, 2011.
- Svetličić, V., Žutić, V., Pletikapić, G., and Mišić Radić, T.: Marine polysaccharide networks and diatoms at the nanometric scale, *Int. J. Mol. Sci.*, 14, 20064-20078, doi:10.3390/ijms141020064, 2013.
- Tas, S. and Erdogan, O.: Effects of oil pollution on the phytoplankton community in the küçükcekece bay (north-east sea of Marmara, Turkey), *Rapp. Comm. Int. Mer. Medit.*, 38, 320-320, 2007.
- Tsekov, R., Kovač, S., and Žutić, V.: Attachment of oil droplets and cells. on dropping mercury electrode, *Langmuir*, 15, 5649–5653, doi:10.1021/la980944q, 1999.



- Urbani, R., Magaletti, E., Sist, P., and Cicero, A. M.: Extracellular carbohydrates released by the marine diatoms *Cylindrotheca closterium*, *Thalassiosira pseudonana* and *Skeletonema costatum*: effect of P-depletion and growth status, *Sci. Total. Environ.*, 353, 300-306, doi:10.1016/j.scitotenv.2005.09.026, 2005.
- Urbani, R.; Sist, P.; Pletikapić, G.; Mišić Radić, T.; Svetličić, V., and Žutić, V.: Diatom polysaccharides: Extracellular  
5 reduction, isolation and molecular characterization, in: *The complex world of polysaccharide*, Karunaratn, D.N. (ed.);  
Intech: Rijeka, Croatia, 346–356, doi: 10.5772/51251, 2012
- Verdugo, P., Alldredge, A. L., Azam, F., Kirchman, D. L., Passow, U., and Santschi, P.H.: The oceanic gel phase: a bridge  
in the DOM-POM continuum, *Mar. Chem.*, 92, 67–85, doi: 2004.
- Verdugo, P., Alldredge, A. L., Azam, F., Kirchman, D. L., Passow, U., and Santschi, P. H.: The oceanic gel phase: a bridge  
10 in the DOM–POM continuum, *Mar. Chem.*, 92, 67–85, doi:10.1016/j.marchem.2004.06.017, 2004.
- Verdugo, P.: Marine microgels, *Annu. Rev. Mar. Sci.*, 4, 375–400, doi: 10.1146/annurev-marine-120709-142759, 2012.
- Villacorte, L.O., Ekowati, Y., Calix-Ponce, H. N., Schippers, J.C., Amy, G. L. and Kennedy, M.D.:  
Improved method for measuring transparent exopolymer particles (TEP) and their precursors in fresh and saline water,  
*Water Res.*, 70, 300-312, doi: 10.1016/j.watres.2014.12.012, 2015.
- 15 Vollenweider, R. and Rinaldi, A.: Marine mucilages., *Sci. Total. Environ.*, 165, 1–235, doi:10.1016/0048-9697(95)90002-0,  
1999.
- Zvonarić, T., Žutić, V., and Branica, M.: Determination of surfactant activity by polarography, *Thalassia Jugosl.*, 9: 65-75,  
1973.
- Žutić, V., Ćosović, B., and Kozarac, Z.: Electrochemical determination of surface active substances in natural waters, *J.*  
20 *Electroanal. Chem.*, 78, 113-121, doi:10.1016/S0022-0728(77)80427-0, 1977.
- Žutić, V., Pleše, T., Tomaić, J., and Legović, T.: Electrochemical characterization of fluid vesicles in natural waters, *Mol.*  
*Cryst. Liq. Cryst.*, 113, 131–145, doi:10.1080/00268948408071678, 1984.
- Žutić, V. and Legović, T.: A film of organic matter at the fresh-water/sea-water interface of an estuary, *Nature*, 328, 612–  
614, doi:10.1038/328612a0, 1987.
- 25 Žutić, V., Svetličić, V., and Tomaić, J.: Dissolved and dispersed organic matter in natural waters: Progress by  
electroanalysis, *Pure Appl. Chem.*, 62, 2269-2276, doi: 10.1351/pac199062122269, 1990.
- Žutić, V., Kovač, S., Tomaić, J., and Svetličić, V.: Heterocoalescence between organic microdroplets and charged  
conductive interface, *J. Electroanal. Chem.*, 349, 173-186, doi:10.1016/0022-0728(93)80171-D, 1993.
- Žutić, V., Ivošević, N., Svetličić, V., Long, R. A., and Azam, F.: Film formation by marine bacteria at a model fluid  
30 interface. *Aquat. Microb. Ecol.*, 17, 231-238, doi:10.3354/ame017231, 1999.
- Žutić, V. and Svetličić, V.: Interfacial processes, in: *The Handbook of environmental chemistry*, Wangersky, P. (ed.),  
Springer-Verlag: Berlin, Germany, 149–165, 2000.
- Žutić, V., Svetličić, V., Ivošević, N., Hozić, A., and Pečar, O.: Northern Adriatic mesocosm experiment Rovinj 2003:  
Dynamics of organic microparticles studied by the electrochemical technique, *Period. Biolog.*, 106, 67-74, 2004.




Wells, M. L. and Goldberg, E. D.: Occurrence of small colloids in sea water, *Nature*, 353, 342-344, doi:10.1038/353342a0, 1991.

Wells, M. L. and Goldberg, E. D.: Colloid aggregation in seawater, *Mar. Chem.*, 41, 353-358, doi: 10.1016/0304-4203(93)90267-R, 1993.

5 Wells, M. L. and Goldberg, E. D.: The distribution of colloids in the North Atlantic and Southern Oceans. *Limnol. Oceanogr.*, 39, 286-302, doi: 10.4319/lo.1994.39.2.0286, 1994.


10



**Figure 1: Macroaggregation phenomena in the Northern Adriatic (a) giant gel aggregate at 10 m depth as captured by a scuba-diver in August 1997 (courtesy of Gerald Müller-Niklas), and (b) map of the study area (inset) and sampling stations in the**


15 Northern Adriatic.





**Figure 2: Illustration of different classes of organic constituents in seawater and their electrochemical signals at the electrode: biopolymer and small colloids, surface-active particles and gel particles (Svetličič et al., 2006).**

5



10

**Figure 3: Basic principle of electrochemical adhesion based detection: Interaction of oil droplets with the charged electrode interface.**




Figure 4: Annual variation of concentrations of: surface-active particles (SAP), gel microparticles (GeP) and surfactant activity determined by electrochemical analysis of raw seawater samples taken at sampling station SJ 105 in the Northern Adriatic (Project Jadran).






Figure 5: Basic principle of atomic force microscopy.

5




10 Figure 6: AFM topographic images of marine gel representing networks with (a) a low degree of fibril crosslinking and with (b) a high degree of crosslinking with the corresponding cross-sections along the indicated lines. Marine gel samples were collected in the Northern Adriatic (SJ 105) in June 23, 2003, at a depth of 10-13 m (Mišić Radić et al., 2011).



**Figure 7:** Large marine vesicle of 1.5  $\mu\text{m}$  diameter imaged by AFM with the corresponding cross-section along the indicated line. Samples were collected in the Northern Adriatic (SJ 101) in May 18, 2010, at a depth of 0 m.


5



10

**Figure 8:** Sampling station of Rt Murva (●) in the bay of Boka Kotorska (Montenegro, inset: study area, southern Adriatic Sea, adopted from Ivošević DeNardis et al., 2014).

15



**Figure 9:** Dispersed fuel oil droplets observed (a) under an optical microscope with DAPI staining, (b) three-dimensional views of typical nanometer sized fuel oil droplets in the seawater sample taken at the Rt Murva depth of 0 m and (c) depth of 5 m visualized by AFM with the corresponding vertical profiles. Vertical profile enables to calculate oil droplet diameter in seawater (adopted from Ivošević DeNardis et al., 2014).