

Meteorological factors driving glacial till variation and the associated periglacial debris flows in Tianmo Valley, southeast Tibetan Plateau

M. F. Deng^{1,2}, N. S. Chen^{1*}, and M. Liu^{1,2}

(¹Key Laboratory of Mountain Hazards and Surface Process, Institute of Mountain Hazards and Environment, Chinese Academy of Sciences, Chengdu 610041, China;

²University of Chinese Academic of Sciences, Beijing 100049, China)

Abstract: Meteorological studies have indicated that high alpine environments are strongly affected by climate warming, and periglacial debris flows are frequent in deglaciated regions. The combination of rainfall and air temperature controls the initiation of periglacial debris flows, and the addition of meltwater due to higher air temperatures enhances the complexity of the triggering mechanism compared to that of storm-induced debris flows. On the southeastern Tibetan Plateau, where temperate glaciers are widely distributed, numerous periglacial debris flows have occurred over the past 100 years, but none occurred in the Tianmo watershed until 2007. In 2007 and 2010, three large-scale debris flows occurred in the Tianmo Valley. In this study, these three debris flow events were chosen to analyse the impacts of the annual meteorological conditions, including the antecedent air temperature and meteorological triggers. The remote sensing images and field measurements of the adjacent glacier suggested that sharp glacier retreats occurred in the one to two years preceding the events, which coincided with spikes in the mean annual air temperature. Glacial till changes providing enough active sediment driven by a prolonged increase in the air temperature are a prerequisite of periglacial debris flows. Different factors can trigger periglacial debris flows, and they may include high intensity rainfall, as in the first and third debris flows, or continuous, long-term increases in air temperature, as in the second debris flow event.

Key words: glacial till variation; meteorological factors; periglacial debris flows; southeast Tibetan Plateau

1. Introduction

Alpine environments are vulnerable to climate changes, and alpine glaciers and permafrost are the most sensitive to degradation (Harris et al., 2009; IPCC, 2013). Glacier and permafrost retreat can induce mass movements such as landslides, shallow slides, debris slides, moraine collapses, etc. (Cruden and Hu, 1993; Korup and Clague, 2009; McColl, 2012; Stoffel and Huggel, 2012; Fischer et al., 2012). These movements would bring the material out of the watersheds in the form of debris

批注 [admin1]: Answer to the comment:
1. The acronym TM (line 16) should be defined before its use. Moreover, its use in the abstract should be avoided as much as possible.

删除的内容: TM

批注 [admin2]: Answer to the comment:
2. Line 17 the nearby??????

删除的内容: nearby

批注 [admin3]: Answer to the comment:
3. Sentence lines 18-20: the glacial till change is a pre-requisite for debris flow formation because it provides sediments. Therefore, the beginning word, "Moreover", can be omitted, and a brief explanation should be introduced.

删除的内容: Moreover, g

批注 [admin4]: Answer to the comment:
4. Line 26: "changes,and": the missing of a space after points and commas is seen in all the paper. Please correct.

35 | flows or sediment fluxes. Debris flows in alpine regions often bury residential areas,
36 | cut off main roads, block rivers (Shang et al., 2003; Cheng et al., 2005; Deng et al.,
37 | 2013) and destroy basic facilities downstream; thus, they pose a considerable threat to
38 | the local economy and social development. In undeveloped alpine areas where the
39 | transportation system is particularly poor or limited, such as in southeastern Tibet, the
40 | negative effects produced by debris flows, such as cutting off main roads, can be
41 | serious (Cheng et al., 2005).

42 | Periglacial debris flows occur in high alpine areas with large areas of glaciers,
43 | such as on the Tibetan Plateau in China (Shang et al., 2003; Ge et al., 2014), in the
44 | Alps in Europe (Sattler et al., 2011; Stoffel and Huggel, 2012), in the Caucasus
45 | Mountains in Russia (Evans et al., 2009) and in northern Canada (Lewkowitz and
46 | Harris, 2005). Periglacial debris flows can be initiated by rainfall (Stoffel et al., 2011;
47 | Schneuwly-Bollschweiler and Stoffel, 2012), glacial meltwater flow or ice particle
48 | ablation (Arenson and Springman, 2005; Decaulne et al., 2005) or outburst floods
49 | from glacier lakes (Chiarle et al., 2007) in different parts of the world; however,
50 | multiple triggers of a single event have rarely been studied. Because debris flows are
51 | commonly triggered by rainfall (Sassa and Wang, 2005; Decaulne et al., 2007; Kean
52 | et al., 2013; Takahashi, 2014), the rainfall threshold, intensity and duration have been
53 | widely used for debris flow monitoring and to provide event warnings in non-glacier
54 | areas (Guzzetti et al., 2008).

55 | In deglaciated areas, the debris flow threshold can be more difficult to determine.
56 | Periglacial debris flows tend to occur in the summer when the thawing of glaciers and
57 | glacial tills predominates and meltwater penetrates the glacial tills at a constant and
58 | successive flow rate. The effect of meltwater is similar to that of antecedent rainfall
59 | (Rahardjo et al., 2008) and is variable in different periods, considering snow and
60 | glacier shrinkage and air temperature fluctuations. In the Swiss Alps, the meltwater
61 | volume is high in early summer, and debris flows can be initiated by low intensity
62 | rainfall. However, larger rainstorms are required to produce debris flows in late
63 | summer and early autumn when the meltwater volume is low (Stoffel et al., 2011;
64 | Schneuwly-Bollschweiler and Stoffel, 2012). On the southeastern Tibetan Plateau, the

批注 [admin5]: Answer to the comment:
5. Line 32: "Debris flowsin": the missing of
space between two words is seen in all the
paper. Please correct.

65 rainfall threshold given by Chen et al. (2011) is relatively wide (0.2~2.0 mm/10min,
66 0.6~6.3 mm/h or 3.0~19.4 mm/24h), the small rainfall threshold of which is likely
67 affected by the air temperature. Moreover, periglacial debris flows induced by sudden
68 releases of water from glacier lakes are closely related to increasing air temperature
69 (Liu et al., 2014).

70 Air temperature fluctuations are likely important triggers of periglacial debris
71 flows. Compared to storm-induced debris flows, the addition of meltwater due to
72 increased air temperature can greatly enhance the complexity of the initiation
73 mechanism of periglacial debris flows. It is difficult to simulate the triggering process
74 via experiments or mathematical simulation; thus, case studies of natural debris flows
75 must be explored. In this study, three debris flow events in the Tianmo watershed on
76 the southeastern Tibetan Plateau are used as examples after a debris flow-free period
77 of nearly 100 years as deglaciation continues. The annual meteorological conditions,
78 antecedent air temperature and triggering conditions prior to debris flows are analysed
79 to further understand the meteorological triggers and their roles in glacier retreat,
80 glacial till variation and debris flow initiation.

81 2. Background

82 (1) Study area

83 Temperate glaciers on the Tibetan Plateau are primarily distributed in the Parlung
84 Zangbo Basin, and they covered a total landmass of 2381.47 km² in 2010 based on
85 TM images (taken by the No. 4 or 5 thematic mappers on the Landsat satellite with a
86 spatial resolution of 30 m) (Liu, 2013). Historically, the movement of temperate
87 glaciers has produced numerous moraines, the depth of which can reach 500 m locally
88 (Yuan et al., 2012). In recent decades, a significant temperature increase has occurred,
89 and the temperature at the Bomi meteorological station (central Parlung Zangbo Basin)
90 increased by 0.23°C/10a from 1969 to 2007, resulting in remarkable glacial shrinkage
91 (Yang et al., 2010).

92 Tianmo Valley, which is located in Bomi County and to the south of the Parlung

批注 [admin6]: Answer to the comment:
6. Line 69; maybe "possibility" is better
than "complexity". ____may complexity is
better as the author want to say than the
mechanism is complexity as the combined
effect of rainfall and ice melt water

批注 [admin7]: answer to comment :
14. Line 216: mapperson??????

93 Zangbo River, covers an area of 17.76 km² (29°59'N/95°19'E; Figure 1). This valley
94 has a northeast-southeast orientation and is surrounded by high mountains reaching
95 5590 m a.s.l. at the southernmost location and 2460 m a.s.l. at the fork in the Parlung
96 Zangbo River. The TM image from 2013 illustrated the presence of a hanging glacier
97 with an area of 1.42 km² in the upper concave area at an elevation of 4246 m to 4934
98 m. Bare rock, dipping at an angle of approximately 60°, emerges below and above the
99 hanging glacier and is often covered by snow. Below 3800 m a.s.l., vegetation,
100 including forest and shrubs, occupies most of the area (Table 1).

101 The river channel in the watershed is sheltered by shade and not directly affected
102 by sunlight, resulting in less solar radiation and a location at which a small trough
103 glacier can form. In the main channel, the trough glacier extended to 2966 m a.s.l. in
104 2006. The lower part of the trough glacier has been eroded by glacier meltwater flow,
105 and an arch glacier that is vulnerable to high pressure was formed (Figure 2). The
106 remnants of the landslide deposits are approximately 10 metres high can be observed
107 on both sides of the channel. These deposits consist of low-stability sediment and can
108 be easily entrained by debris flows.

109 Tianmo Valley is located on the north side of the bend in the Yarlung Zangbo
110 River and is strongly affected by new tectonic movement. An inferred normal fault
111 vertical to the channel cuts through the valley and is only 30 km from the Yarlung
112 Zangbo fault. In 1950, a rather significant earthquake (Ms. 8.6) hit Zayu, which is
113 only 200 km away, and local records reported that a large amount of rock collapsed
114 and landslides were produced at that time. The whole valley is located in a strong
115 ductile deformation zone and is dominated by gneissic lithology belonging to a
116 Presinian System.

117 (2) Disaster history

118 According to our field interviews with local residents, there were no debris flows
119 in the approximately 100 years prior to 2007 in Tianmo Valley. The channel was
120 relatively narrow before 2007, and the local people could walk across via a wooden
121 bridge to live and farm on the terrace on the west side. On the morning of September

122 4th, 2007, a rainfall even hit this area and it ceased around 7:00. On the evening of
 123 approximately 18:00, the local forest guard heard a loud noise coming from the
 124 upstream area and rainfall later began in the upstream area at approximately 19:00.
 125 Then, debris flows occurred in the Tianmo Channel after the second rainfall event,
 126 and they subsequently blocked the Parlung Zangbo River. It is told by the local citizen
 127 that several debris flows occurred during that entire night while we cannot separate
 128 them according to the field measurements, and approximately 1,340,000 m³ of
 129 sediment was transported during this event, resulting in 8 missing persons and deaths.
 130 This debris flow event is listed as DF1 in this paper, which contained the first debris
 131 flows and the following waves. Concurrently, debris flows occurred in the four
 132 adjacent 4 valleys (Table 2). According to the size classification proposed by Jakob
 133 (2005), which is based on the total volume, peak discharge and inundated area, the
 134 size classes of the debris flows in the five valleys are given in Table 2.

135 At 11:30 on July 25th, 2010, debris flows were again triggered in Tianmo Valley
 136 that traced the path of the preceding debris flow deposits and reached the other side of
 137 the Parlung Zangbo River. According to Ge et al. (2014), a solid sediment mass of
 138 approximately 500,000 m³ was carried to (Table 1) and deposited in the channel and
 139 blocked the main river. A barrier lake was formed, and the rising water destroyed the
 140 roadbed of G318. Dozens of small-magnitude debris flows occurred in the following
 141 week. This debris flow event is listed as DF2 in this paper and it contained several
 142 waves.

143 Debris flows occurred again two months later on Sep. 6th (The Ministry of Land
 144 and Resources P. R. C., 2010), although we could not determine the exact time
 145 sequence of the events. According to speculation, the debris flows could have
 146 occurred in the early morning before dawn when the rainfall intensity reached its
 147 maximum (Figure 9). This theory agrees with the findings of Chen (1991), who found
 148 that periglacial debris flows historically occurred between 18:00~24:00 in this area.
 149 The debris barrier in the main river was consequently increased by an additional
 150 450,000 m³, and the barrier lake was enlarged to hold 9,000,000 m³ of water. This
 151 debris flow event is listed as DF3 in this paper and waves of which cannot be

- 删除的内容: after
- 批注 [admin8]: Answer to the comment: 7. Lines 118-121. "On the morning" is not coherent with "18.00" and add t to even (line 119); I suggest to rewrite the sentence as:the triggering area was hit by a rainfall event and after that some loud noise were heard about 18:00..... (... [2])
- 删除的内容: a rainfall even that did not hit the downstream area ceased, the local forest guard heard a loud noise coming from the upstream area at approximately 18:00 (... [1])
- 删除的内容: Rainfall
- 批注 [admin9]: Answer to the comment: 8. Lines 121-123 Rewrite the sentences as: a debris flow occurred after a second rainfall event that began at 19:00 (... [3])
- 删除的内容: began to
- 已下移 [1]: This debris flow event is listed as DF1 in this paper.
- 删除的内容: According
- 已移动(插入) [1]
- 批注 [admin10]: Answer to the comment: 9. Lines 122, 131 and 138 debris flows? The writer does not understand if for a debris flow event the authors intend different debris flows or a debris flow composed by several waves or something else. Please explain in the paper. (... [4])
- 批注 [admin11]: Answer to the comment: 10. Line 128: Table 2 is not necessary because it deals with debris flows that are not object of present work. ___the other debris flows cases in table 2 is used to show that the metrological condition in these days is favorable for debris flows (... [5])
- 删除的内容: nearby
- 批注 [admin12]: Answer to the comment: 11. Lines 142-143: the finding of Chen (1991) could be due to the increase of (... [6])

167 | determined.

168 | A field investigation revealed that a high percentage of boulders in the
169 | downstream area and glacial tills above the trough glacier are loose and high porosity
170 | rocks (Figure 2); hence, they have low density and can be easily entrained. Our
171 | particle size tests of the glacial tills and debris flow deposits indicate a low clay
172 | ($d < 0.005$ mm) content, whereas the debris flow deposits contain more fine particles
173 | that are smaller than 10 mm (Figure 4), suggesting that entrainment accounted for a
174 | considerable amount of fine particles.

175 | **(3) Meteorological data**

176 | The study area is located in a high alpine area where the economy is relatively
177 | undeveloped and few meteorological stations exist. Before 2011, the Bomi
178 | meteorological station (established in 1955) was the only station in the area. It is
179 | located 54 km from Tianmo Valley at an elevation of 2730 m, and other stations were
180 | located more than 200 km away.

181 | The Tibetan Plateau is a massive terrace that obstructs the Indian monsoon,
182 | causing it to travel through the Yarlung Zangbo Canyon and its tributaries. As the
183 | Indian monsoon is transported to higher altitudes, a rainfall gradient emerges in the
184 | Parlung Zangbo Basin. However, according to the rainfall data from the area, rainfall
185 | often exhibits a similar intensity as that of the long-term rainfall process from
186 | Guxiang to Songzong, which suggests that a large rainfall gradient does not exist
187 | between Tianmo Valley and Bomi meteorological station; therefore, the rainfall data
188 | from Bomi meteorological station could be used in our study. To conduct additional
189 | studies, another meteorological station was built in 2011 near Tianmo Valley.

190 | It has been established that the air temperature decreases with altitude; therefore,
191 | the air temperature in the source area of Tianmo Valley is lower than that in Bomi
192 | County. According to the research by Li and Xie (2006), the air temperature decreases
193 | at a rate of $0.46\text{--}0.69^\circ\text{C}/100$ m over the entire Tibetan Plateau, and the rate in the
194 | study area is $0.54^\circ\text{C}/100$ m. Because the glacier and permafrost in the source area
195 | have planar distributions, the air temperature at the geometric centre of the glacier and

196 permafrost can be used to analyse the temperature process.

197 3. Analysis and results

198 (1) Air temperature and rainfall changes

199 The mean annual air temperature is generally used to reflect the trends of glacier
200 change (Yang et al., 2015). We collected mean annual air temperature and annual
201 rainfall data from 1970 to 2014 from the Bomi meteorological station (Figure 5). The
202 records showed that the mean air temperature has increased by approximately 1.5 °C
203 over the past 45 years at a rate of 0.033 °C/a. This air temperature increase was
204 particularly rapid between 2005 and 2007 at approximately 0.7 °C/3a, which is 7
205 times the average value over the past 45 years. However, the annual rainfall from
206 2000 to 2010 was low and estimated as 828.2 mm per year. From 2000 to 2004, the
207 rainfall during summer (July to September) accounted for approximately 50% of the
208 total annual rainfall; however, only 32% of the rainfall occurred in the summer of
209 2005~2006, even though the annual rainfall exhibited a similar trend. In 2007, rainfall
210 in the summer and the entire year returned to the mean rainfall state.

211 Figure 5 shows similar air temperature and rainfall trends before DF2 and DF3.
212 The air temperature increased in 2009 and reached 10.2 °C which is the maximum of
213 the past 45 years; however, the annual rainfall went down to only 65% of the average
214 amount; and the summer rainfall reached a minimum value. In 2010, rainfall was
215 abundant, and the annual rainfall increased to 1080.6 mm, which is approximately 30%
216 more than the average value and close to the maximum.

217 The following common traits can be identified by comparing the annual
218 meteorological conditions of DF1, DF2 and DF3: 1) one or two years before the
219 debris flows, the mean annual temperature increased and the annual rainfall and
220 summer rainfall decreased. Additionally, the climate was in a "hot-dry" state; 2) As
221 the temperature gradually decreased, the annual rainfall returned to normal or
222 increased, and a "hot-wet" climate state contributed to debris flow initiation (Lu and
223 Li, 1989).

批注 [admin13]: answer to comment:
12. Lines 185-189. The sentences could be substituted by In the periods 2000-2004 and 2007 rainfall precipitated from July to September was about 50% of the total annual rainfall while.....

删除的内容:

删除的内容: temperature over

删除的内容: of 10.2 °C

删除的内容: was

删除的内容: , which was lower than that in 2005 and 2006, reached a minimum value

批注 [admin14]: answer to comment:
13. Lines 190-196: a more schematic and concise presentation of data is required.

230 (2) Changing of glacier in Tianmo Valley

231 In our study, remote images were collected to analyse the glacier changes in the
232 source area in recent years. To eliminate the effect of snow cover, images were taken
233 in the thawing seasons when snow cover is limited, enabling easy detection of glaciers
234 and snow. Moreover, an image taken on a bright, cloudless day is still needed to show
235 the watershed clearly; however, a difficult case is encountered when the rainy season
236 begins during the thawing season, as the atmosphere is often covered by thick clouds.
237 Furthermore, to illustrate glacier retreat and its impact on debris flows properly, the
238 images should be within similar time intervals, such as 3 years, before and after debris
239 flow events. High-resolution images are rare, and we could only collect one SPOT
240 image (taken by the Systeme Probatoired' Observation de la Terre satellite with a
241 spatial resolution of 5 m) in 2008. To achieve image consistency, we collected 5 TM
242 images taken on September 17th, 2000, July 24th, 2003, September 21st, 2006,
243 September 24th, 2009 and August 4th, 2013.

244 Based on the 5 TM images, we classified the area as glacier, snow, bare land,
245 gully deposit and vegetation in time series (Figure 6), and the area of each is given in
246 Table 1. Figure 6 shows that deglaciation occurred in Tianmo Valley; notably, the
247 eastern branch has experienced considerable deglaciation. To clearly show the rapid
248 rate of glacial retreat in the entire basin and eastern branch, a graph of retreat was
249 plotted, as shown in Figure 7.

250 Figure 7 shows that the glacier in Tianmo Valley shrank from 2000 to 2013, with
251 variable rates of glacier retreat. In 2000~2003, 2003~2006, 2006~2009 and
252 2009~2013, the glacier retreat rates in Tianmo Valley were 0.02, 0.06, 0.027 and
253 0.0075 km²/a, respectively, and those of the eastern branch were 0.0033, 0.01, 0.008
254 and 0.002 km²/a, respectively. According to the figure, the largest glacier retreat rate
255 was observed in 2003~2006, followed by that in 2006~2009. The glacier area at the
256 beginning should be noted to assess the rate of change of the glacier. The glacier
257 retreat rate can be normalized, and the relative glacier retreat rate can be calculated
258 based on this change in area.

删除的内容: (taken by the No. 4 or 5
thematic mapperson the Landsat satellite
with a spatial resolution of 30 m)

262 | The relative glacier retreat rates were 11.30, 35.09, 17.43 and 5.17 $10^{-3}\text{km}^2/\text{a}/\text{km}^2$
263 | in 2000~2003, 2003~2006, 2006~2009 and 2009~2013, respectively; and the
264 | corresponding values were 20.83, 66.67, 66.67 and 20.83 $10^{-3}\text{km}^2/\text{a}/\text{km}^2$ for the
265 | eastern branch. The relative glacier retreat rate of the eastern branch decreased sharply
266 | between 2000 ~2013.

267 | In this study, TM images over 3-year intervals were applied to obtain the mean
268 | glacier retreat rate. As glacier retreat rate in the 3 three years could be either high or
269 | low, field measurements of the nearby glacier were used to show the glacier retreat
270 | condition before debris flows occurred. Yang et al. (2015) conducted field
271 | measurements of the No.94 Glacier in the Parlung Zangbo Basin since 2006, and the
272 | field measurements suggest it had a negative balance from 2006~2010 (Figure 7). The
273 | negative balance reached a maximum level in 2009, followed by 2008 and 2006,
274 | indicating rapid deglaciation in these three years.

275 | When we combined the results of the TM image analysis and field measurements
276 | of the No. 94 Glacier, we observed that the glacier in Tianmo Valley experienced the
277 | most rapid deglaciation prior to debris flows in 2006, 2008 and 2009, which coincided
278 | with an increase in the mean annual air temperature (Figure 5). Moreover, the
279 | maximum glacier retreat in 2009 was potentially related to the decline in snowfall in
280 | the preceding winter and early spring.

281 | **(3) Antecedent air temperature and rainfall**

282 | The air temperature in the source area can be obtained based on a vertical rate of
283 | decline ($0.54\text{ }^{\circ}\text{C}/100\text{ m}$). Based on this method, the air temperature in the source area
284 | was $9.8\text{ }^{\circ}\text{C}$ lower than that at the Bomi meteorological station. We collected the
285 | lowest temperature, the mean temperature and daily rainfall from June to September
286 | in 2007 and 2010 (Figure 8).

287 | Figure 8 shows that the lowest air temperature was below 0 at the end of June
288 | 2007. At the beginning of July, the air temperature started to rise quickly, which
289 | continued until early September when DF1 occurred. This trend suggests that the high
290 | air temperatures in July and August contributed to DF1.

291 Additionally, the air temperature was high from early July to late August, and
292 another high air temperature period was observed in early September. When DF2
293 occurred in late July, the air temperature had reached the maximum for that year,
294 which suggests that the air temperature in early and mid-July was responsible for DF2.
295 After DF2 occurred, the air temperature in August varied towards the conditions that
296 caused DF3.

297 Antecedent air temperature fluctuations include the air temperature and the
298 duration of variations. The air temperatures and durations before debris flows are
299 variable and difficult to evaluate. The accumulation of positive air temperature is
300 often used to analyse the effect of air temperature on glacier melting (Rango and
301 Martinec, 1995) and can be expressed as follows:

$$302 \quad T_{PT} = \sum_{i=n}^0 T_i (T_i > 0) \quad (1)$$

303 where T_{PT} is positive air temperature accumulation (°C) and T_i is the average
304 daily air temperature (only $T_i > 0$ is included).

305 Because air temperature is successive, it is difficult to determine the beginning of
306 positive air temperature accumulation. Glacial tills can decrease the heat that
307 penetrates into them, and the low air temperature is only observed in the upper thin
308 layer. Moreover, freeze-thaw cycles exist when the lowest air temperature is less than
309 0°C. From this perspective, the beginning of positive air temperature accumulation is
310 defined as the time at which the lowest air temperature exceeds 0°C for two or three
311 successive days or since the last debris flow.

312 Based on the above method, we can deduce that positive air temperature
313 accumulation began when the lowest air temperature exceeded 0°C for several
314 successive days beginning on June 28th, 2007, June 9th, 2010, and July 26th, 2010,
315 which correspond to DF1, DF2 and DF3, respectively. The duration and T_{PT} were
316 calculated for each debris flow event. The results were 69 days and 517.9 °C, 47 days
317 and 332.1 °C and 42 days and 320.4 °C (Figure 8) for DF1, DF2 and DF3,

批注 [admin15]: answer to comment:
15. Line 261: explain in the caption of
Figure 8 the meaning of PT. PT is T_{PT}

318 respectively. The results showed that T_{PT} for DF1 was much larger than the other
319 two T_{PT} values, which coincides with the fact that there ~~were~~ no debris flows in the
320 past dozens of years, and extraordinary external forces such as large T_{PT} are required
321 to disrupt the long-term balance.

322 (4) Triggering conditions

323 ~~Rainfall in a short can trigger debris flows while it cannot be~~ triggered by a sole
324 abrupt increase in air temperature ~~as the continuous and limited nature of air~~
325 ~~temperature, instead, air temperature of longer term should be included.~~ In our
326 analysis, the rainfall over the three days preceding a debris flow event is given in
327 Figure 9.

328 Before DF1, the air temperature was high, which continued through July and
329 August. Notably, the T_{PT} reached ~~517.9°C~~. According to the local forest guard, an
330 isolated convective storm occurred prior to DF1, although no rainfall was recorded at
331 the Bomi meteorological station or in the downstream area at that time. In Figure 9, as
332 the rainfall right before DF1 occurred was not recorded by the Bomi meteorological
333 station, we added approximately 5 mm/h of rainfall intensity (according to the
334 description provided by the forest guard) before DF1 to account for the storm, which
335 might not reflect the real rainfall process. We can therefore conclude that this isolated
336 convective storm initiated DF1, while the long-term high air temperature trend paved
337 the way for DF1. Considering a large deglaciaded area, several other periglacial debris
338 flows simultaneously occurred near Tianmo Valley (Deng et al., 2013), which
339 suggests the advantageous meteorological conditions for debris flow initiation.

340 DF2 occurred when the air temperature reached a peak in 2010. The thaw season
341 began in the middle of June, and T_{PT} reached 332.1 °C. On July 24th, one day before
342 DF2, the air temperature reached a maximum value for that year. ~~No rainfall event hit~~
343 ~~this area preceding DF2 according to the record of~~ Bomi meteorological station, and
344 the local citizens also observed no rain. ~~The trigger of DF2 was likely the continuous~~

批注 [admin16]: answer to comment:
16. Line 291: were instead of "was"

删除的内容: was

删除的内容: The continuous nature of air
temperature limits the possibility for debris
flows

删除的内容: . Since previous air
temperature trends cannot be neglected, it is
of no sense to study air temperature
triggers. .
Antecedent rainfall is a factor that favours
debris flows.

批注 [admin17]: answer to comment: 17.
Lines 295-297. The writer does not
understand the first sentence and therefore
its link and the sense of the second
sentence.

批注 [admin18]: answer to comment:
18. Lines 301, 313 and 321: 517.9°C: this
high value (in centigrade) is not possible.
___517.9°C is the positive air temperature
accumulation which can be calculated by
the equation (1).

删除的内容: The rainfall record at the

删除的内容: shows that there had been
no rainfall several days preceding DF2

批注 [admin19]: answer to comment :
19. Line 315-316: from that there had been
no..... the sentence becomes unclear.

358 | percolation of meltwater due to the long-term increase in air temperature.

359 | According to field interviews, several debris flows of small magnitude occurred
360 | before DF3. The air temperature decreased in late August but increased to another
361 | high value before DF3, and the T_{PT} reached 320.4 °C. Rainfall began 2 days prior to
362 | DF3 and lasted the entire day before DF3. According to the rainfall trend at the Bomi
363 | meteorological station, the rapid increase in rainfall intensity started 4 hours before
364 | DF3 and reached 3.8 mm/h, which was responsible for the initiation of DF3.

批注 [admin20]: answer to comment:
20. Line 322 What does it mean a steady
rainfall?

删除的内容: was steady

删除的内容:

365 | 4. Discussion

366 | In this study, we found that the triggering factors of the three debris flows were
367 | high air temperature and rainfall for DF1, high air temperature for DF2 and storm for
368 | DF3, respectively. When we analysed the dates and triggers of these events, various
369 | questions should be settled first: 1) why did debris flows not occur in 2006 or 2009
370 | when deglaciation reached its peak and more ice meltwater was present; 2) why did
371 | DF1 and DF3 occur in September when the air temperature and volume of ice
372 | meltwater were decreasing; and 3) why there were no large-scale debris flows
373 | triggered by previous heavy storms. Based on our results, we believe that the impact
374 | of the water source on the magnitude and frequency of debris flows is relatively small,
375 | or more debris flows would form during the early larger storm; instead, the sediment
376 | source, including the associated magnitude and activity, may be the predominant
377 | control, as reported by Jakob et al. (2005), who noted that channel recharge is a
378 | prerequisite for debris flows. However, in most situations, we cannot reach the source
379 | area to detect the soil source, and high-tech remote sensing can only distinguish the
380 | boundary of the soil source. In the periglacial area where glacial till is often covered
381 | by glacier or everlasting snow, a change in the soil source would be highly difficult to
382 | detect. In this study, we combine the meteorological conditions and literature reports
383 | to discuss the likely variations in glacial tills before debris flows.

批注 [admin21]: answer to comment :
21. Line 332 write were after "there"

删除的内容: were

删除的内容: soil

批注 [admin22]: answer to comment :
22. Line 335 and following: write sediment
source instead of "soil source"

384 | (1) Annual variations in glacial till

385 | Climate warming is a global trend (IPCC, 2013), and the Tibetan Plateau, as the

390 third pole, is no exception to climate change. According to our statistics, the air
391 temperature in Bomi County has increased by 1.5 °C over the past 45 years
392 (1970~2014). Glacier retreat induced by climate warming has been widely accepted,
393 and recent research suggests that the weaker Indian monsoon could be another reason
394 for such retreat (Yao et al., 2012). Glaciers are always located in concave ground
395 areas and cover large volumes of glacial tills. Glacial pressure can generate normal
396 stress vertical to the slope, which can strengthen the slope stability. The effect of
397 glaciers on slope stability is called glacial debuttressing (Cossart et al., 2008). As
398 deglaciation continues, the result could lead to the exposure of the frozen glacial tills
399 (Figure 10 A to B) and smaller glacial debuttressing.

400 The retreats of glaciers and glacial tills due to climate warming are quite
401 different. The newly formed bare glacial till is frozen with a high ice content. The
402 cohesion of the ice particles creates a bare glacial till with high shearing strength and
403 stability. Deglaciation is accompanied by the melting of internal ice particles, which
404 can greatly enhance the activity. This process first occurred at the surface layer of
405 glacial till, followed the layers below, resulting in enlargement of active debris. As the
406 debris obstruct heat fluxes from penetrating into the layer below, so the melting rate of
407 internal ice particles is quite slower than that of glacial retreat (Takeuchi et al., 2000),
408 result into a strong heat gradient at the surface while limited in deep layers, which
409 means the activity of the debris decline with depth and long term high air temperature
410 is required to enhance the activity in a deeper layer. As the ablation rates is quite low,
411 only the surface layer is highly active and the sediment is relatively limited. Therefore,
412 no debris flows of large magnitude could occurred in 2006 and 2009 when glacier
413 retreat reached a maximum while the active glacial till is restricted to the surface
414 layer.

415 (2) Variation in glacial till on antecedent days

416 After the long, cold winter, glacial tills become frozen. If a regressive glacier
417 does not recover in the winter, glacial tills are covered by snow. As the air temperature
418 increases again, the surface snow melts first, followed by the internal ice particles.

删除的内容: produce an active surface layer that can

删除的内容: deep

删除的内容: and result in

删除的内容: at a

删除的内容: rate

删除的内容: slower

删除的内容: . Because a strong

删除的内容: occurs

删除的内容: but

删除的内容: is

删除的内容: glacial tills with thicker coverage always have relatively thinner thawed layers, and the ablation rates of glaciers and internal ice particles are similar to that at the glacier surface and close to the moraine slope. The newly formed bare glacial till is frozen with a high ice content. The cohesion of the ice particles creates a bare glacial till with high shearing strength and stability,

删除的内容: and

删除的内容: were observed

删除的内容: and

批注 [admin23]: answer to comment: 24. Lines 356-368. The link between the thaw process within the till, its duration and the absence of debris flow in 2006 and 2009 is ill explained or missing.

删除的内容: quite limited

删除的内容: .

445 The thawing of internal ice particles induces a series of changes in the glacial till,
446 which include the following: 1) the thawing will break the bonds of ice particles and
447 increase the instability between ice cracks (Ryzhkin and Petrenko, 1997; Davies et al.,
448 2001); 2) the sharp air temperature fluctuations in high alpine, mountainous areas
449 induce a repeated cycle of expansion and contraction in the glacial till that can destroy
450 the mass structure to some extent; 3) the seepage of ice meltwater can transport
451 fine-grained sediments that were formerly frozen in the ice matrix (Rist, 2007); and 4)
452 the ice meltwater can result in a higher water content and pore water pressure
453 (Christian et al., 2012). These changes in glacial till can sharply decrease the soil
454 strength, shifting to an active mass from an uncovered and frozen moraine (Figure
455 10B to C). Because heat conduction in glacial till is relatively slow, this process may
456 last for a very long time and require a high antecedent air temperature.

457 Heat conduction via the percolation of rainfall and ice meltwater can amplify the
458 depth of active glacial till (Gruber and Haeberli, 2007), whereas covering the surface
459 glacial till can hinder a heat flux from penetrating into the deep layers (Noetzli et al.,
460 2007). At a low air temperature, the heat flux should be constrained to the surface
461 layer, and a large heat gradient due to a high air temperature would contribute much
462 more to the heat flux and ice melt in the deep mass. Thus, the long-term effect of a
463 high air temperature can amplify the active glacial till (Noetzli et al., 2007; Åkerman
464 and Johansson, 2008), under which lies frozen glacial till with a high ice content. The
465 activity of glacial till varies with depth from high at the surface to low in the deep
466 layers, and landslide failure can take place on glacial till slopes in a retrogressive
467 manner, coinciding with long-term air temperature fluctuations, as active glacial till is
468 relatively limited in deglaciated areas.

469 **(3) Failure of glacial till**

470 Different factors can lead to glacial till failure. Active glacial till slopes with low
471 strength are usually vulnerable, and their failure can occur when the air temperature is
472 above 0 °C (Arenson and Springman, 2005). Rainfall or ice melt water induced by air
473 temperature can trigger the failure (Figure 10 C to D). This type of event is called a

474 shallow landslide, and the failure mechanism lies in the ablation of internal ice
475 particles and the percolation of meltwater, which can initially decrease the soil
476 strength (Arenson and Springman, 2005; Decaulne et al., 2005). Later, the subsequent
477 rapid percolation of ice meltwater or heavy rainfall can saturate the debris, decrease
478 soil suction and shearing strength, and form seepage flows that can trigger the shallow
479 landslide failure (Springman et al., 2003; Decaulne and Sæmundsson, 2007; Chiarle et
480 al., 2007). Whether the failure can induce debris flows is still dependent on its ability
481 to entrain the debris layer as the flow moves through the channel.

482 Another type of failure might take place when peaked runoff flows over and
483 entrains debris deposits in the charged channel and reach a critical discharge (Berti
484 and Simoni, 2005; Gregoretto and Dalla Fontana, 2008; Kean et al., 2012, 2013;
485 Takahashi, 2014; Rengers et al., 2016, Gregoretto et al., 2016), which is more
486 determined by channel bed slope and grain size of debris (Tognacca et al., 2000;
487 Gregoretto, 2000; Theule et al., 2012; Hurlimann et al., 2014; Degetto et al., 2015).

488 This type of channelized runoff could be a combination of three factors: rainfall,
489 melting ice or the overflow that forms when a glacier collapses downward into a
490 water pool. Mechanism of this process lies in the hydrodynamic forces exerted on the
491 surface elements of debris layers and surpassing sediment resistance (Gregoretto and
492 Dalla Fontana, 2008; Recking et al., 2009; Prancevic et al., 2014). FaThe
493 concentration of runoff in the channel bottom causes the erosion of the debris surface
494 layer forming a solid-liquid current at first, then extends to the layers below with
495 whole or partial mobilization and debris flows was generated (Gregoretto and Dalla
496 Fontana, 2008). Therefore, debris flows initiated by landslide failure caused by
497 seepage flow or channelized runoff that entrain sediments in the periglacial area is
498 similar with the mechanism of debris flows initiation in non-glacier areas (Iverson et
499 al., 1997; Springman et al., 2003; Sassa and Wang, 2005; Gregoretto and Dalla
500 Fontana, 2008; Kean et al., 2013), while the difference lies in the activity of debris
501 and the source of water. In the European Alps, periglacial debris flows are mainly
502 provoked by rainfall, which is also related to air temperature fluxes (Stoffel et
503 al., 2011). Additionally, the values of rainfall and air temperature required to trigger

批注 [admin24]: answer to comment:
25. Line 409 in which care????

删除的内容: , in which care the debris is deposited

删除的内容: a

批注 [admin25]: answer to comment:
26. Line 411 Delete "a" before "peaked"
and write "peaked runoff flows (Kean et al.,
2012, Rengers et al., 2016, Gregoretto et al.,
20016) "

批注 [admin26]: answer to comment:
27. Line 414 add the references Theule et
al., 2012, Hurlimann et al., 2014 and
Degetto et al., 2015.

删除的内容: Armanini and Gregoretto,
2005;

删除的内容: Kean et al., 2013

带格式的: 字体: Times New Roman, 小
四, 字体颜色: 文字 1, 英语(英国)

带格式的: 字体: Times New Roman, 小
四, 字体颜色: 文字 1, 英语(英国)

批注 [admin27]: answer to comment :
28. Line 415 delete the reference Armanini
and Gregoretto, 2005 and Kean et al., 2013.

批注 [admin28]: answer to comment: 29.
Line 418-421 Rewrite the sentence as:
Mechanism of this process lies in the
hydrodynamics forces exerted on the
surface elements of debris layers and
surpassing sediment resistance (Gregoretto,
2008; Recking et al., 2009; Prancevic et al.,
2014).

删除的内容: , created by the channelized
runoff, acting on the surface elements of the
debris layer and surpassing resistance of the
(... [7])

删除的内容: 5; Prancevic et al., 2014).

批注 [admin29]: answer to comment:
Theule, J.I., Liebault, F., Loye, A. Laigle
(... [8])

删除的内容: from

删除的内容: that leads to a landslide
failure

522 debris flows could be inversely correlated. Air temperature increases can cause
523 melting and water runoff; thus, the rainfall required to create percolation flows or
524 critical discharge to trigger a debris flow would be much less. In addition, the
525 intensity and duration of the required rainfall may require other preconditions, such as
526 those associated with the distributions of glaciers and frozen glacial tills and the
527 terrain of the source area, to enhance the debris flow (Lewkowicz and Harris, 2005).

528 The three debris flow events were associated with similar annual meteorological
529 conditions, except that the positive air temperature accumulation prior to DF1 was the
530 largest. DF1 occurred at the end of a prolonged period of high air temperature, prior to
531 this, there were instances of small failures but no large-scale debris flows. On July
532 25th, 2007, the daily rainfall reached 20.7 mm, while no debris flows were generated
533 because the active glacial till is restricted in the shallow surface layer after a short
534 term of air temperature increase.

535 In 2010, the largest daily rainfall occurred on June 7th, accounting for 37.5 mm,
536 at the beginning of an air temperature increase when the glacial till was frozen with
537 low activity. The lack of glacial till with high activity was the likely cause of the
538 absence of debris flows. The similar situation could be found on August 23rd, when the
539 daily rainfall was 20.3 mm and the positive air temperature accumulation is low since
540 DF2, which had produced quite limited active glacial till. Besides, a low and positive
541 air temperature was observed prior to September 6th when DF3 occurred, the
542 boundary of active glacial till had been enlarged before; moreover, the high rainfall
543 intensity could supplemented this lack of prolonged high air temperature and trigger
544 debris flows.

545 5. Conclusion

546 Climate changes have serious effects in high mountainous areas, and the mass
547 movement of sediments such as periglacial debris flows has become increasingly
548 frequent. Prolonged increases in the mean annual air temperature are regarded as very
549 favourable for periglacial debris flows. In particular, the annual “hot-dry” weather
550 conditions one or two years prior were responsible for three debris flow events in

删除的内容: 2010

删除的内容: when

删除的内容: thick active glacial till was still lacking after small failure events.

删除的内容:

删除的内容: and had

删除的内容: O

删除的内容: .

删除的内容: , the antecedent air temperature accumulation had remained stable since DF2 and active glacial till was still developing

删除的内容: On

删除的内容: , the antecedent positive air temperature accumulation was small, and a low air temperature was observed previously; however,

删除的内容: lack

批注 [admin30]: answer to comment: 31. Lines 442-452. These sentences should be resumed in a more concise form. The link of the amount of rainfalls and the triggering of debris flow should be clearly explained. Moreover, authors should consider that the two debris flows triggered by rainfall (DF1 and DF2) when the areas not covered by glacier should have reached the largest extension of the year and therefore, runoff in the downstream area should increase.

569 Tianmo Valley. Debris flows are generally not initiated in the year when the mean
570 annual air temperature spikes, as the melting of internal ice particles lags behind the
571 rate of glacial retreat resulting from a prolonged increase in air temperature.

572 Glacial till is unlimited in deglaciated areas, and its activity relies on glacial
573 retreat and internal ice particle melting. Glacial till changes induced by increased air
574 temperature are the first steps in forming periglacial debris flows compared to
575 storm-induced debris flows in non-glacier areas. Glacial tills require a four-phase
576 process prior to debris flow occurrence. In this process, the variation in air
577 temperature drives the glacial till change, including causing glacier recession,
578 producing bare glacial till and enhancing the glacial till activity. Debris flows can
579 occur when a sufficient amount of active glacial till exists and rainfall-induced
580 seepage or runoff is more likely to generate debris flows.

581 It is difficult to observe glacial till changes in source areas of debris flows, and
582 the analysis of the phase conversion of glacial till in this study is based on the
583 triggering conditions and other literature findings. Indeed, the meteorological
584 conditions, such as the antecedent air temperature and meteorological triggers that
585 drive the phase conversion, are partly coupled and difficult to distinguish.

586 **Acknowledgements:** This research was supported by the National Natural Science Foundation
587 of China (grant nos. 41190084, 41402283 and 41371038) and the “135” project of IMHE, CAS.
588 We wish to acknowledge the editors of the Natural Hazards and Earth System Science Editorial
589 Office and the anonymous reviewers for their constructive comments, which helped us improve
590 the contents and presentation of the manuscript.

591 References

592 Åkerman, H. J. and Johansson, M.: Thawing permafrost and thicker active layers in
593 sub-Arctic Sweden, Permafrost Periglac. Process., 19, 279-292, 2008.
594 Arenson, L. U. and Springman, S. M.: Mathematical descriptions for the behaviour of
595 ice-rich frozen soils at temperatures close to 0 °C, Can. Geotech. J., 42, 431-442,
596 2005.

删除的内容: variable

批注 [admin31]: answer to comment:
32. Line 466-467: what is the meaning of
variable air temperature condition?

删除的内容: condition

删除的内容: due to different factors

删除的内容: s

删除的内容: and temperature increases can

删除的内容: cause

删除的内容: produce

删除的内容: e

批注 [admin32]: answer to comment: 33.
Lines 475-477 Last part of the sentence is
not well linked to the previous part.

删除的内容: In future studies, we hope to
determine the effect of each meteorological
condition, and more detailed studies should
be performed.

609 | Berti, M., and Simoni, A.: Experimental evidences and numerical modelling of debris
610 | flow initiated by channel runoff. *Landslides*, 2(3), 171-182, 2005.

611 | Bommer, C., Fitze, P. and Schneider, H.: Thaw-consolidation effects on the stability
612 | of Alpine talus slopes in permafrost, *Permafrost Periglac. Process.*, 23, 267-276,
613 | 2012. doi:10.1002/ppp.1751.

614 | Chen, N. S., Zhou, H. B., and Hu, G. S.: Development rules of debris flow under the
615 | influence of climate change in Nyingchi, *Adv. Clim. Change Res.*, 7, 412-417,
616 | 2011. (In Chinese).

617 | Chen, R.: Initiation and the Critical Condition of Glacial Debris Flow, Master thesis,
618 | Institute of Mountain Hazards and Environment, Chinese Academic of Sciences,
619 | P19, 1991.

620 | Cheng, Z. L., Wu, J. S., and Geng, X.: Debris flow dam formation in southeast Tibet,
621 | *J. Mt Sci.*, 2, 155-163, 2005.

622 | Chiarle, M., Iannotti, S., Mortara, G., and Deline, P.: Recent debris flow occurrences
623 | associated with glaciers in the Alps, *Glob. Planet. Change*, 56, 123-136, 2007.

624 | Cossart, E., Braucher, R., Fort, M., Bourlès, D. L., and Carcaillet, J.: Slope instability
625 | in relation to glacial debuttressing in alpine areas (Upper Durance catchment,
626 | southeastern France): evidence from field data and ¹⁰Be cosmic ray exposure
627 | ages, *Geomorphology*, 95, 3-26, 2008.

628 | Cruden, D. M. and Hu, X. Q.: Exhaustion and steady state models for predicting
629 | landslide hazards in the Canadian Rocky mountains, *Geomorphology*, 8,
630 | 279-285, 1993.

631 | Davies, M. C. R., Hamza, O., and Harris, C.: The effect of rise in mean annual
632 | temperature on the stability of rock slopes containing ice-filled discontinuities,
633 | *Permafrost Periglac. Process.*, 12, 137-144, 2001.

634 | Decaulne, A. and Sæmundsson, T.: Spatial and temporal diversity for debris-flow
635 | meteorological control in subarctic oceanic periglacial environments in Iceland,
636 | *Earth Surf. Process. Landf.*, 32, 1971-1983, 2007.

删除的内容: Armanini, A., and
Gregoretti, C.: Incipient sediment motion
at high slopes in uniform flow condition.
Water Res Res., 41: W12431, 2005. .

删除的内容: -
删除的内容: doi:10.1002/esp.1509. .

643 Decaulne, A., Sæmundsson, T., and Petursson, O.: Debris flows triggered by rapid
644 snowmelt in the Gleidarhjalli area, northwestern Iceland, Geografiska Annaler,
645 87A, 487-500, 2005 .

646 Degetto, M., Gregoretti, C and Bernard, M. Comparative analysis of the differences
647 between using LiDAR contour-based DEMs for hydrological modeling of
648 runoff generating debris flows in the Dolomites. Front. Earth Sci., 3,21, 2015.

带格式的: 字体: (中文) +中文正文,
(中文) 中文(简体, 中国)

649 Deng, M. F., Chen, N. S., Ding, H. T., and Zhou, C. C.: The hydrothermal condition
650 of 2007 group-occurring debris flows and its triggering mechanism in Southeast
651 Tibet, J. Nat. Dis., 22, 128-134, 2013 (In Chinese).

652 Evans, S. G., Tutubalina, O. V., Drobyshev, V. N., Chernomorets, S. S., McDougall,
653 S., Petrakov, D. A., and Hungr, O.: Catastrophic detachment and high-velocity
654 long-runout flow of Kolka Glacier, Caucasus Mountains, Russia in 2002,
655 Geomorphology, 105, 314-321, 2009.

656 Fischer, L., Purves, R. S., Huggel, C., Noetzli, J., and Haeberli, W.: On the influence
657 of topographic, geological and cryospheric factors on rock avalanches and
658 rockfalls in high-mountain areas, Nat. Hazards Earth Syst. Sci., 12, 241-254,
659 2012.

660 Ge, Y. G., Cui, P., Su, F. H., Zhang, J. Q., and Chen, X. Z.: Case history of the
661 disastrous debris flows of Tianmo watershed in Bomi County, Tibet, China:
662 some mitigation suggestions, J. Mt. Sci., 11, 1253-1265, 2014.

删除的内容: n

663 Gregoretti, C.: The initiation of debris flow at high slopes: experimental results, J.
664 Hydraulic Res., 38, 83-88, 2000.

665 Gregoretti, C., and Dalla Fontana, G.: The triggering of debris flow due to
666 channel-bed failure in some alpine headwater basins of the Dolomites: analyses
667 of critical runoff, Hydrol. Process., 22, 2248-2263, 2008.

668 Gregoretti, C., Degetto, M., Bernard, M., Crucil, G., Pimazzoni, A., De Vido G., Berti,
669 M., Simoni, A. Lanzoni, S. Runoff of small rocky headwater catchments: Field
670 observations and hydrological modeling. Water Resources Research. 52, 10,
671 8138-8158, 2016.

带格式的: 字体: 非倾斜

带格式的: 字体: (默认) Times New Roman, 小四, 字体颜色: 自动设置, 图案: 清除

带格式的: 字体: (默认) Times New Roman, 小四, 字体颜色: 自动设置, 图案: 清除

删除的内容: .

带格式的: 字体: (中文) +中文正文,
(中文) 中文(简体, 中国)

674 Gruber, S., and Haeberli, W.: Permafrost in steep bedrock slopes and its
675 temperature-related destabilization following climate change. *J. Geophys. Res.*,
676 112, (F02S18), 2007.

677 Guzzetti, F., Peruccacci, S., Rossi, M., and Stark, C. P.: The rainfall intensity-duration
678 control of shallow landslides and debris flows: an update, *Landslides*, 5, 3-17,
679 2008.

680 Harris, C., Arenson, L. U., Christiansen, H. H., Etzelmüller, B., Frauenfelder, R.,
681 Gruber, S., Haeberli, W., Hauck, C., Hölzle, M., Humlum, O., Isaksen, K., Kääb,
682 A., Kern-Lütschg, M. A., Lehning, M., Matsuoka, N., Murton, J. B., Nötzli, J.,
683 Phillips, M., Ross, N., Seppälä, M., Springman, S. M., and Vonder Mühll, D.:
684 Permafrost and climate in Europe: monitoring and modelling thermal,
685 geomorphological and geotechnical responses, *Earth Sci. Rev.*, 92, 117-171,
686 2009.

687 Hurlimann, M., Abanco, C., Moya, J., Vilajosana, I. Results and experiences gathered
688 at the Rebaixader debris-flow monitoring site, Central Pyrenees, Spain.
689 *Landslides*. 11, 6, 939-953, 2014.

690 Intergovernmental Panel of Climate Change, Summary for Policymakers. Working
691 Group I Contribution to the IPCC Fifth Assessment Report Climate Change
692 2013: The Physical Science Basis, Cambridge University Press, Cambridge, UK,
693 2013.

694 Iverson, R. M., Reid, M. E., and LaHusen, R. G., Debris-flow mobilization from
695 landslides, *Annu. Rev. Earth Planet Sci.*, 25, 85-138, 1997.

696 Jakob, M., Bovis, M., and Oden, M.: The significance of channel recharge rates for
697 estimating debris-flow magnitude and frequency, *Earth Surf. Proc. Land*, 30,
698 755-766, 2005.

699 Jakob, M.: A size classification for debris flows, *Eng. Geol.*, 79, 151-161, 2005.

700 Kean, J. W., McCoy, S. W., Tucker, G. E., Staley, D. M., and Coe, J. A.:
701 Runoff-generated debris flows: observations and modeling of surge initiation,
702 magnitude, and frequency, *J. Geophys. Res. Earth Surf.*, 118, 2190-2207, 2013.

删除的内容: -

带格式的: 字体: 非倾斜

删除的内容: .

带格式的: 字体: (中文) +中文正文,
(中文) 中文(简体, 中国)

带格式的: 字体: (中文) +中文正文,
(中文) 中文(简体, 中国)

705 Kean J.W., Staley D.M., Leeper R.J., Schmidt K.M., Gartner J.E. A low-cost method
706 to measure the timing of postfire flash floods and debris flows relative to
707 rainfall. *Water Resources Research*, 48,5, W05516, 2012.

708 Korup, O. and Clague, J. J.: Natural hazards, extreme events, and mountain
709 topography, *Quat. Sci. Rev.*, 28, 977-990, 2009.

710 Lewkowicz, A. G. and Harris, C.: Frequency and magnitude of active-layer
711 detachment failures in discontinuous and continuous permafrost, northern
712 Canada, *Permafrost Periglac. Process.*, 16, 115-130, 2005.

713 Li, Q. Y. and Xie, Z. C.: Analysis on the characteristics of the vertical lapse rates of
714 temperature. Taking Tibetan Plateau and its adjacent area as an example, *J*
715 *Shihezi University (Natural Science)*, 24, 719-723, 2006 (In Chinese).

716 Liu, J. J., Cheng, Z. L., and Su, P. C.: The relationship between air temperature
717 fluctuation and Glacial Lake outburst floods in Tibet, China, *Quat. Int.*, 321,
718 78-87, 2014.

719 Liu, Y.: Research on the Typical Debris Flows Chain Based On RS in Palongzangbu
720 Basin of Tibet, Master thesis, Chengdu University of Science And Technology,
721 2013 (In Chinese).

722 Lu, R.R. and Li, D.J.: Ice-snow-melt debris flows in the Dongru Longba, Bomi
723 county, Xizang, *J. Glaciol. Geocryol.*, 11, 148-160, 1989. (In Chinese).

724 McColl, S. T.: Paraglacial rock-slope stability, *Geomorphology*, 153-154, 1-16, 2012.

725 Noetzli, J., Gruber, S., Kohl, T., Salzmann, N., and Haeberli, W.: Three-dimensional
726 distribution and evolution of permafrost temperatures in idealized
727 high-mountain topography, *J. Geophys. Res.*, 112, F02S13, 2007.

728 Prancevic, J. P., Lamb, M. P., and Fuller, B. M.: Incipient sediment motion across the
729 river to debris-flow transition. *Geology*, 42(3), 191-194, 2014.

730 Rahardjo, H., Leong, E. C., and Rezaur, R. B.: Effect of antecedent rainfall on
731 pore-water pressure distribution characteristics in residual soil slopes under
732 tropical rainfall, *Hydrol. Process.*, 22, 506-523, 2008.

733 Rango, A. and Martinec, J.: Revisiting the degree-day method for snowmelt
734 computations, *JAWRA, J. Am. Water Resour. Assoc.*, 31, 657-669, 1995.

带格式的: 字体: 非倾斜

带格式的: 字体: (中文) +中文正文,
(中文) 中文(简体, 中国)

735 Recking A. Theoretical development on the effect of changing flow hydraulics on
736 incipient bed load motion. Water Resources Research. 45,4, W04401, 2009.
737 Rengers, F.K., L.A. McGuire, J.W. Kean and D.E. Hobley, Model simulations of
738 flood and debris flow timing in steep catchments after wildfire. Water
739 Resources Research, 52, 8, 6041-6061, 2016.
740 Rist, A.: Hydrothermal Processes within the Active Layer above Alpine Permafrost in
741 Steep Scree Slopes and Their Influence on Slope Stability, PhD Thesis, Swiss
742 Federal Institute for Snow and Avalanche Research and University of Zurich,
743 168 pp., 2007.
744 Ryzhkin, I. A. and Petrenko, V. F.: Physical mechanisms responsible for ice adhesion,
745 J. Phys. Chem. B, 101, 6267-6270, 1997.
746 Sassa, K. and Wang, G. H., Mechanism of Landslide-Triggered Debris Flows:
747 Liquefaction Phenomena due to the Undrained Loading of Torrent Deposits.
748 Debris-flow Hazards and Related Phenomena, Springer, Berlin Heidelberg,
749 81-104, 2005.
750 Sattler, K., Keiler, M., Zischg, A., and Schrott, L.: On the connection between debris
751 flow activity and permafrost degradation: a case study from the Schnalstal,
752 South Tyrolean Alps, Italy, Permafrost Periglac. Process., 22, 254-265, 2011.
753 Schneuwly-Bollschweiler, M. and Stoffel, M.: Hydrometeorological triggers of
754 periglacial debris flows in the Zermatt valley (Switzerland) since 1864, J.
755 Geophys. Res., 117, F02033, 2012.
756 Shang, Y. J., Yang, Z. F., Li, L., Liu, D. A., Liao, Q., and Wang, Y.: A super-large
757 landslide in Tibet in 2000: background, occurrence, disaster, and origin,
758 Geomorphology, 54, 225-243, 2003.
759 Springman, S. M., Jommi, C., and Teyssere, P.: Instabilities on moraine slopes
760 induced by loss of suction: a case history, Géotechnique, 53, 3-10, 2003.
761 Stoffel, M. and Huggel, C.: Effects of climate change on mass movements in
762 mountain environments, Prog. Phys. Geog., 36, 421-439, 2012.

带格式的: 字体: 非倾斜

删除的内容: .

带格式的: 字体: (中文) +中文正文,
(中文) 中文(简体, 中国)

764 Stoffel, M., Bollschweiler, M., and Beniston, M.: Rainfall characteristics for
765 periglacial debris flows in the Swiss Alps: past incidences, potential future
766 evolutions, *Clim. Change*, 105, 263-280, 2011.

767 Takahashi, T., Debris Flow: Mechanics, Prediction and Countermeasures, CRC Press,
768 Boca Raton, FL, 2014.

769 Takeuchi, Y., Kayastha, R. B., and Nakawo, M.: Characteristics of ablation and heat
770 balance in debris-free and debris-covered areas on Khumbu Glacier, Nepal
771 Himalayas, in the pre-monsoon season. *IAHS PUBLICATION*, 53-62, 2000.

772 The Ministry of Land and Resources P. R. C.: China geological hazard
773 Bulletin(September edition), 2010.

774 Theule, J.I., Liebault, F., Loye, A., Laigle, D., and Jaboyedoff, M. Sediment budget
775 monitoring of debris flow and bedload transport in the Manival Torrent, SE
776 France. *Nat. Hazards Earth Syst. Sci.*, 12, 731-749, 2012.

777 Tognacca, C., Bezzola, G. R., and Minor, H. E.: Threshold criterion for debris-flow
778 initiation due to channel bed failure, in: *Wieczoreck, G. F. (Ed.), Proceedings*
779 *Second International Conference on Debris Flow Hazards Mitigation,*
780 *Prediction and Assessment, Taipei, 89-97, 2000.*

781 Yang, W., Guo, X., Yao, T., Zhu, M. and Wang, Y.: Recent accelerating mass loss of
782 southeast Tibetan glaciers and the relationship with changes in macroscale
783 atmospheric circulations, *Clim. Dynam.*, 47, 805-815, 2016.

784 Yang, W., Yao, T., Xu, B., Ma, L., Wang, Z., and Wan, M.: Characteristics of recent
785 temperate glacier fluctuations in the Parlung Zangbo River basin, southeast
786 Tibetan Plateau, *Chin. Sci. Bull.*, 55, 2097-2102, 2010.

787 Yao, T.D., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K.,
788 Zhao, H., Xu, B., Pu, J., Lu, A., Xiang, Y., Kattel, D.B., and Joswiak, D.:
789 Different glacier status with atmospheric circulations in Tibetan Plateau and
790 surroundings, *Nat. Clim. Change*, 2, 663-667, 2012.

791 Yuan, G.X., Ding, R.W., Shang, Y.J., and Zeng, Q.L.: Genesis of the Quaternary
792 accumulations along the Palong section of the Sichuan-Tibet Highway and

删除的内容: -

带格式的: 字体: (中文) +中文正文,
(中文) 中文(简体, 中国)

带格式的: 字体: (中文) +中文正文,
(中文) 中文(简体, 中国)

删除的内容: -

795 Their distribution regularities, *Geology and Exploration*, 48, 170-176, 2012 (In
796 Chinese).
797

798

799

Table 1 Changes in glacier, snow, bare land, gully deposition and vegetation in Tianmo Valley


Year	Glacier (km ²)	Glacier(eastern branch) (km ²)	Snow (km ²)	Bare land (km ²)	Gully deposition (km ²)	Vegetation (km ²)
2000	1.77	0.16	2.13	2.80	0.44	10.46
2003	1.71	0.15	2.44	2.54	0.44	10.48
2006	1.53	0.12	2.68	2.44	0.44	10.55
2009	1.45	0.096	2.81	3.03	0.47	9.90
2013	1.42	0.088	1.74	3.83	0.51	10.17

800

801

Table 2 Basic information regarding the debris flows in Tianmo Valley and nearby valleys

No.	Name	Coordinates	Basin area (km ²)	Glacier area (in 2006) (km ²)	Date	Size class
1	Tianmo Valley	29°59'N 95°19'E	17.74	1.53	4 th Sep. 2007	6
					25 th Jul. 2010	5
					6 th Sep. 2010	5
2	Kangbu Valley	30°16'N 94°48'E	48.7	1.06	4 th Sep. 2007	3
3	Xuewa Valley	29°57'N 95°23'E	33.22	0.95	4 th Sep. 2007	2
4	Baka Valley	29°53'N 95°33'E	22.15	2.46	7 th Sep. 2007	3
5	Jiaqing Valley	30°16'N 94°49'E	15.51	1.12	9 th Sep. 2007	3




802
803

Figure 1 Location of Tianmo Valley and related information



804
805

Figure 2 Overview of the valley from the channel(in 2014)




806

807 Figure 3 DF1 in 2007(A. Overview of the Tianmo debris flows from the downstream area; B& C.

808


Boulder and debris flow deposits on the north side of the Parlung Zangbo River)



809

810


Figure 4 Particle size distributions of the glacial tills and debris flow deposits



811

812

Figure 5 Variation in the mean annual air temperature and rainfall at Bomi from 1970 to 2014




813

814

815


816

Figure 6 Distribution and changes in glacier, snow, bare land, gully deposition and vegetation in Tianmo Valley



817
818
819


Figure 7 Changes in glacier over time and the measured annual mass balance of the Parlung No. 94 Glacier (mass balance was edited from Yang et al.(2015))



820
821

Figure 8 Air temperature and rainfall before and after DF1, DF2 and DF3

带格式的: 字体: Times New Roman, 字体颜色: 文字 1



删除的内容:




Figure 9 Variations in rainfall accumulation prior to DF1 and DF3 (no rainfall before DF2)





Figure 10 Changes in glacier and frozen glacial till before periglacial debris flow initiation (A: glacial-covered glacial tills; B: uncovered and frozen glacial tills; C: active glacial tills; D: failure of glacial tills)

带格式的: 字体: Times New Roman



删除的内容: glacier glacial tills with ice

带格式的: 字体: Times New Roman, 小四, 字体颜色: 文字 1

批注 [admin33]: answer to comment: 23. Figure 10: perhaps panels C and D should be inverted. __C and D had been inverted.

823

824

825

826

827

828

页 5: [1] 删除的内容 administrator 2017-1-27 13:42:00

a rainfall even that did not hit the downstream area ceased, the local forest guard heard a loud noise coming from the upstream area at approximately 18:00

页 5: [2] 批注 [admin8] administrator 2017-1-27 15:33:00

Answer to the comment:

7. Lines 118-121. "On the morning" is not coherent with "18.00" and add t to even (line 119); I suggest to rewrite the sentence as:the triggering area was hit by a rainfall event and after that some loud noise were heard about 18:00.....

页 5: [3] 批注 [admin9] administrator 2017-1-27 15:33:00

Answer to the comment:

8. Lines 121-123 Rewrite the sentences as: a debris flow occurred after a second rainfall event that began at 19:00

页 5: [4] 批注 [admin10] administrator 2017-1-27 15:33:00

Answer to the comment:

9. Lines 122, 131 and 138 debris flows? The writer does not understand if for a debris flow event the authors intend different debris flows or a debris flow composed by several waves or something else. Please explain in the paper.

页 5: [5] 批注 [admin11] administrator 2017-1-27 15:33:00

Answer to the comment: 10. Line 128: Table 2 is not necessary because it deals with debris flows that are not object of present work. ___the other debris flows cases in table 2 is used to show that the metrological condition in these days is favorable for debris flows formation.

页 5: [6] 批注 [admin12] administrator 2017-1-27 15:33:00

Answer to the comment: 11. Lines 142-143: the finding of Chen (1991) could be due to the increase of melting water, while in present case debris flow has been triggered by rain storm.

11. Lines 142-143: the finding of Chen (1991) could be due to the increase of melting water, while in present case debris flow has been triggered by rain storm.

11. Lines 142-143: the finding of Chen (1991) could be due to the increase of melting water, while in present case debris flow has been triggered by rain storm.

页 15: [7] 删除的内容

administrator

2017-1-27 15:27:00

, created by the channelized runoff, acting on the surface elements of the debris layer and surpassing resistance of the sediment (Tognaccaet al., 2000; Gregoretti, 2000; Armanini and Gregoretti, 2000

页 15: [8] 批注 [admin29]

administrator

2017-1-27 15:38:00

answer to comment: Theule, J.I., Liebault, F., Loye, A., Laigle, D., and Jaboyedoff, M., 2012. Sediment budget monitoring of debris flow and bedload transport in the Manival Torrent, SE France. *Natural Hazard Earth Sciences*, 12, 731-749.