

Final response to the Referees' comments

First, we would like to thank the editor and the referees for the useful suggestions and constructive comments. To respond to these comments, we have made several modifications to the manuscript and we added some new refs. Please find below our response to their comments and a marked-up manuscript version showing the changes made.

Editor

Comment 1: Figure 1 has no legend for the terrestrial topography

Answer: This is a fine point by the editor. We have changed Figure 1 as suggested (section 2.2, page 5).

Referee #1

Comment 1: I would like the authors to clarify in the manuscript the drivers/components of the short-term sea level increases. Do they refer to extreme sea levels due to storm surges and waves (e.g. is wave setup included) or to storm surge alone?

Answer: This is a fine point by the reviewer. The drivers/components of the short-term sea level increases refer to extreme sea levels due to the combined effect of storm surges and wave set up. We agree with the reviewer that it needs to be better clarified in the text and we have made the necessary changes in the revised manuscript (section 2.3, page 6, lines 12-13, section 3.2, page 9, lines 10-11 and 15-17, section 4.2.2, page 19, lines 8-9)

Comment 2: Pg.3, Lines 18 – 20 There are refs to US\$ and then to C It is confusing. Please clarify; this may be confusing for the reader.

Answer: Changed as suggested (section 2.1, page 3, line 22)

Comment 3: Also Pg. 8, Lines 21 - 22 12 SLR scenarios are mentioned, but only 11 are detailed.

Answer: The examined SLR scenarios are eleven. We made the necessary correction in the revised manuscript (section 3.2, page 9, line 8)

Referee #2

Comment 1: The width of the beach has been extracted from Google Earth images and 4 operators obtained consistent results on 400 beaches, and considers irrelevant the influence of the tide (0,15 m) on its position. But it does not take into account the fact that the baric tides can have a much greater value and add up to the astronomical one.

Answer: We certainly agree with the referee that beach width estimations from satellite snapshots may not represent mean conditions, as we have stated in the text (Section 3.1, beginning of page 7). Beach width estimations are controlled by the shoreline positions, which are dynamic coastal feature, showing large spatial and temporal variability; and being strongly controlled by the beach morphodynamic processes. Recent detailed research on one of the Aegean Archipelago beaches (Ammoudara, N. Crete) for which data of high spatio-temporal resolution are available (i.e. 10 month period hourly shoreline positions), has shown high shoreline position variability (up to 6 m, or about 10-12 % of the maximum width (Velegrakis et al., 2016). We believe that any method to record beach widths using satellite imagery snapshots, one-off land based topographic and/or LIDAR surveys or, even, video-imaging of limited duration may not provide synoptic information on the 'mean' beach conditions. Accurate estimation of mean shoreline positions requires long time series of high temporal

resolution, from which estimations of the mean shoreline position could be obtained. However, such information is rarely available, particularly at the basin/Archipelago scale. Therefore, satellite imagery information appears to be the only alternative, which we deem adequate for a first assessment of the beach exposure to sea level rise over larger spatial scales. We have modified the text in order to clarify further the above issue (section 3.1, page 7, lines 3-11).

Comment 2: Sediment texture cannot be retrieved from satellite images for pixel size at ground.

Answer: The sediment texture (e.g. sand or gravel) was not retrieved from satellite images, it was assessed on the basis of the available photos on the Google Earth application and other available information collated from scientific literature/reports. To address this comment we modified the text to make it clearer to the readership (section 3.1, page 7, lines 24-27).

Comment 3: No bathymetry data are presented for beaches, which affects the distance from the shore of the depth of closure, value that enters the erosion evaluation resulting in some SLR some models (e.g., Bruun).

Answer: Given the large (Archipelago) scale of the application, the input data of the models for the evaluation of beach retreat, could not be based on in situ measurements. So we used linear profiles of a wide range of beach slopes. The distance from the shore of the depth of closure and the surf zone width, values necessary for the use of the analytical models, were estimated on the basis of the beach slope. The lack of accurate bathymetry data may introduce some uncertainty. However the validation of the models showed that the results of the models set with the equivalent linear profile were reasonably close to those of the physical experiments, and that the use of the models in an ensemble mode gave improved projections, with differences between models and experiments ranging from about 3 % to 11 % (see section 4.2.1). The aim of the exercise has not been to replace detailed modeling studies for individual beaches, but to provide ranges of beach erosion and flooding at a large (Archipelago) scale using minimum environmental information. The necessary clarification is made in the revised manuscript (section 3.2, page 8, lines 10-13).

Comment 4: Well-sorted sand was simulated, but data provided is D50, not sorting.

Answer: One of the input data needed for the models is the median sediment size D50 (not sorting), this is the reason that D50 data are provided and not sorting. We modified the text in order to make it clearer to the readership (section 4.2.1, page 13, line 24).

Comment 5: Beach rock exposure do not degrade beach aesthetics, its presence is considered a positive factor in Coastal Scenery Assessment.

Answer: In this point we do not agree with the referee. Beachrocks not only can affect the actual widths of a sandy beach as they can promote beach sediment erosion and outcropping of the initially buried beachrocks (see Vousdoukas et al., 2007, Vousdoukas et al. 2009a), but also affect perceptions regarding the beach. The presence of weathered/deformed beachrock outcrops at the beachface, commonly colonised by assemblages of epilithic and borrowing organisms (Brattström, 1992) that form a 'slippery' mat, can make the access to the sea difficult, or even dangerous, and degrade the aesthetics and amenity value of the beach and, thus, affect its touristic potential (Vousdoukas et al. 2009b). Beachrock formation/outcropping may also change the biodiversity, since beachrock outcrops create habitats suitable for colonisation by hard substrate species (e.g. corals, molluscs, algae and annelid worms) (Brattström, 1992; Vousdoukas et al., 2007; 2012). However, it is questionable whether the overall effect on the coastal ecology would be beneficial, particularly in view of the biodiversity losses in soft-substrate species (Brown, 1982). Beach aesthetics/scenery may also suffer by the presence of beachrocks, as beachrock beaches do not comply with the widely-recognisable beach model (long and wide beaches consisting of light coloured sands). With regard to the tourists' perceptions on this subject, a contingent valuation study among European tourists showed that although the majority of tourists were not previously aware of beachrock phenomenon, half of them paid notice to the hard coastal sedimentary formations. Survey respondents believe that the authorities should undertake precautionary measures and that European Union should increase research funding in order to avoid further beachrock expansion. Actually,

almost half of the respondents would be willing to pay an annual tax in the range of 13.2-16.4 € per household in order to contribute to this effort (Kontogianni et al., 2014).

We clarified further the issue by adding some refs in order to address the comment but due to space limitations we do not think that detailed documentation would be beneficial to the manuscript (section 5, page 22, lines 17-19).

References

- Brattström, H., 1992. Marine biological investigations in the Bahamas. 22. Littoral zonation at three Bahamian beachrock localities. *Sarsia* 77, 81-109.
- Brown, B., 1982. Spatial and temporal distribution of a deposit-feeding polychaete on a heterogeneous tidal flat. *Journal of Experimental Marine Biology and Ecology* 65(3), 213-227.
- Kontogianni, A., Damigos, D., Tourkolias, C., Vousdoukas, M., Velegrakis, A., Zanou, B., & Skourtos, M., 2014. Eliciting beach users' willingness to pay for protecting European beaches from beachrock processes. *Ocean & Coastal Management* 98, 167-175.
- Velegrakis, A.F., Trygonis, V., Chatzipavlis, A.E., Karambas, Th., Vousdoukas, M.I., Ghionis, G., Monioudi I.N., Hasiotis, Th., Andreadis, O., Psarros, F., 2016. Shoreline variability of an urban beach fronted by a beachrock reef from video imagery. *Natural Hazards* DOI: 10.1007/s11069-016-2415-9.
- Vousdoukas, M.I., Velegrakis, A.F., & Plomaritis, T.A., 2007. Beachrock occurrence, characteristics, formation mechanisms and impacts. *Earth-Science Reviews* 85, 23-Vousdoukas, M., Velegrakis, A.F. and Karambas, Th., 2009. Morphology and sedimentology of a beachrock-infected beach: Vatera Beach, Lesbos, Greece. *Continental Shelf Research* 29, 1937–1947.
- Vousdoukas, M.I., Velegrakis, A.F., Karambas, T.V., 2009a. Morphology and sedimentology of a microtidal beach with beachrocks: Vatera, Lesbos, NE Mediterranean. *Continental Shelf Research* 29(16), 1937-1947.
- Vousdoukas, M.I., Velegrakis, A.F., Kontogianni, A., Makrykosta, E.N., 2009b. Implications of the cementation of beach sediments for the recreational use of the beach. *Tourism Management*, 30 (4), 544-552.
- Vousdoukas, M.I., A.F. Velegrakis, M. Paul, C. Dimitriadis, E. Makrykosta, D. Koutsoubas, 2012. Field observations and modeling of wave attenuation over colonized beachrocks. *Continental Shelf Research* 48, 100-109.

Assessment of island beach erosion due to sea level rise: The case of the Aegean Archipelago (Eastern Mediterranean)

Isavela N. Monioudi¹, Adonis F. Velegrakis¹, Antonis E. Chatzipavlis¹, Anastasios Rigos^{1,2}, Theophanis Karambas³, Michalis I. Vousdoukas^{4,1}, Thomas Hasiotis¹, Nikoletta Koukourouvli⁵, Pascal Peduzzi⁶, Eva Manoutsoglou¹, Serafim E. Poulos⁷, and Michael B. Collins⁸

¹Department of Marine Sciences, University of the Aegean, University Hill, Mytilene, GR-81100, Greece

²Department of Cultural Technology and Communication, University of the Aegean, University Hill, Mytilene GR-81100, Greece

³Department of Civil Engineering, Aristotle University of Thessaloniki, University Campus, GR-54124 Thessaloniki, Greece

⁴European Commission, Joint Research Centre (JRC), Directorate for Space, Security & Migration, Disaster Risk Management Unit, Via E. Fermi 2749, Ispra (VA), I-21027, Italy

⁵Department of Geography, University of the Aegean, University Hill, GR-81100 Mytilene, Greece

⁶UNEP/DEWA/GRID-Geneva, International Environment House, 11 Chemin des Anemones, CH-1219 Châtelaine, Switzerland

⁷Faculty of Geology and Geoenvironment, National and Kapodistrian University of Athens, Panepistimioupoli Zografou, 15784 Athens, Greece

⁸Plentziako Itsas Estazioa, University of the Basque Country, Areatza z/g. E-48620, Plentzia- Bizkaia, Spain

Correspondence to: Michalis I. Vousdoukas (michalis.vousdoukas@jrc.ec.europa.eu)

Abstract. The present contribution constitutes the first comprehensive attempt to (a) record the spatial characteristics of the beaches of the Aegean Archipelago (Greece), a critical resource for both the local and national economy; and (b) provide a rapid assessment of the impacts of the long- term and episodic sea level rise (SLR), under different scenarios. Spatial information and other attributes (e.g. presence of coastal protection works and backshore development) of the beaches of the 58 largest islands of the Archipelago were obtained on the basis of remote-sensed images available in the web. Ranges of SLR-induced beach retreats under different morphological, sedimentological and hydrodynamic forcing and SLR scenarios were estimated, using suitable ensembles of cross-shore (1-D) morphodynamic models. These ranges, combined with empirically-derived estimations of wave run up-induced flooding, were then compared with the recorded maximum beach widths, to provide ranges of retreat/erosion and flooding at the Archipelago scale. The spatial information shows that the Aegean ‘pocket’ beaches may be particularly vulnerable to mean (MSLR) and episodic SLRs due to: (i) their narrow widths (about 59 % of the beaches have maximum widths < 20 m); (ii) their limited terrestrial sediment supply; (iii) the substantial coastal development and (iv) the limited existing coastal protection. Modeling results indeed project severe impacts under mean and episodic SLRs, which by 2100 could be devastating. For example, under MSLR of 0.5 m (RCP4.5), a storm-induced sea level rise of 0.6 m is projected to result in complete erosion of between 31 and 88 % of all beaches (29 - 87 % of beaches currently fronting coastal infrastructure and assets), at least temporarily. Our results suggest a very considerable risk which appears that, in addition to, will require the significant effort, and financial resources and policies/regulation in

~~order required to protect/maintain the critical economic resource of the Aegean Archipelago; appropriate coastal 'set-back zone' policies should be adopted and implemented.~~

1 Introduction

Beaches are critical components of the coastal zone; not only are they significant habitats in their own right (e.g. Defeo and McLachlan, 2013), but also provide protection from marine flooding to other transitional ecosystems and the coastal assets, infrastructure and activities they front (e.g. Neumann et al., 2015). At the same time, tourism has been increasingly associated with beach recreational activities according to the dominant 'Sun, Sea and Sand-3S' tourism model (Phillips and Jones, 2006). Consequently, beaches have become very important economic resources (Ghermandi and Nunes, 2013), forming one of the pillars of tourism, an economic sector that contributes an estimated 5 % of Global Gross Product - GGP, and about 6 – 7 % of global employment (directly and indirectly) (Hall et al., 2013).

Beaches are also very dynamic environments, controlled by complex forcing-response processes that operate at various spatio-temporal scales (Short and Jackson, 2013). They are generally under erosion (EuroSION, 2004; IPCC SREX, 2012; IPCC, 2013), which can be differentiated into (a) long-term erosion, i.e. irreversible retreat of the shoreline, due to mean sea level rise (MSLR) and/or negative coastal sedimentary budgets that force either beach landward migration or drowning (Nicholls and Cazenave, 2010); and (b) short-term erosion, caused by storm surges and waves, which may, or may not, result in permanent shoreline retreats but can be nevertheless devastating (e.g. Smith and Katz, 2012; UNECE, 2013). The accelerating MSLR coupled with episodic storm events will aggravate the already significant beach erosion with severe impacts on coastal activities, infrastructure and assets (e.g. Jiménez et al., 2012) and the beach carrying capacity for recreation/tourism (Valdemoro and Jiménez, 2006; McArthur, 2015).

Beach erosion appears to be particularly alarming in islands. Island beaches are increasingly vulnerable to erosion due to their (generally) limited dimensions and diminishing sediment supply (e.g. Velegrakis et al., 2008; Peduzzi et al., 2013). At the same time, island beaches are amongst the most significant 3S tourism destinations. For example, 3S tourism accounts for more than 23 % of the Gross Domestic Product - GDP in many Caribbean Small Island States - SIDS and, in some cases, e.g. Antigua and Barbuda, for more than 75 % (ECLAC, 2011). Mediterranean islands are also major tourism destinations; in Greece, most of the hotel capacity and foreign tourist arrivals and earnings are associated with the Greek islands (SETE, 2016).

Under a variable and changing climate, projections on the future evolution of beach morphology are not easy, due to uncertainties regarding both forcing and beach response (e.g. Short and Jackson, 2013). Nevertheless, beach erosion is amongst the first issues to consider when planning for the sustainable development of the coastal zone, particularly in areas where beaches function as natural 'armor' to valuable coastal infrastructure and assets and/or as significant environments of leisure (e.g. Paula et al., 2013). Therefore, assessments of the beach morphological evolution at different spatio-temporal scales are required, based on advanced numerical, analytical, and/or empirical models constructed and applied by

experienced operators, set up/validated using appropriate field data and backed by expert analysis (e.g. Roelvink et al., 2009; Bosom and Jiménez, 2010; Ding et al., 2013). However, such efforts are usually hampered by the (a) scarcity of relevant information in many coastal areas, and (b) dearth in the necessary human and financial resources (e.g. Parker et al., 2013); this is particularly true when assessments of beach erosion are carried out over larger spatial scales. All the same, it is necessary to assess future beach retreat/erosion and flood risk at large spatial scales, in order to identify ‘hot spots’ and plan for effective adaptation policies and efficient allocation of resources.

Against this background, the objective of the present study is to assess the erosion and temporary inundation/flood risks of the beaches of the Aegean Archipelago islands of the Aegean Archipelago (Greece) under different scenarios of SLR. Towards this objective, spatial characteristics such as the area, length, maximum width, orientation, sediments and the presence of coastal works and backshore development of the Aegean beaches were recorded. This information was then used in conjunction with projections from ensembles of cross-shore morphodynamic models to obtain estimates of the ranges of potential beach retreat/erosion and flooding under different MSLRs and storm condition events.

2 Aegean Archipelago beaches: Significance, environmental setting and sea level rise

2.1 Significance of the Aegean Archipelago beaches

Aegean Archipelago (Fig. 1) consists of several thousand islands and rock islets, with a combined area of 17550 km² and total coastline length of about 5880 km (Eurosion, 2004). Few of these islands and islets are populated; less than 70 islands have more than 100 and 45 more than 1000 permanent inhabitants (http://www.statistics.gr/portal/page/portal/ESYE/PAGE-themes?p_param=A2001). Yet, Aegean islands form very significant tourist destinations. 50 % of all Greek hotel beds (and > 60 % of all 5 star hotel beds) are located in the Aegean Archipelago with 43 % of the foreign arrivals to Greece in 2015 (7.4 out of a total of 17.1 million) arriving at its 11 international airports (SETE, 2016).

In recent years, tourism has become a most significant economic activity in Greece. In 2013, foreign earnings of the Greek tourist industry were about €15.5 billion (€15.5 billion). As recent studies suggest that for each 1 € generated by tourism in Greece, an additional 1.2 - 1.65 € is created by related economic activity (a multiplier of, at least, 2.2 see IOBE (2012)), it follows that direct and indirect earnings from tourism may account for up to about 20 % of the country's GDP (and 30 % of the private sector employment). Tourism is even more important for the island (local) economies. For example, in 2012 tourism accounted for about 48 % of the GDP of Crete and 60 % of the GDP of the Cyclades and Dodecanese island complexes (SETE, 2016).

In Aegean Archipelago, 3S tourism is the dominant model. A most critical component of 3S tourism is the availability of beaches that are aesthetically and environmentally and aesthetically sound and retain adequate carrying capacity (e.g. McArthur, 2015; Cisneros et al., 2016). Therefore, the assessment/management of beach erosion which could constitute a major threat risk for the sustainable development of the Aegean islands beaches should be prioritized; a decade-

Σχόλιο [I1]: Changed as suggested by Referee #1, Comment 2.

old approximation had suggested that about 25 % of the total coastline of the Aegean islands was already under erosion (Eurosion, 2004).

2.2 Environmental Setting

Aegean Archipelago is located at the Aegean Sea, a peripheral sea of the Eastern Mediterranean that covers an area of some 160 x 103 km², drains high relief basins with a total area of 200 × 103 km² and is connected to Black Sea through the Dardanelles Straits and to Eastern Mediterranean through the Cretan Arc Straits. Aegean Sea ~~has a high relief is characterized by irregular morphology~~ due to complex regional tectonics and comprises different geomorphological units (Poulos, 2009), including: an extensive shelf (N. Aegean Shelf), a tectonic trough (N. Aegean Trough), a central platform (Cyclades Plateau) with large concentration of islands, as well as deep basins (~~some > 2500 m deep~~) ~~mainly in the South Aegean~~ (Fig.1).

Aegean Sea shows complex hydrographic patterns and circulation (e.g. Theocharis et al., 1993) which are ~~mainly partly~~ controlled by the cold and low salinity water inputs from the Black Sea through the Dardanelles Strait and the warm and saline water inputs from the Levantine Sea through the Eastern Cretan Arc Straits (Skliris et al., 2011). Under certain conditions (the Eastern Mediterranean Transient -EMT), the ~~deep~~-Aegean basins ~~can be important~~ ~~have been observed to be~~ the locations of deep water formation in the Eastern Mediterranean (e.g. Zervakis et al., 2000; Androulidakis et al., 2012).

The complex physiography of the Aegean Archipelago controls its wind and wave climate, which is generally relatively mild due to the short fetches and durations. Northerly winds (44% frequency of occurrence, Androulidakis et al. (2015)) and waves (Soukissian et al., 2007; 2008) appear to prevail, ~~and, although~~ Although waves are generally more energetic in winter, there are also energetic events in summer, forced by N-NE winds ('the Etesians'). Highly energetic wave events of relatively short duration may also occur, particularly along island straits. Soukissian et al. (2008) suggested ~~as the most energetic areas of the Aegean Archipelago~~ (i) the area to the N-NE of the Cyclades platform (particularly the Mykonos-Ikaria Strait) and (ii) the western and eastern Cretan Arc Straits (Fig.1) as the most energetic areas of the Aegean Archipelago; for example, maximum wave heights of about 11 m (Tp of 13.3 s and direction of 345^{±0} N) have been reported for the Mykonos-Ikaria Strait ~~in~~ (22/01/2004). Analysis of ERA-INTERIM wave information (1979-2013) from different representative areas of the Aegean Archipelago carried out as part of the present study shows: (a) mean significant wave heights (Hs) of about 1 m in all areas, apart from an area to the northeast of eastern Cretan Arc Straits (mean Hs of about 0.8 m); (b) mean maximum wave heights of about 2.4 m; and (c) significant interannual variability.

Recent studies on the future wave climate of the Aegean Archipelago, project small changes in significant wave heights for the 21st century. For the period 2001-2049, significant wave heights in the N. Aegean are projected to slightly increase for the SW waves relative to the 1950 - 2000 reference period, whereas for the end of the century (2050 – 2099), wave occurrence and heights ~~patterns~~ are projected to show high spatio-temporal variability (e.g. Prinos, 2014; Tsoukala et al., 2016).




Figure 1: The Aegean Archipelago

Σχόλιο [I2]: The Figure was changed as suggested by the editor

2.3 Mean ~~and Extreme~~ Sea Levels

Mediterranean MSLR rates were $1.1 - 1.3 \text{ mm yr}^{-1}$ for the most part of the 20th century. Since the late 1990s, however, much higher rates ($2.4 - 3.8 \text{ mm yr}^{-1}$) have been recorded, an increase attributed mainly to additional water mass inputs rather than to steric contributions; ~~these changes have been also related to being also correlated with~~ North Atlantic Oscillation-NAO modulations (Tsimplis et al., 2013). For the Aegean Archipelago, in particular, satellite altimetry suggests recent MSLR rates of $4.3 - 4.6 \text{ mm yr}^{-1}$ (Mamoutos et al., 2014), with some periods characterized by even higher rates (e.g. ~~up to 15.3 mm yr^{-1} , see Tsimplis et al., (2009).~~). In terms of future projections, recent studies project decreases in the MSLR rates in the Aegean Sea for the 21st century; ~~although this might benevertheless, such trends could be an~~ underestimations due to the uncertainties regarding ~~water the~~ mass exchanges particularly between the Black and Aegean Seas (Mamoutos et al., 2014). Hinkel et al. (2014) using an approach ~~that accounts accounting~~ for changes in ice mass and ocean circulation ~~projected the following suggested the following likely future~~ MSLRs for the region of the Aegean Archipelago ($33.5^{\circ} - 40^{\circ} \text{ N}$,

18.5⁰-28.5⁰ E). In 2050, ~~mean sea level~~MSL was projected to be 0.13 - 0.15 m and 0.14 - 16 m higher than that of the 1985 – 2005 reference period for a medium land-ice scenario and RCPs 4.5 and 8.5, respectively; in 2100, under the same scenarios, ~~mean sea level is~~MSL was projected to be 0.46 - 0.48 m and 0.66 - 0.72 m higher than that of the ~~1985–2005~~ reference period, respectively.

In addition to MSLR, changes in the intensity, frequency and/or patterns of extreme storm surges and waves can, at least temporarily, induce beach erosion and flooding, particularly when combined with increasing mean sea levels (e.g. Xu and Huang, 2013). Extreme sea levels (ESLs) in the Mediterranean have a seasonal footprint with extreme positives levels occurring mostly in winter and under certain North Atlantic Oscillation (NAO) modulations (Tsimplis and Shaw, 2010). ~~In the Aegean Archipelago, extreme sea levels are relatively low (heights of up to about 0.5 m) (Tsimplis and Shaw, 2010; Krestenitis et al., 2011), increasing slightly towards the north (Androulidakis et al., 2015).~~

Future ESLs are projected to show~~extreme sea levels will be associated with~~ high spatial variability, being sensitive to the evolution of the thermohaline circulation and the Black Sea buoyant inputs (e.g. Mamoutos et al., 2014). Generally, ~~e~~Extreme levels due to storm surges and (waves) are projected to decrease in terms of magnitude~~show generally decreasing trends~~ over the Mediterranean basin towards the end of the 21st century (e.g. Conte and Lionello, 2013). although ; nevertheless, model choice and resolution (Marcos et al., 2011; Androulidakis et al., 2015; Vousdoukas et al., 2016) and the quality/resolution of the ~~available~~ coastal observations available for model validation (~~Calafat et al., 2014~~) may have influenced ~~thesesuch~~ projections (e.g. Calafat et al., 2014). For example, a recent study has suggested substantial changes in the return periods of ESLs for E. Mediterranean for the end of the 21st century, with the current 1000-year event projected to occur every 5 years (Vousdoukas et al., under review). Regarding the storm surges in the Aegean Archipelago, these are relatively small (heights of up to about 0.5 m, Krestenitis et al. (2011)) and~~Storm surges are generally~~ projected to show (generally) small height increases until 2050, as well as changes in their temporal distribution (e.g. Marcos et al., 2011; Androulidakis et al., 2015; Vousdoukas et al., 2016a).

Σχόλιο [I3]: This part is slightly changed to account for the Comment 1 of Referee #1. There are other clarifications on this issue later

3 Materials and Methods

3.1 Geo-spatial characteristics of the Aegean Archipelago beaches

The geo-spatial characteristics of Aegean Archipelago ('dry') beaches have been recorded, on the basis of the ~~images~~ imagery and other related optical information available in the Google Earth Pro application. In this study, 'dry' beaches were defined as the low-lying coastal sedimentary bodies bounded on their landward side by either natural boundaries (vegetated dunes and/or cliffs) or permanent artificial structures (e.g. coastal embankments, seawalls, roads, and buildings) and on their seaward side by the shoreline, i.e. the median line of the foaming swash zone shown on the imagery. Regarding the lateral extent of individual beaches, these were delimited by natural barriers, such as rock promontories. Tiny beaches (areas less than about 20 m²) were ignored/not included in the data set. Digitization of the remote-sensed imagery was carried out by few (4) analysts, who followed consistently the above beach delimitation rules. To assess inconsistencies, 400 beaches from

different islands (about 12 % of the recorded beaches) were processed by all 4 analysts and the standard deviation of the extracted shoreline positions was estimated less than 0.3 m; this was considered acceptable for the scope of the study.

However, it should be noted that the primary information used i.e. satellite imagery snapshots cannot provide synoptic information at the Aegean Archipelago scale, as the available images have been collected in different periods and under variable hydrodynamic conditions. Regarding the beach width estimations, although astronomical tidal effects are small due to the microtidal regime of the Aegean Archipelago (tidal ranges in most areas less than about 0.15 m) and the generally increased beach slopes, there are still uncertainties, as the estimations were based on ‘snapshots’ of the shoreline positions which are very dynamic coastal features controlled, amongst others, by the spatio-temporal variability of the nearshore waves and currents (e.g. Velegrakis et al., 2016). Satellite imagery, as well as other ‘one-off’ land-based, airborne or LIDAR surveys provide only ‘snapshots’ of beach widths which may not represent ‘mean’ conditions; nevertheless, this can not be avoided when working at large spatial scales (e.g. Allenbach et al., 2015; Vousdoukas et al., 2016a).

Beaches were digitized as polygons and exported to a GIS for further analysis. There has been no geo-rectification, as the aim of the exercise has not been to provide definitive locations and elevations of beach features, but to extract/record (horizontal) geo-spatial characteristics. To this end, a custom-made AML (ARC Macro Language, proprietary language for ArcInfo applications in ESRI software) script was used to estimate beach areas, lengths, maximum widths and orientations (Altenbach et al., 2015). It should be noted that the satellite imagery used cannot provide synoptic information at the Aegean Archipelago scale, as images have been collected in different years, seasons and hydrodynamic conditions. Although tidal effects are small due to the microtidal regime of the Aegean Archipelago (tidal ranges in most areas less than about 0.15 m) and the (generally) increased beach slopes, geo-spatial characteristics controlled by the shoreline position and obtained from remotely sensed snapshots may not represent mean conditions (Velegrakis et al., 2016). Nevertheless, this cannot be avoided when working at a basin/Archipelago scale (Altenbach et al., 2015).

In addition to beach dimensions, other relevant information was recorded and codified, including: the presence of (a) natural features, such as river mouths, back-barrier lagoons and cliffs and beachrock outcrops; and (b) artificial features such as coastal protection schemes and backshore infrastructure/assets. Classification Assessment of the beach sediments of the Aegean island beaches in terms of sediment texture classes (e.g. sand or gravel) was also carried out on the basis of all available web-based optical information from the scientific literature/reports (e.g. Monioudi et al., 2016) as well as web-based photographic material and other available information collated from scientific literature/reports.

3.2 Beach retreat predictions due to sea level rise

Sea level rise represents a most significant hazard for threat to beaches, forcing their retreat/erosion; a sea level rise α will result in a shoreline retreat S due to erosion of the beach face, the sediments of which are transported/deposited offshore, with the extent/rates of the cross-shore retreat controlled (amongst others) by bed slope, the texture and supply of beach sediments and the hydrodynamic conditions (e.g. Dean, 2002).

Σχόλιο [a4]: Response to Comment 1 of Referee #2

Σχόλιο [a5]: Response to Comment 2 of Referee #2

In the present study, seven cross-shore (1-D) morphodynamic models were used to project beach response to SLR: the Bruun (Bruun, 1988), Edelman (Edelman, 1972) and Dean (Dean, 1991) analytical models and the numerical models SBEACH (Larson and Kraus, 1989), Leont'yev (Leont'yev, 1996), XBEACH (Roelvink et al. 2010) and a model, the hydrodynamic component of which involves high-order Boussinesq equations - Boussinesq model (Karambas and Koutitas, 2002). The Bruun model is a widely-used (e.g. Hinkel et al., 2010; Ranasinghe et al., 2013) analytical morphodynamic model that estimates long-term coastal retreat S under a SLR a on the basis of the equilibrium profile concept; its results are controlled by the height of the beach face and the cross-shore distance between the beach closure depth and the shoreline. The Edelman model estimates beach erosion/retreat using the initial height of the beach face, the water depth at wave breaking and the surf zone width, whereas the Dean model estimates retreats on the basis of the wave height, the water depth at wave breaking and the surf zone width. It should be noted that due to lack of accurate bathymetry data, the necessary input values of the analytical models i.e. the distance of the shoreline from the closure depth and the width of the surf zone were estimated on the basis of the beach slope.

The SBEACH model (Larson and Kraus, 1989) is a numerical morphodynamic model, consisting of 3 modules: a hydrodynamic, a sediment transport and a morphological evolution module. It can describe wave transformation in shoaling waters, with the coastal sediment transport controlled by the coastal wave energy fluxes; the sediment continuity equation in a finite difference scheme and a 'stair-step' beach profile discretization is used in its morphological module. The numerical model based on Leont'yev (1996) uses the energetic approach, with the cross-shore wave energy balance controlled by wave propagation angle and dissipation; sediment transport rates are estimated separately for the surf and swash zones. The XBEACH model (Roelvink et al., 2010) is an open-source, widely used numerical model of the nearshore processes intended to estimate the effects of time-varying storm conditions; it contains a time-dependent wave action balance solver and allows for variations in the wave action over time and over the directional space. Finally, the Boussinesq model used computes non-linear wave transformation in the surf and swash zone, based on a wave propagation module involving high-order Boussinesq equations (Karambas and Koutitas, 2002); its sediment transport module can estimate sheet flow as well as bed and suspended load over uneven sea beds (e.g. Karambas, 2006). Detailed descriptions of the models used can be found elsewhere (e.g. Vousdoukas et al., 2009a; Monioudi et al., ~~2014~~2016).

In the present contribution, all models were used in a stationary mode. Validation of model results was provided through comparisons with the results of physical experiments in the GWK wave flume, Hanover, Germany (~~see~~ Section 4.2.1). The models were used in an ensemble mode in order to assess the range of long- and short-term beach retreats/erosion for different beach slopes, sediment ~~textures~~ (grain size) ~~s~~ and wave conditions, and under different scenarios of MSL changes and/or ~~storm sea levels~~ extreme sea levels caused by storm surges/waves. Two model ensembles were created, a 'long-term' ensemble consisting of the analytical models Bruun, Dean and Edelman and a 'short-term' ensemble comprising the numerical SBEACH, Leont'yev, XBEACH and Boussinesq models. ~~‡~~ The former is used to assess beach retreat/erosion under MSLR, ~~whereas~~ and the latter retreat due to temporary SLR (i.e. ~~from due to~~ storm surges ~~and~~ waves. The adopted

Σχόλιο [I6]: Response to Comment 3 of Referee #2

approach was based on the proposition that as models have differential sensitivity to the controlling environmental factors, ensemble applications may provide ‘tighter’ prediction ranges than the individual models (Section 4.2.1).

Experiments were carried out using various plausible energetic wave conditions in the Aegean Archipelago (Section 2.2), i.e. waves with offshore heights (H_s) of 1, 1.5, 2, 3 and 4 m and periods (T) of 4, 5, 6, 7 and 8 s. Likewise, in order to address the sediment texture variability over the Archipelago beaches, experiments were carried out for seven different median (d_{50}) grain sizes (d_{50} of 0.2, 0.33, 0.50, 0.80, 1, 2 and 5 mm); note that the results of the analytical models are not controlled by beach sediment size. Five (5) different linear profile slopes (bed slopes of 1/10, 1/15, 1/20, 1/25 and 1/30) and ~~twelve~~ eleven (11) SLR scenarios (0.05, 0.15, 0.22, 0.30, 0.40, 0.50, 0.75, 1, 1.25, 1.50 and 2 m) were examined.

Experiments were carried out for all combinations (about 5500 experiments), and the means (best fits) of the lowest and highest projections by all models of the two ensembles were estimated. With regard to combined SLRs (i.e. coastal ESLs due to storm surges and wave set ups (e.g. Vousdoukas et al., 2016b) superimposed on the MSLRs), the long-term and short-term ensembles were used consecutively.

Recent MSLR projections for the area that take into account ie mass ~~the~~ contributions of ice mass were used (Hinkel et al., 2014): (i) 0.15 m, average of the RCP 4.5 and RCP 8.5 scenarios in 2040; and (ii) 0.5 m and 0.7 m under RCP 4.5 and RCP 8.5 (2100), respectively. With regard to short-term SLR, recent trends ~~projections~~ for Extreme Sea Levels (ESLs) storm surge heights in the Aegean Archipelago (up to 0.5 - 0.6 m high) (Tsimplis and Shaw, 2010) are used; ESLs in the region are projected to only slightly increase in the future according to recent projections regarding their storm surge component (Androulidakis et al., 2015; Vousdoukas et al., 2016a) ~~-were used~~.

The above approach is designed to project beach retreat/erosion, but not temporary inundation/flooding due to wave run-up. Although the wave run-up is dealt within the numerical models of the ensemble, its effects are manifested in the results only if it induces sediment transport that forces morphological changes (e.g. Leont'yev (1996)). Yet, wave run up-induced temporary flooding that does not result in beach retreats might ~~is~~ also be a significant beach management issue (e.g. Jiménez et al., 2012; Hoeke et al., 2013). Therefore, estimations of wave run up excursion/inundation were also undertaken on the basis of run up heights; these were estimated for all tested conditions, using the expressions of Stockdon et al. (2006) which have been validated for the beaches of the Aegean Archipelago (Vousdoukas et al. 2009b):

$$R_{2\%} = 1.1 \left(0.35\beta(H_o/L_o)^{1/2} + \frac{[H_o L_o (0.563\beta^2 + 0.004)]^{1/2}}{2} \right), \quad (\text{all data}) \quad (1)$$

$$R_{2\%} = 0.043(H_o/L_o)^{1/2}, \text{ for dissipative beaches } (\xi < 0.3) \quad (2)$$

where $R_{2\%}$, the 2% exceedence of the peak run-up height, H_o , L_o are the deep water wave height and length, β the beach slope and ξ the Iribarren number ($\xi = \beta/(H_o/L_o)^{1/2}$).

Σχόλιο [17]: Corrected as recommended in Comment 3 of Referee #1.

Σχόλιο [18]: This is again to highlight the case of the coastal ESLs being the sum of storm surge and wave set up (Comment 1 of Referee #1).

Σχόλιο [19]: These changes were made to clarify further the issue posed in Comment 1 of Referee #1

Wave run up excursions were then calculated from the wave run up heights ($R_{2\%}$) for all tested bed slopes and wave conditions and added to the beach erosion/retreat projections of the seven 1-D cross-shore morphodynamic models to project final flooding excursions ($S(i)$). The best fits of the lower and upper limits of the final projections of flooding by all models were then estimated.

5 4 Results

4.1 Characteristics of the beaches of the Aegean Archipelago

3234 beaches were recorded along the coasts of the 58 larger islands of the Aegean Archipelago. These beaches were found to be 'pocket' beaches, having a total area of about 21.35 km²; ~~indicating this indicates~~ that the total carrying capacity of the Aegean Archipelago beaches (i.e. the number of visitors that can be simultaneously hosted), is about 2135 thousand according to the Rajan et al. (2013) criterion (10 m² for each beach user). A rough estimation on the basis of the average number of tourist days spent in the Aegean islands suggests that, with the exception of the busiest months (July and August) during which about 7000 thousand tourist days per month are recorded (SETE, 2016) and many beaches operate close to their full carrying capacity, the Archipelago beaches in their present condition still have (as a total) potential for development as environments of leisure.

15 Most of the Aegean beaches are narrow, with about 59 % of them having recorded maximum widths of less than 20 m (Fig. 2). Regarding beach sediment type, three different ~~beach types~~ classes were recorded: (i) sandy beaches (36.4 %); (ii) gravel/pebble beaches (44.4 %); and (iii) cobble/rocky beaches (about 2 %). For the remainder of the beaches, no sediment type could be assigned on the basis of the available information (Fig. 2).




Figure 2: Statistics of the Aegean Archipelago beaches. (a) Pie diagrams of major characteristics; (b) relationship between the beach maximum width and the presence of river mouth; and (c) boxplots showing relationships between beach widths and sediment types.

Beach orientation was examined by: (i) taking into consideration the 8 main compass directions; and (ii) considering the 8 main compass directions for each of the 8 island complexes of the Aegean Sea (N. Aegean, Dodecanese, Cyclades, Crete, SW Aegean, Argosaronikos, Evia, and Sporades, Fig. 1). In the first case, occurrence of beaches with S, SE and SW orientations appears more prevalent (16.1 %, 13.9 % and 12.5 % respectively) than that of beaches facing towards the other 5 compass directions; this was found to be statistically significant (Table 1). Prevailing beach orientations were also found to differ over the different island complexes. For example, the prevailing (statistically significant) beach orientations in the Dodecanese islands were found to be towards the SE (18 %) and in Crete towards the S (22%). There is also significant correlation between beach orientation and recorded maximum width. Statistical tests (Table 1) showed that W and SW facing beaches are associated with greater maximum widths (24.5 and 24.0 m, respectively).

Table 1: Statistical analysis of the characteristics of the Aegean Archipelago beaches.

Comparison	Null Hypothesis (H_0)	Statistical test results	Observations
Beach orientation and occurrence	H_0 : The data are uniformly distributed around the compass rose	<i>Rayleigh's Test</i> : H_0 rejected ($p < 0.001$). Beach orientation not uniformly distributed over the compass rose	Prevailing beach orientation towards the south (S) (occurrence 16.1 %)
Beach orientation and maximum width	H_0 : The 8 compass orientations have equal mean values regarding maximum width ($\mu_1 = \mu_2 = \dots = \mu_8$) H_0 : $\rho_M = 0$, No correlation in the data population	<i>Kruskal-Wallis (Analysis of variance by ranks)</i> : H_0 rejected ($p < 0.001$). Mean beach maximum widths differ with orientation <i>Mardia's correlation coefficient (Zar, 2010)</i> : H_0 rejected ($p < 0.01$). Significant correlation between orientation and maximum width	W and SW facing beaches have greater maximum widths (mean maximum widths of 24.5 and 24 m)
Beach orientation and beachrock occurrence	H_0 : the two variables are independent	χ^2 tests for (i) 4 and (ii) 8 main compass orientations: H_0 rejected ($p < 0.001$). Small association's power Cramer's (i) $V = 0.13$ and (ii) $V = 0.17$	No significant trend
Sediment type and presence of river mouth	H_0 : Beach sediment type independent of the river mouth presence	χ^2 test: H_0 rejected ($p < 0.05$). Significant correlation, but small power of association (Cramer's $V = 0.064$)	Sand-sized beaches are more Beaches with coarse grained sediments are less associated with river mouths
Beach maximum width and presence of river mouth	H_0 : the sets are independent (mean values ($\mu_1 = \mu_2$))	<i>Student's t-test</i> (two-tailed test): H_0 rejected ($p < 0.001$)	Beaches with large maximum widths are associated with river mouths (Fig. 2b)
Beach sediment type and maximum width	H_0 : Mean values of the 3 categories are equal ($\mu_1 = \mu_2 = \mu_3$)	<i>ANOVA</i> , after splitting the data set into the 3 sediment types: H_0 rejected ($p < 0.001$).	Beaches with coarse sediments are associated with smaller maximum widths (Fig. 2c)

The above results indicate some hydrodynamic control in the development and maintenance of the Aegean beaches: there are more and wider beaches along the island coasts that are (at least) partially protected from the prevailing northerly wind/waves (Soukissian et al., 2007; 2008). However, the correlation is weak (Table 1) due to other important factors controlling island beach development ~~and maintenance, such as (e.g. the antecedent topography and geological history, the~~ 5 ~~terrestrial sediment supply, and the human development and the wave fetch/duration at each individual beach).~~

In terms of terrestrial sediment supply, few beaches (about 18 %) were found to be associated with intermittent (very rarely permanent) flow river mouths; most of those were found in the large islands of North Aegean (314 out of a total of 728 beaches). Riverine supply appears to be a significant control for both beach width and sediment type; beaches with river 10 mouths are more likely to be wider and built on finer sediments (sands) (Fig. 2 and Table 1).

Another interesting finding is that perched beaches (Gallop et al., 2012) form a significant fraction of the Aegean Archipelago beaches; beachrock outcrops (e.g. Vousdoukas et al., 2009a) were recorded on the beach face of 23 % of all beaches. In terms of orientation, although there appears that beaches with beachrocks are more prevalent at the southern coasts of islands (192 out of 744 total occurrences), statistical testing did not show any significant trends (Table 1). As there

could be a significant number of beaches which may contain buried beachrocks (i.e. not outcropping at the time of the analysed imagery), it seems that beachrocks are quite widespread along the Aegean island beaches.

With regard to coastal protection schemes ~~have been recorded, these are present only~~ at about 15 % of the Aegean island beaches (Fig. 2a). In comparison, many beaches are associated with coastal development: 80.8 % of all beaches front public and private assets such as coastal roads, housing and tourist infrastructure. The density of these assets is variable, ranging from a coastal road and few houses found in remote island beaches to the plethora of valuable public and private assets found behind the urban beaches of e.g. Heraklion (Crete) and Rhodes. Generally, about 32.7 % of the Aegean beaches front coastal infrastructure and assets with moderate/high density. The considerable coastal development, coupled with the narrow widths of the Aegean island beaches increase exposure under a variable and changing climate.

4.2 Predictions of erosion/retreat and flooding for the Aegean island beaches

4.2.1 Model sensitivity and validation

Model sensitivity tests undertaken within the present study have shown that beach retreats are controlled by beach typology. All models show higher retreats with decreasing Iribarren numbers ξ (i.e. for beaches with milder beach slopes β and/or offshore waves of increased steepness (H_o/L_o), with the exception of the Bruun model the results of which are independent of ξ for linear profiles. The most sensitive models to beach slope are the Edelman and SBEACH models and the least sensitive the Xbeach model; with regard to the offshore wave climate, Xbeach appears to be the most and Leont'yev and Boussinesq the least sensitive models. The effect of sediment texture is not always clear, although a weak negative correlation between beach retreat and the median sediment size (d_{50}) might be discerned in the numerical models; analytical model results are independent of the beach sediment texture. Generally, models showed differential sensitivity to initial conditions and forcing, which justifies their collective use in ensembles.

Model results have been compared with those by physical experiments conducted at the wave flume (GWK, Hanover, Germany) in early 2013 (details in Voudoukas et al. (2014)). In these experiments, the initial slope of the beach was set to about 1/15, tested waves had an offshore height (H) of 1 m and a period (T) of 5 s and the seabed consisted of well-sorted sand of median grain size (d_{50}) equal to 0.3 mm (~~$d_{50} = 0.3$ mm~~). Three SLR scenarios were tested (rises of 0.2, 0.4 and 0.6 m), with the initial profiles of these experiments controlled by the final profile (wave forcing only) of the first experiment (i.e. the test without level rise). Simulation times were set to 3000 s.

In Fig. 3, profiles by the numerical models Leont'yev, SBEACH, Xbeach and Boussinesq under the baseline level as well as 3 increased sea levels (+0.2, +0.4, and +0.6 m) are compared against those resulted from the physical experiments. It appears that there are some discrepancies, particularly with regard to the dynamics of the offshore bar and trough. The Leont'yev, and Xbeach models seem to considerably 'smooth' these features, whereas the SBEACH and Boussinesq results tend to represent better the results of the physical experiments; the best performance was by the Boussinesq model (Fig. 3). Nonetheless, there appears to be a good agreement between the beach retreats estimated by the models and those recorded

Σχόλιο [I10]: A small change in order to address Comment 4 of Referee #2.

during the physical experiments (Fig. 3 and Table 2). Leonty'ev, SBEACH and Boussinesq models appear to slightly underestimate and Xbeach to overestimate beach retreats. Bruun and Edelman models appear to overestimate beach retreats under higher SLRs whereas the Dean model under all SLRs tested.




Figure 3: Profiles by numerical models plotted against results from physical experiments at the GWK wave flume (Hanover): (a) initial/present water level; (b) water level rise of 0.2 m; (c) rise of 0.4 m; and (d) rise of 0.6 m. Both numerical and physical experiments were set up for the same initial (non-linear) profile (stippled line). Simulation times for the numerical experiments set to 3000 s to match those of the physical experiment.

When the models are used in an ensemble mode, the comparison between the retreats projected by the numerical models (short-term retreat) and those recorded in the physical experiments improves (Table 2). Conversely, analytical models tend to overestimate beach retreats by up to 20 %. Interestingly, when the results of all (7) analytical and numerical models are combined in a unified ensemble, then retreat overestimations decrease to less than 10 % for the tested SLRs.

Table 2: Comparison of results from models and physical experiments. Negative values represent erosion. Key: SLR, sea level rise; Exper, physical experiment; Leo, Leont'yev model; SB, SBEACH model; Xb, Xbeach model; Bous, Boussinesq model; Br, Bruun model; Edel, Edelman model; Dean, Dean model; ST Ens, Short-term ensemble; and LT Ens, Long-term ensemble.

SLR (m)	Shoreline retreat/erosion or advance/accretion (m)									
	Exper	Leo	SB	Xb	Bous	ST Ens	Br	Edel	Dean	LT Ens
0	-0.35	-0.01	0.02	-2.32	-0.06	-0.59	---	---	---	---
0.2	-3.57	-3.2	-3.04	-5.63	-3.0	-3.72	-3.01	-3.09	-6.7	-4.27
0.4	-6.23	-5.94	-5.83	-8.35	-5.6	-6.43	-6.13	-6.43	-8.82	7.13
0.6	-8.66	-8.22	-8.02	-10.27	-7.9	-8.60	-9.25	-9.97	-11.5	-10.24

The above results refer to natural profiles. In order to relate the validation to the linear profiles used here, further tests were undertaken. In these tests, the models were run using an ‘equivalent’ linear profile having a slope of 1/15 to compare with the results of the physical experiments. The ‘equivalent’ profile was estimated using the best linear fitting to the natural profiles used in the physical experiment (between water depths of 3.5 and elevations of 1.5 m).

The results of this exercise are shown in Table 3. Generally, the comparison showed that, at least for the conditions tested, the results of the models set with the equivalent linear profile were reasonably close to those of the physical experiments, with the comparison worsening with increasing water levels. When the numerical models are used as a (short-term) ensemble, beach retreat forced by waves only is ~~substantially~~ overestimated; under SLR, however, ensemble projections improve (overestimations of up to 19 %, Table 3). Regarding the analytical models, the long-term ensemble gave improved projections, with differences between models and experiments ranging from about 3 % to 11 %.

Table 3: Comparison of results by models and physical experiments. Models used an ‘equivalent’ linear profile slope of 1/15 to represent the natural profile of the physical experiment (see text). Negative values represent beach retreat/erosion. Key: SLR, sea level rise; Exper, physical experiment; Leo, Leont’yeu model; SB, SBEACH model; Xb, Xbeach model; Bous, Boussinesq model; Br, Bruun model; Edel, Edelman model; Dean, Dean model; ST Ens, Short-term ensemble; and LT Ens, Long-term ensemble.

SLR (m)	Shoreline retreat/erosion or advance/accretion (m)									
	Exper	Leo	SB	Xb	Bous	ST Ens	Br	Edel	Dean	LT Ens
0	-0.35	-0.34	-0.02	-3.23	0.0	-0.9	---	---	---	---
0.2	-3.57	-3.58	-3.2	-7.00	-3.03	-4.20	-3	-3.07	-4.05	-3.37
0.4	-6.23	-6.67	-6.25	-10.47	-5.9	-7.32	-6	-6.3	-7.05	-6.45
0.6	-8.66	-9.73	-9.25	-13.35	-8.8	-10.28	-9	-9.69	-10.05	-9.58

4.2.2 Beach erosion and inundation/flooding projections

Modeling results show that sea level rise will cause shoreline retreats as well as significant morphological changes (Fig. 4). Numerical models showed differential profile changes in both breaker and surf zones, supporting their use in an ensemble mode.




Figure 4: Examples of morphodynamic changes in the profile of the upper part of the beach (see inset) and mean retreats of the 2 ensembles for SLR of 0.4 m. (a) Offshore (at 20 m water depth) wave height H of 1 m and period T of 5 s, and median (d50) sediment grain size of 0.8 mm; linear beach profile with 1/10 slope; (b) Offshore (at 20 m water depth) wave height H of 2 m and period T of 6 s, and median (d50) sediment grain size of 0.33 mm; linear profile with 1/30 slope. Origin of X axis at 20 m water depth.

Models displayed differential behaviour for almost all tested conditions, showing as expected significant ranges of results (Fig. 5) due to the varying initial conditions and forcing used i.e. different bed slopes, sediment sizes, wave conditions and SLRs. The means (best fits) of the lowest and highest projections of all models were calculated. It was found that the 'low' mean of the beach retreat projections by the short-term ensemble (i.e. the best fit of the lowest projections from the 4 numerical models) is given by $S = 0.1 \alpha^2 + 9.7 \alpha + 0.4$ and the 'high' mean by $S = 0.7 \alpha^2 + 28.5 \alpha + 4.8$ (Fig. 5a), where S is the beach retreat and α is the SLR. Also, the low projection mean of the long-term ensemble is given by $S = 0.1 \alpha^2 + 10 \alpha + 0.3$ and the high projection mean by $S = 1.6 \alpha^2 + 29.8 \alpha + 2.3$ (Fig. 5b). Ranges in beach temporary inundation/flooding due to wave run-up combined with (a) episodic (short term) SLRs (Fig. 5c) were estimated as $S(i) = 0.1 \alpha^2 + 9.7 \alpha + 4$ (minimum) and $S(i) = -0.7 \alpha^2 + 31.2 \alpha + 27.2$ (maximum) and with long-term SLRs (Fig. 5d) as $S(i) = 0.4 \alpha^2 + 10.1 \alpha + 3.7$ (minimum) and $S(i) = 0.4 \alpha^2 + 30 \alpha + 28.1$ (maximum).

Ranges of decreases in 'dry' beach widths were projected through the comparison between the ranges of beach retreat/erosion (S) and the recorded maximum widths of the 3234 pocket beaches. Ranges in beach temporary inundation/flooding were estimated by the comparison between the ranges of combined beach retreat and wave run-up excursions (S(i)) and the beach maximum widths. In Table 4, estimations are presented for 9 SLR scenarios: (i) 0.15, 0.5 and 0.7 m MSLRs according to recent projections for the area (Hinkel et al. 2014) using the long-term ensemble; (ii) short-term SLRs due to storm surges and waves of 0.2, 0.4 and 0.6 m using the short-term ensemble; and (iii) combined MSLRs and short-term SLRs of 0.55, 1.1 and 1.3 m, using consecutively the long- and short-term ensembles.




Figure 5: Projections of beach retreat/erosion (a and b) and temporary inundation/flooding (c and d) due to short-term (a and c) and long-term (b and d) SLR. Projections are for different beach slopes, sediment sizes and wave conditions. The means of the highest and lowest projections of all models in the ensembles are shown as yellow stippled lines.

According to the projections of the long-term ensemble, MSLRs of 0.15, 0.5 and 0.7 m will result to beach retreats/erosion by about 1.8 - 6.8, 5.3 - 17.6 and 7.3 - 24 m, respectively, whereas short-term ensemble estimates show that ~~storm-induced episodic increased~~ levels of +0.2, +0.4, +0.6 m could result in ranges of beach retreat/erosion of about 2.3 - 10.6, 4.3 - 16.3, 5.3 - 19.3 and 6.2 - 22.2 m, respectively (Table 4). For the considered SLR scenarios, there will be significant impacts on the Aegean Archipelago beaches as shown by the percentages of beaches that are projected to be eroded/shifted landward to a distance equal to 50 % and 100 % of their maximum width (Table 4).

Even under a MSLR of 0.15 m (RCPs 4.5 and 8.5 in 2040), there could be substantial impacts on the basis of the mean minimum and maximum projections of the long-term ensemble (Table 4). Temporary inundation (S(i)) could overwhelm between 4.9 and 81 % of the beaches, flooding (occasionally) 4.6 - 80 % of the beaches fronting currently existing coastal infrastructure/assets. Under a MSLR of 0.5 m (RCP 4.5, 2100), projected impacts will be severe. 5 to 54 % of all Aegean beaches will be completely eroded in the absence of appropriate coastal defences, whereas between about 18 and 90 % of all beaches are projected to be occasionally overwhelmed by flooding. For 2100, under the high emission scenario (RCP8.5, MSLR of 0.7 m), impacts could be catastrophic (Table 4 and Fig. 6): 12.4 - 68 % and 28 - 93 % of Aegean island beaches

will be completely eroded and occasionally overwhelmed by flooding, respectively. Associated infrastructure/assets are also projected to be greatly impacted, with 12 - 66 % and 26 – 92 % of all beaches fronting currently existing assets projected to be lost to beach erosion and occasionally overwhelmed by flooding, respectively.

Table 4: Minimum and maximum mean estimates of beach retreat (S) and inundation/flooding (S(i)) by the long-term, short-term and combined ensembles and the empirical model of Stockdon et al. (2006) Ranges of cross-shore retreat/erosion (R) and temporary inundation/flooding (F) for the beaches of the Aegean Archipelago are projected by comparing the highest and lowest mean S and S(i) with the maximum width of the 3234 Aegean Archipelago beaches under different SLRs. Numbers (N) and percentages of beaches where backshore infrastructure/assets are projected to be affected by beach retreat/erosion and flooding are also shown.

SLR (m)			S (m)		R		F		R		F		R		F	
					Equal to 50 % of max. width (%)		Equal to max. width (%)		Beaches with assets affected		Beaches with assets affected		N	%	N	%
Long-term	0.15	Min	1.8	5.2	0.6	27.4	0.0	4.9	0	0.0	119	4.6				
		Max	6.8	32.6	39.1	96.4	8.2	81.1	203	7.8	2086	79.8				
	0.5	Min	5.3	8.8	27.5	54.2	5.0	17.6	120	4.6	437	16.7				
		Max	17.6	43.2	84.4	98.3	54.2	89.7	1368	52.4	2320	88.8				
	0.7	Min	7.3	10.9	43.2	64.3	12.4	27.6	306	11.7	688	26.3				
		Max	24	49.3	91.5	99.1	68.0	92.6	1732	66.3	2400	91.8				
Short-term	0.2	Min	2.3	5.9	2.4	30.9	0.1	5.0	2	0.1	122	4.7				
		Max	10.6	33.4	64.3	96.6	27.5	82.1	684	26.2	2113	80.9				
	0.4	Min	4.3	7.9	17.4	47.9	2.3	12.7	49	1.9	315	12.1				
		Max	16.3	39.6	81.1	97.8	51.3	87.2	1293	49.5	2249	86.1				
	0.6	Min	6.2	9.8	34.7	59.1	7.9	21.5	195	7.5	537	20.6				
		Max	22.2	45.7	90.2	98.8	66.5	90.7	1695	64.9	2343	89.7				
MSLR + Short-term	0.55	Min	6	9.6	34.7	59.0	7.9	21.4	195	7.5	535	20.5				
		Max	23.2	46.4	91.0	98.8	68.0	91.0	1730	66.2	2354	90.1				
	1.1	Min	11.6	15.2	68.0	78.8	30.8	47.8	763	29.2	1204	46.1				
		Max	40.3	63.6	97.9	99.7	88.0	96.2	2270	86.9	2506	95.9				
	1.3	Min	13.5	17.2	74.6	82.8	39.1	54.1	977	37.4	1364	52.2				
		Max	46.2	69.7	98.8	99.7	91.0	96.8	2354	90.1	2526	96.7				

According to the mean low projections of the short-term ensemble, ~~storm coastal sea levels increased sea levels due to storm surges/waves~~ of ± 0.2 and ± 0.4 m will result in moderate (temporary) beach retreats (0.1 - 2.3 % of all beaches will retreat more than their maximum width) and temporary flooding (5 – 12.7 % of the beaches will be occasionally completely flooded). On the basis of the mean high projections (forced by the high wave conditions expected in storms, see also Tsoukala et al, (2016)), beach retreats and flooding will be substantial with severe reductions in ‘dry’ beach widths and potential damages of assets located at the back of the beach. About 27 and 51 % of all Aegean beaches will retreat by more than their maximum width and 82 and 87 % will be completely overwhelmed by temporary flooding under short-term SLRs of 0.2 and 0.4 m, respectively (Table 4).

In the case of a 0.4 m short-term SLR, up to 50 % and 86 % of all beaches fronting assets will be affected by beach retreat/erosion and temporary flooding, respectively (Fig. 6), whereas impacts will be more severe under higher coastal sea

levels (+0.6 m) (Table 4). The picture does not change much, even when projections are adjusted for the maximum overestimation observed in the results of the numerical models (by 19 %) when linear profiles were used (Section 4.2.1); for example, for a 0.4 m level rise, up to 37 and 84 % of all beaches fronting assets will be fully eroded and flooded, respectively.

5 | The worst impacts are projected from the combined mean and short-term SLRs. In 2040, in the case that storm-~~induced~~ sea levels of ± 0.4 m are combined with a projected MSLR of 0.15 m (combined SLR of 0.55 m), 8 - 68 % of beaches are projected to be (at least temporarily) eroded and 21 - 91 % of beaches flooded. In 2100, superimposition of storm levels on the projected MSLRs will have devastating effects. A combined sea level rise of 1.1 m (e.g. a storm-~~induced~~ **extreme level** ~~of +surge/wave set-up of~~ 0.6 m superimposed on a MSLR of 0.5 m (RCP4.5)) will have very severe impacts, indeed (Figs 6

10 | and 7): 31 – 88 % of all beaches will be completely (at least temporarily) eroded (29 - 87 % of all beaches fronting assets) under the low and high mean projections of the ensemble, respectively, with 48 - 96 % of all beaches occasionally overwhelmed by flooding (Table 4).

A combined SLR of 1.3 m (RCP8.5, a MSLR of 0.7 m combined with a storm-induced coastal sea level of +0.6 m) represents a ‘doom’ scenario for the beaches of the Aegean Archipelago. Based on the low projections of the combined

15 | ensembles, about 75 % of beaches will be shifted landward (and/or drowned) to a distance equal to 50 % of their maximum ‘dry’ width, whereas about 39 % of all beaches will be (at least temporarily) completely eroded; 37 % of beaches fronting existing assets will be fully eroded and 54 % fully flooded.

Σχόλιο [I11]: This part is slightly changed to account for the Comment 1 of Referee #1.





Figure 6: Minimum and maximum retreat and flooding of Aegean Archipelago beaches for different SLR scenarios on the basis of the low and high means respectively of the ensemble projections. a) and b) Minimum and maximum retreat and flooding under a MSLR of 0.7 m. c) and d) Minimum and maximum retreat and flooding under a storm-induced level of +0.4 m. e) and f) Minimum and maximum retreat and flooding under a combined lon- and short-term SLR of 1.1 m (see text). Initial and final (after SLR) maximum beach widths are shown. Final widths < 0 show beaches lost or shifted landward or flooded by their entire maximum width. Black bars show currently existing infrastructure/assets fronted by beaches.

10 According to the high ensemble estimates, it is projected that almost 91 % of all beaches will be completely eroded, at least for the time of the storm event (90 % of all beaches fronting assets); about 97 % of all beaches will also be flooded temporarily (Table 4).



5 Figure 7: Projections of minimum beach retreat under a combined SLR of 1.1 m (see text), showing Aegean beaches projected to retreat by distances equal to different percentages of their initial maximum widths. In the inset, projections for the beaches of Naxos (Cyclades) are shown in larger scale for detail.

Σχόλιο [I12]: Has been slightly improved

When projections are adjusted for overestimations observed in the validation exercise due to the use of linear profiles (Section 4.2.1), projected impacts are still devastating: 35 -89 % of all beaches will be (at least temporarily) completely eroded and 48 – 86 % flooded, whereas according to the high estimates, 88 % and 96 % of all beaches fronting assets will be completely eroded and flooded, respectively.

It should be noted that projected impacts do not account for vertical land motions (Poulos et al., 2009). Moreover, since other significant erosion factors such as the diminishing coastal sediment supply are not considered, the above projections may actually underestimate future beach retreat/erosion and flooding at the Aegean Archipelago beaches.

5 Discussion and Conclusions

Analysis of the first systematic record of the geo-spatial characteristics of the beaches of the Aegean Archipelago has provided interesting findings. Aegean island beaches are narrow pocket beaches (59 % ~~were recorded having have~~ recorded widths < 20 m), ~~characterised by have~~ characterised by limited terrestrial sediment supply and fronting a moderate (as a total) load of backshore development; about 81 % of the Aegean beaches currently front coastal infrastructure/assets with 33 % fronting moderate to high coastal development. At the same time, only 15 % of beaches are associated with some form of existing coastal protection. Moreover, at least 23 % of the Aegean beaches exhibit outcropping beachrocks at their beach faces. In addition to aesthetic considerations and the safety risks that slippery beachrock surfaces might pose to beach users (Vousdoukas et al., 2009c), beach face beachrocks can also ~~degrade aesthetics,~~ induce ecological changes and promote beach sediment removal (e.g. Vousdoukas et al. 2009b; 2012) ~~erosion,~~ decreasing the carrying capacity of the Archipelago beaches and constraining their potential as environments of leisure (Kontogianni et al., 2014).

Development and maintenance of the Aegean beaches may be (at least partially) controlled by hydrodynamics, as more and wider beaches occur along the southern and western coasts of the islands that are relatively protected from the prevailing N-NE waves. Maximum beach widths were found to correlate with sediment type and terrestrial sediment supply; sandy beaches are wider as are those associated with river outlets. With regard to the carrying capacity of the Archipelago beaches, these appear to have potential (as a total) for further development as environments of leisure; nevertheless, better spatial distribution and a lengthening of the period of beach tourism should be considered.

~~The geo-spatial characteristics of the~~ Aegean island beaches ~~appear particularly indicate that these may be considerably exposed vulnerable~~ appear particularly indicate that these may be considerably exposed vulnerable to SLR. Our modeling results ~~indeed~~ indeed project severe ~~erosion and flooding impacts~~ erosion and flooding impacts under SLR from as early as in 2040, particularly under the combined effects of the projected MSLR and storm-induced sea levels (Table 4). By 2100, ~~beach erosion impacts~~ beach erosion impacts from combined MSLR and storm events ~~is projected to be very substantial, with potentially devastating impacts on both coastal infrastructure and tourism could be devastating.~~ is projected to be very substantial, with potentially devastating impacts on both coastal infrastructure and tourism could be devastating. For example, under a MSLR of 0.5 m (RCP4.5), a ~~short-term coastal sea level rise due to storm surges and waves of storm event inducing an additional~~ short-term coastal sea level rise due to storm surges and waves of storm event inducing an additional 0.6 m ~~rise~~ rise

Σχόλιο [I13]: Some refs were added in order to address the Comment 5 of Referee #2.

is projected to result in complete erosion of between 31 (minimum) and 88 % (maximum) of all beaches (29 - 87 % of beaches currently fronting coastal infrastructure and assets), at least temporarily. ~~In addition, between 46 and 96 % of the coastal infrastructure and assets behind the Aegean island beaches are projected to be temporarily flooded during such an event. Regarding the Aegean island tourism, this will be negatively affected by the projected large decreases in 'dry' beach width, a critical control of the beach recreational use, value and resilience (e.g. Yang et al., 2012). Under increasing beach erosion/retreat and flooding, the long-term recreational value of the Aegean Archipelago beaches as well as the value of associated assets may fall considerably (e.g. Gopalakrishnan et al., 2011).~~ As a recent study projects substantial changes in the return periods of ~~ESL~~~~extreme sea levels~~ for E. Mediterranean ~~at for~~ the end of the 21st century (~~the current 1000 year event is projected to occur once every 5 years,~~ Voudoukas et al. (~~submitted under review~~)), ~~the~~ projected impacts on both beaches and coastal infrastructure/assets are worrying.

~~In the present study, SLR impacts have been estimated on the basis of decreases in 'dry' beach width, a critical parameter for the estimation of beach resilience and recreational use and value (e.g. Yang et al., 2012). Under increasing beach erosion/retreat and flooding, the long-term recreational value of Aegean Archipelago beaches as well as the value of associated assets may fall considerably (e.g. Gopalakrishnan et al., 2011).~~

Against this background, it appears that plans to respond effectively to the projected beach erosion risk should be urgently drawn up with different adaptation options ~~analysed~~~~considered~~. Options based on the ecosystem approach should be the first to consider in order to protect both beaches and backshore ecosystems and infrastructure/assets (e.g. Peduzzi et al., 2013), although 'hard' works might, in some cases, be deemed necessary. However, the significance of beaches as critical economic resources and the low effectiveness of 'hard' coastal works (e.g. breakwaters) to protect beaches from MSLR indicate that beach nourishment schemes will be also required, at least for the most economically important beaches. As marine aggregates constitute the most suitable, but often scarce (Peduzzi, 2014) material for beach nourishment, particular care should be taken to ensure sustainability of marine aggregate deposits (Velegrakis et al., 2010). The significance of such deposits should be certainly considered in future marine spatial plans as a matter of priority (see also EU Directive 2014/89/EU).

A primary tool to manage assets and economic activities at risk is the introduction of effective policies and regulation. An underpinning principle of coastal management under SLR should be the introduction of effective precautionary controls on future coastal development, e.g. through regulation that allocates buffer zones ('set-back' zones) behind retreating coastlines. It is submitted that, on the basis of our results, there is an urgent need to adopt/implement relevant regulation for the Aegean Archipelago.

A way forward could be the ratification by Greece of the 2009 ICZM Protocol to the Barcelona Convention, which prescribes set-back zones (Art. 8.2(a)) and has been already ratified by the EU (Council Decision 2010/631/EU). However, there are also challenges as e.g. those related to the set-back zone demarcation and risk allocation, (e.g. Sano et al., 2010; Gibbs et al., 2013) that can be only alleviated by publicly transparent criteria and decisions (e.g. Abbott, 2013). Assessments of beach retreat/erosion and flooding are required at various scales that are science-based and, at the same time, accessible by

coastal planners, managers, stakeholders and the wider public for the planning and smooth implementation of ‘set-back’ zones as well as the prioritization in the allocation of resources for adaptation.

The approach adopted in the present study to assess impacts of the SLR presents certain advantages. Existing methodologies/tools for rapid assessment of coastal/beach erosion ~~due to and corresponding vulnerabilities to~~ MSLR and extreme events at large scales (e.g. Hinkel et al., 2010; Ramieri et al., 2011; Khouakhi et al., 2013) have limitations stemming from (amongst others): (a) their requirements for coastal Digital Elevation Models (DEMs) of high resolution/accuracy; and (b) the limited generally consideration for major controls (e.g. hydrodynamics). At the same time, advanced modeling approaches (e.g. Vousdoukas et al., 2016b) in addition to detailed environmental information commonly require experienced operators and high computation costs that ~~may makemakes~~ them impractical to coastal planners/managers (e.g. McLeod et al., 2010).

The present approach, which compares ranges of SLR induced beach retreat and flooding under different initial conditions and hydrodynamic forcing with beach maximum widths, is not limited by the resolution/accuracy of available coastal DEMs or the availability of detailed environmental information (e.g. Jiménez et al., 2012), and can be easily incorporated in other beach vulnerability tools (e.g. Alexandrakis et al., 2015) and used in areas with limited human resources. Nevertheless, there are also constraints. Projections are based on the assumption that beaches comprise inexhaustible sediment reservoirs, with no lateral sediment losses; cross-shore modeling obviously cannot resolve such issues. In addition, the approach is not designed to account for other erosion-controlling factors, such as: geological controls, coastal sedimentary budgets, and extreme event duration and sequencing (e.g. Gallop et al., 2012; Corbella and Stretch, 2012); the presence of artificial beach protection schemes and/or protecting nearshore ecosystems (e.g. Peduzzi et al., 2013); and the effects of coastal use (e.g. Bi et al., 2013). However, the aim of the exercise has not been to replace detailed modeling studies for individual beaches, but to provide ranges of beach erosion and flooding at a large (Archipelago) scale.

Acknowledgements

This research has been co-financed by the European Union (European Social Fund – ESF) and Greek national funds through the Operational Program “Education and Lifelong Learning” of the National Strategic Reference Framework (NSRF) – Research Funding Program: THALES (Project ISLA). ~~The senior authors (INM, AFV and AEC) also acknowledge the support from ERA-Beach Project during the writing-up of the [research]; the programme is co-funded in 85% by the EEA GRANTS, 2009-2014, and 15% by the Public Investments Programme (PIP) of the Hellenic Republic.~~ M.I. Vousdoukas acknowledges funding from the European Union Seventh Framework Programme FP7/2007–2013 under Grant Agreement No. 603864 (HELIX: ‘High-End cLimate Impacts and eXTremes’; www.helixclimate.eu), as well as by the JRC institutional project Coastalrisk.

Σχόλιο [a14]: Updated acknowledgements

References

- Abbott, T., 2013. Shifting shorelines and political winds The complexities of implementing the simple idea of shoreline setbacks for ocean front developments in Maui, Hawaii. *Ocean & Coastal Management* 73, 13–21.
- Alexandrakis, G., Manasakis, C., Kampanis, N.A., 2015. Valuating the effects of beach erosion to tourism revenue. A management perspective. *Ocean & Coastal Management* 111, 1-11.
- Allenbach, K., Garonna, I., Herold, C., Monioudi, I., Giuliani, G., Lehmann, A.F. Velegrakis, 2015. Black Sea beach vulnerability to sea level rise. *Environmental Science and Policy* 46, 95–109.
- Androulidakis, Y.S., Kourafalou, V.H., Krestenitis, Y.N., Zervakis, V., 2012. Variability of deep water mass characteristics in the North Aegean Sea: The role of lateral inputs and atmospheric conditions. *Deep-Sea Research I* 67, 55–72.
- Androulidakis, Y.S., Kombiadou, K.D., Makris, C.V., Baltikas, V.N., Krestenitis, Y.N., 2015. Storm surges in the Mediterranean Sea: Variability and trends under future climatic conditions. *Dynamics of Atmospheres and Oceans* 71, 56–82.
- Bi, X., Lu, Q., Pan, X., 2013. Coastal use accelerated the regional sea-level rise. *Ocean & Coastal Management* 82, 1-6.
- Bosom, E., Jiménez, J.A., 2010. Storm-induced coastal hazard assessment at regional scale: application to Catalonia (NW Mediterranean). *Advances in Geosciences*. 26, 83–87.
- Bruun, P., 1988. The Bruun Rule of erosion by sea level rise: A discussion on large-scale two- and three-dimensional usages. *Journal of Coastal Research* 4(4), 622-648.
- Calafat, F.M., Chambers, D.P., Tsimplis, M.N., 2014. On the ability of global sea level reconstructions to determine trends and variability. *Journal of Geophysical Research* 119 7840-7853.
- Cisneros, H.M.A., Revollo, Sarmiento, N.V., Delrieux, C.A., Piccolo, M.C., Perillo, G.M.E., 2016. Beach carrying capacity assessment through image processing tools for coastal management. *Ocean & Coastal Management* 130, 138-147.
- Conte, D., Lionello, P., 2013. Characteristics of large positive and negative surges in the mediterranean sea and their attenuation in future climate scenarios. *Global Planetary Change* 111, 159–173.
- Corbella, S., Stretch, D.D., 2012. Predicting coastal erosion trends using non-stationary statistics and process-based models. *Coastal Engineering* 70, 40–49.
- Dean, R.G., 1991. Equilibrium beach profiles: characteristics and applications. *Journal of Coastal Research* 7(1), 53–84.
- Dean, R.G., 2002. Beach nourishment: Theory and practice. In: *Advanced Series on Ocean Engineering*, Singapore, World Scientific Publishing Company.
- Defeo, O., McLachlan, A., 2013. Global patterns in sandy beach macrofauna: Species richness, abundance, biomass and body size. *Geomorphology* 199, 106–114.
- Ding, Y., Kuiry, S.N., Elgohry, M., Jia, Y., Altinakar, M.S., Yeh, K.-C., 2013. [Impact assessment of sea-level rise and hazardous storms on coasts and estuaries using integrated processes model](#). *Ocean Engineering* 71, 74–95.

- ECLAC, 2011. An assessment of the economic impact of climate change on the transportation sector in Barbados. UN Economic Commission for Latin America and the Caribbean ECLAC Technical Report LC/CAR/L309. 44 pp
- Edelman, T., 1972. Dune erosion during storm conditions. In: Proceedings of the 13th International Conference on Coastal Engineering ASCE, pp. 1305-1312.
- 5 Eurosion, 2004. Living with coastal erosion in Europe: Sediment and Space for Sustainability. Part II. DG Environment, EC (<http://www.eurosion.org/reports-online/part2.pdf>).
- Gallop, S.L., Bosserelle, C., Eliot, I., Pattiaratchi, C.B., 2012. The influence of limestone reefs on storm erosion and recovery of a perched beach. *Continental Shelf Research* 47, 16–27.
- Ghermandi, A., Nunes, P.A.L.D., 2013. A global map of coastal recreation values: Results from a spatially explicit meta-analysis. *Ecological Economics* 86, 1–15.
- 10 Gibbs, M.T., Thébaud, O., Lorenz, D., 2013. A risk model to describe the behaviours of actors in the houses falling into the sea problem. *Ocean & Coastal Management* 80, 73-79.
- Gopalakrishnan, S., Smith, M.D., Slott, J.M., Brad Murray, A., 2011. The value of disappearing beaches: A hedonic pricing model with endogenous beach width. *Journal of Environmental Economics and Management* 61, 297-310.
- 15 Hall, C.M., Scott, D., and Gössling, S., 2013. The Primacy of Climate Change for Sustainable International Tourism. *Sustainable Development* 21, 112–121.
- Hinkel, J., Nicholls, R.J., Vafeidis, A.T., Tol, R.S.J., Avagianou, T., 2010. Assessing risk of and adaptation to sea-level rise in the European Union: An application of DIVA. *Mitigation and Adaptation Strategies for Global Change* 15(7), 703-719.
- Hinkel, J., Lincke, D., Vafeidis, A.T., Perrette, M., Nicholls, R.G., Tol, R.S., Marzeion, B., Fettweis, X., Ionescu, C., 20 Levermann, A., 2014. Coastal flood damages and adaptation costs under 21st century sea-level rise. *Proceedings of the National Academy of Sciences*, 111, 3292-3297.
- Hoeke, R.K., McInnes, K.L., Kruger, J.C., McNaught, R.J., Hunter, J.R. and Smithers, S.G., 2013. Widespread inundation of Pacific islands triggered by distant-source wind-waves. *Global and Planetary Change* 108, 128-138.
- IOBE, 2012. The impact of tourism on Greek Economy (in Greek). Foundation for Economic & Industrial Research, pp. 43, 25 http://iobe.gr/docs/research/RES_05_E_01092012REP_GR.pdf
- IPCC, 2013. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)], Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- 30 IPCC SREX, 2012. Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation. A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change IPCC [C.B. Field et al. (eds.)], Cambridge University Press, Cambridge, UK, and New York, NY, USA, 582 pp.
- Jiménez, J.A., Bosom, E., Valdemoro, H.I., Guillén, J., 2012. Storm-induced damages along the Catalan coast (NW Mediterranean) during the period 1958–2008. *Geomorphology* 143-144, 24-33.

- Karambas, T.V., Koutitas, C., 2002. Surf and swash zone morphology evolution induced by nonlinear waves. *Journal of Waterway, Port, Coastal and Ocean Engineering*, ASCE 128 (3).
- Karambas, Th., 2006. Prediction of sediment transport in the swash zone by using a nonlinear wave model. *Continental Shelf Research* 26, 599–609.
- 5 Khouakhi, A., Snoussi, M., Niazi, S., Raji, O., 2013. Vulnerability assessment of Al Hoceima bay (Moroccan Mediterranean coast): a coastal management tool to reduce potential impacts of sea-level rise and storm surges. *Journal of Coastal Research* SI 65, 968-973.
- Kontogianni, A.D., Damigos, D., Tourkolias C., Vousdoukas M.I., Velegrakis A.F., Zanou B., Skourtos, M., 2014. The social cost of coastal erosion: eliciting beach users' willingness to pay for protecting European beaches from beachrock processes. *Ocean & Coastal Management* 98, 167-175.
- 10 Krestenitis, Y.N., Androulidakis, Y.S., Kontos, Y.N., Georgakopoulos, G., 2011. Coastal inundation in the north-eastern Mediterranean coastal zone due to storm surge events. *Journal of Coastal Conservation* 15 (3), 353-368.
- Larson, M., Kraus, N.C., 1989. SBEACH: numerical model for simulating storm-induced beach change; report 1: empirical foundation and model development, Technical Report - US Army Coastal Engineering Research Center 89-9.
- 15 Leont'ev, I.O., 1996. Numerical modelling of beach erosion during storm event. *Coastal Engineering* 29(1-2), 187-200.
- Mamoutos, G., Tragou, E., Kakagiannis, G., 2014. Mean sea level changes in the Greek Seas. In 6th Panhellenic Conference in Management and Improvement of Coastal Zones, November 2014, Athens, Greece, pp. 335-344.
- Marcos, M., Jordà, G., Gomis, D., Pérez, B., 2011. Changes in storm surges in Southern Europe from a regional model under climate change scenarios. *Global Planetary Change* 77, 116–128.
- 20 McArthur, S., 2015. Why Carrying Capacity Should be a Last Resort, 2015. In: T.Vir Singh (Ed.), *Challenges in Tourism Research*, Channel View Publications, pp. 288-302.
- McLeod, E., Poulter, B., Hinkel, J., Reyes, E., and Salm, R., 2010. Sea-level rise impact models and environmental conservation: A review of models and their applications. *Ocean & Coastal Management* 53, 507-517.
- Monioudi, I.N., Karditsa, A., Chatzipavlis, A., Alexandrakis, G., Andreadis, O.P., Velegrakis, A.F., Poulos, S.E., Ghionis, G., Petrakis, S., Sifnioti, D., Hasiotis, Th., Lipakis, M., Kampanis, N., Karambas, Th., Marinos, E. (2016) Assessment of the vulnerability of the eastern Cretan beaches (Greece) to sea level rise. *Regional Environmental Change* 16 (7), 1951–1962. DOI 10.1007/s10113-014-0730-9.
- Neumann, B., Vafeidis, A.T., Zimmerman, J., and Nicholls, R.J., 2015. Future Coastal Population Growth and Exposure to Sea-Level Rise and Coastal Flooding - A Global Assessment. *Plos One* <http://dx.doi.org/10.1371/journal.pone.0118571>
- 30 Nicholls, R.J., Cazenave, A., 2010. Sea-level rise and its impact on coastal zones. *Science* 328, 1517-1520.
- Parker, A., Saad Saleem, M., Lawson, M., 2013. Sea-level trend analysis for coastal management. *Ocean & Coastal Management* 73, 63-81.
- Paula, D.P., Dias, J.M.A., Ferreira, O. and Morais, J.O., 2013. High-rise development of the sea-front at Fortaleza (Brazil): Perspectives on its valuation and consequences. *Ocean & Coastal Management* 77, 14-23.

- Peduzzi, P., Velegrakis, A.F., Estrella, M., Chatenoux, B., 2013. Integrating the role of ecosystems in disaster risk and vulnerability assessments: Lessons from the Risk and Vulnerability Assessment Methodology Development Project (RiVAMP) in Negril Jamaica, in: FG Renaud et al. (eds), *The role of Ecosystems in Disaster Risk Reduction*. United Nations University Press (ISBN 978-9280812213), 109-139.
- 5 Peduzzi, P., 2014. Sand, rarer than one thinks. Article reproduced from United Nations Environment Programme (UNEP) Global Environmental Alert Service (GEAS). *Environmental Development* 11 (2014), 208–218
- Phillips, M.R., Jones A.L., 2006. Erosion and tourism infrastructure in the coastal zone: Problems, consequences and management. *Tourism Management* 27, 517-524.
- Poulos, S.E., 2009. Origin and distribution of the terrigenous component of the unconsolidated surface sediment of the Aegean floor: A synthesis. *Continental Shelf Research* 29, 2045–2060.
- 10 Poulos, S.E., Ghionis G., Maroukian, H., 2009. Sea-level rise trends in the Attico–Cycladic region (Aegean Sea) during the last 5000 years. *Geomorphology* 107, 10-17.
- Prinos, P., 2014. Climate change effects on the Greek seas and coastal areas — the research project THALIS-CCSEAWAVS. In: 6th Panhellenic Conference on Coastal Zones Management and Improvement, 24—27 November 2014, Athens, 315-324.
- 15 Rajan, B., Varghese, V.M., Pradeepkumar, A.P., 2013. Beach Carrying Capacity Analysis for Sustainable Tourism Development in the South West Coast of India. *Environmental Research, Engineering and Management* 1 (63), 67-73.
- Ramieri, E., Hartley, A., Barbanti, A., Duarte Santos, F., Laihonon, P., Marinova, N., Santini, M., 2011. Methods for assessing coastal vulnerability to climate change. ETC CCA Background Paper, Copenhagen (DK) 8-9 June 2011. European Topic Centre on Climate Change Impacts, Vulnerability and Adaptation European Environment Agency, 76pp. <http://www.oannes.org.pe/>
- 20 Ranasinghe, R., Duong, T.M., Uhlenbrook, S., Roelvink, D., Stive, M., 2013. Climate-change impact assessment for inlet-interrupted coastlines. *Nature Climate Change* 3, 83-87.
- Roelvink, D., Reniers, A., Van Dongeren, A., Van Thiel de Vries, J., McCall, R., Lescinski, J., 2009. Modeling storm impacts on beaches, dunes and barrier islands. *Coastal Engineering* 56, 1133–1152.
- 25 Roelvink, D., Reniers, A., van Dongeren, A., van Thiel de Vries, J., Lescinski, J., McCall, R., 2010. Xbeach Model Description and Manual, Unesco-IHE Institute for Water Education. Delft and Delft University of Technology, 106.
- Sano, M., Marchard, M., Medina, R., 2010. Coastal set backs for the Mediterranean: a challenge for ICZM. *Journal of Coastal Conservation* 14, 33-39.
- 30 SETE, 2016. Association of the Greek Touristic Businesses. (<http://sete.gr/el/statistika-vivliothiki/statistika/?c=&cat=&key=%CE%9A%CE%9B%CE%99%CE%9D%CE%95%CF%82&tags=%CE%BE%CE%B5%CE%BD%CE%BF%CE%B4%CE%BF%CF%87%CE%B5%CE%AF%CE%B1>)
- Short, A.D., Jackson, D.W.T., 2013. Beach Morphodynamics. In: *Treatise on Geomorphology*, Vol. 10, pp 106-129.

- Skiridis, N., Sofianos, S.S., Gkanasos, A., Axaopoulos, P., Mantziadou, A., Vervatis, V., 2011. Long-term sea surface temperature variability in the Aegean Sea. *Advances in Oceanography and Limnology* 2:2, 125-139.
- Smith, A.B., Katz, R.W., 2012. U.S. Billion-dollar Weather and Climate Disasters: Data Sources, Trends, Accuracy and Biases. *Natural Hazards* 67(2), 387-410.
- 5 Soukissian, T., Hatzinaki, M., Korres, G., Papadopoulos, A., Kallos, G., Anadranistakis, E., 2007. Wind and wave atlas of the Hellenic seas. Hellenic Centre for Marine Research Publications, HCMR.
- Soukissian, T., Prospathopoulos, A., Hatzinaki, M., Kabouridou, M., 2008. Assessment of the wind and wave climate of the Hellenic Seas using 10-Year hindcast results. *The Open Ocean Engineering* 1, 1-12.
- Stockdon, H.F., Holman, R.A., Howd, P.A., Sallenger, J.A.H., 2006. Empirical parameterization of setup, swash, and runup. *Coastal Engineering* 53, 573-588.
- 10 Theocharis, A., Georgopoulos, D., Lascaratos, A., Nittis, K., 1993. Water masses and circulation in the central region of the Eastern Mediterranean: Eastern Ionian, South Aegean and north west Levantine, 1986-1987. *Deep Sea Research II*, 40(6), 1121-1142.
- Tsimplis, M., Marcos, M., Colin, J., Somot, S., Pascual, A., Shaw, A.G.P., 2009. Sea level variability in the Mediterranean Sea during the 1990 on the basis of two 2D and one 3D model. *Journal of Marine Systems* 78 (1), 109-123.
- 15 Tsimplis, M.N., Shaw, A., 2010. Seasonal sea level extremes in the Mediterranean Sea and at the Atlantic European coasts. *Natural Hazards and Earth System Sciences* 10, 1457-1475.
- Tsimplis, M.N., Calafat, F.M., Marcos, M., Jorda, G., Gomis, D., Fenoglio-Marc, L., Struglia, M.V., Josey, S.A., Chambers, D.P., 2013. The effect of the NAO on sea level and on mass changes in the Mediterranean Sea. *Journal of Geophysical Research: Oceans* 118, 944-952.
- 20 Tsoukala, V.K., Chondros, M., Kapelonis, Z.G., Martzikos, N., Lykou, A., Belibassakis, K., Makropoulos, Ch., 2016. An integrated wave modelling framework for extreme and rare events for climate change in coastal areas - the case of Rethymno, Crete. *Oceanologia* 58, 71-89.
- UNECE, 2013. Climate Change Impacts and Adaptation for International Transport Networks. Expert Group Report, ITC
- 25 UN Economic Commission for Europe ECE/TRANS/238, 223 pp.
http://www.unece.org/fileadmin/DAM/trans/main/wp5/publications/climate_change_2014.pdf
- Valdemoro, H.I., Jiménez, J.A., 2006. The influence of shoreline dynamics on the use and exploitation of Mediterranean tourist beaches. *Coastal Management* 34, 405-423.
- Velegrakis, A.F., Voudoukas, M.I., Andreadis, O., Pasakalidou, E., Adamakis, G., Meligonitis, R., 2008. Impacts of dams on their downstream beaches: A case study from Eresos coastal basin, Island of Lesbos, Greece. *Marine Georesources and Geotechnology* 26, 350-371.
- 30 Velegrakis, A.F., Ballay, A., Poulos, S., Radzevičius, R., Bellec, V., Manso, F., 2010. European marine aggregates resources: Origins, usage, and mining techniques. *Journal of Coastal Research* SI 51, 1-14.

- Velegrakis, A.F., Trygonis, V., Chatzipavlis, A.E., Karambas, Th., Vousdoukas, M.I., Ghionis, G., Monioudi I.N., Hasiotis, Th., Andreadis, O., Psarros, F., 2016. Shoreline variability of an urban beach fronted by a beachrock reef from video imagery. *Natural Hazards* DOI: 10.1007/s11069-016-2415-9.
- Vousdoukas, M.I., Velegrakis, A.F., Karambas, T.V., 2009a. Morphology and sedimentology of a microtidal beach with beachrocks: Vatera, Lesbos, NE Mediterranean. *Continental Shelf Research* 29(16), 1937-1947.
- Vousdoukas, M.I., Velegrakis, A.F., Dimou, K., Zervakis, V., Conley, D.C., 2009b. Wave run-up observations in microtidal, sediment-starved beaches of the Eastern Mediterranean. *Journal of Marine Systems* 78, 537-547.
- Vousdoukas, M.I., Velegrakis, A.F., Kontogianni, A. and Makrykosta, N.E. 2009c. Implications of the cementation of beach sediments (beachrock formation) for the recreational use of the beach. *Tourism Management* 30, 544-552.
- Vousdoukas, M.I., A.F. Velegrakis, M. Paul, C. Dimitriadis, E. Makrykosta, D. Koutsoubas, 2012. Field observations and modeling of wave attenuation over colonized beachrocks. *Continental Shelf Research* 48, [100-109]
- Vousdoukas, M.I., Kirupakaramoorthy, T., Oumeraci, H., de la Torre, M., Wübbold, F., Wagner, B., Schimmels, S., 2014. The role of combined laser scanning and video techniques in monitoring wave-by-wave swash zone processes. *Coastal Engineering* 83, 150-165.
- Vousdoukas, M.I., Voukouvalas, E., Annunziato, A., Giardino, A., Feyen, L., 2016a. Projections of extreme storm surge levels along Europe. *Climate Dynamics* DOI: 10.1007/s00382-016-3019-5.
- Vousdoukas, M.I., Voukouvalas, E., Mentaschi, L., Dottori F., Giardino A., Bouziotas D., Bianchi A., Salamon P., and Feyen L., 2016b. Developments in large-scale coastal flood hazard mapping. *Natural Hazards Earth System Science* 16, [1841]-1853.
- Vousdoukas, M.I., Mentaschi, L., Voukouvalas, E., Verlaan, M., Feyen, L., Extreme sea levels on the rise along Europe's coasts. Earth's Future Nature Climate Change (submitted under review).
- Xu, S., Huang, W., 2013. Effects of sea level rise on frequency analysis of 1% annual maximum water levels in the coast of Florida. *Ocean Engineering* 71, 96-102.
- Yang, B., Madden, M., Kim, J., Jordan, T.R., 2012. Geospatial analysis of barrier island beach availability to tourists. *Tourism Management* 33, 840-854.
- Zar, J.H., 2010. Biostatistical analysis. 5th ed., Prentice Hall PTR, 658 pp. ISBN-13: 978-0321656865.
- Zervakis, V., Georgopoulos, D., Drakopoulos, P., 2000. The role of the North Aegean in triggering the recent Eastern Mediterranean climatic changes. *Journal of Geophysical Research* 105, 26103-26116.

Σχόλιο [I15]: Refs added in order to address the Comment 5 of Referee #2.

Σχόλιο [a16]: Added that reference to address further the Comment 1 of Referee #1.