

1 Spatial and temporal analysis of fatal off-piste and
2 backcountry avalanche accidents in Austria with a
3 comparison of results in Switzerland, France, Italy and
4 the United States.

5 Christian Pfeifer, Dr.

6 Institute of Basic Sciences in Engineering Science
7 Unit for Engineering Mathematics, University of Innsbruck

8 Peter Höller, Dr.

9 Austrian Research Centre for Forests
10 Institute for Natural Hazards, Innsbruck

11 Achim Zeileis, Dr.

12 Department of Statistics, University of Innsbruck

13 **Address for correspondence:**

14 Christian Pfeifer

15 Institut für Grundlagen der Technischen Wissenschaften

16 Arbeitsbereich für Technische Mathematik, Universität Innsbruck

17 Technikerstraße 13, A-6020 Innsbruck

18 E-Mail: christian.pfeifer@uibk.ac.at

Abstract

19
20 In this article we analyzed spatial and temporal patterns of fa-
21 tal Austrian avalanche accidents caused by backcountry and off-piste
22 skiers and snowboarders within the winter periods 1967/68–2015/16.
23 The data were based on reports of the Austrian Board for Alpine
24 Safety and reports of the information services of the federal states.

25 Using the date and the location of the recorded avalanche accidents
26 we were able to carry out spatial and temporal analyses applying
27 generalized additive models and Markov random field models.

28 As the result of the trend analysis we noticed an increasing trend of
29 backcountry and off-piste avalanche fatalities within the winter periods
30 from 1967/68 to 2015/16 (although slightly decreasing in recent years),
31 which is in contradiction to the widespread opinion in Austria that the
32 number of fatalities is constant over time. Additionally, we compared
33 Austrian results with results of Switzerland, France, Italy and the
34 United States based on data from the International Commission of
35 Alpine Rescue (ICAR). As the result of the spatial analysis we noticed
36 two hotspots of avalanche fatalities (‘Arlberg-Silvretta’ and ‘Sölden’).

37 Because of the increasing trend and the rather ‘narrow’ regional
38 distribution of the fatalities consequences on prevention of avalanche
39 accidents were highly recommended.

40 **Keywords:** Snow, Avalanches, Accidents

41 1 Introduction

42 In the Alps, backcountry skiing has become very popular in the last 50
43 years. Unfortunately, there are a lot of fatal accidents due to snow avalanches
44 caused by skiers and/or snowboarders. They are of special public interest
45 (Januskovecz, 1989).

46 In Austria, about 25–30 fatalities caused by snow avalanches are expected
47 every year (Neuhold, 2012; Höller, 2009). Furthermore, it is reported that
48 in Alpine countries (such as Austria) the number of fatalities is more or
49 less constant over the time (Brugger et al., 2001; Valt & Pivot 2013; Roth,
50 2013) and that there is some sort of seasonality in the data in terms of
51 higher frequencies of accidents within a distance of 5 or 6 years (Höller,
52 2009; Tschirky et al., 2000). Harvey and Zweifel (2008) even denote that
53 fatalities are decreasing over time in Switzerland. In a most recent paper
54 Techel et al. (2016) investigated avalanche fatalities in the European Alps (in
55 addition to Switzerland–Austria–Slovenia) over time stratified for controlled
56 and uncontrolled terrain, concluding that in the case of uncontrolled terrain
57 the trend seems to be constant over time from the 1980s up to now.

58 Usually trend information for Austrian avalanche fatalities is given in
59 the annual reports of the Austrian Board for Alpine Safety (Kuratorium
60 für alpine Sicherheit, 2016). Considering these profiles, we notice higher
61 frequencies of fatalities in the 1980s. However, the highest frequency in winter
62 1998/99 is due to avalanche fatalities in villages (Galtür, Ischgl), also affecting
63 buildings. This is because the statistics in the reports do not distinguish
64 between fatal avalanches in buildings, on roads, outdoors without skiing,
65 fatalities due to skiing on slopes and backcountry skiing.

66 In this paper our focus is on accidents caused by backcountry (using no
67 ascent support) and off-piste ('leaving the ski resort in order to travel in areas
68 that were not controlled for avalanches', see (Silverton et al., 2009)) skiers or
69 snowboarders. Until now there has not been an investigation for this special
70 group of avalanche incidents in Austria keeping in mind that accidents due
71 to off-piste and backcountry skiing are by far the most common way to be
72 involved in avalanche accidents.

73 Our task in this paper is to carry out a spatial and temporal analysis,
74 identifying (potentially nonlinear) trends over time and regional patterns.
75 In the case of trend analysis, we compare Austrian results with results of
76 Switzerland, France, Italy and the United States.

77 **2 Materials and methods**

78 **2.1 Data**

79 For our study we built a data base of fatal avalanche accidents recording the

- 80 1. date,
- 81 2. municipal area where the accident took place,
- 82 3. federal state of the municipality,
- 83 4. number of persons involved,
- 84 5. number of fatalities,
- 85 6. type of activity (on/off-piste, backcountry skiing, etc.)

86 of fatal accident events in Austria within the winter periods
87 1980/81–2015/16, which are available from the annual reports of the
88 Austrian Board for Alpine Safety (Kuratorium für alpine Sicherheit, 2016)
89 and the annual reports of the information services of the federal states
90 (Amt der Tiroler Landesregierung, 2009). In order to check the reliability of
91 the accident data, we made a cross-check between those reported in the two
92 sources. Looking at winter season 1986/87 we figured out that the reports
93 were incomplete. However, we were able to fill this gap using records of the
94 BFW (Austrian Research Centre for Forests, Institute for Natural Hazards,
95 Innsbruck), e.g. see (Schaffhauser et al., 1988).

96 For the period 1967/68–1979/80 we used aggregated information pub-
97 lished in the annual reports of the Austrian Board for Alpine Safety (Ku-
98 ratorium für alpine Sicherheit, 2016). Starting from 1977/78 we were able
99 to distinguish between backcountry and off-piste fatalities. Finally, further
100 annual reports of the BFW were helpful in order to resolve classification
101 problems of avalanche events.

102 Keeping in mind aspects of data quality, it seems to be that avalanche
103 information back to the period 1967/68 is reliable for our purposes. In general
104 information relating to fatal avalanches seems to be much more reliable than
105 information relating only to avalanches with injured or uninjured persons.
106 Most notably, in the case of fatal avalanches we do not expect that there are
107 records missing.

108 In order to compare Austrian results with international results we use
109 data from the International Commission of Alpine Rescue (ICAR) which
110 was kindly made available for us by the ICAR.

111 The data are annual count data of fatal avalanche events (‘Statistique
112 d’accidents d’avalanche’) based on 21 countries within the periods 1983/84–
113 2015/16 which are categorized by the type of fatalities (backcountry skiing or
114 snowboarding, off-piste, on-piste, alpinist without ski/snowboard, on road,
115 buildings, snowmobile, other).

116 In case of the international data (Switzerland: Frank Techel, ‘Auszug aus
117 der Lawinenschadensdatenbank des SLF’ (SLF 2017); Italy: Mauro Valt,
118 Associazione Interregionale Neve e Valanghe, Trento; France: Frederic Jarry,
119 ANENA; United States: Ethan Greene, Colorado Avalanche Information
120 Center) a crosscheck was carried out.

121 For looking at the regional distribution of avalanche fatalities we built
122 small area maps based on Austrian municipalities. For this purpose
123 we use polygon boundaries of the small-scaled areas provided by the
124 ‘Bundesamt für Eich- und Vermessungswesen’ (BEV) in a shapefile.
125 In order to get a regional overview of the alpine terrain ($\geq 1500m$
126 above sea level) for discussion, we use digital elevation model (DEM)
127 data from the BEV at an $250m$ resolution. Further on, we use data
128 of overnight stays in the winter season 2015/16 at community level
129 provided by the ‘Statistik Austria’ as an additional approach for
130 discussion ([https://www.statistik.gv.at/web_de/statistiken/wirtschaft/
131 tourismus/beherbergung/ankuenfte_naechtigungen/index.html](https://www.statistik.gv.at/web_de/statistiken/wirtschaft/tourismus/beherbergung/ankuenfte_naechtigungen/index.html)).

132 **2.2 Statistical methods**

133 After aggregating the spatio-temporal data y_{st} (denoting the observed fatal-
134 ities at time t and location s) in terms of location, which means summing up

135 over the locations, $\sum_s y_{st}$, we propose the following model for capturing the
 136 trend over time:

$$\log(\mu_t) = f(t) + x_t \quad (2.1)$$

137 where μ_t denotes the expectation of the Poisson distributed number of annual
 138 avalanche fatalities over time t (in our case: winter periods). The logarithms
 139 of these values are modelled as the sum of potentially nonlinear trend function
 140 $f(t)$ and a stationary remainder x_t . We use the Aikake information criterion
 141 (AIC) and the Bayesian information criterion (BIC) in order to compare
 142 the constant, linear and nonlinear model (which is in our opinion the better
 143 choice than reporting pairwise comparisons of p-values for potentially non-
 144 parametric trend functions, see e.g. Venables & Ripley, 2002). To account
 145 for potential serial correlation and periodic variation in the remainder, we
 146 consider autoregressive moving-average (ARMA) effects.

147 After aggregating the spatio-temporal data y_{st} in terms of time, which
 148 means summing up over the time, $\sum_t y_{st}$, we propose a Markov random
 149 field approach modelling the expected number of avalanche fatalities μ_s (s ,
 150 $s \in \{1, \dots, S\}$, denoting the region which are municipalities in our case) as
 151 follows:

$$\log(\mu_s) = Z\beta_s \quad (2.2)$$

152 where the $S \times S$ design matrix Z depends on the specific form of the spatial
 153 layout. The coefficients β_s are conditionally Gaussian distributed (Markov

154 random fields) according to:

$$\beta_s | \beta_{-s} \sim N \left\{ \frac{1}{n_s} \sum_{r \sim s} \beta_r, \frac{\tau^2}{n_s} \right\} \quad (2.3)$$

155 where β_{-s} denotes the vector of parameters without its s th component, n_s is
 156 equal to the number of neighboring regions with reference to region s , $s \sim r$
 157 indexes all units adjacent to region s and τ^2 denotes a (unknown) variance
 158 parameter.

159 For fitting these models we use the R package `mgcv` (R Development Core
 160 Team, 2012; Wood, 2006) which applies the smoothing spline approach for
 161 fitting generalized additive models (GAM).

162 Further on, for looking at the regional distribution of avalanche fatalities
 163 (and subsequently at the regional distribution of alpine terrain and overnight
 164 stays) we build small area maps based on Austrian municipalities using the
 165 geographic information system (GIS) ArcMap. We, of course, use Markov
 166 random field estimates as described above which helps us to identify regional
 167 hot spots of avalanche fatalities.

168 **3 Results**

169 **3.1 Temporal results**

170 In the following, we give the plots of temporal estimated functions of
 171 avalanche fatalities at first plotting the function for Austria in total within
 172 the winter periods 1967/68–2015/16 (see Figure 1). Additionally, we plot
 173 the trend function of exclusively off-piste fatalities starting from the winter

174 season 1977/78 (see Figure 2). Further on, we calculate 90% confidence
175 bands of the estimated functions in both cases as shown in the plots.

176 For reasons of comparison Table 1 gives the frequencies of backcountry,
177 off-piste and total fatalities of Austria and the Austrian neighboring countries
178 Italy and Switzerland within the winter periods 1983/84–2015/16. Addition-
179 ally the off-piste percentages are reported. Furthermore, we report the results
180 of fatalities in France, which turns out to be the country with the highest
181 counts of fatalities in Europe, and the results of the United States, which is
182 probably the most important country outside of Europe in terms of avalanche
183 fatalities. For this purpose, however, we use ICAR data as described above.

184 For further international comparison we consider estimated functions of
185 off-piste and backcountry avalanche fatalities (and off-piste fatalities de-
186 tached) of Switzerland, France, Italy and the United States in Figures 3–6.

187 Finally, the Aikake information criterion (AIC) and the Bayesian infor-
188 mation criterion (BIC) of the constant (no trend effect), linear and nonlinear
189 models are reported for model comparison – see Table 2. Lower AIC- and
190 BIC-values, however, indicate significantly better fits when comparing the
191 different models.

192 **3.2 Regional results**

193 Figures 7 and 8 show the regional distribution of fatal avalanche events (Fig-
194 ure 7 in total and Figure 8 off-piste only) using colored maps based on small
195 areas, which are the Austrian municipalities in our case. The coloring, how-
196 ever, is based on Markov random field estimates of avalanche fatalities as
197 described in the previous Section; the number corresponding with each spa-

208 tial unit in the plot is equal to the original count.

209 In addition to Figures 7–8, Table 3 gives a list of those municipalities with
200 the most avalanche fatalities in Austria. Further on, we list those avalanche
201 events in Austria with the highest counts of fatalities in Table 4 which turns
202 out to be useful for the discussion section.

203 Finally, Figure 9 shows the distribution of Alpine terrain ($\geq 1500m$ above
204 sea level) and the distribution of the overnight stays in the winter season
205 2016 at municipal level (restricted to the 130 municipalities with more than
206 100,000 overnight stays in Austria) which allows us to discuss possible reasons
207 for the observed distribution of avalanche fatalities in Figure 7 and Figure 8.

208 4 Discussion

209 4.1 Temporal analysis with an international overview

210 If we look at the trend function of Austria in total (see Figure 1) we notice
211 an increasing trend having its maximum at winter period 2005/06 (1969/70:
212 approx. 12, 2005/06 approx. 22). In recent years we, however, notice that
213 the number of annual fatalities is slightly decreasing.

214 Additionally we take notice of a peak in the 1980s ranging between
215 1981/82 and 1987/88. But keeping in mind that increased snowfall has an es-
216 sential effect on the number of accidents (Harvey, 2008; Harvey et al., 2012;
217 Höller, 2012), increased solid precipitation in the 1980s during wintertime
218 (Laternser & Schneebeli, 2003; Abegg, 1996) could give some evidence for
219 this pattern.

220 Looking at the off-piste trend function (see Figure 2), we notice an in-

221 creasing (linear) trend without any peak in the 1980s. As in the ‘total’ case,
222 the off-piste fatalities are slightly decreasing from the mid 2000s on.

223 Lower AIC- and BIC-values (see Table 2) indicate that the nonlinear
224 model is preferable to the constant or linear model – although in case of
225 ‘Austria off-piste’ the BIC-value indicates that the linear model seems to be
226 preferable.

227 Considering ARMA effects, we did not find any substantial serial correla-
228 tion or any sort of periodicity in the remainder x_t . Further on, we notice that
229 there is a lot of variation of the observed counts around estimated function(s).

230 Comparing Austrian fatal backcountry and off-piste counts within
231 1983/84 – 2015/16 with results of counts in Switzerland, France, Italy and
232 the United States (see Table 1) we notice, led by France (787 fatalities in
233 total, 23.85 fatalities per year), the second largest number of total avalanche
234 fatalities (680, 20.61) in Austria. Having a focus on backcountry fatalities
235 only, Austria is leading (458, 13.88) followed by France (433,13.12) and
236 Switzerland (395, 11.97). In Austria a share of 32.65% of total fatalities are
237 due to off-piste accidents (largest value France: 44.98%; smallest: United
238 States 29.23%).

239 Comparisons with total fatality profiles of France, Switzerland and Italy
240 result in:

- 241 1. high frequencies in the 1980s,
- 242 2. low counts in the 1990s,
- 243 3. increasing trend beginning in 2000
- 244 4. to some extent decreasing in recent years,

245 which in turn is rather similar to the results of Austria.

246 However, if we consider the results of the United States in Figure 6 (284
247 total fatalities, 8.61 fatalities per year) we note a positive almost linear trend
248 without any peaks in the 1980s. The AIC- and BIC-values indicate that,
249 with the exception of the United States (linear model), nonlinear models are
250 preferable (whereas the BIC-values of France almost indicate that there is
251 no effect at all in case of France).

252 If we compare the off-piste trends of the countries we notice quite different
253 shapes to those of Austria (positive trend without peak in the 1980s):

- 254 1. Italy: similar to shape as seen in case of total counts.
- 255 2. Switzerland: difference to total trend function, peak of off-piste trend
256 around year 2000.
- 257 3. France: decrease of off-piste counts in recent years.
- 258 4. United States: almost no increase; because of the lowest AIC-value,
259 the constant model turns out to be the best one.

260 Such as in the ‘total’ case above, lower AIC- and BIC-values indicate that,
261 with the exception of the United States (constant model), nonlinear models
262 are best-performing. Usually trend information is given as a linear function
263 in the literature for avalanche data, see e.g. (Tschirky et al., 2000; Harvey &
264 Zweifel, 2008; Spencer & Ashley, 2011; Page et al., 1999). Our investigations
265 - see AIC- and BIC-values in Table 2 - showed that (with the exception of the
266 US-data) linear models are not appropriate – see also the results of (Techel
267 et al, 2016) and (SLF, 2016) in the recent research.

268 At the beginning and the end of the longitudinal profiles we observe larger
 269 confidence bands indicating less precise estimates due to missing data in their
 270 neighbourhoods. As a result of this extreme estimates at the beginning of
 271 the temporal profiles could be less reliable (e.g. in case of Switzerland ‘total’,
 272 if we compare the results with those of SLF, 2016)

273 The temporal profiles could also be seen as an indicator for low/high
 274 frequency temporal clusters, which are: Austria total (6 larger values) in the
 275 mid 1980s; Switzerland off-piste, France and Italy total (5 smaller values for
 276 each) in the early and mid 1990s.

277 4.2 Regional analysis

278 In Figure 7 we explore the regional or spatial distribution of avalanche fatal-
 279 ities in Austria within the years 1981–2016. Here the total area of Austria
 280 is divided into small areas, equal to the areas of the Austrian municipalities
 281 (211 municipalities with at least one reported fatality). Looking at Table
 282 3, we notice that the municipalities with highest numbers are ‘Sölden’ and
 283 ‘St. Anton a. Arlberg’ Around the municipalities ‘St. Anton a. Arlberg’ and
 284 ‘Sölden’ in the western part of the Austrian federal state Tyrol we observe 2
 285 clusters or hot spots of increased fatalities:

286 The first cluster (CL1), centered around the regions Arlberg and Silvretta,
 287 is including the municipalities St. Anton a. Arlberg (number of avalanche
 288 fatalities: 31), Kaisers (10), Klösterle (9), Lech (22) in Arlberg, and the
 289 municipalities St. Gallenkirch (8), Gaschurn (8), Galtür (21), Ischgl (9) in
 290 Silvretta.

291 The second cluster (CL2), located in the southern part of Ötztal, Kühtai

292 and Stubai, is including the municipalities Sölden (50), St. Leonhard i. Pitztal
 293 (18), Längenfeld (9) in the Ötztal Alps, and the municipalities St. Sigmund
 294 i. Sellrain (11), Silz (14), Sellrain (5), Neustift i. Stubaital (11) in Kühtai-
 295 Stubai.

296 Further on, we observe some smaller spots in the federal states:

- 297 – Tyrol (Tuxer Alpen): Navis (9), Wattenberg (9), Schmirn (5), Tux (10)
- 298 – Salzburg (Saalbach): Saalbach-Hinterglemm (10), Niedernsill (13)
- 299 – Styria (Triebener Tauern – Seckauer Tauern): Gaal (6), Wald am
 300 Schoberpaß (6), Hohentauern (6).

301 Finally we notice some single areas with increased frequency such as:

302 Mittelberg Vorarlberg (10), Heiligenblut Carinthia (11), Werfenweng
 303 Salzburg (15), Pusterwald Styria (10). Some single areas with increased
 304 frequencies, e.g. Werfenweng (15) and Niedernsill (13), are due to disastrous
 305 single avalanche events, see e.g. Table 4.

306 Figure 8 plots the distribution of the off-piste fatalities (without back-
 307 country fatalities; 77 municipalities with at least one reported off-piste fa-
 308 tality). As a conclusion we notice 2 hot spots of off-piste fatalities which
 309 are: ‘St. Anton a. Arlberg’ - ‘Lech’ - ‘Ischgl’ (Arlberg, Ischgl) and ‘Sölden’
 310 (southern part of Ötztal).

311 Furthermore, there are some single spots or small clusters such as: Tux
 312 Tyrol (5), Jochberg Tyrol (5), Saalbach-Hinterglemm Salzburg (6), Niedern-
 313 sill Salzburg (12).

314 If we compare Figure 7 and Figure 8 (or if we have a look at Table 3)
 315 we notice centres of off-piste avalanche fatalities in CL1 such as Lech (20 off-
 316 piste fatalities out of 22 total, 90.91% off-piste), St. Anton a. Arberg (26 out

317 of 31 total, 83.87%) and Ischgl (9 out of 9 total, 100%) while the accidents
318 of Galtür (0% off-piste), St. Gallenkirch (2 out of 8 total) and Gaschurn (0%
319 off-piste) are mainly due to backcountry skiers.

320 Looking at CL2, the fatal accidents of our interest are mainly caused by
321 backcountry skiers except Sölden which off-piste rate is about 50% (25 out
322 of 50 total, $\geq 31.24\%$ in case of Austria).

323 Figure 9 (distribution of alpine terrain and overnight stays in the win-
324 ter season 2015/16) tries to give some idea in order to explain the spatial
325 distribution of avalanche fatalities of Figure 7 and Figure 8. Obviously, the
326 percentage of alpine terrain at municipal level coincides with the number of
327 fatalities. However, there are alpine areas with less number of fatalities than
328 in those in the western part of Tyrol, see e.g. East Tyrol. The majority of
329 fatalities are restricted to 2 clusters, which is more or less only a small part
330 of the terrain of our interest.

331 Looking at the overnight stays in Figure 9, we notice that the largest
332 counts of overnight stays coincide with the largest counts of in total and
333 off-piste fatalities (Sölden, St. Anton a. Arlberg, Lech), but there are winter
334 tourist regions with less number of avalanche fatalities, see e.g. the Tauern
335 region or the northeastern part of Tyrol. In the case of total number of
336 fatalities, overnight stays are partly misleading because they e.g. do not take
337 into account the considerable number of native backcountry skiers around
338 Innsbruck.

339 This is more or less in agreement with considering the size of Austrian
340 ski resorts, see (Fleischhacker, 2016), instead of overnight stays (Spencer and
341 Ashley (2011) stated that areas with higher winter sports activity are those

342 with higher number of avalanche fatalities).

343 Finally, if we consider the spatial patterns of buildings exposed to snow
344 avalanches in Austria (Fuchs et al., 2015) we could find some remarkable
345 congruences (looking at CL1 and CL2) if we compare them with avalanche
346 fatalities at municipal level.

347 5 Conclusion

348 As the result of the trend analysis we notice an increasing trend (although
349 decreasing in recent years) of off-piste and backcountry avalanche fatalities
350 within the winter periods from 1967/68 to 2015/16. This clearly contradicts
351 the widespread opinion that the number of fatalities is constant over time.

352 Comparing results of off-piste and backcountry avalanche fatalities in
353 Austria with other relevant countries we notice the second highest number
354 of off-piste and backcountry fatalities in Austria and the largest number of
355 backcountry fatalities in Austria. We notice similar estimated functions if
356 we compare Austrian results with results of the relevant European countries.
357 However, the off-piste trend function of Austria is quite different to those
358 of the other relevant European countries (but similar to those of the United
359 States).

360 As the result of the regional analysis we notice two hot spots of avalanche
361 fatalities in Figure 7: ‘St. Anton a. Arlberg (29)’ (Arlberg-Silvretta) and
362 ‘Sölden (43)’ (southern part of Ötztal, Stubai-Kühtai).

363 Because of the increasing trend (although decreasing in recent years) and
364 the rather ‘narrow’ regional distribution of the fatalities, consequences on

365 prevention of avalanche accidents are highly recommended, e.g. starting a
366 ‘campaign against avalanche accidents’ in the centers of the clusters St. Anton
367 and Sölden. This should especially be done in order to prevent the large
368 number of off-piste (freerider) fatalities in St. Anton-Lech-Ischgl and Sölden.

369 Unfortunately, we are not able to verify the influence of increased num-
370 ber of backcountry and off-piste skiers over time because there is no valid
371 information about frequencies of backcountry and off-piste skiers in general.
372 However, we find some evidence that increased winter overnight stays (which
373 could be seen as an evidence for increased winter sports activity) has an effect
374 on higher number avalanche fatalities, see Figure 9.

375 Finally, we do not hesitate to mention that further research is needed, e.g.
376 to explore the influence of new fallen snow, temperature, etc. on the number
377 of fatalities. For this purpose, further and more precise data are necessary.

378 **Acknowledgement:** The authors kindly acknowledge data provision
379 from the International Commission of Alpine Rescue (ICAR) by Hans-Jürg
380 Etter. Additionally, the authors are grateful to Frank Techel (SLF, Davos
381 Switzerland), Mauro Valt (Associazione Interregionale Neve e Valanghe,
382 Trento Italy), Frederic Jarry (Association Nationale pour l’Étude de la
383 Neige et des Avalanches, Grenoble France) and Ethan Greene (Colorado
384 Avalanche Information Center, Boulder United States) which allowed us to
385 check the data provided from the ICAR.

386 Financial/Material Support: None

387 Disclosures: None

388 References

- 389 Abegg B. Klimaänderung und Tourismus: Klimafolgenforschung am
390 Beispiel der Wintertourismus in den Schweizer Alpen. Zurich,
391 Switzerland: vdf Hochschulverlag AG; 1996.
- 392 Amt der Tiroler Landesregierung. Schnee und Lawinen. Jahresberichte.
393 Innsbruck, Austria: Raggl; 1994–2009.
- 394 Brugger H, Durrer B, Adler-Kastner L, Falk M, Tschirky F. Field manage-
395 ment of avalanche victims. *Resuscitation* 2001; 51(1):7–15.
- 396 Fleischhacker V. Aktuelle Nachfragetrends im Winter-
397 sporttourismus in Österreich. 2016. Available from:
398 http://netoffice.salzburgerland.com/downloads/salzburgerland_aktuell/0916
- 399 Fuchs S, Keiler M, Zischg A. A spatiotemporal multi-hazard exposure as-
400 sessment based on property data. *Nat Hazards Earth Syst Sci* 2015;
401 15:2127—2142.
- 402 Harvey S, Zweifel B. New trends of recreational avalanche accidents in
403 Switzerland. Proceedings of the ISSW2008. Whistler, Canada; 2008.
- 404 Harvey S. Mustererkennung in der Lawinenkunde. Sicherheit im Bergland.
405 Jahrbuch des Kuratoriums für alpine Sicherheit. Vienna, Austria: Ku-
406 ratorium; 2008:53–72.
- 407 Harvey S, Rhyner H, Schweizer J. Lawinenkunde: Praxiswissen für Ein-
408 steiger und Profis zu Gefahren, Risiken und Strategien. Munich, Ger-
409 many: Bruckmann Verlag; 2012.

- 410 Höller P. Avalanche cycles in Austria: an analysis of the major events in
411 the last 50 year. *Nat Hazards* 2009; 48(3):399–424.
- 412 Höller P. The cumulation of avalanche accidents in certain periods - an anal-
413 ysis of backcountry events in Austria. *Proceedings of the ISSW2012*.
414 Anchorage, USA; 2012:32–339.
- 415 Januskovecz A. Zeitungsberichterstattung über Naturkatastrophen,
416 Ansätze für die forstliche Öffentlichkeitsarbeit zum Theme
417 Lawinen–Hochwasser–Muren. Vienna, Austria: Hochschulschrift ,
418 Universität für Bodenkultur; 1989.
- 419 Kuratorium für alpine Sicherheit. Sicherheit im Bergland. Jahrbücher
420 des Kuratoriums für alpine Sicherheit. Vienna, Austria: Kuratorium;
421 1973–2016.
- 422 Laternser M, Schneebeli M. Long-term snow climate trends of the Swiss
423 Alps. *Int J Climatol* 2003; 23(7):733–750.
- 424 Neuhold T. 25 Lawinentote werden akzeptiert. Pressebericht zum Law-
425 inenkolloquium 2012 Salzburg. *Der Standard Printausgabe* 16.1.2012.
426 Vienna, Austria: Standard Verlagsgesellschaft m. b. H.; 2012.
- 427 Page C, Atkins D, Shockley L, Yaron M. Avalanche deaths in the United
428 States: A 45-year analysis. *Wilderness Environ Med* 1999; 10:146–151.
- 429 R Development Core Team. R: A language and environment for statistical
430 computing. Vienna, Austria; 2012. Available from: [http://www.r-](http://www.r-project.org/)
431 [project.org/](http://www.r-project.org/)

432 Roth E. Lawinen: verstehen–vermeiden–Praxistipps. Munich, Germany:
433 Bergverlag Rother; 2013.

434 Schaffhauser H. Lawinenereignisse und Witterungsverlauf in Österreich.
435 Winter 1986/87. FBVA-Berichte 35. Vienna, Austria: Österreichischer
436 Agrarverlag; 1988.

437 Silvertown NA, McIntosh SE, Kim HS. Risk Assessment in Winter Backcoun-
438 try Travel. Wilderness Environ Med 2009; 20:269–274.

439 SLF. Lawinen in der Schweiz: Entwicklung in den letzten 80 Jahren. Avail-
440 able from: <http://www.slf.ch/praevention/lawinenunfaelle/Publikationen/lawinenunfaelle/Sp>

441 Spencer J, Ashley. Avalanche fatalities in the western United States: a
442 comparison of three databases. Nat Hazards 2011; 58(1):31–44.

443 Techel F, Jarry F, Kronthaler G, Mitterer S, Nairz P, Pavšek M, Valt M,
444 Darms G. Avalanche fatalities in the European Alps: long-term trends
445 and statistics. Geographica Helvetica 2016; 71:147–159.

446 Tschirky F, Brabec B, Kern M. Lawinenunfälle in den Schweizer Alpen
447 – eine statistische Zusammenstellung mit den Schwerpunkten Ver-
448 schüttung, Rettungsmethoden und Rettungsgeräte. 2000. Available
449 from: [http://www.slf.ch/praevention/lawinenunfaelle/unfallstatistik-](http://www.slf.ch/praevention/lawinenunfaelle/unfallstatistik-de.pdf)
450 [de.pdf](http://www.slf.ch/praevention/lawinenunfaelle/unfallstatistik-de.pdf)

451 Valt M, Pivot S. Avalanche accident documentation is of fundamental
452 importance to understand the dynamics, taking place in snow, of
453 risky activities in order to implement the best possible prevention

454 strategies. Proceedings of the ISSW2013. Grenoble-Chamonix,
455 France; 2013:1142–1147.

456 Venables WN, Ripley BD. Modern applied Statistics with S. New York,
457 United States: Springer; 2002.

458 Wood SN. Generalized Additive Models: An Introduction with R. Boca
459 Raton, USA: Chapman and Hall/CRC; 2006.




Figure 1: Observed (○) and estimated (●) annual total avalanche fatalities (off-piste and backcountry) with 90% confidence band (grey) in Austria within 1967/68–2015/16.




Figure 2: Observed (o) and estimated (•) annual off-piste avalanche fatalities with 90% confidence band (grey) in Austria within 1977/78–2015/16.




Figure 3: Observed (\circ) and estimated (\bullet) annual avalanche fatalities (off-piste and backcountry, i.e. total, on the left and off-piste on the right) with 90% confidence bands (grey) in Switzerland within 1983/84–2015/16.




Figure 4: Observed (○) and estimated (●) annual avalanche fatalities (off-piste and backcountry, i.e. total, on the left and off-piste on the right) with 90% confidence bands (grey) in France within 1983/84–2015/16.




Figure 5: Observed (o) and estimated (●) annual avalanche fatalities (off-piste and backcountry, i.e. total, on the left and off-piste on the right) with 90% confidence bands (grey) in Italy within 1983/84–2015/16.




Figure 6: Observed (○) and estimated (●) annual avalanche fatalities (off-piste and backcountry, i.e. total, on the left and off-piste on the right) with 90% confidence bands (grey) in the United States within 1983/84–




Figure 7: Regional distribution of avalanche fatalities (off-piste and backcountry) in Austria within 1980/81–2015/16.




Figure 8: Regional distribution of avalanche fatalities (off-piste) in Austria within 1980/81–2015/16.




Figure 9: Distribution of Alpine terrain ($\geq 1500m$ above sea level) and number of overnight stays in the winter season 2016 at community level.

Country	Backcountry		Off-piste		Total		% off-piste
	number	per year	number	per year	number	per year	
Austria	458	13.88	222	6.73	680	20.61	32.65%
Switzerland	395	11.97	222	6.73	617	18.70	35.98%
France	433	13.12	354	10.73	787	23.85	44.98%
Italy	322	9.76	138	4.18	460	13.94	30.00%
USA	201	6.09	83	2.52	284	8.61	29.23%
sum	1785	54.09	1033	31.30	2818	85.39	36.66%

Table 1: Number of avalanche fatalities and annual average (off-piste, back-country and total) of 5 countries within the winter periods 1983/84–2015/16.

		Total			Off-piste		
		const	linear	nonlin.	const	linear	nonlin.
Austria	AIC	550.87	543.14	530.3	241.35	238.17	236.46
	BIC	552.76	546.93	539.46	243.02	241.5	243.03
Switzerland	AIC	256.47	254.29	242.79	189.9	191.87	186.79
	BIC	257.96	257.29	250.1	191.4	194.87	192.83
France	AIC	268.9	270.9	267.74	251.2	253.09	245.7
	BIC	270.39	273.89	275.39	252.7	256.08	252.28
Italy	AIC	285.01	286.78	250.69	189.79	191.62	175.23
	BIC	286.5	289.77	257.59	191.29	194.61	180.78
United States	AIC	188.64	182.43	186.33	147.45	148.92	151.33
	BIC	190.13	185.42	192.65	148.95	151.92	156.4

Table 2: AIC and BIC of the constant, linear and nonlinear trend model considering data of Austria total and off-piste (Figure 1, Figure 2), Switzerland total and off-piste (Figure 3), France total and off-piste (Figure 4), Italy total and off-piste (Figure 5) and United States total and off-piste (Figure 6).

Communitiy	Backcountry	Off-piste	Total
Sölden	25	25	50
St. Anton am Arlberg	5	26	31
Lech	2	20	22
Galtür	21	0	21
St. Leonhard im Pitztal	14	4	18
Werfenweng	13	2	15
Silz	14	0	14
Niedersill	1	12	13
Neustift im Stubaital	7	4	11
Heiligenblut am Großglockner	9	2	11
St. Sigmund im Sellrain	11	0	11
Tux	5	5	10
Kaisers	6	4	10
Mittelberg	6	4	10
Saalbach-Hinterglemm	4	6	10
Pusterwald	9	1	10
Klösterle	4	5	9
Navis	9	0	9
Ischgl	0	9	9
Längenfeld	9	0	9
Wattenberg	9	0	9
Gaschurn	8	0	8
St. Gallenkirch	6	2	8
Fügenberg	5	2	7
Jochberg	1	5	6
Axams	3	3	6
Gaal	6	0	6
Häselgehr	6	0	6
Wald am Schoberpaß	6	0	6
Hohentauern	4	2	6
Mallnitz	6	0	6
Prägraten am Großvenediger	5	0	5
Tweng	2	3	5
Nauders	2	3	5
Kitzbühel	3	2	5
Serfaus	0	5	5
Sellrain	5	0	5
Schmirn	5	0	5
Fusch an der Großglocknerstraße	5	0	5
Alpbach	4	1	5
Bad Gastein	3	2	5
Rohrmoos-Untertal	5	0	5
Untertauern	3	2	5

Table 3: Number of avalanche fatalities (off-piste, backcountry and total) in Austria within 1980/81–2015/16 stratified for communities with more than 4 fatalities in the observation period.

Date	Location	Municip.	Fatalities
1982-01-31	Werfenweng	Werfenweng	13
2000-03-28	Schmiedinger Kogel	Niedersill	12
1999-12-28	Jamtalhütte - Gde. Galtür	Galtür	9
1987-04-05	Idalpe	Ischgl	6
1988-03-28	Jamtal	Galtür	6
2009-05-02	SchalFKogel	Sölden	6
2016-02-06	Wattener Lizum	Wattenberg	5
1985-03-21	Sonntagkarzinken, Schladm. Tauern	Rohrmoos-Untertal	4
1988-02-14	Hühnereggen, Stubai Alpen	Sellrain	4
1993-04-12	Querkogeljoch, Ötztaler Alpen	Sölden	4
1997-02-18	Luxnacher Sattel	Häselgehr	4
2005-01-22	Rendl	St. Anton a. Arlberg	4
1981-03-01	Hohe Veitsch	Mürzsteg	3
1984-02-19	Hoher Gleirsch, Karwendelgebirge	Scharnitz	3
1985-05-04	Speikogel, Kitzbüheler Alpen	Westendorf	3
1986-01-08	Kühkarkopf, Hohe Tauern	Fusch a. d. Großglocknerstr.	3
1986-04-01	Tschambreuspitze, Silvretta	Gaschurn	3
1986-04-07	Windachscharte, Stubai Alpen	Sölden	3
1986-12-21	Lattenberg Triebener Tauern	Wald a. Schoberpaß	3
1987-01-06	Fluchtalpe, Kleines Walsertal	Mittelberg	3
1987-04-18	Scharkogel	Uttendorf	3
1991-12-21	Scharnitzfeld, Wölzer Tauern	Pusterwald	3
1995-01-03	Schöngraben/Törli	St. Anton a. Arlberg	3
1995-02-11	Scheibenspitze	Navis	3
1996-03-09	Frommerkogel, Tennengebirge	Hüttau	3
1996-04-03	Murkarspitze, Gde. Längenfeld	Längenfeld	3
2000-03-16	Wasserradkopf	Heiligenblut	3
2000-11-19	Roßkarschneid	Sölden	3
2003-01-30	Scharnitzalm, Scharnitzfeld	Pusterwald	3
2004-12-20	Mohnenfluh	Lech	3
2005-02-22	Sulzkogel	Silz	3
2005-03-05	Rotschrofenspitze	Kaisers	3
2013-01-18	Mittagskofel, Karnische Alpen	Lesachtal	3

Table 4: List of avalanche events (off-piste or backcountry) in Austria within 1980/81–2015/16 with more than 2 fatalities in each event.