

## ***Interactive comment on “Geomorphological evolution of landslides near an active normal fault in Northern Taiwan, as revealed by LiDAR and unmanned aircraft system data” by Kuo-Jen Chang et al.***

**Kuo-Jen Chang et al.**

epidote@ntut.edu.tw

Received and published: 10 October 2017

Thank you for the helpful comments on our manuscript. Please find below our response and modifications that we have revised in the manuscript following the referee's comments and suggestions.

Anonymous Referee #2 The study by Chang et al. investigates two large landslides developed along an active normal fault in a volcanic environment. Starting from previous knowledge about two large landslides in the area, the authors build their study on

C1

mapping the two landslides from visual interpretation of UAS imagery, as well as the interpretation of high-resolution digital topography (1 x 1 m LiDAR DEM). Based on their mapping, they estimate the volume of the two landslides by subtracting the present day topography from a reconstructed pre-failure topography. They conclude that the volume obtained is six times higher than the reported largest landslide volume in Taiwan. They further postulate that an active normal fault controlled the morphological evolution of the two landslides, and that ongoing faulting is responsible for maintaining landslide hazard condition in the study area. While it is interesting the attempt of the authors to relate landslide evolution directly to fault activity, I'm not fully convinced by the story they want to tell. I identified many issues and problems with the data (1), methods (2), and interpretations (3) that preclude this from being a convincing study. These include lack of clarity in data and methods and what was actually measured, issues with the interpretations and what the data mean, and a lack of depth in the interpretations and implications that are drawn from the data.

Referee #2-1. I have reservations about some of the assumptions that the authors have gone into their dataset. In particular, I don't know where their slip surfaces position estimates have come from. These are critical, because it is the postulated spatial coincidence between the slip surfaces and the present-day topography that provides the condition to calculate the landslides volume according to the method presented in the paper. The authors are not clear at this point: only short and general shrift are done at lines 15-20 page 8, but without any geological evidence or examples, it's hard to know what, exactly, they have considered for their assumption. Geology of the area is presented in figure 1, but the figure is not informative enough to support the assumption of the authors. Clearly, the present day topography is somehow related to the movement along the slip surfaces, but I think the authors need to be a lot more careful about what they say, and do a better job of documenting why the present day topography can be considered the slip surface of an old landslide. I also have reservation about the landslide detection, mapping and classification. Figure 5 illustrate the detection of zones affected by mass movements highlighted by ridges and scarps, which

C2

are commonly interpreted as the topographic response to movements along the slip surfaces at depth. However, the evidences strongly contrast with the assumption done by the authors about the coincidence between the slip surface and the present-day topography. This is a main issue that the authors should address to be their contribution convincing. In addition, I have reservations about the mapping itself. Landslide mapping should include the definition of the scarp area, deposit area, and both the flanks (see for instance Santangelo et al. 2015 NHESS, 15, 2111–2126; Guzzetti et al. 2012, Earth Science Reviews, 112, 42-66; Ambrosi and Crosta, 2006, Engineering Geology, 83, 183-200). Looking Figure 5, I really don't know where the limits (even supposed) of the two landslides are positioned. The circumstance undermine the possibility to visually appreciate and to quantitatively measure landslide area in map. Furthermore, the paper is not informative enough about the landslide type, landslide age (even relative age) and different generation of landslides recognized inside the old landslides. The information is necessary to characterize the landslide morphology, evolution and hazard, which are specific purposes of the paper. I think a more detailed mapping using the high quality materials (UAS imagery and LiDAR DEM) available to the authors should be add to the paper.

Response #2-1: We have reconstructed the above section into 5 separate questions (a to e), and responded these questions accordingly:. a) The slip surfaces and the present-day topography that provides the condition to calculate the landslides volume according to the method presented in the paper, but without any geological evidence or examples. Response #2-1a: Indeed, to estimate the landslide volume, the original topography and slip surface are the key issues. However, with regarding to an old landslide, the original surface is unknown. On the other hand, slip surface is usually covered by the slid mass, and is not easily exposed. Therefore, in this study we try to propose one of the methods to reasonably construct reasonably by the original ground surface and assume the location of the slip surface that likely located at the interface between the volcanic cover and the underlying sedimentary rocks. The original ground surface is constructed from ideal volcano cone edifice. The sedimentary rock basement

C3

and the volcanic rock cover have been well mapped both on the geologic maps (Fig. 1 and the new Fig. 5) and in field survey in the region. Based on the distribution of rock types, it is supposed that the contact between the volcanic cover and the underneath sedimentary rocks may serve as a weak plane for the slip surface. The slip surface consists from the difference of material and the exposed different lithology.

We have revised and newly improved the paragraph in the manuscript. b) Geology of the area is not informative enough to support the assumption of the authors, and do a better job of documenting why the present day topography can be considered the slip surface of an old landslide. Response #2-1b: We have now added a figure to show more detailed local geologic conditions. In the new geological map, many landslides that occurred in the study area and in the Tatum Volcano region were attached to demonstrate the distribution of landslides. Compare Comparing the size, distribution and classification, etc. the two largest landslides (XSL and CSL) were thus chosen as the target for this study. We have newly revised and improved the paragraph in the manuscript. On the other hand, the 2D hillshade map (the original Fig. 5, now change to Fig. 6) has been modified with the azimuth of shade illumination to being 315° so as to better illustrate the landslide geomorphologic features.

c) Figure 5 illustrate the detection of zones affected by mass movements highlighted by ridges and scarps, which are commonly interpreted as the topographic response to movements along the slip surfaces at depth. However, the evidences strongly contrast with the assumption done by the authors about the coincidence between the slip surface and the present-day topography. This is a main issue that the authors should address to be their contribution convincing. Response #2-1c: The original Fig. 5 is now modified as Fig. 6. Indeed, ridges and scarps of a landslide are commonly interpreted as the topographic response of the movements along the slip surfaces at depth. However, the topographic feature responses reflect only the ground subsidence actually. Thus if the slid mass glides with a long run out distance or the displaced mass has been eroded away, both processes will preserved topographic remain relicts by

C4

distinct shutter ridges and scarps. In consequence, in this study we interpret that most of the material has been eroded away, and discussed from the perspectives of normal faulting and tectonic setting of the study area. We have newly improved the manuscript to better illustrate the full overall framework of the study.

d) Looking Figure 5, I really don't know where the limits (even supposed) of the two landslides are positioned. The circumstance undermine the possibility to visually appreciate and to quantitatively measure landslide area in map. Response #2-1d: The original figure Fig. 5 (now modified as Fig. 6) is has been modified with the azimuth of shade illumination to being  $315^\circ$  so as to better illustrate the landslide geomorphologic features. This new hillshade image should shall improve essentially the identification of landslide region visually, since because not necessary that all readers are familiar with the landslide morphology.

e) The paper is not informative enough about the landslide type, landslide age (even relative age) and different generation of landslides recognized inside the old landslides. Response #2-1e: The normal faulting in the region started since from 400Ka, and is activated continuously ever since. The faulting being was identified in the Taipei basin area and northeastern offshore Taiwan, with the fault line situated on both two sides of the study area. And the fault line was recently identified and linked together as only one normal fault in Tatun Volcano region (near and surrounding the study area) by the authors. In conclusion, for the relative age of the landslide, we interpret that the landslide has been triggered since right after normal faulting started and the formation of Tatun Volcano, say which is far later than 200 Ka. Regarding to the different generation of landslide, the geomorphologic components show different degrees of preservation within the two observed landslides. Furthermore, the CSL is interpreted to have occurred from a combination of multiple landslide events. We have newly improved revised the manuscript to denote the relative age of the landslide, and the different generation of landslides as well.

Referee #2-2. Although the method seems to be reasonable in theory, too many is-

C5

sues remain unexplained. For instance: I disagree with the assumption that detailed UAV imagery are better than aerial photographs and/or satellite images to detect and characterized large landslides. My own experience suggest quite the opposite. Indeed, UAV imagery and detailed LiDAR DEM are very useful to perform detailed studies. As a matter of fact, one of the more interesting piece of work in the paper is related to the characterization of the micro-topography of the landslides and the discussion about the possibility to apply the method to the study of gully erosion. However, gully erosion appear to be as a minor complication compared to the estimation of the landslide volume of a giant landslide. Complication is irrelevant here if the authors focus their paper on the calculation of the total landslide volume.

Response #2-2: We have reconstructed divided the above section into 2 questions (a to b), and response responded the questions accordingly: a) I disagree with the assumption that detailed UAV imagery are better than aerial photographs and/or satellite images to detect and characterized large landslides. Response #2-2a: In Taiwan, on the one hand heavy precipitation induced by the annual northeast monsoon markedly modifies easily the landslide topography. On the other hand, the study region is situated within a national park and preserves well dense forest very well. Both effects conceal detailed topography and nearly impossible to study directly from aerial photographs and/or satellite images. The same situation can be found in the Tsaoling and Jiufengershan two giant landslides (namely, Tsaoling and Jiufengershan) triggered by the Chi-Chi earthquakes, where the vegetation colonization cover and concealed almost all the topographic features details, especially for the zone of accumulation, in just only ten years after the landslides has been occurred. That is why we deploy employed two high-resolution and high-precision data datasets/methods, the UAV and the airborne LiDAR, to decipher the landslide features of the study area. And that is why we insist assert the quality and different levels of the data sets, and illustrated and denoted them in Figs. 2 and 4. We have newly improved revised and clarified the documentation in the manuscript.

C6

b) Gully erosion appears to be as a minor complication compared to the estimation of the landslide volume of a giant landslide. Complication is irrelevant here if the authors focus their paper on the calculation of the total landslide volume. Response #2-2b: Yes, the gully incision is a minor factor to estimate the overall landslide volume. The method is used only to assess the landslide morphology and evolution. We have clarified the documentation in the manuscript.

Referee #2-3. The final interpretation is not convincing and rise many question: Why just such two landslides developed along a regional normal fault? What about other places along the fault? There is somethings peculiar in the specific location of the two landslides? (i.e. relative relief higher respect to other places along the fault?) geo-structural setting different respect to other places along the fault and prone to landslides? cluster of strong earthquakes? evidence of high vertical deformation rates? what else?) In the scheme proposed by the authors the fault is the main factor controlling both the onset and the disruption of the landslides, but no analysis support their conclusion. I have also reservation about the idea that normal fault activity has the effect of cancel the landslide signature (third diagram in the final scheme). I think quite the opposite; fault activity sustain relief formation, maintaining the condition for landslide development (see Bucci et al. 2016, ESPL, 41, 711-720; and Densmore et al. 1997, Science 275, 369-72). The authors conclude somethings similar at lines 27-29 page 12, but their statement conflict with the idea illustrated in the scheme. Finally, the authors never explicitly address time scales of the considered landslides and fault, as well as the probable mismatch in timescale of the landsliding and faulting processes.

Response #2-3: We have reconstructed divided the paragraph by into 3 questions (a to c) and responded se accordingly: a) Why just such two landslides developed along a regional normal fault? What about other places along the fault? There is somethings peculiar in the specific location of the two landslides? (i.e. relative relief higher respect to other places along the fault?) geo-structural setting different respect to other places along the fault and prone to landslides? cluster of strong earthquakes? evidence of

C7

high vertical deformation rates? what else?) Response #2-3a: In northern Taiwan, the tectonic activity of the region is in extensional regionregime, thus dominated by normal faulting in the study area nowadays. The Jinshan fault (JSF), and Shanchiao fault (SCF, also known as the Jinshan Fault through with normal faulting mechanism), both of the faulting was were being identified longtime ago in Taipei Basin area (Southwest to the study area) and in northeastern offshore Taiwan (northeast of the study area). And recently these two faults were identified and to have linked together as only by only one normal fault in the Tatun Volcano region (around and across the study area ) by the authors. (tThe result was published in the Central Geological Survey CGS project report written in Chinese, and the paper for international journal is now in preparation). On the other hand, there are many landslides within the study area and in the Tatun Volcano region, as shown in the newly added figure Fig. 5. Compare Comparing the size, distribution and classification, etc. the two largest landslides (XSL and CSL) were thus chosen as the target for this study. We have newly revised and improved the paragraph in the manuscript.

b) I have also reservation about the idea that normal fault activity has the effect of cancel the landslide signature (third diagram in the final scheme). I think quite the opposite; fault activity sustain relief formation, maintaining the condition for landslide development. The authors conclude somethings similar at lines 27-29 page 12, but their statement conflict with the idea illustrated in the scheme. Response #2-3b: In northern Taiwan, the tectonic activity of the region is in extensional regionregime. The Jinshan normal faulting cause resulted in the formation of Taipei basin by over one thousand meter throw of the fault separation. The normal faulting is has been very well documented recently, e.g. Teng et al., (2001); Shyu et al., (2005); C.T. Chen et al., (2007, 2010); Huang et al., (2007); and K.C. Chen et al., (2010). And this normal faulting caused the formation of the Taipei basin and may also cause also the continuous eruption of the Tatun Volcano. The evidence of normal faulting being has been recently identified in Taipei basin area and northeastern offshore Taiwan. And recently identified and linked together as only one normal fault in Tatun Volcano region

C8

(near and around the study area) by the authors. And two original normal faults are considered to be linked together as a long stretched normal fault that may provide significant earthquake faulting. Finally, the total length of the Jinshan normal fault is over more than as long as 130 Km long. In conclusions, we thus interpret that the normal faulting leads has led to the formation of the slope daylight, as well as the volcano subsidence in southern the south of the study area. This process may likely leads to the formation of the landslide. As Because the normal faulting activated continuously, the sliding mass may be transporting continuously to the Jinshan Delta. The fig original Fig. 13 (now modified as Fig. 14) demonstrates ideally (of course not fully adapted the current topography) the general geomorphologic evolution ideally, and so as to explain the wear off of the landslide deposits, especially in XSL.

c) The authors never explicitly address time scales of the considered landslides and fault, as well as the probable mismatch in timescale of the landsliding and faulting. Response #2-3c: The normal faulting started since from 400 Ka, and activated continuously ever since. The age of the Tatun volcano is smaller than 200 Ka. So the relative age of the landslide is been triggered sincemost probably after the normal faulting and the formation of the Tatun Volcano, say farwhich is later than 200 Ka. On the other hand, the CSL and XSL preserve different degrees of landslide geomorphologic components, showing the sitesa combination of multiple landslide events. Furthermore, part of the fault branches is identified on the lower slope within the sliding area, express prompting the faulting behavior truncates and enhances the erosion process. In conclusion, based on many aspects, thus the authors thus interpret propose one model to access highlight the possiblethe landslide evolution that will be useful for further testing.

Referee #2-4. Finally, I have reservation about the general organization of the paper. The chapter Introduction is a blend (sometime confused) of general issues about landslide identification and characterization. I suggest to restructure the text, developing a sharper motivation with some clearer objectives. Also, quote the pertinent literature

C9

addressing the mapping and analysis of large landslides. Pertinent local literature help understanding the state of the art at local scale. The authors are not clear enough at this point. For instance at line 25 page 2 the authors acknowledge that the two landslides were already recognized. So why the authors define the two landslides as "obscure" if they were already recognized? I think additional information should be provided, and a comparison of previous and new results should be done. Similarly, the manuscript lacks of references to international literature addressing mapping and analysis of large landslide in active regions. Pertinent international literature help defining the framework of the study and it should be quoted along the paper (see for instance Bucci et al. 2016, ESPL, 41, 711-720; Scheingross et al. 2013, Geological Society of America Bulletin, 125, 473-489; Bucci et al. 2013, Physics and Chemistry of the Earth, 63, 12-24; Strecker M.R. and Marret R. 1999, Geology, 27, 307-310) The chapter geological background (lines 14-23 page 3) is confused: it is hard to follow and to understand the polygenic history of the faults of the area. The chapter contain information negligible for the aim of the paper. At the same time, the chapter lack of potentially useful information about the age and deformation rate of active structures, seismicity, landslide events. Finally, lines 3-11 page 4 belong to method, not to geological background. The chapters 3 and 4 mix up methods, results and discussion, which is also included in the following chapter: Discussion. This writing setting makes reading hard to follow and to understand. Please change the text of the manuscript including the following chapters: Methods (include here technical issues regarding UAS imagery, digital topography (1 x 1 m LiDAR DEM), how you define landslides, what do you map using conventional approach (i.e. stereoscopic aerial photo-interpretation), what new using UAS imagery and LiDAR DEM (would be good to see in map the differences), how you estimate the landslide dimension, how you carried out the morphological reconstruction); Results (includes the new data and maps); and then Discussion (what can we learn from the new data and what is the meaning also comparing to other works) and Conclusions (take home messages in short). The chapters Discussion and Conclusion focus on the evolution of the two landslides, stressing the role of tectonics. However,

C10

the paper do not contain any new information/analysis/result related to tectonics. The evolution scheme drawn by the authors remain poorly constrained also by the lacks of geological evidences supporting the supposed coincidence of the slip surfaces and the present day topography. I suggest to reconsider in depth (or to drop) the part of the analysis related to the volume calculation of the two landslides, because it simply raises too many questions.

Response #2-4: We have reconstructed divided the above paragraph by into 4 questions (a to d), and response responded accordingly:

a) I suggest to restructure the text, developing a sharper motivation with some clearer objectives. Also, quote the pertinent literature addressing the mapping and analysis of large landslides. Pertinent local literature help understanding the state of the art at local scale. The authors are not clear enough at this point. For instance at line 25 page 2 the authors acknowledge that the two landslides were already recognized. So why the authors define the two landslides as "obscure" if they were already recognized? I think additional information should be provided, and a comparison of previous and new results should be done. Similarly, the manuscript lacks of references to international literature addressing mapping and analysis of large landslide in active regions. Pertinent international literature help defining the framework of the study and it should be quoted along the paper. Response #2-4a: Some of the p Pertinent literatures were are now added into the manuscript. The two landslides were already recognized from geomorphologic 40 m DTM by Prof. C. T. Lee of the National Central University (from only personal communication). However, due to the lake lack of existed/available datasets and no districtwithout distinct features, the landslides was were not been analyzed deeply in depth till this study. From climatologic point of view, the annual rainfall is more than 2500 mm in this area, thus a vast portion of the study area is covered by vegetation. Dense forest thus partially conceals morphological features and has prevented detailed geomorphic studies in the past. On the other hand, the heavy rainfall also increases enhances the surface processes, e.g., incision and erosion. As a con-

C11

sequence, the erosion effect also obscures the landslide features in addition. To all of the points, weWe have newly improved the documentation in the manuscript based on the abovementioned points.

b) The chapter geological background (lines 14-23 page 3) is confused: it is hard to follow and to understand the polygenic history of the faults of the area. The chapter contain information negligible for the aim of the paper. At the same time, the chapter lack of potentially useful information about the age and deformation rate of active structures, seismicity, landslide events. Response #2-4b: To discuss the landslide evolution, especially for an old landslide, the geologic and regional tectonics must be included, even for the factor of surface process, e.g. climate etc. So theThe polygenic history of the study area must be taken into account. In the study area, so we consider many factors, including, : lithology, normal fault, climate, vegetation, erosion and human agriculture activity etc., so asin order to access the landslide geomorphologic evolution. Regarding to the slip rate of Jinshan normal faulting, it is shown between 8.2-1.8 mm/yr subsiding rate in at different sites and in time intervals (, e.g., Rau et al., 2006; Huang et al., 2007; Chen et al., 2010). This high slip rate creates the Taipei Basin, and may significantly affects importantly the landslide evolution as well. But unfortunately, these slip rate studiesy is were focused only on the Taipei Basin, and not on the study area. The manuscript is now newly reinforced and improved to clarify the tectonic factor and the interaction.

c) This writing setting makes reading hard to follow and to understand. Please change the text of the manuscript including the following chapters: Methods; Results; and then Discussion and Conclusions. Response #2-4c: We have newly improved the manuscript according to the comment.

d) The chapters Discussion and Conclusion focus on the evolution of the two landslides, stressing the role of tectonics. However, the paper do not contain any new information/analysis/result related to tectonics. The evolution scheme drawn by the authors remain poorly constrained also by the lacks of geological evidences support-

C12

ing the supposed coincidence of the slip surfaces and the present day topography. Response #2-4d: One geological map (Fig. 5) have has been added to demonstrate the geological background of study area, and to better linked the relationship between the regional tectonics and landslide geology and evolution. The manuscript has been improved accordingly.

Please also note the supplement to this comment:

<https://www.nat-hazards-earth-syst-sci-discuss.net/nhess-2017-227/nhess-2017-227-AC2-supplement.pdf>

---

Interactive comment on Nat. Hazards Earth Syst. Sci. Discuss., <https://doi.org/10.5194/nhess-2017-227>, 2017.