



Modeling anthropogenic and natural fire ignitions in an inner-alpine valley

Giorgio Vacchiano¹, Cristiano Foderi², Roberta Berretti¹, Enrico Marchi², Renzo Motta¹

¹Università degli Studi di Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari, Grugliasco TO, 10095, Italy

5 ²Università degli Studi di Firenze, Dipartimento di Gestione dei Sistemi Agrari, Alimentari e Forestali, Firenze, 50145, Italy

Correspondence to: Giorgio Vacchiano (gvacchiano@gmail.com)

Abstract. Modelling and assessing the factors that drive forest fire ignitions is critical for fire prevention and sustainable ecosystem management. In southern Europe, the anthropogenic component of wildland fire ignitions is especially relevant. In the Alps, however, the role of fire as a component of disturbance regimes in forest and grassland ecosystems is poorly known. The aim of this work is to model the probability of fire ignition for an alpine region in Italy using a regional wildfire archive (1995-2009) and MaxEnt modeling. We analyzed separately: i) winter forest fires; ii) winter fires on grasslands and fallow land; iii) summer fires. Predictors were related to morphology, climate, and land use; distance from infrastructures, number of farms, and number of grazing animals were used as proxies for the anthropogenic component; collinearity among predictors was reduced by a Principal Component Analysis. 30% of ignitions occurred in agricultural areas, 24% in forests. Ignitions peaked in the late winter–early spring. Negligence from agro-silvicultural activities was the main cause of ignition (64%); lightning accounted for 9% of causes across the study timeframe, but increased from 6% to 10% between the first and second period of analysis. Models for all groups of fire had a high goodness-of-fit (AUC 0.90-0.95). Temperature was proportional to the probability of ignition, and precipitation inverse-proportional. Proximity from infrastructures had an effect only on winter fires, while the density of grazing animals had a remarkably different on summer (positive correlation) and winter (negative) fires. Implications are discussed regarding climate change, fire regime changes, and silvicultural prevention. Such spatially-explicit approach allows to carry out spatially targeted fire management strategies, and may assist in developing better fire management plans.

25

Keywords: Disturbance regime, Forest fire, MaxEnt, risk model, PCA, ignition



1 Introduction

Wildland fires drive the dynamics of vegetation in many parts of the world (Thonicke et al., 2001; Bond and Keeley, 2005), shaping landscape patterns and influencing the provision of productive and regulatory ecosystem services (Cannon, 2001; Conedera et al., 2003). Fire is part of the natural dynamics of ecosystems, but it may become a nuisance or a hazard for human settlements and activities. In this context, knowing where wildland fire hazard is higher is a fundamental support to distribute and make prevention actions more efficient. Spatially explicit fire databases are used routinely for hazard zoning (Zumbrunnen et al., 2011; Vacchiano et al., 2016). However, in a context of changing anthropogenic and natural drivers of fire occurrence – such as the increase of mean and extreme temperatures, altered precipitation patterns (Beniston, 2006), land use change (abandonment and/or higher tourist and urban pressures) in the wildland-urban interface (WUI) (Moreira et al., 2011) – fire hazard maps and or danger indices based on past conditions may not be able to forecast the future influence of environmental and social drivers onto fire ignition. These processes could imply higher risk even in areas where fires were not traditionally considered a frequent and/or severe disturbance and risk agent, such as in the Alps (Bebi et al., 2017). In such context, modeling the factors that drive ignitions (Ganteaume et al., 2013) can help to choose the most appropriate fire prevention actions even under changing environmental and social conditions (Chuvienco et al., 2014).

Considerable research has been carried out to quantify the influence of natural (climate, vegetation, topography, and landscape connectivity) and anthropogenic drivers of wildland fire ignitions (Cardille and Ventura, 2001; Yang et al., 2007; Martínez et al., 2009; Renard et al., 2012), but the relative importance of each factor is still debated. The prevailing paradigm at the planetary scale is that climate (Carcaillet et al., 2001; Whitlock et al., 2003), fire weather (Schoennagel et al., 2004), or fuel (Krawchuk et al., 2006) are the most important factors. Moreover, forest management and human actions may strongly influence the spatio-temporal pattern of fire ignitions (DeWilde and Chapin, 2006), especially in highly developed regions such as southern Europe, where most of the wildland fires are human-induced. Here, several studies have shown the significance of the distance of forest fires to roads, settlements and infrastructure, or specific land uses or even its abandonment as predisposition for fire ignition (Vega-Garcia et al., 1995; Goldammer, 2003; Catry et al., 2009; Martinez et al., 2009).

In the Alps, however, the role of fire as a component of disturbance regimes in forest and grassland ecosystems is still poorly known. Historically, some ecosystems such as forests dominated by Scots pine or European larch might have evolved under medium- to high- frequency fire regimes (Vazquez Moris et al., 2017). However, the frequency and severity of fires has been heavily influenced by human activities, such as widespread pastoral fires, and a century-long strict fire suppression policy. Such lack of knowledge is particularly upsetting in an area where wildland-urban interfaces are expanding under the pressure of tourism, and where fire danger may already be experiencing phase transitions induced by climate change (e.g., +26 % increase in total area burned by fires larger than 10 ha in 1981–2000 relative to 1961–1980, or the unprecedented occurrence of summer stand-replacing fires) (Vacchiano et al., 2014; Vacchiano and Motta, 2015).



Since both sustainable ecosystem management and effective fire prevention must rely on understanding the characteristics and effects of fire regimes, it is important to improve the understanding of spatio-temporal patterns of natural and anthropogenic fire in the Alpine region, the most important drivers and agents of change, and their potential effects on fire severity and danger. In this paper, we fit spatially-explicit multivariate models for fire ignition in Valle d'Aosta, a dry, inner-
5 alpine region of the western Italian Alps. In doing so, we set out to answer to the following research questions: 1) What are the current fire regimes in the study area? 2) What are the most important environmental drivers of fire ignition in areas with different fire regimes (e.g., summer crown fires in forests vs. winter surface fires in grasslands and fallow lands)? 3) How to solve the “curse of dimensionality” when trying to fit fire ignition models while exploring a high-dimensional space of many potentially collinear predictors (Bar Massada et al., 2013)? 4) How sensitive are current fire ignitions to climate vs
10 anthropogenic drivers in the study area?

Using a 15-year database of temporally- and spatially-explicit fire ignitions allowed us to answer to such questions without being hindered by current fire suppression, which may bias fire behavior data when trying to reconstruct fire regimes from observed burned areas and wildfire frequencies. The results of our analysis can be used by land managers to inform fire prevention actions, and by ecologists to parameterize models of fire ignition for the simulation of future landscape dynamics
15 as a function of natural disturbances.

2. Methods

2.2 Study area

Our study area was Aosta Valley region in northwest Italy (Figure 1). The region covers 3262 km² and is shaped by an east–west main valley with several north–south lateral arms. More than 90% of the land has an elevation of at least 1000 m a.s.l.,
20 and 60% is above 2000 m. Mean annual temperature in Aosta (45°26' N, 7°11' E, 583 m) is 10.9 °C (years 1961–1990; Tetrarca et al., 1999). Climate is warm-summer continental (Dfb) according to the Köppen classification (Peel et al., 2007); mean temperatures of July and January may differ by as much as 22°C. Annual precipitation in Aosta is much lower than in other central Alpine valleys (494 mm, years 1961–1990; Biancotti et al., 1998), with a period of water deficit extending from June to September, and winter precipitation in the form of snow.

25 Forests cover 98,439 ha (Gasparini and Tabacchi, 2011), a third of the region's area. The dominant forest types are conifers—mixed larch (*Larix decidua* Mill.) and Swiss stone pine (*Pinus cembra* L.) (45% of total forest cover), Norway spruce (*Picea abies* [L.] Karst.) (19%), and Scots pine (*Pinus sylvestris* L.) (13%). Broadleaf forest types such as chestnut (*Castanea sativa* Mill.), downy oak (*Quercus pubescens*) (4% each), and early-seral secondary woodlands (12%) are common at lower elevations. Other forest cover types in the region include silver fir (*Abies alba* Mill.) and beech (*Fagus sylvatica* L.) (1% each), Austrian pine (*Pinus nigra* J. F. Arnold) (0.5%), and riparian forests (0.5%) (Gasparini and
30 Tabacchi, 2011). The maximum elevation of the tree line is about 2300 m a.s.l. (Viglietti et al., 2010).



Wildfires occur mainly in late winter and early spring and are 95% anthropogenic in origin (Vacchiano et al., 2015). Surface fires usually start at the bottom of the valley and spread upward, often transitioning into crown fires in conifers due to the low moisture content of live foliage during the dormant season. Most wildfires are small (average size =5.8 ha); however, 70% of the burned area is due to sporadic but relatively large wildfires (up to 400 ha; Vacchiano et al., 2016). This fire regime is common to several inner Alpine valleys (Moser et al., 2010; Zumbrennen et al., 2011) and reflects the influence of strict fire suppression policies adopted in the last decades throughout the Alpine region (Pezzatti et al., 2013; Valese et al., 2014).

2.2 Data analysis

We georeferenced all ignition points (N= 1118, excluding re-ignitions from previously active fires) from a regional forest fire archive for the years 1995–2009, maintained by Regional Forest Service staff (Figure 2). Information available for each point included the date and time of the ignition, duration, cause of ignition, and main vegetation type. Since the study area was covered by widely different vegetation types, with presumably heterogeneous fire regimes, after preliminary scrutiny of the frequency distribution of ignition months and vegetation cover types we grouped fire ignitions in three main categories, indicative of three broadly distinct fire regimes:

- i) 246 summer ignitions (May to September);
- ii) 316 winter ignitions (October to April) in forested areas
- iii) 556 winter ignitions in meadows and fallow lands

For each group, we analyzed the effect of topography, climate, and land use (vegetation). We opted to study the relationship between fire ignition and mean climate, as opposed to fire weather, because this allowed us to assess the long-term relative fire danger in the study area during an “average” fire season. Climate predictors (mean annual temperature BIO1; maximum temperature of the warmest month BIO5; minimum temperature of the coldest month BIO6; yearly temperature range BIO7; average temperature of the wettest quarter BIO8; average temperature of the driest quarter BIO9; average temperature of the warmest quarter BIO10; average temperature of the coldest quarter BIO11; yearly precipitation sum BIO12; precipitation sum of the wettest month BIO13; precipitation sum of the driest month BIO14; precipitation sum of the wettest quarter BIO16; precipitation sum of the driest quarter BIO17; precipitation sum of the warmest quarter BIO18; precipitation sum of the coldest quarter BIO19), with a spatial resolution of 1 km, were obtained from the WorldClim dataset (Hijmans et al., 2005) and resampled at a 30 m resolution by linear interpolation. We used a digital elevation model (DEM) with a resolution of 10m to calculate elevation slope, southwesteriness as $\cos(\text{aspect} - 225^\circ)$ as a proxy for evapotranspiration (Franklin and Tolonen, 2000), and Heat Load Index (McCune and Keon, 2002) as a proxy of solar radiation. Vegetation was described using Corine Land Cover for year 2006 level 3, reclassified in 5 types at 30 m spatial resolution: 1) urban areas; 2) water bodies; 3) pastures and grazing land; 4) agricultural areas; 5) forest areas.



Human activities were factored in using three predictors: the distance of the ignition point from a main road or building (DIST_INFR), as mapped by a regional vector cartography; the number of grazing domestic animals (N_GRAZ_ANIMALS) and the number of dairy and meat enterprises with grazing animals (N_FARMS). The latter two variables were extracted for each municipality from the most recent national agricultural survey (ISTAT 2010) and scaled on
5 the area of pasture land in each municipality to derive unit-area values (D_GRAZ_ANIMALS and D_FARMS).

Potential issues of collinearity between predictors were addressed by running a Principal Component Analysis (PCA) – based clustering (Booth, 1994) of all independent variables for the ignition subset. From each principal component, we selected only those variables that were correlated to the component by a loading higher than 0.32, i.e., explaining at least 10% of the variance of that component (Booth, 1994); in case more than one variable satisfied this rule, we selected only the
10 one with the highest loading. In order to accommodate non-linear effects, each ignition group was modeled as a function of the selected drivers by using MaxEnt v. 3.3.3.k (Phillips et al., 2006), i.e., a maximum entropy algorithm that produces a spatially-explicit probability of ignition, under the null hypothesis of uniform spatial distribution. Being based on presence-only data, MaxEnt maximizes the information available, without the need to draw assumptions about unknown absence or pseudo-absence points (Jaynes, 1990). The fitting algorithm in MaxEnt finds the best set of predictors after attempting to
15 carry out several mathematical transformations, i.e., linear, product, power, hinge, threshold, and categorical. The regularization coefficient was set to 1.5.

For each subset, a model was calibrated on 70% of observation and validated on the remaining 30%; data splitting was carried out by 5 bootstrap samples. Model performance was assessed by the area under the Receiver Operating Characteristic curve (AUC) (Fielding and Bell, 1997), ranging from 0 to 1, with 0.5 indicating the absence of correlation between
20 predictors and observation, and 0.8 commonly considered the minimum threshold for a good fit (Manel et al., 2001). Finally, we measured the effect size of each individual predictor by a leave one out procedure assisted by jackknife repeated replication (JRR) (Quenouille, 1949). Using JRR, we recalculated AUC of models that excluded one predictor at a time, and ranked predictors by their Percent Contribution (%C), i.e., the cumulative relative increase in gain of the model produced by the MaxEnt algorithm while modifying the coefficient of that predictor, and the Percent Permutation Importance (PPI), i.e.,
25 the normalized drop in AUC resulting from randomly permuting the values of that predictor among the training points (Phillips et al., 2006). Contrary to %C, PPI does not depend on the particular path followed by the MaxEnt algorithm in fitting the model.

3. Results

Of the 1118 ignitions analyzed for year 1995-2009, most (30.0%) occurred in agricultural areas, 23.7% in forests, 20.5% in
30 permanent meadows, and 10.6% in pastures and fallow lands. The average yearly ignition density in each of these vegetation classes was 0.18, 0.16, 0.03 and 0.01 fires km⁻¹, respectively. The frequency of ignitions peaked in the late winter – early spring (February to April), with a secondary peak in the summer (June to September) (Figure 3).



Most ignitions could not be assigned a definite cause (Figure 4). However, among known causes, negligence from agricultural and forest activities was the main one (64% of all ignitions with a known cause). Arson accounted for only 7% of all ignitions with a known cause, and lightning for 9%. The time window analyzed is too short, and the share of unknown causes too high, to make a robust analysis of temporal trends; however, among known causes, lightning-induced fires showed the highest increase between the first and second half of the study period (relative frequency of lightning fires was 6% and 10% in 1995-2001 and 2002-2009, respectively), while arson was almost unchanged (5% and 7%) and negligence-related factors decreased (81% and 75%). The relative incidence of summer fires increased with altitude (up to about 40% of all fires above 1500 m) (Figure 5).

Variable selection by PCA loadings was carried out the analysis of the first 13 principal components (PC), which explained 99% of the cumulative proportion of variance. In all three subsets, climatic/topographic and anthropogenic variables were clearly orthogonal to each other in the most important principal components. DIST_INFR and D_GRAZ_ANIMALS were selected (loading >0.32) in all three datasets, as were elevation, slope, land cover, and at least one between heatload index and BIO1 and BIO10 were always collinear to other climatic predictors and were discarded; however, in all subsets, at least one temperature- and one precipitation-related variable were always selected (Table 1).

All models had a high goodness-of-fit, with AUC ranging between 0.90 (summer fires) to 0.94 (winter agro-pastoral fires) and 0.95 (winter forest fires) (mean of five bootstrap iterations). The most important predictors (IPP>10%) were: BIO5 and BIO13 for summer fires; BIO14, elevation, BIO9, and DIST_INFR for winter forest fires; and elevation and DIST_INFR for winter agro-pastoral fires (Table 2).

The effect of climatic predictors conformed to expectations, with temperature variables being monotonically proportional to the probability of ignition, and precipitation variables inverse-proportional. Elevation effects were consistent with temperature ones. The most important anthropogenic driver, i.e., distance from infrastructures, had a large effect only on winter fires, where it contributed to a high probability of ignition at distances lower than about 1000 m (Figure 6); however, its relative importance was never higher than 28%. Variables related to the density of grazing animals were not particularly informative (IPP <2%), but their effect on ignition probability was remarkably different between summer (direct correlation) and winter fires (inverse correlation) (Figure 7).

The spatially-explicit description of ignition probability showed a pattern driven primarily by temperature and human presence, with valley bottoms and areas closed to developed settlements having the highest ignition probabilities, especially in the summer (Figure 8).

4. Discussion

4.1 Natural drivers of fire ignition

The sources of wildland fire ignition can be natural (lightning) or anthropogenic, either accidental or voluntary. In the study area, fires exhibited a marked distinction between climate-driven summer fires and anthropogenic winter fires. In this study,



lighting represented on average 33% of all summer ignitions among those with a known cause, and 38% between June and August (comparable to other Alpine regions, e.g., 40% in Austria: Müller et al., 2012). Such high rate is indicative of the fact that summer fires in this study area can still be considered a partially “natural” disturbance regime, except for the fact that suppression policies are still quite aggressive, and prevent a more complete restoration of fire as an agent potentially useful for regeneration and ecosystem functioning (Vazquez Moris et al., 2017). In turn, in the study area the frequency of lightning-induced fires is highly correlated with the summer Standardized Precipitation and Evaporation Index in Aosta (Castagneri et al., 2015) ($R = 0.55$ for all months between May and September, and 0.69 for the month of July). Two consequences of such correlation are an increase of lightning-induced fire from the first to the second half of the study period (+130% after vs. before year 2001), and the extraordinary number of lightning-induced fires in the summer 2003 (which accounted for one-third of all lightning ignitions in the period). Both can be an indicator of ongoing climate change, with year 2003 showing extreme drought that could become the new norm in the middle or end of this century (Schar and Jendritzky, 2004). The consequences of ongoing climate change on fire regimes can be pervasive, not only for an expected increase in frequency and intensity of fires in a warmer world (Kulakowski et al., 2017; Seidl et al., 2017), but also because of a likely shift in fire regime, from a winter-dominated to an all-year-round seasonality.

Lightning-induced forest fires are ignited when lightning reaches the forest floor, and they can smoulder in the organic soil layer until this is dry enough to carry a surface fire (Wotton et al., 2003; Pezzatti et al., 2009). In turn, the properties of fuel depend on forest type, weather conditions, and vegetation structure (Schoennagel et al., 2004). Wall-to-wall information on forest biomass, structure, and flammability (e.g., Corona et al., 2014) was not available for the study area; even without such factors, fire prediction models had a very high accuracy. The effect of weather on ignition confirmed our expectations and previous literature (e.g., Vilar et al., 2010; Naraynaraj and Wimberly, 2012; Nunes, 2012), with summer temperature and spring precipitations driving summer ignitions, and temperature and precipitation of the driest quarter (i.e., often winter), together with southwesternness (i.e., stands located on drier, sunnier slopes), driving winter fires. The main driver of fire density distribution in southern Europe is precipitation, both off-season (which affects vegetation growth and fuel accumulation) and during the fire season, which limits fire ignition and spread (Oliveira et al., 2012).

Concerning topography, we found a negative effect of elevation on fires. This is likely a result of the correlations between increasing elevation and decreasing human activity density (Xu et al., 2006), and increasing elevation and more positive evapotranspiration balance (Padilla and Vega-Garcia, 2011). Moreover, when altitude increases, fuels are less abundant due to the abundance of non-vegetated areas (rocks, sparse vegetation, snow).

4.2 Anthropogenic fires

In the study area, winter fires were mainly driven by proximity to human infrastructure, as opposed to summer ones. Winter (October to April) ignitions followed an anthropogenic cause in 99.6% of cases. Human-induced ignitions mostly depend on the moisture of fine fuels, e.g., cured needles or leaves and small woody residues on the forest floor (Wotton et al., 2003).



Fuel structure and continuity then control the eventual spread and intensity of fire in the post-ignition phase (e.g., Ruffault and Mouillot, 2017).

Human presence usually directly influences fire density and burned area where fire regimes are anthropogenic (Bar Massada et al., 2013), while human-induced landscape fragmentation has a reverse effect on fire where regimes are mostly natural
5 (Parisien et al., 2004). In the Alps, the density of railroads, forest roads, and trails, together with agricultural and forestry developments, contribute significantly to fire danger (Arndt et al., 2013), due to accidental or negligent fires (Catty et al., 2009; Martinez et al., 2009; Vilar et al., 2010; Narayananaraj and Wimberly, 2012; Oliveira et al., 2012).

Reineking et al. (2010) assessed the importance of weather, forest type, and human actions on forest fire ignitions in a fire-prone area of Switzerland over a 37-year period. Drought was the most important driver of lightning-induced fires, with a
10 negligible effect of distance to human infrastructures. Anthropogenic fires, in contrast, were significantly associated to both distance to infrastructure and weather. Also, anthropogenic fires occurred across a wider range of weather conditions relative to naturally ignited ones, since human activities may provide enough energy to ignite a fire even under sub-optimal fuel moisture conditions.

From the point of view of land cover, areas with higher agricultural population were more prone to fire. In the region, fire is
15 one of the preferred tools to eliminate harvest and agricultural residues, clean field margins, and eliminate excess litter in orchards (Leone et al., 2009). Conversely, pastures have the lowest fire density, probably due to the effectiveness of grazing animals in decreasing fuel load (Bovio and Marchi, 2010).

4.3 Analytical approach

The relationship between fire ignitions and environmental drivers at the regional scale has been historically analyzed using
20 many different statistical models. Due to the need to accommodate non-linearities, most works used logistic regression based on negative binomial or Poisson error distributions (e.g., Martinez et al., 2009; Chuvieco et al., 2009; Prasad et al., 2008). Other authors have proposed regression trees (Amatulli et al., 2006), neural networks (Vega-García et al., 2007), Bayesian probability techniques (Romero-Calcerrada et al., 2008), or Generalized Additive Models (Vilar et al., 2010). Compared to
25 such algorithms, most of which are parametric, the machine learning approach adopted by MaxEnt has been performing equally, if not better (Bar Massada et al., 2013). Being based on presence-only data, MaxEnt avoids the problem of dealing with false, unreliable, or pseudo-absences (Jiménez-Valverde et al., 2008).

The other characteristic of this study is the selection of predictors before fitting the final model. Collinearity between spatial predictors is common, especially when spatially-calibrated climate grids are used (Vacchiano and Motta, 2015), which can lead to unstable or unreliable model fit. Even if MaxEnt is much less sensitive to collinearity (Kuemmerle et al., 2010), its
30 results would be less clearly interpretable in the presence of strongly collinear predictors. The use of PCA and PC loading thresholds to select the most important predictors allowed a reduction in collinearity of predictors and a less ambiguous interpretation of model results (Cruz-Cárdenas et al., 2014).



5. Conclusions

This study showed that fire regime in an alpine region has distinct patterns depending on the ecosystem (forest or grassland/cropland) and season involved (winter or summer). Anthropogenic drivers, mainly from negligence-related reasons, are the main causes of ignition, but the incidence of fires starting from lightning has been recently increasing. From a management point of view, the spatially-explicit approach we describe here allows to carry out spatially targeted fire management strategies, and may be integrated into future fire management plans at regional and local scale. Spatially-explicit hazard assessments can support managers in carrying out appropriate preventive and pre-suppression activities, to locate helicopter water points, to plan fuel management interventions, as well as to parameterise fire behaviour and landscape dynamics models and simulate different fire scenarios with different fire-fighting tactics. Furthermore, by combining the fire danger with the vulnerability to fire (i.e., the ecological and economic fire impact), a cell-by-cell assessment the resulting fire risk can be carried out. As a consequence, in less vulnerable areas the strict fire suppression approach may be reconsidered, and fires allowed to burn a certain share of the land to restore historical disturbance regimes and improve the functionality of those forest ecosystems that had co-evolved with fire (e.g., regeneration of light-demanding species).

From the point of view of the drivers of fire ignition, the present study highlights the importance of urban areas and roads as potential sources of ignition, and of policies that regulate development in the urban-wildland interface, and potentially hazardous human activities close to hotspot fire locations and during periods of increased weather risk. Under the current scenarios of climate change and more frequent drought and lightning, silvicultural prevention should be carried out more frequently to reduce the load and continuity of forest fuels for an effective reduction of wildfire risk.

20 Author contributions

GV co-wrote the paper and created the figures, CF conceived and co-wrote the paper and carried out data analysis, RB digitized fire ignition input data, EM and RM contributed to the discussion.

Acknowledgements

25 We acknowledge Giancarlo Cesti and the forest fire service of Corpo Forestale Regione Autonoma Valle d'Aosta for data provision.

References

- Amatulli, G., Rodrigues, M. J., Trombetti, M. and Lovreglio, R.: Assessing long-term fire risk at local scale by means of decision tree technique, *J. Geophys. Res.*, 111, G4, doi:10.1029/2005JG000133, 2006.
- 30 Arndt, N., Vacik, H., Koch, V., Arpaci, A. and Gossow, H.: Modeling human-caused forest fire ignition for assessing forest fire danger in Austria, *iForest*, 6, 315-325, doi:10.3832/ifor0936-006, 2013.



- Bar Massada, A., Syphard, A. D., Stewart, S. I. and Radeloff, V. C.: Wildfire ignition-distribution modelling: a comparative study in the Huron-Manistee National Forest, Michigan, USA, *Int. J. Wildl. Fire*, 22(2), 174–183, doi:10.1071/WF11178, 2013.
- Bebi, P., Seidl, R., Motta, R., Fuhr, M., Firm, D., Krumm, F., Conedera, M., Ginzler, C., Wohlgemuth, T. and Kulakowski, D.: Changes of forest cover and disturbance regimes in the mountain forests of the Alps, *For. Ecol. Manage.*, 388, 43–56, doi:10.1016/j.foreco.2016.10.028, 2017.
- Beniston, M.: Mountain weather and climate: A general overview and a focus on climatic change in the Alps, *Hydrobiologia*, 562, 3–16, doi:10.1007/s10750-005-1802-0, 2006.
- Biancotti, A., Bellardone, G., Bovo, S., Cagnazzi, B., Giacomelli, L. and Marchisio, C.: Distribuzione regionale di piogge e temperature, Regione Piemonte, Torino, 1998.
- Bond, W. J. and Keeley, J. E.: Fire as a global “herbivore”: the ecology and evolution of flammable ecosystems, *TRENDS Ecol. Evol.*, 20(7), 387–394, doi:10.1016/j.tree.2005.04.025, 2005.
- Booth, G. D., Niccolucci, M. J., and Schuster, E. G.: Identifying proxy sets in multiple linear regression - an aid to better coefficient interpretation, USDA Forest Service, Intermountain Research Station, Ogden UT, 1994.
- Bovio, G. and Marchi, E.: Selvicoltura: incendi, pascolo, *It. J. For. Mount. Env.*, 65, 113–119, 2010.
- Cannon, S. H.: Debris-flow generation from recently burned watersheds, *Environ. Eng. Geosci.*, 7(4), 321–341, 2001.
- Carcaillet, C., Bergeron, Y., Richard, P. J. H., Frechette, B., Gauthier, S. and Prairie, Y. T.: Change of fire frequency in the eastern Canadian boreal forests during the Holocene: does vegetation composition or climate trigger the fire regime?, *J. Ecol.*, 89(6), 930–946, doi:10.1111/j.1365-2745.2001.00614.x, 2001.
- Cardille, J. A. and Ventura, S. J.: Occurrence of wildfire in the northern Great Lakes Region: Effects of land cover and land ownership assessed at multiple scales, *Int. J. Wildl. Fire*, 10(2), 145–154, doi:10.1071/WF01010, 2001.
- Castagneri, D., Bottero, A., Motta, R. and Vacchiano, G.: Repeated spring precipitation shortage alters individual growth patterns in Scots pine forests in the Western Alps, *Trees*, 29(6), 1699–1712, doi:10.1007/s00468-015-1250-z, 2015.
- Catry, F. X., Rego, F. C., Bacao, F. and Moreira, F.: Modeling and mapping wildfire ignition risk in Portugal, *Int. J. Wildl. Fire*, 18(8), 921–931, doi:10.1071/WF07123, 2009.
- Chuvieco, E., Gonzalez, I., Verdu, F., Aguado, I. and Yebra, M.: Prediction of fire occurrence from live fuel moisture content measurements in a Mediterranean ecosystem, *Int. J. Wildl. Fire*, 18(4), 430–441, doi:10.1071/WF08020, 2009.



- Chuvieco, E., Martinez, S., Victoria Roman, M., Hantson, S. and Lucrecia Pettinari, M.: Integration of ecological and socio-economic factors to assess global vulnerability to wildfire, *Glob. Ecol. Biogeogr.*, 23(2), 245–258, doi:10.1111/geb.12095, 2014.
- Conedera, M., Peter, L., Marxer, P., Forster, F., Rickenmann, D. and Re, L.: Consequences of forest fires on the hydrogeological response of mountain catchments: A case study of the Riale Buffaga, Ticino, Switzerland, *Earth Surf. Process. Landforms*, 28(2), 117–129, doi:10.1002/esp.425, 2003.
- Corona, P., Ferrari, B., Cartisano, R. and Barbati, A.: Calibration assessment of forest flammability potential in Italy, *iForest*, 7, 300–305, doi:10.3832/ifor1123-007, 2014.
- Cruz-Cardenas, G., Lopez-Mata, L., Luis Villasenor, J. and Ortiz, E.: Potential species distribution modeling and the use of principal component analysis as predictor variables, *Rev. Mex. Biodivers.*, 85(1), 189–199, doi:10.7550/rmb.36723, 2014.
- DeWilde, L. and Chapin III, F. S.: Human impacts on the fire regime of interior Alaska: Interactions among fuels, ignition sources, and fire suppression, *Ecosystems*, 9(8), 1342–1353, doi:10.1007/s10021-006-0095-0, 2006.
- Dong, X., Shao, G., Limin, D., Zhanqing, H., Lei, T. and Hui, W.: Mapping forest fire risk zones with spatial data and principal component analysis, *Sci. China Ser. E-Tech. Sci.*, 49(1), 140–149, doi:10.1007/s11434-006-8115-1, 2006.
- Fielding, A. H. and Bell, J. F.: A review of methods for the assessment of prediction errors in conservation presence/absence models, *Environ. Conserv.*, 24(1), 38–49, doi:10.1017/S0376892997000088, 1997.
- Franklin, S. B. and Tolonen, M.: Temporally-explicit models of fire and forest, *Plant Ecol.*, 146(2), 145–168, doi:10.1023/A:1009891610140, 2000.
- Ganteaume, A., Camia, A., Jappiot, M., San-Miguel-Ayanz, J., Long-Fournel, M. and Lampin, C.: A review of the main driving factors of forest fire ignition over Europe, *Environ. Manage.*, 51(3), 651–662, doi:10.1007/s00267-012-9961-z, 2013.
- Gasparini, P. and Tabacchi, G.: L’Inventario Nazionale delle Foreste e dei serbatoi forestali di Carbonio INFC 2005 - Secondo inventario forestale nazionale italiano - Metodi e risultati, Edagricole, Bologna, 2011.
- Goldammer, J. G.: Towards international cooperation in managing forest fire disasters in the Mediterranean region, in *Security and environment in the Mediterranean: conceptualising security and environmental conflicts*, edited by H. G. Brauch, P. H. Liotta, A. Marquina, P. F. Rogers and M. E. Selim, pp. 907–915, Springer, Berlin, 2003.
- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. and Jarvis, A.: Very high resolution interpolated climate surfaces for global land areas, *Int. J. Climatol.*, 25(15), 1965–1978, doi:10.1002/joc.1276, 2005
- Jaynes, E. T.: Probability-theory as logic, in *Maximum entropy and bayesian methods*, edited by P. H. Fougere, pp. 1–16, Kluwer, Dordrecht, 1990.



- Jiménez-Valverde, A., Lobo, J. M. and Hortal, J.: Not as good as they seem: The importance of concepts in species distribution modelling, *Divers. Distrib.*, 14(6):885–890, 2008.
- Krawchuk, M. A., Cumming, S. G., Flannigan, M. D. and Wein, R. W.: Biotic and abiotic regulation of lightning fire initiation in the mixedwood boreal forest, *Ecology*, 87(2), 458–468, doi:10.1890/05-1021, 2006.
- 5 Krivtsov, V., Vigy, O., Legg, C., Curt, T., Rigolot, E., Lecomte, I., Jappiot, M., Lampin-Maillet, C., Fernandes, P. and Pezzatti, G. B.: Fuel modelling in terrestrial ecosystems: An overview in the context of the development of an object-orientated database for wild fire analysis, *Ecol. Modell.*, 220(21), 2915–2926, doi:10.1016/j.ecolmodel.2009.08.019, 2009.
- Kuemmerle, T., Perzanowski, K., Chaskovskyy, O., Ostapowicz, K., Halada, L., Bashta, A.-T., Kruhlov, I., Hostert, P., Waller, D. M. and Radeloff, V. C.: European bison habitat in the Carpathian mountains, *Biol. Conserv.*, 143(4), 908–916, doi:10.1016/j.biocon.2009.12.038, 2010.
- 10 Kulakowski, D., Seidl, R., Holeksa, J., Kuuluvainen, T., Nagel, T. A., Panayotov, M., Svoboda, M., Thorn, S., Vacchiano, G., Whitlock, C., Wohlgemuth, T. and Bebi, P.: A walk on the wild side: Disturbance dynamics and the conservation and management of European mountain forest ecosystems, *For. Ecol. Manage.*, 388, 120–131, doi:10.1016/j.foreco.2016.07.037, 2017.
- 15 Leone, V., Lovreglio, R., Pilar Martin, M., Martinez, J. and Vilar, L.: Human factors of fire occurrence in the Mediterranean, in *Earth observation of wildland fires in Mediterranean ecosystems*, edited by E. Chuvieco, pp. 149-170, Springer, New York, 2009.
- Manel, S., Williams, H. C. and Ormerod, S. J.: Evaluating presence-absence models in ecology: the need to account for prevalence, *J. Appl. Ecol.*, 38(5), 921–931, doi:10.1046/j.1365-2664.2001.00647.x, 2001.
- 20 Martinez, J., Vega-Garcia, C. and Chuvieco, E.: Human-caused wildfire risk rating for prevention planning in Spain, *J. Environ. Manage.*, 90(2), 1241–1252, doi:10.1016/j.jenvman.2008.07.005, 2009.
- McCune, B. and Keon, D.: Equations for potential annual direct incident radiation and heat load, *J. Veg. Sci.*, 13(4), 603–606, doi: 10.1111/j.1654-1103.2002.tb02087.x, 2002.
- Moreira, F., Viedma, O., Arianoutsou, M., Curt, T., Koutsias, N., Rigolot, E., Barbati, A., Corona, P., Vaz, P., Xanthopoulos, G., Mouillot, F. and Bilgili, E.: Landscape - wildfire interactions in southern Europe: Implications for landscape management, *J. Environ. Manage.*, 92(10), 2389–2402, doi:10.1016/j.jenvman.2011.06.028, 2011.
- Moser, B., Temperli, C., Schneiter, G. and Wohlgemuth, T.: Potential shift in tree species composition after interaction of fire and drought in the Central Alps, *Eur. J. For. Res.*, 129(4), 625–633, doi:10.1007/s10342-010-0363-6, 2010.
- Müller, M. M., Vacik, H., Diendorfer, G., Arpaci, A., Formayer, H. and Gossow, H.: Analysis of lightning-induced forest fires in Austria, *Theor. Appl. Clim.*, 111, 183-193, 2012.
- 30



- Narayananaraj, G. and Wimberly, M. C.: Influences of forest roads on the spatial patterns of human- and lightning-caused wildfire ignitions, *Appl. Geogr.*, 32(2), 878–888, doi:10.1016/j.apgeog.2011.09.004, 2012.
- Nunes, A. N.: Regional variability and driving forces behind forest fires in Portugal an overview of the last three decades (1980–2009), *Appl. Geogr.*, 34, 576–586, doi:10.1016/j.apgeog.2012.03.002, 2012.
- 5 Oliveira, S., Oehler, F., San-Miguel-Ayanz, J., Camia, A. and Pereira, J. M. C.: Modeling spatial patterns of fire occurrence in Mediterranean Europe using multiple regression and random forest, *For. Ecol. Manage.*, 275, 117–129, doi:10.1016/j.foreco.2012.03.003, 2012.
- Padilla, M. and Vega-Garcia, C.: On the comparative importance of fire danger rating indices and their integration with spatial and temporal variables for predicting daily human-caused fire occurrences in Spain, *Int. J. Wildl. Fire*, 20(1), 46–58, doi:10.1071/WF09139, 2011.
- 10 Parisien, M. A., Hirsch, K. G., Lavoie, S. G., Todd, J. B. and Kafka, V. G.: Saskatchewan fire regime analysis. Canadian Forest Service, Northern Forestry Centre, Edmonton AB, 2004.
- Peel, M. C., Finlayson, B. L. and McMahon, T. A.: Updated world map of the Koppen-Geiger climate classification, *Hydrol. Earth Syst. Sci.*, 11(5), 1633–1644, 2007.
- 15 Pezzatti, G. B., Zumbrunnen, T., Buergi, M., Ambrosetti, P. and Conedera, M.: Fire regime shifts as a consequence of fire policy and socio-economic development: An analysis based on the change point approach, *For. Policy Econ.*, 29(SI), 7–18, doi:10.1016/j.forpol.2011.07.002, 2013.
- Phillips, S. J., Anderson, R. P. and Schapire, R. E.: Maximum entropy modeling of species geographic distributions, *Ecol. Modell.*, 190(3–4), 231–259, doi:10.1016/j.ecolmodel.2005.03.026, 2006.
- 20 Prasad, V. K., Badarinath, K. S. and Eaturu, A.: Biophysical and anthropogenic controls of forest fires in the Deccan Plateau, India, *J. Environ. Manage.*, 86(1), 1–13, doi:10.1016/j.jenvman.2006.11.017, 2008.
- Quenouille, M. H.: Problems in plane sampling, *Ann. Math. Stat.*, 20, 355–375, 1949.
- Reineking, B., Weibel, P., Conedera, M. and Bugmann, H.: Environmental determinants of lightning- v. human-induced forest fire ignitions differ in a temperate mountain region of Switzerland, *Int. J. Wildl. Fire*, 19(5), 541–557, doi:10.1071/WF08206, 2010.
- 25 Renard, Q., Pelissier, R., Ramesh, B. R. and Kodandapani, N.: Environmental susceptibility model for predicting forest fire occurrence in the Western Ghats of India, *Int. J. Wildl. Fire*, 21(4), 368–379, doi:10.1071/WF10109, 2012.



- Romero-Calcerrada, R., Novillo, C. J., Millington, J. D. A. and Gomez-Jimenez, I.: GIS analysis of spatial patterns of human-caused wildfire ignition risk in the SW of Madrid (Central Spain), *Landscape Ecol.*, 23(3), 341–354, doi:10.1007/s10980-008-9190-2, 2008.
- Ruffault, J. and Mouillot, F.: Contribution of human and biophysical factors to the spatial distribution of forest fire ignitions and large wildfires in a French Mediterranean region, *Int. J. Wildl. Fire*, 26(6), 498–508, doi:10.1071/WF16181, 2017.
- Schar, C. and Jendritzky, G.: Climate change: Hot news from summer 2003, *Nature*, 432(7017), 559–560, doi:10.1038/432559a, 2004.
- Schoennagel, T., Veblen, T. T. and Romme, W. H.: The interaction of fire, fuels, and climate across rocky mountain forests, *Bioscience*, 54(7), 661–676, doi:10.1641/0006-3568(2004)054[0661:TIOFFA]2.0.CO;2, 2004.
- Seidl, R., Thom, D., Kautz, M., Martin-Benito, D., Peltoniemi, M., Vacchiano, G., Wild, J., Ascoli, D., Petr, M., Honkaniemi, J., Lexer, M. J., Trotsiuk, V., Mairota, P., Svoboda, M., Fabrika, M., Nagel, T. A. and Reyser, C. P. O.: Forest disturbances under climate change, *Nat. Clim. Chang.*, 7(6), 395–402, doi:10.1038/NCLIMATE3303, 2017.
- Tetarca, S., Spinelli, F., Cogliani, E. and Mancini, M.: *Profilo climatico dell'Italia*, ENEA, Roma, 1999.
- Thonicke, K., Venevsky, S., Sitch, S. and Cramer, W.: The role of fire disturbance for global vegetation dynamics: coupling fire into a Dynamic Global Vegetation Model, *Glob. Ecol. Biogeogr.*, 10(6), 661–677, doi:10.1046/j.1466-822x.2001.00175.x, 2001.
- Vacchiano, G. and Motta, R.: An improved species distribution model for Scots pine and downy oak under future climate change in the NW Italian Alps, *Ann. For. Sci.*, 72(3), doi:10.1007/s13595-014-0439-4, 2015.
- Vacchiano, G., Berretti, R., Mondino, E. B., Meloni, F. and Motta, R.: Assessing the effect of disturbances on the functionality of direct protection forests, *Mt. Res. Dev.*, 36(1), 41–55, doi:10.1659/MRD-JOURNAL-D-15-00075.1, 2016.
- Vacchiano, G., Lonati, M., Berretti, R. and Motta, R.: Drivers of *Pinus sylvestris* L. regeneration following small, high-severity fire in a dry, inner-alpine valley, *Plant Biosyst.*, 149(2), 354–363, doi:10.1080/11263504.2013.819821, 2015.
- Vacchiano, G., Stanchi, S., Marinari, G., Ascoli, D., Zanini, E. and Motta, R.: Fire severity, residuals and soil legacies affect regeneration of Scots pine in the Southern Alps, *Sci. Total Environ.*, 472, 778–788, doi:10.1016/j.scitotenv.2013.11.101, 2014.
- Valese, E., Conedera, M., Held, A. C. and Ascoli, D.: Fire, humans and landscape in the European Alpine region during the Holocene, *Anthropocene*, 6, 63–74, doi:10.1016/j.ancene.2014.06.006, 2014.
- Vazquez Moris, J., Vacchiano, G., Enri, S. R., Lonati, M., Motta, R. and Ascoli, D.: Resilience of European larch (*Larix decidua* Mill.) forests to wildfires in the western Alps, *New For.*, 48(5), 663–683, doi:10.1007/s11056-017-9591-7, 2017.



- Vega Garcia, C., Woodard, P. M., Titus, S. J., Adamowicz, W. L. And Lee, B.: A logit model for predicting the daily occurrence of human caused forest-fires, *Int. J. Wildl. Fire*, 5(2), 101–111, doi:10.1071/WF9950101, 1995.
- Vega-García, C.: Propuesta metodológica para la predicción diaria de incendios forestales, in *Wildfire 2007 - IV International Wildfire Conference*, Seville, Spain, 13-17 May, 2007.
- 5 Viglietti, D., Letey, S., Motta, R., Maggioni, M. and Freppaz, M.: Snow avalanche release in forest ecosystems: A case study in the Aosta Valley Region (NW-Italy), *Cold Reg. Sci. Technol.*, 64(2, SI), 167–173, doi:10.1016/j.coldregions.2010.08.007, 2010.
- Vilar, L., Woolford, D. G., Martell, D. L. and Pilar Martin, M.: A model for predicting human-caused wildfire occurrence in the region of Madrid, Spain, *Int. J. Wildl. Fire*, 19(3), 325–337, doi:10.1071/WF09030, 2010.
- 10 Whitlock, C., Shafer, S. L. and Marlon, J.: The role of climate and vegetation change in shaping past and future fire regimes in the northwestern US and the implications for ecosystem management, *For. Ecol. Manage.*, 178(1–2), 5–21, doi:10.1016/S0378-1127(03)00051-3, 2003.
- Wotton, B. M., Martell, D. L. and Logan, K. A.: Climate change and people-caused forest fire occurrence in Ontario, *Clim. Change*, 60(3), 275–295, doi:10.1023/A:1026075919710, 2003.
- 15 Yang, J., He, H. S., Shifley, S. R. and Gustafson, E. J.: Spatial patterns of modern period human-caused fire occurrence in the Missouri Ozark Highlands, *For. Sci.*, 53(1), 1–15, 2007.
- Zumbrunnen, T., Pezzatti, G. B., Menendez, P., Bugmann, H., Buergi, M. and Conedera, M.: Weather and human impacts on forest fires: 100 years of fire history in two climatic regions of Switzerland, *For. Ecol. Manage.*, 261(12, SI), 2188–2199, doi:10.1016/j.foreco.2010.10.009, 2011.



Tables

Table 1 – Principal components analysis and selected predictors

PC	Retained variable	Loading
1	-	-
	BIO8	0.335
2	BIO1	0.359
	BIO10	0.447
	BIO5	0.469
3	D_GRAZ_ANIM	0.691
	D_FARMS	0.691
4	Heatload	-0.637
	Southwesternness	-0.690
	Slope	-0.582
5	DIST_INFR	-0.770
6	-	-
7	-	-
8	-	-
9	-	-
10	-	-
	BIO17	-0.362
11	BIO11	-0.409
	BIO19	-0.448
	BIO6	-0.475
12	-	-
	BIO17	-0.327
13	BIO18	-0.725



Table 2 – Predictor effect size (PC: percent contribution; PPI: percent permutation importance) in the models for ignition of summer, winter forest, and winter agropastoral fires

Predictor	Summer fires		Winter forest fires		Winter agro-pastoral fires	
	%C	IPP	%C	IPP	%C	IPP
Bio16	29.41	0.76	0.91	0.22	5.87	1.19
Elevation	19.99	2.78	42.99	20.06	35.51	41.20
Bio5	11.29	38.59	8.42	5.69	1.94	2.76
Bio6	9.31	1.61	11.86	0.44	3.37	0.23
Bio12	7.55	0.01	11.85	0.39	9.66	1.52
Bio14	4.48	1.12	5.41	30.56	10.54	6.35
Southwesternness	3.47	2.05	4.96	6.97	2.28	4.21
Bio18	3.23	9.67	4.49	3.36	2.46	0.95
Land cover	2.96	3.18	1.27	3.7	4.54	4.17
DIST_INFR	2.77	2.58	4.86	10.59	20.6	28.13
Bio9	2.14	1.91	0.99	12.59	1.10	5.95
D_GRAZ_ANIM	1.80	1.91	0.44	1.12	1.21	1.55
Bio13	1.59	33.85	1.56	5.29	0.93	1.79



Figure captions

Figure 1: Map of the study area

5 Figure 2: Fire ignitions in years 1995-2009 (red dots) and Corine Land Cover (year 1990) vegetation types in the study area (yellow: agricultural areas, light green: permanent meadows, dark green: forests, blue: pastures and shrubs, grey: unvegetated, white: urban)

Figure 3: Frequency of fire ignitions by month and vegetation cover type

Figure 4: Fire ignitions by cause in the study period

Figure 5: Absolute and relative frequency of summer (black) vs. winter (grey) fire ignitions in the study area by elevation class

10 Figure 6: Effect of distance from infrastructures on ignition probability in winter forest (a) and agro-pastoral fires (b).


Figure 7: Effect of density of grazing domestic animals on ignition probability in summer fires (a), winter forest (b) and winter agro-pastoral fires (c).

Figure 8: Ignition probability in the study area for summer fires (a), winter forest (b) and winter agro-pastoral fires (c).



Figures

Figure 1



5



Figure 2






Figure 3






Figure 4






Figure 5








Figure 6

a



b




5




Figure 7

a



b



5



c








Figure 8

a





b





c

