

1 Dear Editor and the reviewers,
2 We do appreciate your constructive, thoughtful, careful, and helpful comments and suggestions.
3 After careful discussions, calculations, and analyses, we finished the preparation of responses to
4 you. There are totally four parts: “Response to Reviewer 1”, “Response to Reviewer 2”,
5 “Response to Short Comments” and a clean version of the manuscript “Risk Zoning of Typhoon
6 Disasters in Zhejiang Province, China”.
7 If there are any new comments or suggestions, please let us know.
8 Best Regards
9 Yi Lu and the coauthors

10
11

12 **Response to Reviewer 1**

13 1. In the "2. Study Area" section, it is better to give a background introduction of typhoon disasters
14 in the study area; otherwise, it is not unstoodable why you use Zhejiang Province as a case.

15 Reply: We do appreciate the helpful suggestion.

16 Mainly according to the suggestion, some modifications have been done. The modifications
17 include:

18 (1) The structure of the paper has been changed. Considering that Section 2 is so thin and
19 unbalanced to other sections, “2 Study Area” is merged into “3 Data and Methods” and then the
20 Section numbers hereafter are changed.

21 (2) In the last paragraph of “Introduction”, a sentence “In this study, Zhejiang province, which is
22 frequently affected by the strongest landfall typhoons (Ren et al., 2008) and experiences most
23 serious typhoon disasters (Liu and Gu, 2002) in the mainland of China, is selected as the study
24 area.” has been added.

25 The following two new references have been added:

26 Liu, T. J. and Gu, J. Q.: A statistical analysis of typhoon disasters in Zhejiang province, Journal of
27 Catastrophology, 17 (4): 64-71, 2002. (in Chinese)

28 Ren, F. M., Wang, X. L., Chen, L. S., and Wang, Y. m.: Tropical cyclones landfalling on mainland
29 China, Hainan and Taiwan and their correlations, Acta Meteorologica Sinica, 66 (2): 224-235,
30 2008. (in Chinese)

31

32 2. For the meteorological data, how many stations are there in Zhejiang Province? The authors said
33 2419 stations provided by NMIC; it is not clear if they are distributed throughout China or only in
34 Zhejiang province. It needs to be clarified.

35 Reply: Thanks so much for the comment. In this paper, the OSAT method need to identify typhoon
36 wind and precipitation from wide range than Zhejiang province, so 2419 stations of precipitation
37 data and 2479 stations of wind data over the mainland of China are used, which all contained 71
38 stations corresponding to counties in Zhejiang province.

39 We add following sentences at the end of L80 in original section 3.1.1.

40 “In addition, the OSAT method need to identify typhoon wind and precipitation from a wider
41 range than Zhejiang province (please see detail in section 2.2.1), so 2419 stations of precipitation
42 data and 2479 stations of wind data over the mainland of China are used, which all contained 71
43 stations corresponding to counties in Zhejiang province.”

44

45 3. Canonical Correlation Analysis is the main tool for this study, which is however not introduced
46 to readers at all. It is necessary to give an introduction of this method and how it is applied in this
47 study.

48 Reply: Thanks for your suggestion. According to the suggestion, we have added introduction and
49 application of Canonical Correlation Analysis (CCA) in this paper.

50 We add following sentences at the end of L102 in original section 3.2.2.

51 “In statistics, canonical correlation analysis (CCA) is a way of inferring information from
52 cross-covariance matrices. If we have two vectors $X = (X_1, \dots, X_n)$ and $Y = (Y_1, \dots, Y_m)$ of
53 random variables, and there are correlations among the variables, then CCA can find linear
54 combinations of the X_i and Y_j which have maximum correlation with each other (Hardoon et al.,
55 2014). The method was first introduced by Hotelling in 1936 (Hotelling, 1936). The main point of
56 CCA is to separate linear combination of new variables from the two sets of variables. In this case,
57 the correlation coefficient between new variables reaches the maximum. In this paper, we chose
58 factors causing typhoon disasters as a set of variables, and typhoon disaster as another. Under the
59 maximum canonical correlation coefficient, the linear combination coefficients (typical variable
60 coefficients) of factors causing typhoon disasters can be used as weight coefficients of this group
61 of variables. Then we can determine the impact of factors causing typhoon disasters.”

62 The added references are as follows:

63 Hardoon, D. R., Szedmak, S., and Shawetaylor, J.: Canonical Correlation Analysis: An Overview
64 with Application to Learning Methods, *Neural Computation*, 16(12):2639-2664, 2014.

65 Hotelling, H.: Relations between two sets of variates, *Biometrika*, 28(3/4), 321-377, 1936.

66

67 4. In Section 3.2.4, it is necessary to introduce how the so-called SoVI is used to calculate the
68 population vulnerability index.

69 Reply: Thanks very much for the suggestion. According to the suggestion, we have added
70 introduction and application of SoVI in this paper.

71 To introduce how we used the SoVI method, we add following definition of SoVI at the beginning
72 of L109 in original section 3.2.4.

73 “County-level socioeconomic and demographic data are used to construct an index of social
74 vulnerability to environmental hazards named the Social Vulnerability Index (SoVI). Principal
75 Component Analysis (PCA) is the primary statistical technique for constructing the SoVI. The
76 PCA method captures multi-dimensionality by transforming the raw dataset to a new set of
77 independent variables. Then a few components can represent the dimensional data, and underlying
78 factors can be identified easily. These new factors are placed in an additive model to compute a
79 summary score—SoVI (Cutter et al., 2003).”

80 The added reference is as follows:

81 Cutter, S. L., Boruff, B. J., and Shirley, W. L.: Social Vulnerability to Environmental Hazards,
82 *Social Science Quarterly*, 84(2):242-261, 2003.

83 In addition, we have supplemented the specific calculation of the SoVI method in original section
84 5.2, and the detail can be seen in the response to the question 13 of Reviewer 2.

85

86 5. The data source of typhoon disaster losses?

87 Reply: Typhoon disaster losses used in this paper are obtained from the National Climate Center

88 who collected these disaster data from local meteorological departments. The data source can be
89 seen in lines 82-83 of original section 3.1.2.

90
91 6. How is "typhoon rainstorm" defined? Its probability is an important issue in this study. How
92 the probability is determined? It is not introduced at all.

93 Reply: Thanks very much for the comment and suggestion.

94 (1) For the first question, according to the suggestion, we have added a definition of "typhoon
95 rainstorm" and "typhoon torrential rainstorm". In original section 4.1, we did some research about
96 risk of typhoon rainstorm. Typhoon rainstorm in this study means daily typhoon precipitation over
97 50mm, and typhoon torrential rainstorm means daily typhoon precipitation over 100mm.

98 (2) For the second question, the probability in this paper is the annual possibility of the occurrence
99 of typhoon rainstorms. The probability denominator is the total number of years, and the
100 numerator is the annual frequency of typhoon precipitation. If a station experiences typhoon
101 precipitation in one year, the numerator increases by one. Then we can obtain probability in each
102 station.

103 We add following definition at the original L133 after "higher risk of typhoon disasters."

104 "Typhoon rainstorm in this study means daily typhoon precipitation over 50mm, and typhoon
105 torrential rainstorm means daily typhoon precipitation over 100mm. "

106 We add following definition at the original L133 before "Based on".

107 "The probability is the annual possibility of the occurrence of typhoon rainstorms. The probability
108 denominator is the total number of years, and the numerator is the annual frequency of typhoon
109 precipitation. If a station experiences typhoon precipitation in one year, the numerator increases by
110 one. "

111
112 7. In Eq(1), it is not clear how population and economic loss are both included. Only for
113 population or economic loss? Why in the form of "Ax+By", not a "multiplying" form?

114 Reply: Thanks for the comment.

115 (1) For the first question, as we explained in question 3, the main point of CCA is to separate
116 linear combination of new variables from the two sets of variables. In this paper, we chose factors
117 causing typhoon disasters as a set of variables, and typhoon disasters as another. Factors causing
118 typhoon disasters (typhoon wind and precipitation) and typhoon disasters (affected population and
119 economic loss) are both contained. As explained in original section 5.1, taking the county as a unit,
120 we select all the typhoons having affected population in Zhejiang province from 2004 to 2012.
121 The total precipitation and daily maximum wind speed during typhoons measured in each county
122 are used to describe the factors causing typhoon disasters. Using CCA, we determine the impact of
123 typhoon precipitation and winds on the population. Similarly, we determine the impact of typhoon
124 precipitation and winds on economic loss. In this method, population and economic loss are both
125 included.

126 (2) For the second question, both typhoon rainstorm and high wind will bring certain disasters.
127 When reaching a certain critical value, they will have a superposition effect. However, the effects
128 of wind and rain on disaster are different. We want to explore the contribution of wind and rain to
129 typhoon disasters, which can be shown by different weighting factors. Therefore, it is necessary to
130 determine their influence through typical correlation analysis, which is a typical variable
131 coefficient. So the form is "Ax+By", not a multiplier. When the typical correlation coefficient

132 passes the significance test, weight discrimination can be made to determine A and B.
133 We add following sentences at the original L175 after “describe the impact.”.
134 “It is necessary to determine their influence through typical correlation analysis, and then typhoon
135 wind and rain effect are superimposed by the weight coefficients. ”
136

137 8. P12 L205, it says, after "performing factor analysis"..... How is the factor analysis performed?
138 In the title of Table 3, principal component analysis is mentioned. How is the principal component
139 analysis performed? Factor analysis is the principal component analysis? If these methods are
140 used, they need to be introduced in the methodology section.

141 Reply: Thanks very much for the comment and suggestion. The factor analysis performed here is
142 Principal Component Analysis (PCA), which is the primary statistical procedure for constructing
143 the SoVI. According to the suggestion, we have added detailed explanations of PCA in original
144 section 3.2.4, which have been answered in question 4.

145 We add following sentences at the beginning of L109 in original section 3.2.4.

146 “County-level socioeconomic and demographic data are used to construct an index of social
147 vulnerability to environmental hazards named the Social Vulnerability Index (SoVI). Principal
148 Component Analysis (PCA) is the primary statistical technique for constructing the SoVI. The
149 PCA method captures multi-dimensionality by transforming the raw dataset to a new set of
150 independent variables. Then a few components can represent the dimensional data, and underlying
151 factors can be identified easily. These new factors are placed in an additive model to compute a
152 summary score—SoVI (Cutter et al., 2003). ”
153

154 9. In Table 4, the derived disaster risk index are divided into 5 grades. How are the thresholds are
155 determined? It is not mentioned.

156 Reply: Thanks for the comment. The classification of typhoon disaster risk index is based on the
157 natural breaks method (Jenks) provided by Arcgis. Then we divide disaster risk index into 5
158 grades, which represent five risk zones for typhoon disasters in Zhejiang province.

159 We add following sentences at the end of original L248.

160 “The classification of typhoon disaster risk index is based on the natural breaks method (Jenks)
161 provided by Arcgis.”
162
163

164 **Response to Reviewer 2**

165 1. The English of the manuscript is need be improved.

166 Thanks for your comment. According to the suggestion, modifications include:

167 (1) The first half of Abstract has been rewritten.

168 (2) “Province” has been changed into “province”.

169 (3) “Comprehensive Risk Index for Typhoon Disasters” has been changed into “Typhoon Disaster
170 Comprehensive Risk Index”.

171 (4) The second half of the last paragraph in Introduction has been rewritten.

172 (5) The structure of the paper has been changed. Considering that Section 2 is so thin and
173 unbalanced to other sections, “2 Study Area” is merged into “3 Data and Methods” and then
174 the Section numbers hereafter are changed.

175 (6) Analyses for the figures have been revised especially these for Figure 7.
176 (7) For grammar, all past tense have been changed into present tense.
177 (8) Other specific modifications can be seen in detail in the text with revision-tracing.
178
179 2. Extend data from 2012 to 2016?
180 Thanks very much for the suggestion.
181 As county-level typhoon disaster data is so limited and it's hard to get new data, we can't extend
182 data from 2012 to 2016.
183
184 3. The background of typhoon disaster over Zhejiang province must be introduced.
185 According to the suggestion, modifications include:
186 (1) In the last paragraph of "Introduction", a sentence "In this study, Zhejiang province, which is
187 frequently affected by the strongest landfall typhoons (Ren et al., 2008) and experiences most
188 serious typhoon disasters (Liu and Gu, 2002) in the mainland of China, is selected as the study
189 area." has been added.
190 (2) The following two new references have been added:
191 Liu, T. J. and Gu, J. Q.: A statistical analysis of typhoon disasters in Zhejiang province, *Journal of*
192 *Catastrophology*, 17 (4): 64-71, 2002. (in Chinese)
193 Ren, F. M., Wang, X. L., Chen, L. S., and Wang, Y. m.: Tropical cyclones landfalling on mainland
194 China, Hainan and Taiwan and their correlations, *Acta Meteorologica Sinica*, 66 (2): 224-235,
195 2008. (in Chinese)
196
197 4. L74-75. Add reference about best track data from CMA.
198 Reply: Thanks for your suggestion. According to the suggestion, we have added references about
199 best track data from CMA, and cite them in original section 3.1.1.
200 New references are added in "References".
201 Eunjeong, C. and Ying, M.: Comparison of three western North Pacific tropical cyclone best track
202 datasets in seasonal context, *Journal of the Meteorological Society of Japan*, 89(3):211-224, 2009.
203 Li, S. H. and Hong, H. P.: Use of historical best track data to estimate typhoon wind hazard at
204 selected sites in China, *Natural Hazards*, 76(2):1395-1414, 2015.
205
206 5. Describe more detail about the OSAT methods.
207 Reply: Thanks very much for the suggestion. According to the suggestion, we have added more
208 detail about the OSAT method.
209 We add following sentences in L96 of original section 3.2.1 before "Lu".
210 "The OSAT method is a numerical technique to separate tropical cyclone induced precipitation
211 from adjacent precipitation areas. Based on the structural analysis of precipitation field, it can be
212 divided into different rain belts. Then, according to the distances between a TC center and these
213 rain belts, typhoon center and each station, typhoon precipitation is distinguished."
214
215 6. L103: In this section, authors describe the methods of standardization. But corresponding
216 variables are unknown.
217 Reply: Thanks for the comment. According to the comment, we have added introduction of
218 corresponding variables.

219 We add following sentences at the end of L107 in original section 3.2.3.
220 “Z-score standardization is used for calculating intensity index of factors causing typhoon
221 disasters. Both typhoon precipitation and typhoon maximum wind speed are standardized by this
222 method. When calculating typhoon disaster comprehensive risk index (R), we use MIN-MAX
223 standardization to standardize the intensity index of the factors causing typhoon disasters (I) and
224 the population vulnerability index (SoVI).”

225

226 7. Introduce the method of calculating probability of typhoon rainstorms.

227 Reply: Thanks very much for the suggestion. According to the suggestion, we have added a
228 definition of “typhoon rainstorm” and “typhoon torrential rainstorm”. In original section 4.1, we
229 did some research about risk of typhoon rainstorm. Typhoon rainstorm in this study means daily
230 typhoon precipitation over 50mm, and typhoon torrential rainstorm means daily typhoon
231 precipitation over 100mm. The probability is the annual possibility of the occurrence of typhoon
232 rainstorms. The probability denominator is the total number of years, and the numerator is the
233 annual frequency of typhoon precipitation. When a station has experienced typhoon precipitation
234 in one year, the numerator increases by one.

235 We add following sentences at the end of original L133 before “Based on”.

236 “Typhoon rainstorm in this study means daily typhoon precipitation over 50mm, and typhoon
237 torrential rainstorm means daily typhoon precipitation over 100mm. The probability is the annual
238 possibility of the occurrence of typhoon rainstorms.”

239

240 8. L148: Do you sure about “over six sites”?

241 Reply: Thanks for your question. We feel sorry for that it is a translation error. This should be
242 referred to Typhoon wind over 6 grade (≥ 10.8 m/s). Figure 6 shows average duration (days) of
243 typhoon winds at each station in Zhejiang province from 1980 to 2014. We have modified original
244 L148.

245 Original L148 is modified as follows.

246 “The average duration (days) of typhoon winds (over 6 grade) is calculated in Zhejiang province
247 (Figure 6).”

248

249 9. Durations of data in different parts are different. Could durations of different kinds of data are
250 uniformed?

251 Reply: Thanks for the comment. In the original paper, we used three durations of data, including
252 typhoon precipitation during 1960 - 2013, typhoon wind during 1980 - 2014, and typhoon disaster
253 data during 2004 - 2012. According to the suggestion, durations of daily precipitation and wind
254 have been uniformed with 1980–2014. Duration of typhoon disaster data remains unchanged.
255 Detailed reasons are as follows.

256 (1) First of all, because of limited access to county-level typhoon disaster data, we have only
257 obtained data during 2004 to 2012 from National Climate Center. So all analyses of intensity
258 index of factors causing typhoon disasters are during 2004 to 2012, which remains unchanged.
259 However, this duration is short for risk analyses of typhoon precipitation and typhoon wind.
260 Therefore, longer time-series data are needed.

261 (2) We feel sorry for that it is a expression mistake to say “The statistics showed a rapid increase
262 in the number of automated wind measurement stations from 1980” in original L77. As Lu et al.

263 (2016) mentioned, considering the homogeneity and continuity of wind data, we use daily wind
264 data during 1980 - 2014 to identify typhoon wind.

265 (3) Considering the consistency between wind and precipitation data, 1980 to 2014 is selected as
266 the period of study. In addition, the OSAT method need to identify typhoon wind and precipitation
267 from wide range rain belts, so 2419 stations of precipitation data and 2479 stations of wind data
268 over the mainland of China are used, which all contained 71 stations corresponding to counties in
269 Zhejiang province.

270 According to the suggestion, modifications include:

271 (1) The introduction of daily precipitation and wind data in original section “3.1.1 Typhoon,
272 Precipitation and Wind Data” are rewrote as follow.

273 “Daily precipitation data for 2479 stations and daily wind data for 2419 stations during the time
274 period 1960 - 2014 over the mainland of China are obtained from National Meteorological
275 Information Center. The maximum wind speed is given as the maximum of 10-minute mean. In
276 this paper, two time periods of precipitation and wind data are used.

277 Because of limited access to county-level typhoon disaster data, we have only obtained data
278 during 2004 to 2012. So when calculating intensity index of factors causing typhoon disasters,
279 time period of typhoon precipitation and typhoon wind are the same as typhoon disasters, which is
280 2004 - 2012.

281 For risk analyses of typhoon precipitation and typhoon wind (please see detail in sections 3.1
282 and 3.2), suppose future probability is the same as historical probability, we then select the period
283 of 1980 – 2014. As Lu et al. (2016) mentioned, considering the homogeneity of wind data, we use
284 the period of 1980 - 2014 for wind analysis. To ensure the consistency between wind and
285 precipitation data, 1980 - 2014 is selected as the period. In addition, the OSAT method need to
286 identify typhoon wind and precipitation from a wider range than Zhejiang province (please see
287 detail in section 2.2.1), so 2419 stations of precipitation data and 2479 stations of wind data over
288 the mainland of China are used, which all contained 71 stations corresponding to counties in
289 Zhejiang province.”

290 (2) The analyses in original section “4.1 Risk of Typhoon Rainstorms” are modified. All time
291 periods in this section have been changed to 1980 - 2014, with corresponding changes of
292 calculations and pictures.

293

294 10. L178: What criterions are used in calculating precipitation or wind?

295 Reply: Thanks for the comment. If a typhoon disaster occurs and there is a corresponding typhoon
296 wind or typhoon precipitation, it will be included in the sample.

297

298 11. L181: The distribution of sample sizes?

299 Reply: Good comment. The total valid disaster records of Zhejiang province from 2004 to 2012
300 are 421. To establish an intensity index of typhoon disaster-causing factors, we carry out CCA
301 analysis. Taking the county as a unit, we select all the typhoons that affected the population, which
302 means all records with an affected population greater than 0. The total precipitation and daily
303 maximum wind speed during affected typhoons measured in each county are used. The total
304 sample size is 322. Then, we do CCA analyses for all the typhoons that caused direct economic
305 losses in the same way, and the total sample size is 404.

306 According to the suggestion, modifications include:

307 (1) We add following sentences at original L179 after “factors causing typhoon disasters”.

308 “ The total sample size is 322. ”.

309 (2) We add following sentences at original L182 before “(Table 1)”.

310 “, and the total sample size is 404.”.

311

312 12. L194: the intensity index is calculated for each typhoons and each stations? And how get the
313 distribution of intensity indices in figure 8?

314 Reply: Thanks for the comment. The intensity index is calculated for each typhoon at each station.

315 Then we average all intensity indices at each station, and we can get the distribution of intensity
316 indices in figure 8.

317

318 13. L199-L238. I am confused by the section 5.2. Each factor or component in factor 1 to factor7
319 in equation 2 or seven components in table 3 is a vector with 29 variables? Please describe more
320 details about factors or components. The signs of vectors from PCA maybe opposite of real
321 meaning. Do you do detail analysis on the relations of variables in a vector and between vectors?
322 The equation 2 is must be evaluated carefully.

323 Reply: Thanks very much for the comments and suggestions.

324 (1) For the first question, component 1 to component 7 are vectors in table 3 with 29 variables.
325 After performing PCA of the 29 variables, 7 components with eigenvalue equal to or greater than
326 1 are extracted.

327 (2) For the second question, 7 components are examined manually as to whether they increase (+)
328 or decrease (–) vulnerability and they are assigned a cardinality on that basis. Then the
329 vulnerability index is produced by summing all the components using equal weighting, following
330 the Chen (2013) approach.

331 To conveniently describe details about 7 components, we rename 29 variables (Table 2). After
332 PCA, we obtain 7 components. The signs and contained variables of 7 components are shown in
333 Table 3. We can see 6 components increase (+) vulnerability and a component decrease (–)
334 vulnerability. For example, the first component, which reflects the income of the population and
335 the employment situation, is positive because the more property there is in an area, the higher the
336 vulnerability to damage. The second component, which reflects education level of the population,
337 is negative because if education level is higher, then the population’s awareness of disaster
338 prevention and reduction is greater and their vulnerability is lower.

339 According to the comments and suggestions, modifications include:

340 (1) We rename the 29 variables in Table 2.

341

Table 2. The 29 variables affecting vulnerability in Zhejiang province.

	variables	Name
	Per capita disposable income of urban residents	
1	(yuan)	UBINCM
2	Percentage of female (%)	QFEMALE
3	Percentage of minority (%)	QMINOR
4	Median age	MEDAGE
	Unemployment rate (calculated - unemployed population / (unemployed + total population)	QUNEMP
5		
6	Population density	POPDEN

7	Percentage of urban population (%)	QUBRES
8	Percentage of non-agricultural household population (%)	QNONAGRI
9	Percentage of households that living in rented houses (%)	QRENT
10	Percentage of employees working in primary industries and mining (%)	QAGREMP
11	Percentage of employees working in secondary industries (%)	QMANFEMP
12	Percentage of employees working in tertiary industries (%)	QSEVEMP
13	Household size (person / household)	PPUNIT
14	Percentage of population with college degree (25 years old and older)	QCOLLEGE
15	Percentage of population with high school degree (20 years old and older)	QHISCH
16	Percentage of illiterate people (15 years old and older)	QILLIT
17	Population growth rate (2000-2010)	POPCH
18	Average number of rooms per household (inter / household)	PHROOM
19	Per capita housing construction area (m ² / person)	PPHAREA
20	Percentage of premises without tap water (%)	QNOPIPWT
21	Percentage of premises without a kitchen (%)	QNOKITCH
22	Percentage of premises without a toilet (%)	QNOTOILET
23	Percentage of premises without a bath (%)	QNOBATH
24	Number of beds per 1000 person in health care institutions	HPBED
25	Number of medical personnel per 1000 resident population	MEDPROF
26	Percentage of people under 5	QPOPUD5
27	Percentage of population over 65 years old	QPOPAB65
28	Population dependency ratio (%)	QDEPEND
29	Percentage of population covered by subsistence allowances (%)	QSUBSIST

342 (2) We add contained variables of 7 components with different signs in Table 3.

343 Table 3. The seven components extracted by PCA.

Components	Contained variables	Name	(Sign)
1	QMANFEMP, UBINCM, QAGREMP, QRENT, POPCH, QDEPEND, QSUBSIST, QPOPAB65, POPDEN, MEDAGE, QNOKITCH, QILLIT,	Employment and poverty	(+)

	PHROOM, PPHAREA		
2	QHISCH, QCOLLEGE, QNONAGRI, QSEVEMP, HPBED, MEDTECH	Education	(-)
3	QNOBATH, QNOTOILET, PPUNIT	Number of dilapidated houses	(+)
4	QILLIT, QDEPEND, QPOPUD5, MEDAGE	Illiteracy and juvenile population	(+)
5	QFEMALE, PHROOM, PPHAREA, QSEVEMP	Household size and ratio of women	(+)
6	QMINOR	Ethnic minority	(+)
7	QUNEMP, QNOPIPWT	Unemployment and housing size	(+)

344

345 (3) To distinguish the 7 components and 29 variables more clearly, we replace all “factor” with
346 “component” in this section.

347

348 14. Could you find data of 29 variables in table 2 in early year? If the difference of population
349 vulnerability between 2010 to early year is analyzed, the manuscript will to be more valuable.

350 Reply: Thanks for your thoughtful suggestion. The population data used in this paper is obtained
351 from the sixth national population census of the Population Census Office of the National Bureau
352 of Statistics of China and the 2010 statistical yearbooks of each city in Zhejiang province
353 published by the cities’ statistical bureaus. There exist many missing and abnormal records in the
354 original data, which take a long time to be processed. This article focuses on typhoon disaster risk
355 zoning in Zhejiang province, so we didn’t discuss the difference of population vulnerability
356 between 2010 to early year. The variation of population vulnerability is an interesting topic.
357 Maybe we can discuss it in future work.

358

359

360

Response to Short Comments

361 1. Please use word “Tropical Cycle” instead of Typhoon.

362 Reply: Thanks for your comment. Considering to be consistent with some published researches
363 (Chen, 2007; Chen et al., 2011; Ding et al., 2002; Niu et al., 2011), we tend to introduce a
364 definition of typhoon rather than change the name. At the beginning of “Introduction”, “Tropical
365 cyclones cause” has been changed into “Typhoon, which means tropical cyclone in this paper,
366 often causes”.

367

368 2. In your paper, you mentioned many times risk. In my knowledge, risk is future probability of
369 hazard events. However, your paper studied the disaster events happened during 2004-2012. In
370 this period, I would not like to mention risk. In your paper, you can only name disaster probability.

371 Reply: We do appreciate the comment.

372 We agree and understand that risk is future probability of hazard events. However, suppose future
373 probability is the same as historical probability for a specific period, we can understand risk by
374 learning from past events. We tend to make a specific statement about this rather to change “risk”
375 into “disaster probability”. According to the comments, modifications include:

376 (1) A sentence “As risk is future probability of hazard events, when suppose future probability is
377 the same as historical probability for a specific period, we can understand risk by learning
378 from past events.” is added and becomes the first sentence of “Abstract”.

379 (2) A sentence “For risk analyses of typhoon precipitation and typhoon wind (please see detail in
380 sections 3.1 and 3.2), suppose future probability is the same as historical probability, we then
381 select the period of 1980 – 2014.” is added as the first sentence of the third paragraph in
382 original section 3.1.1.

383

384 3. Wind and precipitation data. Please clarify wind and precipitation data during TC events or not.

385 Reply: Thanks for your suggestion. As described in original section 3.1.1, the original daily
386 precipitation data and maximum wind speed in this paper are from the National Meteorological
387 Information Center. Then we distinguish typhoon precipitation and winds by using the OSAT
388 method (original section 3.2.1). Therefore, all typhoon wind and rain mentioned later in this paper
389 were during TC events.

390 To clarify wind and precipitation data, last sentence in original section 3.2.1 is rewrote as follows.

391 “With the application of the OSAT method, daily precipitation and wind data over the mainland of
392 China during 1980 to 2014 are used for identifying typhoon precipitation and wind data.”.

393

394 4. The figure should be redrawn as figures have no latitude and longitude.

395 Reply: Thanks for your suggestion. Figures have been modified according to the suggestion.

396

397 5. Latest references should be cited.

398 Reply: Thanks for your suggestion. According to the suggestion, we have added some latest
399 references in “1 Introduction ” and “References”.

400 At the end of original line 44, we add following sentences.

401 “Recently, some research built quantitative assessment in some provinces and carried out
402 preliminary studies on pre-evaluating typhoon disasters (Huang and Wang, 2015; Yin and Li,
403 2017).”

404 At the end of original line 56, we add following sentences.

405 “Xu et al. (2015) comprehensively assessed the impact of typhoons across China using the
406 geographical information system. The future direction of tropical cyclone risk management is
407 quantitative risk models (Chen et al., 2017).”

408 Added references:

409 Chen, W. F., Duan, Y. H., and Lu, Y.: Review on Tropical Cyclone Risk Assessment, Journal of
410 Catastrophology, 32(4), 2017. (in Chinese)

411 Huang, W. K. and Wang, J. J.: Typhoon damage assessment model and analysis in Taiwan, *Natural*
412 *Hazards*, 79(1), 497-510, 2015.

413 Xu, X., Sun, D., and Guo, T.: A systemic analysis of typhoon risk across china, *Natural*
414 *Hazards*, 77(1), 461-477, 2015.

415 Yin, Y. Z. and Li, H. L.: Preliminary study on pre-evaluation method of typhoon disaster in China,
416 *Meteorological Monthly*, 43(6):716-723, 2017. (in Chinese)

417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454

Risk Zoning of Typhoon Disasters in Zhejiang Province, China

Yi Lu¹ (陆逸), Fumin Ren² (任福民), Weijun Zhu³ (朱伟军)

¹ Shanghai Typhoon Institute of China Meteorological Administration, Shanghai 200030, China

² State Key Laboratory of severe weather, Chinese Academy of Meteorological Sciences, Beijing 100081, China

³ Key Laboratory of Meteorological Disaster of Ministry of Education, Nanjing University of Information Science & Technology, Nanjing 210044, China

Abstract ~~As risk is future probability of hazard events, when suppose future probability is the same as historical probability for a specific period, we can understand risk by learning from past events. We analyze the characteristics of typhoon disasters and the factors causing them using~~ Based on precipitation and wind data over the mainland of China during 1980 - 2014, disaster and social data at the county level in Zhejiang ~~Province~~ from 2004 to 2012, a study on risk zoning of typhoon disasters is carried out. Firstly, characteristics of typhoon disasters and factors causing typhoon disasters are analyzed. Secondly, ~~Using canonical correlation analysis, we develop~~ an intensity index for the factors causing typhoon disasters of factors causing typhoon disasters and calculate a population vulnerability index are developed. Thirdly, combining these two indexes, a comprehensive risk index for typhoon disasters is obtained and used to zone areas of risk in Zhejiang Province. Above analyses show that, southeastern Zhejiang ~~Province~~ is the area most affected by typhoon disasters. The annual probability of the occurrence of typhoon rainstorms >50 mm decreases from the southeast coast to inland areas and is at, with a maximum in the boundary region between Fujian and Zhejiang, which has the highest risk of rainstorms. Southeastern Zhejiang and the boundary region between Zhejiang and Fujian ~~province~~ and the Hangzhou Bay area are most frequently affected by typhoon extreme winds and have the highest risk of wind damage. The population of southwestern Zhejiang ~~Province~~ is the most vulnerable to typhoons as a result of the relatively undeveloped economy in this region, the mountainous terrain and the high risk of geological disasters in this region. Vulnerability is lower in the cities and coastal areas due to better disaster prevention and reduction strategies and a more highly educated population. The southeast coastal areas face the highest risk of typhoon disasters, especially in the boundary region between Taizhou and Wenzhou Ningbo cities. Although the inland mountainous

Corresponding author: Dr. Fumin Ren, State Key Laboratory of severe weather (LaSW)/CAMS, Beijing, 100081.
E-mail address: fmren@163.com

30 areas are not directly affected by ~~the~~ typhoons, they are in the medium-risk category for vulnerability.

31 **Keywords:** typhoon disasters, ~~disaster-causing~~ factors causing typhoon disasters, vulnerability,

32 comprehensive risk index, risk zoning

33 **1 Introduction**

34 Typhoon, which means typhoon~~stropical cyclone in this paper, often~~ causes some of the most serious
35 natural disasters in China, with an average annual direct economic loss of about \$9 billion. The arrival
36 of ~~a~~-typhoon is often accompanied by heavy rain, high winds and storm surges, with the main impacts
37 in southern coastal areas of China (Zhang et al., 2009). Zhejiang Province~~province~~ is seriously affected
38 by typhoons—for example, in 2006, the super-typhoon Sang Mei caused 153 deaths in Cangnan county
39 of Wenzhou city, with 11.25 billion yuan of direct economic losses. Therefore it would be of practical
40 significance to develop a system for the risk assessment of typhoon disasters in Zhejiang
41 Province~~province~~.

42 Major risk assessment models include the disaster risk index system of the United Nations
43 Development Program (global scale, focusing on human vulnerability), the European multiple risk
44 assessment (with ~~an~~-emphasis on factors causing disasters and vulnerability) and the American
45 HAZUS-MH hurricane module and disaster risk management system. Vickery et al. (2009) and Fang et
46 al. (2012, 2013) ~~have~~d reviewed the factors causing typhoon disasters. Rain and wind are direct causes
47 of typhoon disasters (~~Mille 1958;~~ Emanuel, 1988~~7~~, 1992~~88~~, 1995; Holland, 1997; Kunreuther and
48 ~~Riehardoth~~, 1998); stronger typhoons produce heavier rain and stronger winds, resulting in a greater
49 number of casualties and higher economic losses. ~~Much~~Many of the researches on the ~~causes of~~
50 ~~typhoon~~factors causing typhoon disasters ~~used~~s a grade index and the probability of occurrence (~~Yang~~
51 ~~et al. 2010;~~ Chen et al., 2011; Su et al., 2008; Ding et al., 2002; Chen, 2007). Recently, some research
52 built quantitative assessment in some provinces and carried out preliminary studies on pre-evaluating
53 typhoon disasters (Huang and Wang, 2015; Yin and Li, 2017).

54 In terms of vulnerability, Pielke et al. (1998, 2008) combined the characteristics of typhoons and
55 socioeconomic factors, suggesting that both the vulnerability of the population and economic factors
56 were important in estimating disaster losses. The vulnerability of a population is a pre-existing
57 condition that influences its ability to face typhoon disasters. Among the most widely used indices is


58 the Social Vulnerability Index (SoVI) (Cutter [et al.](#), 2003; Chen [et al.](#), 2014). Other researches have
59 focused on the vulnerability of buildings, obtaining a fragility curve by combining historical loss with
60 the characteristics of buildings and ~~the typhoons~~ (Hendrick [and Friedman](#), 1966; Howard [et al.](#), 1972;
61 Friedman, 1984; Kafali [and Jain](#), 2009; Pita [et al.](#), 2014). Studies in China have assessed vulnerabilities
62 to typhoon disasters (Yin [et al.](#), 2010; Niu [et al.](#), 2011). ~~An e~~Evaluation indexes for the assessment of
63 disaster losses ~~was~~were established based on the number of deaths, direct economic losses, the area of
64 crops affected and the number of collapsed houses. ~~This~~hese indexes ~~was~~were used to construct
65 different disaster assessment models (Liang [and Fan](#), 1999; Lei [et al.](#), 2009; Wang [et al.](#), 2010). [Xu et al.](#)
66 [\(2015\) comprehensively assessed the impact of typhoons across China using the geographical](#)
67 [information system. The future direction of tropical cyclone risk management is quantitative risk](#)
68 [models \(Chen et al., 2017\).](#)

69 Previous studies have concentrated on semi-quantitative, large-scale research, with less emphasis
70 on quantitative research at ~~the~~ county level based on large amounts of accurate data. In addition, ~~the~~
71 ~~studies~~ have paid more attention to disaster losses and few studies have focused on a comprehensive
72 risk assessment of typhoon disasters coupled with ~~the~~ ~~—~~factors causing typhoon disasters and
73 ~~the~~population vulnerability ~~of populations and infrastructure~~. [In this study, Zhejiang province, which is](#)
74 [frequently affected by the strongest landfall typhoons \(Ren et al., 2008\) and experiences most serious](#)
75 [typhoon disasters \(Liu and Gu, 2002\) in the mainland of China, is selected as the study area. Section 2](#)
76 [introduces the data and methods used in this study. Section 3 provides analyses on typhoon disaster](#)
77 [losses and causing factors. Section 4 presents risk assessment and regionalization of typhoon disasters.](#)
78 [Summary and discussions are given in the final section.](#)~~We analyzed the characteristics of typhoon~~
79 ~~disasters, established a comprehensive risk index for typhoon disasters in Zhejiang Province and~~
80 ~~developed risk zoning for typhoon disasters in this region, which may give some reference for future~~
81 ~~disaster prevention.~~


82 **2 Study Area****2 Data and Methods**

83 This study ~~was~~is carried out in Zhejiang ~~Province~~province (Figure 1) ~~and~~including 11 cities along
84 the Yangtze River Delta. Zhejiang ~~Province~~province is in the eastern part of the East China Sea and
85 south ~~of~~to Fujian ~~Province~~province, which is one of the most economically powerful provinces in
86 China.

87



88



89

Figure 1. Maps of Zhejiang Province, China showing location and major cities.

90 **3 Data and Methods**

91 **3.1 Data**

92 **3.1.1 Typhoon, Precipitation and Wind Data**


93 The typhoon data used in this study were the best-track tropical cyclone datasets from the Shanghai
94 Typhoon Institute for the time period 1960–2014 (Eunjeong and Ying, 2009; Li and Hong, 2015).
95 Daily precipitation data for 2479 stations and daily wind data for 2419 stations during the time period
96 1960–2013/2014 over the mainland of China were obtained from the National Meteorological
97 Information Center. The maximum wind speed is given as the maximum of 10-minute mean. In this
98 paper, two time periods of precipitation and wind data are used.

99 Because of limited access to county-level typhoon disaster data, we have only obtained data
100 during 2004 to 2012. So when calculating intensity index of factors causing typhoon disasters, time
101 period of typhoon precipitation and typhoon wind are the same as typhoon disasters, which is 2004 -
102 2012.


103 For risk analyses of typhoon precipitation and typhoon wind (please see detail in sections 3.1 and
104 3.2), suppose future probability is the same as historical probability, we then select the period of 1980 –
105 2014. As Lu et al. (2016) mentioned, considering the homogeneity of wind data, we use the period of
106 1980 - 2014 for wind analysis. To ensure the consistency between wind and precipitation data, 1980 -
107 2014 is selected as the period. In addition, the OSAT method need to identify typhoon wind and
108 precipitation from a wider range than Zhejiang province (please see detail in section 2.2.1), so 2419
109 stations of precipitation data and 2479 stations of wind data over the mainland of China are used,
110 which all contained 71 stations corresponding to counties in Zhejiang province.

111 **3.2.1.2 Disaster and Social Data**

112 Disaster data for each typhoon that affected Zhejiang ~~Province~~ from 2004 to 2012 ~~were~~
113 obtained from the National Climate Center and the number of records for each county is shown in
114 Figure 2. Of the 11 cities in Zhejiang ~~Province~~, Wenzhou and Taizhou recorded the most
115 typhoon disasters, with a maximum ~~being~~ of 17. Fewer typhoon disasters ~~were~~ recorded in the
116 central and western regions of Zhejiang ~~Province~~, particularly in Changshan and Quzhou,
117 which may be because the strength of typhoons ~~weakened~~ after landfall. The population data ~~for~~
118 2010 ~~were~~ obtained from the sixth national population census ~~of the~~ (Population Census Office of
119 the National Bureau of Statistics of China), and the 2010 statistical yearbooks of each city in Zhejiang
120 ~~Province~~ published by the cities' statistical bureaus. Basic geographical data ~~were~~ obtained
121 from the National Geomatics Center of China.



122



123

124 Figure 2. Number of records of typhoon disasters ~~by county~~ in Zhejiang ~~Province~~ ~~province~~ from 2004
 125 to 2012.

126 **2.2 Methods**

127 **2.2.1 Objective Synoptic Analysis Technique**

128 The widely used objective synoptic analysis technique (OSAT) proposed by Ren et al. (2001, 2007,
 129 2011) ~~wasis~~ used to ~~identify~~ precipitation due to typhoons ~~in this study. The~~ ~~—This method has a high-~~
 130 ~~recognition ability. OSAT method is a numerical technique to separate tropical cyclone induced~~
 131 ~~precipitation from adjacent precipitation areas. Based on the structural analysis of precipitation field, it~~
 132 ~~can be divided into different rain belts. Then, according to the distances between a TC center and these~~
 133 ~~rain belts, typhoon center and each station, typhoon precipitation is distinguished.~~ Lu et al. (2016)
 134 improved the OSAT method and applied it to identify typhoon winds. ~~With the application of-~~ ~~We used~~

135 the OSAT method, daily-to-distinguish-typhoon precipitation and wind data over the mainland of China
136 durings from 2004-1980 to 2014-2012, are used for identifying typhoon precipitation and wind data.

137 **32.32.2 Canonical Correlation Analysis (CCA)**

138 We used the canonical correlation analysis method proposed by Hotelling (1992) to determine the
139 relationship between the affected population, the rate of economic damage, and typhoon precipitation
140 and winds. In statistics, canonical correlation analysis (CCA) is a way of inferring information from
141 cross-covariance matrices. If we have two vectors $X = (X_1, \dots, X_n)$ and $Y = (Y_1, \dots, Y_m)$ of random
142 variables, and there are correlations among the variables, then CCA can find linear combinations of the
143 X_i and Y_j which have maximum correlation with each other (Hardoon et al., 2014). The method was
144 first introduced by Hotelling in 1936 (Hotelling, 1936). The main point of CCA is to separate linear
145 combination of new variables from the two sets of variables. In this case, the correlation coefficient
146 between new variables reaches the maximum. In this paper, we chose factors causing typhoon disasters
147 as a set of variables, and typhoon disaster as another. Under the maximum canonical correlation
148 coefficient, the linear combination coefficients (typical variable coefficients) of factors causing
149 typhoon disasters can be used as weight coefficients of this group of variables. Then we can determine
150 the impact of factors causing typhoon disasters.

151 **32.2.3 Data Standardization**

152 We adopted two methods: Z-score standardization and MIN-MAX standardization. The Z-score
153 standardized method is based on the mean and standard deviation of the raw data. The MIN-MAX
154 standardization is a linear transformation of the original data so that the original value maps the interval
155 [0, 1]. Z-score standardization is used for calculating intensity index of factors causing typhoon
156 disasters. Both typhoon precipitation and typhoon maximum wind speed are standardized by this
157 method. When calculating typhoon disaster comprehensive risk index (R), we use MIN-MAX
158 standardization to standardize the intensity index of the factors causing typhoon disasters (I) and the
159 population vulnerability index (SoVI).


160 **32.2.4 Vulnerability Assessment (SoVI, PCA)**


161 County-level socioeconomic and demographic data are used to construct an index of social
162 vulnerability to environmental hazards named the Social Vulnerability Index (SoVI). Principal

163 Component Analysis (PCA) is the primary statistical technique for constructing the SoVI. The PCA
164 method captures multi-dimensionality by transforming the raw dataset to a new set of independent
165 variables. Then a few components can represent the dimensional data, and underlying factors can be
166 identified easily. These new factors are placed in an additive model to compute a summary
167 score—SoVI (Cutter et al., 2003). Based on the SoVI designed for disaster social vulnerability in
168 America, Chen et al. (2014) collected 29 variables as proxies to build a set of vulnerability indexes for
169 the social and economic environment in China. We used this method to calculate the population
170 vulnerability index for Zhejiang Province.


171 **3.4 Typhoon Disaster Losses and ~~Causation~~ Factors in Zhejiang Province**

172 Based on the distribution of typhoon disaster losses in Zhejiang Province from 2004 to 2012
173 (Figure 3), the affected areas were mainly located in the southeast corner of the province. The
174 centers with the largest affected population (Fig. 3a), the largest area of affected crops (Fig. 3c) and the
175 highest direct economic losses (Fig. 3d) were in Wenzhou and Taizhou cities, although the losses in
176 Ningbo City were also relatively high. Only part of the plain area was affected by serious
177 agricultural disasters; the other losses were far lower than in the southeast of Zhejiang
178 Province. Cangnan in Wenzhou City was the most severely affected, with the highest
179 cumulative death toll (Fig. 3b). The losses in the affected counties were associated with the
180 frequency and intensity of ~~the~~ typhoons. We therefore analyzed the risk of typhoon precipitation and
181 winds in every county in Zhejiang Province to provide a reference dataset for the ~~factors~~
182 responsible for typhoon disaster factors causing typhoon disasters.






184



185




186

187 Figure 3. Distribution of typhoon disaster losses in Zhejiang ~~Province~~ from 2004 to 2012. (a)
 188 Affected population (unit: millions); (b) total number of deaths (unit: person); (c) area of affected crops
 189 (unit: hectares); and (d) direct economic losses (unit: millions yuan).

190 **43.1 Risk of Typhoon Rainstorms**

191 The main hazard of typhoon precipitation is concentrated precipitation, so the average ~~number~~

192 of duration (days) of typhoon precipitation at each sitestation in Zhejiang Province was
 193 counted from 1986 to 2013 (Figure 4). The duration of typhoon rainfall was less in inland areas,
 194 especially in Quzhou City. Persistent precipitation was concentrated in Wenzhou, Taizhou and Ningbo
 195 cities, where there may have been a higher risk of typhoon disasters. Typhoon rainstorm in this study
 196 means daily typhoon precipitation over 50mm, and typhoon torrential rainstorm means daily typhoon
 197 precipitation over 100mm. The probability is the annual possibility of the occurrence of typhoon
 198 rainstorms. The probability denominator is the total number of years, and the numerator is the annual
 199 frequency of typhoon precipitation. If a station experiences typhoon precipitation in one year, the
 200 numerator increases by one. Based on the probability of typhoon rainstorms occurring in each county
 201 in Zhejiang Province (Figure 5), we found that the annual probability of the occurrence of
 202 typhoon rainstorms was highest over the southeast coast of Zhejiang Province from 1986 to
 203 2013, especially in Taizhou City, where the annual probability was 17.83%. The annual
 204 probability of typhoon rainstorms with precipitation >100 mm was lower, but the distribution of
 205 probability was consistent with the rainstorms with lower precipitation. The probability of typhoon
 206 torrential rainstorms decreased rapidly in the western and central regions of Zhejiang
 207 Province, although the range increased. There were three centers of high risk: Taizhou,
 208 Wenzhou and Ningbo cities.



209

210 Figure 4. Average duration (days) withof typhoon precipitation at each sitestation in Zhejiang
 211 Province from 1986 to 2013.





Figure 5. Probability of the occurrence of typhoon rainstorms in Zhejiang Province: (a) rainstorms with >50 mm of precipitation >50 mm; and (b) torrential rainstorms with >100 mm of precipitation >100 mm.

4.3.2 Risk of Typhoon Winds

The average duration (days) of typhoon winds (over 6 grade) is calculated in Zhejiang province (Figure 6). The average duration of typhoon winds was calculated over six sites in Zhejiang Province (Figure 6). The duration of typhoon winds was relatively short in the central and western regions and the typhoon winds were concentrated in the coastal areas of Wenzhou, Taizhou and Ningbo cities. The longest duration of typhoon winds occurs red over the offshore islands.


The main hazard from typhoon winds is manifested in the destructive force of strong winds and. Therefore we calculate the probability of the annual occurrence of typhoon winds at or above grades 6 and 12 at each monitoring station from 1980 to 2014 (Figure 7). Typhoon winds at or above grade 6 mainly occurred along the coastal areas of Zhejiang Province, but were with rare occurrence in the mountainous areas. Meanwhile, the probability of typhoon winds at or above grade 8 is generally

228 0.5~0.9 lower along the coast of Zhejiang Province, and below 0.25 although still much higher than in
 229 the inland mountainous areas interior, with a probability of up to 75% in Hangzhou Bay and over some
 230 islands. Typhoons with winds speeds of at or above grade 10 or 12 were much less likely and were
 231 only seen in the coastal areas and islands, with a rapidly decreasing probability of occurrence from the
 232 coastal areas 71% to the inland mountainous areas 29%. The areas at high risk of typhoon winds were
 233 consistent with those with high typhoon rainfall, i.e. Wenzhou, Taizhou and Ningbo cities. The risk of
 234 typhoon extreme winds associated with typhoons is much higher in coastal areas than inland.




235
 236 Figure 6. Average duration (days) of typhoon winds (over 6 grade) at each station in Zhejiang
 237 Province from 1960 to 2013.

238



239



240

241

242

243

244

245

246

247

Figure 7. Probability of the occurrence of typhoon winds in Zhejiang Province at (a) grade 6 or above, (b) grade 8 or above, (c) grade 10 or above and (d) grade 12 or above.

54 Risk Assessment and Regionalization of Typhoon Disasters ~~in Zhejiang Province~~

248 **54.1 Intensity Index of Factors Causing Typhoon Disasters**

249 The main factors causing typhoon disasters are rainstorms, winds and storm surges. The level and
 250 intensity of a single ~~causative~~ factor cannot fully represent and describe the impact. It is necessary to
 251 determine their influence through typical correlation analysis, and then typhoon wind and rain effect
 252 are superimposed by the weight coefficients. Therefore we established a comprehensive intensity index
 253 that included ~~typhoon precipitation and winds~~~~a number of different factors involved in typhoon~~
 254 ~~disasters~~. Taking the county as a unit, we selected all the typhoons that affected the population of
 255 Zhejiang ~~Province~~province from 2004 to 2012. The total precipitation and daily maximum wind speed
 256 during typhoons measured in each county ~~were~~are used to describe the factors causing typhoon
 257 disasters. The total sample size is 322. Using ~~canonical correlation analysis~~CCA, we determined the
 258 impact of typhoon precipitation and winds on the population. We then ~~carried out~~ canonical
 259 ~~correlation analysis~~CCA for all the typhoons that caused direct economic losses in Zhejiang
 260 ~~Province~~province from 2004 to 2012, and the total sample size is 404 (Table 1). The effect of typhoon
 261 precipitation on both the population and direct economic losses ~~was~~ always greater than that of
 262 typhoon winds. By averaging typical coefficients for both precipitation and wind, ~~weighted~~ coefficients
 263 of 0.85 and 0.65 ~~were~~are obtained within the intensity index for precipitation and winds, respectively.

264 Table 1. Canonical correlation analysis of factors causing typhoon disasters.



Disastes	Canonical correlation coefficient	Canonical variable coefficient	
		Typhoon precipitation	Typhoon wind
Affected population	0.45	0.84	0.651
Direct economic losses	0.477	0.863	0.655

265

266 Based on the weight coefficients in Table 1, an intensity index of factors causing typhoon
 267 disasters ~~was~~ established:

268
$$I = Ax + By \tag{1}$$

269 where I is the intensity index of factors causing typhoon disasters, X is the standard typhoon
 270 precipitation and Y is the maximum wind speed of the typhoon. A and B are the weighted coefficients
 271 for typhoon precipitation and typhoon winds, respectively. Using Equation (1), we ~~calculated~~ average
 272 the intensity indexes of typhoons at each station (Figure 8). Based on the distribution of these average
 273 intensity indexes, ~~we found~~ three high value centers, namely Wenzhou, Taizhou and Ningbo cities,
 274 which are consistent with the results of Chen et al. (2011), can be found.



275
 276
 277 Figure 8. Intensity indexes of factors causing typhoon disasters at each station in Zhejiang

278 Province.

279 **5.4.2 Population Vulnerability Index in Zhejiang Province**

280 Natural disasters are social constructions and the basic causes of losses are the attributes of human
 281 beings and their social system (Jiang 2014). ~~We used t~~ The index system of Chen et al. (2011) is used –
 282 to evaluate the vulnerability of Zhejiang Province. Based on the extracted population

283 information, 29 variables ~~we~~are identified that may affect ~~social~~-vulnerability (Table 2).

284 Table 2. The 29 variables affecting ~~social~~-vulnerability in Zhejiang ~~Province~~province.

	<u>variables</u>	<u>Name</u>
	<u>Per capita disposable income of urban residents</u>	
<u>1</u>	<u>(yuan)</u>	<u>UBINCM</u>
<u>2</u>	<u>Percentage of female (%)</u>	<u>QFEMALE</u>
<u>3</u>	<u>Percentage of minority (%)</u>	<u>QMINOR</u>
<u>4</u>	<u>Median age</u>	<u>MEDAGE</u>
	<u>Unemployment rate (calculated - unemployed</u>	
<u>5</u>	<u>population / (unemployed + total population)</u>	<u>QUNEMP</u>
<u>6</u>	<u>Population density</u>	<u>POPDEN</u>
<u>7</u>	<u>Percentage of urban population (%)</u>	<u>QUBRESD</u>
	<u>Percentage of non-agricultural household</u>	
<u>8</u>	<u>population (%)</u>	<u>QNONAGRI</u>
	<u>Percentage of households that living in rented</u>	
<u>9</u>	<u>houses (%)</u>	<u>QRENT</u>
	<u>Percentage of employees working in primary</u>	
<u>10</u>	<u>industries and mining (%)</u>	<u>QAGREMP</u>
	<u>Percentage of employees working in secondary</u>	
<u>11</u>	<u>industries (%)</u>	<u>QMANFEMP</u>
	<u>Percentage of employees working in tertiary</u>	
<u>12</u>	<u>industries (%)</u>	<u>QSEVEMP</u>
<u>13</u>	<u>Household size (person / household)</u>	<u>PPUNIT</u>
	<u>Percentage of population with college degree (25</u>	
<u>14</u>	<u>years old and older)</u>	<u>QCOLLEGE</u>
	<u>Percentage of population with high school degree</u>	
<u>15</u>	<u>(20 years old and older)</u>	<u>QHISCH</u>
	<u>Percentage of illiterate people (15 years old and</u>	
<u>16</u>	<u>older)</u>	<u>QILLIT</u>
<u>17</u>	<u>Population growth rate (2000-2010)</u>	<u>POPCH</u>
	<u>Average number of rooms per household (inter /</u>	
<u>18</u>	<u>household)</u>	<u>PHROOM</u>
<u>19</u>	<u>Per capita housing construction area (m² / person)</u>	<u>PPHAREA</u>
<u>20</u>	<u>Percentage of premises without tap water (%)</u>	<u>QNOPIPWT</u>
<u>21</u>	<u>Percentage of premises without a kitchen (%)</u>	<u>QNOKITCH</u>
<u>22</u>	<u>Percentage of premises without a toilet (%)</u>	<u>QNOTOILET</u>
<u>23</u>	<u>Percentage of premises without a bath (%)</u>	<u>QNOBATH</u>
	<u>Number of beds per 1000 person in health care</u>	
<u>24</u>	<u>institutions</u>	<u>HPBED</u>
	<u>Number of medical personnel per 1000 resident</u>	
<u>25</u>	<u>population</u>	<u>MEDPROF</u>
<u>26</u>	<u>Percentage of people under 5</u>	<u>QPOPUD5</u>

<u>27</u>	<u>Percentage of population over 65 years old</u>	<u>QPOPAB65</u>
<u>28</u>	<u>Population dependency ratio (%)</u>	<u>QDEPEND</u>
<u>29</u>	<u>Percentage of population covered by subsistence allowances (%)</u>	<u>QSUBSIST</u>

	Variable
1	Per-capita disposable income of urban residents (yuan)
2	Percentage of women
3	Percentage of minority ethnic groups
4	Median age
5	Unemployment rate (calculated = unemployed population / (unemployed + total population))
6	Population density
7	Percentage of urban population
8	Percentage of non-agricultural household population
9	Percentage of households living in rented houses
10	Percentage of employees working in primary industries and mining
11	Percentage of employees working in secondary industries
12	Percentage of employees working in tertiary industries
13	Household size (no. of people/household)
14	Percentage of population with college degree (25 years old)
15	Percentage of population with high school degree (20 years)
16	Percentage of illiterate people (15 years)
17	Population growth rate (2000–2010)
18	Average number of rooms per household

19	Per capita housing construction area (m ² /person)
20	Percentage of premises without tap water
21	Percentage of premises without a kitchen
22	Percentage of premises without a toilet
23	Percentage of premises without a bath
24	Number of beds per 1000 people in health care institutions
25	Number of medical personnel per 1000 resident population
26	Percentage of population <5 years
27	Percentage of population 65 years
28	Population dependency ratio (%)
29	Percentage of population covered by subsistence allowances

286 After performing factor Principal Component Analysis (PCA) analysis of the 29 variables, seven
287 components with an eigenvalue >1 are extracted. Based on the variable meanings in each component,
288 these 7 components are named as table 3. The first component, which reflects the income of the
289 population and the employment situation, contributed 30.1% of the total variance. This
290 factor/component is positive because the more property there is in an area, the higher the vulnerability
291 to damage. The second component, which reflects the level of education level of the population,
292 occupies/contributes 15.6% of the total variance. This factor/component is negative because if the level
293 of education level is higher, then the population's awareness of disaster prevention and reduction is
294 greater and their vulnerability will be is lower. The third component, which reflects the number of
295 dilapidated houses, takes up/contributes 8.7% of the total variance. This factor/component plays a
296 positive part in vulnerability. The fourth component, which reflects the amount of illiteracy and the
297 number of young people, is positive and represents 8.4% of the total variance. The fifth component,
298 which reflects the household size and the percentage of women, explains 7.7% of the total variance and
299 is positive. The sixth component, which reflects the number of people from ethnic minorities,
300 contributes/explains 6.1% of the total variance and is positive. The seventh component, which
301 represents 5.3% of the total variance, reflects the unemployment rate and the housing area and is

302 positive.

303 The total variance explained by these seven components is up to 81.9%, which can be used to
 304 represent the ~~vulnerability of the~~ population vulnerability of Zhejiang ~~Province~~ province. The
 305 distributions of the first (positive) component and the second (negative) component are shown in
 306 Figure 9. Areas with a low employment rate have ~~a~~ high vulnerability, but the vulnerability is low in
 307 urban areas with higher levels of education. The seven components thus represent the real situation of
 308 the population vulnerability ~~of the population~~ in Zhejiang ~~Province~~ province to the effect of typhoons.
 309 The population vulnerability ~~index of the population~~ in Zhejiang ~~Province~~ province (SoVI) ~~wais~~
 310 calculated as:

$$311 \quad \text{SoVI} = \text{factorcomponent } 1 - \text{factorcomponent } 2 + \text{factorcomponent } 3 + \text{factorcomponent } 4 + \\
 312 \quad \text{factorcomponent } 5 + \text{factorcomponent } 6 + \text{factorcomponent } 7 \\
 313 \quad (2)$$


314 By calculating the vulnerability indexes of each county, ~~we obtained~~ the distribution of
 315 population vulnerability in Zhejiang ~~Province~~ province ~~is obtained~~ (Figure 10). The areas with high
 316 vulnerabilities are mountainous regions where the economy is relatively undeveloped, whereas the
 317 vulnerability is low in ~~cities~~ coastal areas, such as Hangzhou and Huzhou cities, where there is a greater
 318 awareness of disaster prevention and reduction and houses are of high quality.

319 Table 3. The seven components extracted by ~~principal component analysis~~ PCA.


<u>Components</u>	<u>Contained variables</u>	<u>Name</u>	<u>(Sign)</u>
1	<u>QMANFEMP, UBINCM, QAGREMP,</u> <u>QRENT, POPCH, QDEPEND,</u> <u>QSUBSIST, QPOPAB65, POPDEN,</u> <u>MEDAGE, QNOKITCH, QILLIT,</u> <u>PHROOM, PPHAREA</u>	<u>Employment and</u> <u>poverty</u>	<u>(+)</u>
2	<u>QHISCH, QCOLLEGE, QNONAGRI,</u> <u>QSEVEMP, HPBED, MEDTECH</u>	<u>Education</u>	<u>(-)</u>

<u>3</u>	<u>QNOBATH, QNOTOILET, PPUNIT</u>	<u>Number of dilapidated houses</u>	<u>(+)</u>
<u>4</u>	<u>QILLIT, QDEPEND, QPOPUD5, MEDAGE</u>	<u>Illiteracy and juvenile population</u>	<u>(+)</u>
<u>5</u>	<u>QFEMALE, PHROOM, PPHAREA, QSEVEMP</u>	<u>Household size and ratio of women</u>	<u>(+)</u>
<u>6</u>	<u>QMINOR</u>	<u>Ethnic minority</u>	<u>(+)</u>
<u>7</u>	<u>QUNEMP, QNOPIPWT</u>	<u>Unemployment and housing size</u>	<u>(+)</u>

320



321




322

323

324

Figure 9. Distribution of population vulnerability index of (a) factorcomponent 1 (employment and income) and (b) factorcomponent 2 (education).



325

326

327

Figure 10. Distribution of population vulnerability index of counties in Zhejiang Province.

328

54.3 Typhoon Disaster Comprehensive Risk Index for Typhoon Disasters and Zoning of

329 **Zhejiang Province**

330 The typhoon disaster risk assessment system ~~considers the~~ is mainly composed of the factors causing
331 disasters, ~~the~~ the population vulnerability ~~of the population~~ and the environment. ~~In this paper,~~
332 ~~the typhoon disaster~~ comprehensive risk index ~~for typhoon disasters~~ is obtained by combining the factors
333 causing typhoon disasters and vulnerability, ~~without~~ but does not take ~~ing~~ the sensitivity of the
334 environment into account. After standardizing the intensity index of factors causing typhoon disasters
335 and the population vulnerability index, the typhoon disaster comprehensive risk index (R) ~~in Zhejiang~~
336 ~~Province~~ was obtained as follows:

337
$$R = \text{intensity index of factors causing typhoon disasters } (I) \times \text{vulnerability index (SoVI)} \quad (3)$$

338 Based on the comprehensive risk index, ~~we defined~~ five risk grades ~~zones~~ for typhoon disasters are
339 defined in Zhejiang Province (Table 4). and risk zoning of typhoon disasters in Zhejiang province has
340 been done as shown in Figure 11. The classification of typhoon disaster risk index is based on the
341 natural breaks method (Jenks) provided by Arcgis.

342 Table 4. Disaster risk index and grade.

Risk grade:	High	High–medium	Medium	Medium–low	Low
Risk index:	0.3	0.18–0.3	0.13–0.18	0.07–0.13	0.07

343 Figure 11 shows that, ~~the index~~ presents ~~gave~~ a good reflection of the distribution of typhoon
344 disasters in Zhejiang ~~Province~~ province (Figure 3), especially in the southeastern coastal areas. ~~The risk-~~
345 ~~zoning of typhoon disasters in Zhejiang Province is shown in Figure 11.~~ The southeast coastal areas
346 face the highest risk, especially in the boundary regions between Zhejiang and Fujian province, and
347 Taizhou and Wenzhou ~~Ningbo~~ cities. Overall, the risk of typhoon disasters decreases from the coast to
348 inland areas. Cities are at medium to low risk as a result of their developed economy, high-quality
349 houses and better educated population. The inland mountainous areas have a high vulnerability. ~~and,~~
350 a Although they are not directly affected by typhoons, they are still in the middle risk areas as a result of
351 their poorly developed economy.





Fig. 11. Risk zoning of typhoon disaster areas in Zhejiang Province.

6.5 Discussion and Conclusions

(1) The intensity indexes of factors causing typhoon disasters is developed, with the highest values in Wenzhou, Taizhou and Ningbo cities, consistent with the risk analysis. A comparison between the distributions of the intensity index and actual typhoon disasters in Zhejiang Province from 2004 to 2012 shows that the index is a good reflection of the possibility of typhoon disasters in each county.

(2) Seven components were extracted after performing factor analysis (PCA) of 29 variables affecting social vulnerability. These seven factors represent 81.9% of the total variance and are a good reflection of the index of population vulnerability in Zhejiang Province. Southwestern Zhejiang is the most vulnerable as it has a relatively undeveloped economy, more mountainous areas and a higher risk of geological disasters. Vulnerabilities are lower in cities and coastal areas as a result

366 of better disaster prevention and reduction measures and a better educated population.

367 (3) A-Typhoon disaster comprehensive risk index ~~for typhoon disasters was~~ obtained by
368 combining the factors causing typhoon disasters and population vulnerability. Based on the
369 comprehensive risk index, ~~we developed~~ risk zoning zones of typhoon disasters in Zhejiang
370 Province province is achieved. The southeast coastal areas are at high risk, especially the boundary
371 regions between Zhejiang and Fujian province, and Taizhou and Wenzhou Ningbo cities. The risk of
372 typhoon disasters decreases quickly from coastal areas to inland regions. Cities are at medium to low
373 risk because of their developed economy, high-quality houses and better educated population.

374 Although some interesting results have been obtained in this study, there are still some problems
375 that require further studies. As a result of the limited data on typhoon disasters ~~in Zhejiang Province~~, it
376 is currently impossible to give a long time trend for high-resolution typhoon disaster analysis. It is also
377 unclear whether this methodology can be applied to other regions.

378 **Acknowledgments**

379 This study was supported by the Chinese Ministry of Science and Technology Project No.
380 2015CB452806.

381 **References**

382 Chen, H. Y., Yan, L. N., and Lou, W. P.: On assessment indexes of the strength of comprehensive
383 impacts of tropical cyclone disaster-causing factors, *Journal of Tropical Meteorology*, 27(1), 139-144,
384 2011. (in Chinese)

385 Chen, W. F., Xu, W., and Shi, P. J.: Risk assessment of typhoon disaster at county level in the Yangtze
386 river delta of china, *Journal of Natural Disasters*, 4, 77-83, 2011.

387 Chen, W. F., Duan, Y. H., and Lu, Y.: Review on Tropical Cyclone Risk Assessment, Journal of
388 Catastrophology, 32(4), 2017. (in Chinese)

389 Chen, X.: Vulnerability diagnosis and assessment of typhoon disaster system at coastal regions, a case
390 study of Fujian province, *Journal of Catastrophology*, 22(3), 6-10, 2007. (in Chinese)

391 Cutter, S. L., Boruff, B. J., and Shirley, W. L.: Social Vulnerability to Environmental Hazards, Social
392 Science Quarterly, 84(2):242-261, 2003.

393 Ding, Y. and Shi, P. J.: Fuzzy risk assessment model of typhoon hazard. *Journal of Natural*
394 *Disasters*, 11(1), 34-43, 2002. (in Chinese)

395 Emanuel, K. A.: The maximum intensity of hurricanes, *Journal of the Atmospheric Sciences*, 45(7),
396 1143-1155, 1988.

397 Emanuel, K. A.: The dependence of hurricane intensity on climate, American Institute of Physics, 277,
398 25-33, 1992.

399 Emanuel, K. A.: Sensitivity of tropical cyclones to surface exchange coefficients and a revised
400 steady-state model incorporating eye dynamics, Journal of the Atmospheric Sciences, 52(22),
401 3969-3976, 1995.

402 [Eunjeong, C. and Ying, M.: Comparison of three western North Pacific tropical cyclone best track
403 datasets in seasonal context, Journal of the Meteorological Society of Japan, 89\(3\):211-224, 2009.](#)

404 Fang, W. H. and Lin, W.: A review on typhoon wind field modeling for disaster risk
405 assessment, Progress in Geography, 32(6), 852-867, 2013. [\(in Chinese\)](#)

406 Fang, W. H. and Shi, X. W.: A review of stochastic modeling of tropical cyclone track and intensity for
407 disaster risk assessment, Advances in Earth Science, 27(8), 866-875, 2012. [\(in Chinese\)](#)

408 Friedman, D. G.: Natural hazard risk assessment for an insurance program, The Geneva Papers on Risk
409 and Insurance - Issues and Practice, 9(1), 57-128, 1984.

410 [Hardoon, D. R., Szedmak, S., and Shawetaylor, J.: Canonical Correlation Analysis: An Overview with
411 Application to Learning Methods, Neural Computation, 16\(12\):2639-2664, 2014.](#)

412 Hendrick, R. L. and Friedman, D. G.: Potential impacts of storm modification on the insurance industry,
413 University of Chicago Press, Chicago, 227-248, 1966.

414 Holland, G. J.: The maximum potential intensity of tropical cyclones, Journal of Atmospheric
415 Sciences, 54(21), 2519-2541, 1997.

416 [Hotelling, H.: Relations between two sets of variates, Biometrika, 28\(3/4\), 321-377, 1936.](#)

417 Howard, R. A., Matheson, J. E., and North, D. W.: The decision to seed hurricanes. Science, 176(4040),
418 1191-202, 1972.

419 [Huang, W. K. and Wang, J. J.: Typhoon damage assessment model and analysis in Taiwan, Natural
420 Hazards, 79\(1\), 497-510, 2015.](#)

421 IPCC: Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation,
422 Cambridge, UK, and New York, NY, USA, 2012.

423 Jiang, T., Li, X. C., and Chao, Q. C.: Highlights and understanding of climate change 2014: impacts,
424 adaptation, and vulnerability, Advances in Climate Change Research, 10(3), 157-166, 2014.

425 Kafali, C. and Jain, V.: A Methodology for Estimating Typhoon Losses to Building Environment in
426 Japan, in: Proceedings of the 11th Americas Conference on Wind Engineering, San Juan, Puerto Rico,
427 2009.

428 Kunreuther, H. and Roth, R. J.: The Status and Role of Insurance Against Natural Disaster in the
429 United States, Washington Joseph Henry Press, 1998.

430 Lei, X. T., Chen, P. Y., Yang, Y. H., and Qian, Y. Z.: Characters and objective assessment of disasters
431 caused by typhoons in china, Acta Meteorologica Sinica, 67(5), 875-883, 2009. [\(in Chinese\)](#)

432 [Li, S. H. and Hong, H. P.: Use of historical best track data to estimate typhoon wind hazard at selected
433 sites in China, Natural Hazards, 76\(2\):1395-1414, 2015.](#)

434 Liang, B. Q., Liang, J. P., and Wen, Z. P.: Study of typhoon disasters and its affects in china, Journal of
435 Natural Disasters, 1, 84-91, 1995. [\(in Chinese\)](#)

436 Liang, B. Q. and Fan, Q.: A fuzzy mathematic of the disaster bytropical cyclones, Journal of Tropical
437 Meteorology, 4, 305-311, 1999. [\(in Chinese\)](#)

438 [Liu, T. J. and Gu, J. Q.: A statistical analysis of typhoon disasters in Zhejiang province, Journal of
439 Catastrophology, 17 \(4\): 64-71, 2002. \(in Chinese\)](#)

440 Lu, Y., Zhu, W. J., Ren, F. M., and Wang, X.: Changes of Tropical Cyclone High Winds and Extreme
441 Winds During 1980-2014 over China, Advances in Climate Change Research, 12(5), 413-421, 2016.
442 [\(in Chinese\)](#)

443 Niu, H. Y., Liu, M., and Lu, M.: Risk assessment of typhoon disasters in china coastal area during last
444 20 years, Scientia Geographica Sinica, 31(6), 764-768, 2011. [\(in Chinese\)](#)

445 Pielke, R. A. and Landsea, C. W.: Normalized hurricane damages in the united states: 1925-95, Weather
446 & Forecasting, 13(3), 621--631, 1998.

447 Pielke, R. A, Gratz, J., and Landsea, C. W.: Normalized hurricane damage in the united states:
448 1900–2005, Natural Hazards Review, 9(1), 29-42, 2008.

449 Pita, G., Pinelli, J. P., and Gurley, K.: State of the art of hurricane vulnerability estimation methods: a
450 review, Natural Hazards Review, 16(2), 04014022, 2014.

451 Ren, F. M., Byron, G., and David, E.: A Numerical Technique for Partitioning Cyclone Tropical
452 Precipitation, Journal of Tropical Meteorology, 17(3), 308-313, 2001. [\(in Chinese\)](#)

453 Ren, F. M., Wang, Y., Wang, X., and Li, W.: Estimating Tropical Cyclone Precipitation from Station
454 Observations, Advances in Atmospheric Sciences, 24(4), 700-711, 2007.

455 [Ren, F. M., Wang, X. L., Chen, L. S., and Wang, Y. m.: Tropical cyclones landfalling on mainland
456 China, Hainan and Taiwan and their correlations, Acta Meteorologica Sinica, 66 \(2\): 224-235, 2008. \(in
457 Chinese\)](#)

458 Ren, F. M. and Wu, G. X.: Tropical cyclone over the past 60 years, China Meteorological Press, Beijing,
459 2011. [\(in Chinese\)](#)

460 Su, G. L., Miao, C. M., and Mao, Y. D.: Typhoon hazard in zhejiang province and risk assessment of its
461 influence on agriculture, Journal of Natural Disasters, 17(5), 113-119, 2008. [\(in Chinese\)](#)

462 Vickery, P. J., Masters, F. J., and Powell, M. D.: Hurricane hazard modeling: the past, present, and
463 future, Journal of Wind Engineering & Industrial Aerodynamics, 97(7–8), 392-405, 2009.

464 Wang, X. R., Wang, W. G., and Ma, Q. Y.: Model for general grade division of typhoon disasters and
465 application, Meteorological Monthly, 36(1), 66-71, 2010. [\(in Chinese\)](#)

466 WMO: Seventh International Workshop on Tropical Cyclones (IWTC-VII), available at:
467 <http://www.wmo.int/pages/prog/arep/wwrp/tmr/IWTC-VII.html/>, 2010.

468 [Xu, X., Sun, D., and Guo, T.: A systemic analysis of typhoon risk across china, Natural Hazards, 77\(1\),
469 461-477, 2015.](#)

470 [Yin, Y. Z. and Li, H. L.: Preliminary study on pre-evaluation method of typhoon disaster in China,
471 Meteorological Monthly, 43\(6\):716-723, 2017. \(in Chinese\)](#)

- 472 Yin, Z. E., Xu, S. Y., Yin, J., and Wang, J.: Small-scale Based Scenario Modeling and Disaster Risk
473 Assessment of Urban Rainstorm Water-logging, *Acta Geographica Sinica*, 65(5), 553-562, 2010. [\(in](#)
474 [Chinese\)](#)
- 475 Zhang, Q., Wu, L. G., and Liu, Q. F.: Tropical cyclone damages in china 1983-2006, *Bulletin of the*
476 *American Meteorological Society*, 90(4), 489-495, 2009.