Analysis of Spatiotemporal variations in mid-upper tropospheric methane during the Wenchuan Ms8.0 earthquake by three indices

Jing Cui¹ and Xuhui Shen¹

¹Key Laboratory of Crustal Dynamics, Institute of Crustal Dynamics, China Earthquake Administration, Beijing 100085, China

Correspondence: Jing Cui (jingcui_86@yahoo.com)

Abstract. This research studied the spatiotemporal variation in methane in the mid-upper troposphere during the Wenchuan earthquake (12 May, 2008) using AIRS retrieval data and discussed the methane anomaly mechanism. Three indices were proposed and used for analysis. Our results show that the methane concentration increased significantly in 2008, with an average increase of 5.12×10^{-8} , compared to the average increase of 1.18×10^{-8} in the previous five years. The Alice and Diff indices

- 5 can be used to identify methane concentration anomalies. The two indices showed that the methane concentration distribution before and after the earthquake broke the distribution features of the background field. As the earthquake approached, areas of high methane concentration gradually converged towards the west side of the epicenter from both ends of the Longmenshan fault zone. Moreover, a large anomalous area was centered at the epicenter eight days before the earthquake occurred, and a trend of strengthening, weakening and strengthening appeared over time. The Gradient index showed that the vertical direction
- 10 obviously increased before the main earthquake, and the value was positive. The gradient value is negative during coseismic or postseismic events. The gradient index reflects the gas emission characteristics to some extent. We also determined that the methane release was connected with the deep crust-mantle stress state, as well as microfracture generation and expansion. However, due to the lack of any technical means to accurately identify the source and content of methane in the atmosphere before the earthquake, an in-depth discussion has not been conducted, and further studies on this issue may be needed.

15 Copyright statement. TEXT

1 Introduction

The great Wenchuan Ms8.0 earthquake, on May 12, 2008, occurred in the Longmenshan Fault zone in western Sichuan Province, China. Its epicenter was located at coordinates $30.95^{\circ}N$, $103.40^{\circ}E$. The extent of the earthquake and aftershock affected areas in the northeast along the Longmen Shan fault, a thrust structure along the border of the Tibetan Plateau and the

20 western Sichuan Basin (Figure 1). This earthquake was one of the worst continental earthquake events to have struck China in recent decades, and it killed more than ten thousand people in several cities along the western Sichuan basin. A surface rupture more than 200 km long formed along the Yingxiu-Beichuan fault. The Guanxian-Jiangyou fault was also ruptured by



Figure 1. Map of the active faults and the location of the Wenchuan earthquake. LMS means the Longmenshan fault zone; XSH means the Xianshuihe fault zone; DLS means the Daliangshan fault zone; Anninghe means the Anninghe fault zone.

surface ruptures that were more than 60 km long, as indicated. The Longmenshan fault zone has a high dipping angle (more than $50^{\circ} - 60^{\circ}$) near the surface and a low angle at depth (15 - 20 km). This listric shape favors significant strain or energy accumulation, forming large earthquakes. This earthquake was characterized by slow strain accumulation, a long recurrence interval and significant damage power. It is a new type of earthquake that deserves further study (Zhang et al., 2008).

5

The variation in soil gas concentration serves as a useful tool for monitoring earthquakes. Numerous field investigations have indicated that large amounts of gases are emitted from active fault zones before, during and after great earthquakes. The increased gas emanation from the Earth's crust in the vicinity of active tectonic faults is triggered by a chain of physical processes and chemical reactions from the ground surface. This complex chain leads to geochemical, atmospheric, ionospheric

and magnetospheric anomalies (Dobrovolsky et al., 1979; Pulinets and Ouzounov, 2011; Ouzounov et al., 2007). The number of pre-earthquake thermal, surface latent heat flux and outgoing longwave radiation anomalies apparently results from earthquake-related gas emission from the lithosphere (Tronin et al., 2002; Dey et al., 2004; Tronin, 2006; Ouzounov et al., 2007; Tramutoli et al., 2013). Thermal (Zhang et al., 2010), ionospheric (Lin, 2012, 2013; Zhu et al., 2010), electromagnetic (Zhang et al., 2011a), and aerosol (Qin et al., 2014) anomalies were found before the Wenchuan earthquake of May 12, 2008.

As the second most important greenhouse gas after carbon dioxide (CO_2) , methane (CH_4) is approximately 20 times better at warming the atmosphere than CO_2 by weight and plays an important role in atmospheric chemistry. Some people have speculated that thermal abnormalities before earthquakes are related to the release of CH_4 . The CH_4 release mechanism has been investigated and results indicate strong methane emissions when friction is applied to marl-type rock (Martinelli

5

- 10 and Plescia, 2005; Italiano et al., 2008). Yue indicated that the Wenchuan earthquake of May 12, 2008, was caused by the rapid migration and expansion of a large amount of highly pressurized and dense CH_4 gas in crustal rock masses (Yue, 2013). Sample results indicated that CH_4 was discharged from a shallow reservoir through faults or fractures caused by the earthquake (Zheng et al., 2013). This study choose to study the spatiotemporal variations of CH_4 during the Ms8.0 Wenchuan earthquake from satellite observations.
- 15 CH_4 satellite observations have started to be applied to seismological studies over recent years. The total column of CH_4 associated with the 12 May 2008 Wenchuan earthquake was investigated using satellite data from the AQUA Atmospheric Infrared Sounder (AIRS), and this work indicates that a large amount of CH_4 was emitted from underground into the atmosphere along the Longmenshan Fault Zone from approximately 1.5 months before to 3 months after the earthquake, and the closer to the epicenter, the larger the amount of emitted gas. The peak values were found at intersection areas (Yue, 2013; Cui et al.,
- 20 2017). However, the same column concentration might correspond to this distinct vertical structure. It is hard to say if the CH_4 anomalies are from a ground source or transform. The gas profile can help us understand the terrestrial emission caused by the earthquake. Recently, NASA declared that the total column values reported for CH_4 constituents are dominated by the initial guess of each and should not be used for research purposes (https://docserver.gesdisc.eosdis.nasa.gov/repository/Mission/AIRS/3.7_ScienceDataProductValidation/V6_Data_Disclaimer.pdf). Previous work has shown that an AIRS for tropospheric
- observation can be used to carry out the study with considerable precision (Zhang et al., 2011b; Xiong et al., 2015). Systematic observation of the vertical variation of CH_4 is scarce. The focus of this study is to examine the spatiotemporal variation of CH_4 in the mid-upper troposphere during the Wenchuan earthquake (12 May, 2008) using AIRS retrieval data and to discuss the mechanism of the methane anomaly. Three indices were proposed and used for analysis: the Absolute Local Index of Change of the Environment (ALICE) was used for anomaly detection; the Vertical Concentration Gradient (Gradient)
- 30 was proposed to study the vertical variation; the Successive Differential Value (Diff) can show the time variation. These three indices analyzed the spatial and temporal distribution of CH_4 before and after the earthquake from horizontal, vertical and time scales helping us understand lithospheric and atmospheric interactions during seismic activity.

2 Data and Methods

10

The Version 6 and Level 3 standard gridded product of 8-day CH_4 volume-mixing ratios in a descending model (local nighttime), with 1 * 1 degree of spatial resolution, were obtained from the NASA Goddard Earth Sciences Data and Information Services Center (DISC) (http://disc.gsfc.nasa.gov/AIRS/index.shtml/). The peak sensitivity of the AIRS to methane retrieval

5 occurs at 300 hPa, and the channels near 7.6 μm are most sensitive to the middle to upper troposphere (Xiong et al., 2015). Therefore, the volume mixing ratio of the middle troposphere (400 hPa, approximately 5 km), the upper middle troposphere (300 hPa, approximately 7 km) and the upper troposphere (200 hPa, approximately 11 km) in the descending orbit data was used.

To extract the methane spatiotemporal anomalies before and after the earthquake, three parameters were applied: the Absolute Local Index of Change of the Environment (ALICE) (Tramutoli, 1998; Tramutoli et al., 2013; Cui et al., 2017) the Vertical

- Concentration Gradient (Gradient) and the Successive Differential (Diff), all carried out on the basis of eliminating the multiyear background. The background field can partially remove the influence of natural sources such as seasonal changes and surface vegetation, effectively capturing emergency information such as earthquakes, reducing "nonseismic anomalies" to a certain extent, providing criteria for the extraction of seismic anomalies, and reducing the misjudgment and leakage of seismic
- 15 anomalies. It was calculated by Eq 1:

N7

$$G_{ref}(x,y,t,l) = \sum_{i=1}^{N} G_i(x,y,t,l) / N$$
(1)

$$\sigma(x, y, t, l) = \sqrt{\sum_{i=1}^{N} \left[G_i(x, y, t, l) - G_{ref}(x, y, t, l)\right]^2 / (N - 1)}$$
(2)

where $G_i(x, y, t, l)$ means the gas value at level l, measured at time t, corresponding to a location at (centered on) the coordinates (x, y); $G_{ref}(x, y, t, l)$ means the reference fields for the study area, defined as a time average gas value; $\sigma(x, y, t, l)$ is the standard deviation of historical records collected under the temporal constraint. For this study, N was defined as 5 years,

20 is the standard deviation of historical records collected under the temporal constraint. For this study, N was defined as 5 years, from 2003 to 2007; t is the time of the measurement acquisition with t ∈ τ, where τ defines the homogeneous domain of the satellite imagery collected in the same time-slot of the day and period(month) of the year; l was defined as 3 layers (400 hPa, 300 hPa and 200 hPa).

The Absolute Local Index of Change of the Environment (ALICE) is calculated by Eq. 3 (Tramutoli, 1998; Cui et al., 2017):

$$25 \quad ALICE(x,y,t,l) = [G(x,y,t,l) - G_{ref}(x,y,t,l)/\sigma(x,y,t,l)]$$

$$(3)$$

The Vertical Concentration Gradient (Gradient) is used to characterize the vertical variation of gas and is calculated by Eq 4 to Eq 7 :

$$Gradient(x, y, t) = \left[G(x, y, t, \Delta l) - G_{ref}(x, y, t, \Delta l)\right] / \sigma(x, y, t, \Delta l)$$
(4)

$$G(x, y, t, \Delta l) = G(x, y, t, l) - G(x, y, t, l+1)$$
(5)

$$G_{ref}(x,y,t,\Delta l) = \sum_{i=1}^{N} G_i(x,y,t,\Delta l)/N$$
(6)

$$\sigma(x, y, t, \Delta l) = \sqrt{\sum_{i=1}^{N} \left[G_i(x, y, t, \Delta l) - G_{ref}(x, y, t, \Delta l)\right]^2 / (N - 1)}$$
(7)

where Gradient(x, y, t, l) stands for vertical concentration gradient value at level l, measured at time t, corresponding to 5 a location at (centered on) the coordinates (x, y); G(x, y, t, l) means the gas value at level l. $G(x, y, t, \Delta l)$ means the vertical difference of adjacent layers; $G_{ref}(x, y, t, \Delta l)$ means the reference fields of the vertical difference for the study area, defined as a time average gas value; $\sigma(x, y, t, \Delta l)$ is the standard deviation of historical records collected under the temporal constraint. P_t stands for the atmospheric pressure at l, l was defined as 3 layers (400 hPa, 300 hPa and 200 hPa).

Successive Differential Value (Diff) refers to the difference of the adjacent time gas value and was calculated by Eq. 8 to Eq. 10 11 :

$$Diff(x, y, \Delta t, l) = \left[G(x, y, \Delta t, l) - G_{ref}(x, y, \Delta t, l)\right] / \sigma(x, y, \Delta t, l)$$
(8)

$$G(x, y, \Delta t, l) = G(x, y, t+1, l) - G(x, y, t, l)$$
(9)

$$G_{ref}(x, y, \Delta t, l) = \sum_{i=1}^{N} G_i(x, y, \Delta t, l) / N$$
(10)

$$\sigma(x, y, \Delta t, l) = \sqrt{\sum_{i=1}^{N} \left[G_i(x, y, \Delta t, l) - G_{ref}(x, y, \Delta t, l)\right]^2 / (N - 1)}$$
(11)

15 3 Results

3.1 Reference field

The 8-day average background field at different CH_4 heights in the study area was obtained and is shown in Fig.2, 3, and 4. These figures show that the CH_4 concentration in the study area decreased significantly from the middle troposphere to the upper, and the average methane concentrations (from Mar 9th to Aug 15th, 2008) in each stratum were 1.794 PPM, 1.775 PPM and 1.762 PPM for middle, upper middle, and upper troposphere, respectively. This indicates that the middle troposphere is significantly affected by terrestrial emission sources.

- The distribution of CH_4 has obvious seasonal characteristics and the seasonal cycles at different heights are similar. The CH_4 mixing ratio has a weak high value during Mar, and then decreases during April-May. It starts to increase in June and stays high during July and August, especially over the Sichuan basin. The relatively high values at different times are mostly located at a structural confluence or at tectonic plate boundaries except for late July and August. Earth gas emissions may be the main cause of the high long-term values at that location (http://www.climatechange2013.org/images/report/P36Doc3_WGI-12_Approved-SPM.pdf). Why does the CH_4 begin to increase in early summer? Studies have shown CH_4 emissions
- 10 significantly correlate with temperature. When the temperature is $< 15^{\circ}$ the Earthseldom produces CH_4 , and when the temperature reaches 35° , CH_4 production can be many times higher than at 25° (KazuyukiYagi and KatsuyukiMinami, 1990; Thomas et al., 1996; Holzapfel-Pschorn et al., 1986; Saarnio et al., 2010; Saarnio and Silvola, 1999). Therefore, the CH_4 concentration in the troposphere is higher during July and August. The CH_4 rice paddy emission, local emissions (such as from gas leakage or energy use), and transport (such as the meridional and zonal advection, convection from the lower troposphere,
- 15 or stratosphere-troposphere exchange) also increase in early summer (Xiong et al., 2010).

The results of the background field analysis show that the CH_4 concentration is greatly influenced by the season, structure and underlying surface, and they present a typical spatial and temporal distribution of methane in southwest China (Zhang et al., 2011b). Gas emanates from the earth's crust continuously, even without earthquakes.

3.2 Anomalous Gas

20 The three indices mentioned above have been applied to identify the spatial and temporal variation of mid-upper tropospheric methane anomalies associated with the Wenchuan Ms 8.0 earthquake using AIRS data.

3.2.1 ALICE index result

Figures 5, 6, and 7 show that as the earthquake approached, the methane concentration near the LMS fault zone gradually increased and the range gradually expanded. High values began to appear in the northern section of the LMS fault zone on

- 25 March 25th, and began to increase in large areas on May 4th, showing a significant north-south distribution through the LMS fault zone. From May 12th to the 19th, the high value decreased near the epicenter, but remained in large parts of the study area and persisted in large areas after the earthquake. From July 7th, the northern section of the LMS fault zone again showed high accumulation. From July 23rd to August 7th, the high value zone reached the maximum, and all along the LMS fault zone it showed NE distribution. Then, the high-value areas began to weaken on August 8. This circle was probably caused by the big
- 30 aftershocks on July 24th and August 1st.



Figure 2. Five year (2003 to 2007) average distribution of 8-day nighttime CH_4 VMR at 200 hPa.



Figure 3. Five year (2003 to 2007) average distribution of 8-day nighttime CH_4 VMR at 300 hPa



Figure 4. Five year (2003 to 2007) average distribution of 8-day nighttime CH_4 VMR at 400 hPa



Figure 5. ALICE index of 8-day CH_4 VMR at 200 hPa



Figure 6. ALICE index of 8-day CH_4 VMR at 300 hPa



Figure 7. ALICE index of 8-day CH_4 VMR at 400 hPa

3.2.2 Vertical Concentration Gradient index result

The Vertical Concentration Gradient stands for the vertical rate of change. Figures 8 and 9 show the value changing obviously at the Bayan har block and plate boundary. The highest values appeared at the Bayan har block, the XSH, ANH and DLS faults and lasted until May 12th before the earthquake. The values were positive before May 12th. They weaken and even turn

5 negative after May 12th, especially during July 23rd to August 7th. The overall shape is still closely related to the fault and basin edge. We will discusses this later.

3.2.3 Successive Differential Value

The trend of the reflected gas concentration over time was reflected by the difference change, and the result of 20080222 represented the difference between February 22nd and the 14th. The places with rapid changes mainly occur at plate boundaries

10 and tectonic intersections. Figure 10 shows that the LMS fault zone began to show an abnormally high value on March 25th, and then the high value area gradually increased. On April 10th, high values appeared at the Bayan har block, the XSH, ANH and DLS faults and lasted until May 12th before the earthquake. The value of the entire LMS fault zone was high during May 12th. The next cycle began on July 7th, when high values first appeared in the northern section of the LMS fault zone, followed by the entire LMS, XSH, ANH and DLS fault zones on July 23rd. Two large aftershocks were reported on July 24th and August 1st.

4 Discussion

Background field analysis can help us understand the gas distribution in the study area and has a certain reference role for the extraction and analysis of late abnormal changes. We have found that the CH_4 concentration is greatly influenced by season, structure and underlying surface, and presents a typical spatial and temporal distribution in southwest China. Comparing our

20 work with the GPS velocity field, the high value area of the background field corresponds to the area of high GPS velocity (Wu et al., 2015). Long-term stress accumulation causes deep earth gases to be expelled along fissures or structural weaknesses (Wang et al., 2017).

Three indices have been used to see if anomalous CH_4 could be identified. The CH_4 spatial variations were most likely caused by geological processes and/or the action of crustal stress in the lithosphere resulting in the Wenchuan earthquake. 25 A large amount of CH_4 was emitted from underground into the atmosphere along plate boundaries and tectonic belts from approximately 1.5 months before to 3 months after the earthquake, and the closer to the epicenter, the larger the amount of emitted gas. A large anomalous area occurs centered at the epicenter eight days before the earthquake. A crucial question in this field of research refers to how can we link an individual precursor with a distinctive stage of the earthquake preparation.

The generation of such a seismic anomaly requires physical and chemical transformations which occur in a spatially extended preparation (activation) zone of an impending earthquake. Earthquakes exhibit in general complex correlations in time, space and magnitude. It is widely accepted that the observed earthquake scaling laws indicate the existence of phenomena closely



Figure 8. Gradient of 400 hPa to 300 hPa



Figure 9. Gradient of 300 hPa to 200 hPa



Figure 10. Distribution of the successive differential value at 300 hPa

associated with the proximity of the system to a critical point (Varotsos et al., 2019). Therefore, such a requirement is satisfied during the appearance of the "critical window", i.e., the epoch during which the short-range correlations have evolved to longrange ones in an extended area, where the "critical radius R" is given by the empirical relation $log R \approx 0.5M$, where M is the earthquake magnitude (Bowman et al., 1998). Notice, based on the recently introduced concept of the "natural time" by

Varotsos and his colleagues (Varotsos et al., 2011). it has been shown that the foreshock seismic activity that occurs in the 5 region around the epicentre of the upcoming significant shock a few days up to one week before the main shock occurrence, behaves as critical phenomenon.

Therefore, the hypothesis that the large anomaly in methane eight days before the earthquake occurred corresponds to the critical point- window of the earthquake preparation process cannot be excluded. Accumulated experimental evidence supports

the aforementioned hypothesis as follows: 10

The earthquake preparatory process has various facets which reflect correspondingly different precursors. Importantly, precursors emerge during the same period. A few days up to one week before the main shock occurrence, while they behave as critical phenomena, as well. Characteristically, such as precursors are: (i) ULF magnetic field variations recorded by groundbased magnetic observatories before significant earthquakes, e.g., (Contoviannis et al., 2016; Potirakis et al., 2018). (ii) MHz

fracture induced MHz EM anomalies (Eftaxias et al., 2018). The generation of such a seismic anomaly also requires physical 15 and chemical transformations which occur in a spatially extended preparation (activation) zone of an impending earthquake.

Characteristic precursors are athe short-lived seismo-ionospheric EM precursors and EM anomalies rooted in preseismic LAI-coupling (Pulinets et al., 2003; Pulinets and Boyarchuk, 2004). Pulinets et al. (Pulinets et al., 2003) have provided strong evidence for the occurrence of ionospheric precursors well before the main shock: ionospheric precursors within a 5 days before the seismic shock were registered in 73% of the cases for earthquakes with a magnitude 5, and in 100% of the cases for



earthquakes with a magnitude 6.

To further verify that the increase in CH_4 concentration is related to earthquakes, we compared and analyzed the variation of overall methane concentration and anomaly index in the study area for nonseismic years (Figures 11 and 12, there was no large earthquake during 2003 to 2007). Compared with 2003-2007 (Figure 11), the CH_4 concentration increased significantly

in 2008 with an average increase of 5.12×10^{-8} , compared with the average increase of 1.18×10^{-8} in the past five years (Figure 25 11). Especially between Mar 25th and June 21st. The results from the past five years showed that the concentration of CH_4 increased significantly due to seasonal influence from June 21st, showing a rapid increase in the slope of the curve (Table 1). Although there was a certain increase in 2008, the curve slope was relatively small (Table 1), which may be due to the large amount of CH_4 released by the aftershock earthquake. The increasing trend of CH_4 concentration caused by seasonal change

weakened between Mar 25th and June 21st, 2008. 30

We also compare the ALICE result with 2008 and 2007 (Figures 12 and 13). The distribution of the ALICE index of 8-day CH_4 VMR at 300 hPa during 2007 is very different from 2008. In 2007, the index results were similar on the whole to stable features with no significant differences. The average value is almost between -2 to 2 except Feb 7th, 2007. However, the average value for 2008 is over 2 after April 10th, 2008, one month before the earthquake. Although gaseous anomalies can be

affected by climate change and weather patterns, human activity, vegetation coverage and many other factors beyond seismic 35



Figure 11. Comparison of the average CH_4 VMR in the study area from 2003 to 2008 (300 hPa)

| Table 1. Linear fitting betw | een June 21st and S | eptember 1st | t for di | fferent years |
|------------------------------|---------------------|--------------|----------|---------------|
|------------------------------|---------------------|--------------|----------|---------------|

| Year | Linear equation | R-square |
|------|----------------------|----------|
| 2003 | y = 0.9272x + 174.15 | 0.9788 |
| 2004 | y = 0.6044x + 176.53 | 0.9051 |
| 2005 | y = 0.6916x + 175.51 | 0.9404 |
| 2006 | y = 0.715x + 176.76 | 0.9847 |
| 2007 | y = 0.7058x + 175.25 | 0.9484 |
| 2008 | y = 0.4387x + 181.43 | 0.8066 |

activity, these effects may be local and can be eliminated by calculation with Equations 1 to 11. Therefore, the identified anomalies of CH_4 VMR can be considered earthquake-related (Cui et al., 2017). CH_4 is an important greenhouse gas next

to carbon dioxide, and previous work shows that the obvious anomalies of thermal infrared brightness temperature appeared on April 25th and were mainly distributed in the LMS fault zone and its southern region (Zhang et al., 2010). These results correspond well with our work.

- Compared with the nonseismic years from 2003 to 2007 (no earthquake >Ms 5.0), the study area before and after the 5 Wenchuan earthquake showed a significantly increased CH_4 anomaly. It is well known that CH_4 is in a supercritical or critical state at high temperature and pressure 10-20 km below the surface, and its physical and chemical properties are relatively stable during periods of weak tectonic activity. However, with active geological activity, this state is also very easy to break. For example, the sudden expansion of the volume of cracks or cavities caused by the change of tectonic stress field can easily change the internal fluid state and cause the supercritical fluid phase transition (Liu and Du, 2000). As soon as the phase
- 10 transition process begins, fluid volume and fracture expansion are triggered, resulting in a large amount of fluid spilling over the surface and spreading into the atmosphere, which is then observed by ground-based instruments or satellites (Martinelli and Plescia, 2005; Italiano et al., 2008).

Before the Wenchuan earthquake, various GPS observation points and level observations showed accelerated activity of different degrees. The LMS Fault activity is characterized by a transition from tensile to compressive, and acceleration occlusion

- 15 occurs near the fault field. Accelerated uplift and extension of the crust appeared in the Bayan har block (Fang et al., 2009; Jiao et al., 2008) indicating that when the main section of the LMS fault zone is still in the state of friction lock, the relatively weak Bayan Har block shows the uplift and expansion of a large number of tensile cracks under long-term tectonic stress. It is with this process that the underground fluid surges up and overflows the surface on a large scale, and then spreads and rises to the upper troposphere, resulting in the high CH_4 concentration region on the western side of the LMS in Fig. 5-10 before
- 20 the earthquake. Many NW fracture surfaces were found on the western side of the LMS after the Wenchuan earthquake. With the occurrence of such cracks, microcracks or preslip, supercritical CH_4 inside the crust can well up and overflow, resulting in increased atmospheric CH_4 concentration.

Figures 5 to 10 show that the CH_4 concentration in the inner area of the Sichuan Basin on the eastern side of the LMS Fault zone is at a low level before and after the earthquake, and occasionally increases temporarily. The occasional short rise may be due to CH_4 release and diffusion from the surrounding fault zone. Although there are a large number of oil and gas deposits in the sedimentary cover of the Sichuan Basin, the present tectonic activity is relatively weak, and it is difficult to produce such phenomena as rupture and expansion in the upper crust, so that large-scale release of CH_4 and other fluids will not occur.

The migration or release of underground fluids such as CH_4 is inevitably completed by other seismic processes in the preearthquake period, which is also an important way for energy conversion and accumulation from the deep part of the crust and

- 30 mantle to the shallow part of the crust. Therefore, it can be inferred that the phase transition process of underground fluids such as CH_4 was mainly completed before the earthquake, so there was an abnormal increase before the earthquake. During the earthquake and coseismic activity, the abnormal increase of fluid concentration such as CH_4 may also occur, which is mainly caused by local fluid release in the brittle layer of the upper crust (Wang et al., 2017). The range is mainly limited to the vicinity of the earthquake rupture zone, which is not on the same order of magnitude as the regional CH_4 release from the deep
- 35 part. For example, the gradient shows that the values are positive before May 12th and become weak and even turn negative



Figure 12. ALICE index of 8-day CH₄ VMR at 300 hPa during 2007



Figure 13. Comparison of the average ALICE index from 2007 and 2008

after May 12th (Figures 8 and 9). The CH_4 in the upper layer may be deep gas released before the earthquake, and locally released after the earthquake decreases, so the concentration of the lower layer decreases and the gradient becomes negative. The iteration of positive and negative values can also indicate geogenic emissions.

The aforementioned results seem to support the hypothesis that the observed anomaly in terms of spatiotemporal variation in methane is rooted in the stage of critical point-epoch of the earthquake preparation process.

5 Conclusions

5

This study has examined the spatiotemporal variation in CH_4 in the mid-upper troposphere during the Wenchuan earthquake (May 12th, 2008) using AIRS retrieval data and discusses the mechanism of the methane anomaly. The CH_4 concentration increased significantly in 2008 with an average increase of 5.12×10^{-8} , compared with 1.18×10^{-8} in the past five years. Three

indices were proposed and used. The Absolute Local Index of Change of the Environment (ALICE) was used for anomaly detection; the Vertical Concentration Gradient (Gradient) was proposed to study vertical variation; the Successive Differential Value (Diff) shows time variation. The three indices analyzed the spatial and temporal distribution of CH_4 before and after the earthquake from horizontal, vertical and time scales. Through this work, the following conclusions are obtained:

5 The background field of CH_4 is greatly influenced by season, structure and underlying surface and presents a typical spatial and temporal distribution of methane in southwest China. Gas emanates from the earth's crust continuously, even without earthquakes.

The CH_4 concentrations in the upper and middle atmosphere before and after the Wenchuan earthquake have a certain temporal and spatial variation. The ALICE and Diff indices could be used to identify the CH_4 concentration anomaly. The research

- 10 results indicate that the CH_4 concentration distribution before and after the earthquake breaks the distribution features of the background field. The CH_4 concentration distribution starts from both ends and gradually gathers around the epicenter. Moreover, a large anomalous area occurs centered at the epicenter eight days before the earthquake, and a trend of strengthening, weakening and strengthening appears over time.
- The Gradient method can reflect the change in gas concentration in the vertical direction. The results show that the vertical direction obviously increases before the main earthquake, and the value is positive. The gradient value is negative during coseismic or postseismic events. It may be that the gas before the main earthquake mainly comes from deep fluid phase transitions, whereas the coseismic or postseismic gas may mainly come from the locally closed fluid in the brittle layer of the crust. However, due to the lack of any technical means to accurately identify the source and content of methane in the atmosphere before the earthquake, an in-depth discussion has not been conducted, and further studies on this issue may be needed. The gradient index can reflect the characteristics of gas emission to some extent.

Appendix A

A1

Author contributions. Jing Cui designed this study, performed most of the data analysis and wrote the paper. Xuhui Shen contributed to the discussions.

25 Competing interests. The authors declare no competing interests.

Acknowledgements. This research was supported by grants from the National Natural Science Foundation of China (Grant No. 41602223) and the Institute of Crustal Dynamics, China Earthquake Administration (Grant No. ZDJ2018-18). The authors would like to thank NASA

for making available the AIRS dataset. The authors wish to thank the anonymous reviewers for their constructive comments that helped improve the scholarly quality of the paper.

References

15

Bowman, D., Ouillon, G., Sammis, C., Sornette, A., and Sornette, D.: An observational test of the critical earthquake concept, Journal of Geophysical Research: Solid Earth, 103, 24 359–24 372, 1998.

Contoyiannis, Y., Potirakis, S., Eftaxias, K., Hayakawa, M., and Schekotov, A.: Intermittent criticality revealed in ULF magnetic fields prior

- 5 to the 11 March 2011 Tohoku earthquake (Mw= 9), Physica A: Statistical Mechanics and its Applications, 452, 19–28, 2016.
 - Cui, Y., Ouzounov, D., Hatzopoulos, N., Sun, K., Zou, Z., and Du, J.: Satellite observation of CH 4 and CO anomalies associated with the Wenchuan M S 8.0 and Lushan M S 7.0 earthquakes in China, Chemical Geology, 469, 2017.
 - Dey, S., Sarkar, S., and Singh, R. P.: Anomalous changes in column water vapor after Gujarat earthquake, Advances in Space Research, 33, 274–278, 2004.
- 10 Dobrovolsky, I. P., Zubkov, S. I., and Miachkin, V. I.: Estimation of the size of earthquake preparation zones, Pure & Applied Geophysics, 117, 1025–1044, 1979.
 - Eftaxias, K., Potirakis, S. M., and Contoyiannis, Y.: Four-Stage Model of Earthquake Generation in Terms of Fracture-Induced Electromagnetic Emissions: A Review, in: Complexity of Seismic Time Series, pp. 437–502, Elsevier, 2018.

Fang, D. U., Wen, X. Z., Zhang, P. Z., Beijing, Chengdu, Beijing, Beijing, and Xi, A.: Interseismic deformation across the Longmenshan fault zone before the 2008 M8. 0 Wenchuan earthquake, Chinese Journal of Geophysics, 52, 2729–2738, 2009.

- Holzapfel-Pschorn, A., Conrad, R., and Seiler, W.: Effects of vegetation on the emission of methane from submerged paddy soil, Plant & Soil, 92, 223–233, 1986.
 - Italiano, F., Martinelli, G., and Plescia, P.: CO2 Degassing over Seismic Areas: The Role of Mechanochemical Production at the Study Case of Central Apennines, Birkhäuser Basel, 2008.
- 20 Jiao, Q., Yang, X., Xu, L., and Wang, B.: PRELIMINARY STUDY ON MOTION CHARACTERISTICS OF LONGMENSHAN FAULT BEFORE AND AFTER Ms8.0 WENCHUAN EARTHQUAKE, Journal of Geodesy & Geodynamics, 28, 7–11, 2008.

KazuyukiYagi and KatsuyukiMinami: Effect of organic matter application on methane emission from some Japanese paddy fields, Soil Science and Plant Nutrition, 36, 599–610, 1990.

Lin, J. W.: Potential reasons for ionospheric anomalies immediately prior to China's Wenchuan earthquake on 12 May 2008 detected by

- 25 nonlinear principal component analysis, International Journal of Applied Earth Observation & Geoinformation, 14, 178–191, 2012.
 - Lin, J. W.: Ionospheric anomaly at the occurring time of China's May 12, 2008, M = 7.9 Wenchuan Earthquake using nonlinear principal component analyses and image decoding, Egyptian Journal of Remote Sensing & Space Sciences, 16, 53–61, 2013.
 - Liu, W. and Du, Jianguo.and Bai, L.: A review on the role of supercritical fluids in the earthquake generation, Seismology and Geology, 22, 339–444, 2000.
- 30 Martinelli, G. and Plescia, P.: Carbon dioxide and methane emissions from calcareous-marly rock under stress: experimental tests results, Annals of Geophysics, 48, 167–173, 2005.
 - Ouzounov, D., Liu, D., Kang, C., Cervone, G., Kafatos, M., and Taylor, P.: Outgoing long wave radiation variability from IR satellite data prior to major earthquakes, Tectonophysics, 431, 211–220, 2007.
 - Potirakis, S. M., Contoyiannis, Y., Asano, T., and Hayakawa, M.: Intermittency-induced criticality in the lower ionosphere prior to the 2016
- Kumamoto earthquakes as embedded in the VLF propagation data observed at multiple stations, Tectonophysics, 722, 422–431, 2018.
 Pulinets, S. and Boyarchuk, K.: Ionospheric precursors of earthquakes, Springer Science & Business Media, 2004.

Pulinets, S. and Ouzounov, D.: Lithosphere–Atmosphere–Ionosphere Coupling (LAIC) model – An unified concept for earthquake precursors validation, Journal of Asian Earth Sciences, 41, 371–382, 2011.

- Pulinets, S., Legen'Ka, A., Gaivoronskaya, T., and Depuev, V. K.: Main phenomenological features of ionospheric precursors of strong earthquakes, journal of atmospheric and solar-terrestrial physics, 65, 1337–1347, 2003.
- 5 Qin, K., Wu, L. X., Zheng, S., Bai, Y., and Lv, X.: Is there an abnormal enhancement of atmospheric aerosol before the 2008 Wenchuan earthquake?, Advances in Space Research, 54, 1029–1034, 2014.
 - Saarnio, S. and Silvola, J.: Effects of increased CO 2 and N on CH 4 efflux from a boreal mire: a growth chamber experiment, Oecologia, 119, 349–356, 1999.

Saarnio, S., Alm, J., Martikainen, P. J., and Silvola, J.: Effects of Raised CO2on Potential CH4Production and Oxidation in, and CH4Emission

- 10 from, a Boreal Mire, Journal of Ecology, 86, 261–268, 2010.
 - Thomas, K. L., Benstead, J., Davies, K. L., and Lloyd, D.: Role of wetland plants in the diurnal control of CH 4 and CO 2 fluxes in peat, Soil Biology & Biochemistry, 28, 17–23, 1996.
 - Tramutoli, V.: Robust AVHRR techniques (RAT) for environmental monitoring: theory and applications, Proc Spie, 3496, 101–113, 1998. Tramutoli, V., Aliano, C., Corrado, R., Filizzola, C., Genzano, N., Lisi, M., Martinelli, G., and Pergola, N.: On the possible origin of thermal
- 15 infrared radiation (TIR) anomalies in earthquake-prone areas observed using robust satellite techniques (RST), Chemical Geology, 339, 157–168, 2013.

Tronin, A. A.: Remote sensing and earthquakes: A review, Physics & Chemistry of the Earth, 31, 138–142, 2006.

Tronin, A. A., Hayakawa, M., and Molchanov, O. A.: Thermal IR satellite data application for earthquake research in Japan and China, Journal of Geodynamics, 33, 519–534, 2002.

20 Varotsos, P., Sarlis, N. V., and Skordas, E. S.: Natural time analysis: the new view of time: precursory seismic electric signals, earthquakes and other complex time series, Springer Science & Business Media, 2011.

Varotsos, P., Sarlis, N., and Skordas, E.: Natural time analysis: Important changes of the order parameter of seismicity preceding the 2011
 M9 Tohoku earthquake in Japan, EPL (Europhysics Letters), 125, 69 001, 2019.

Wang, J., Xianrui, L. I., Chengchen, D. U., and Zeng, Z.: Aerial methane concentration anomaly and air temperature increase before the Wenchuan earthquake, Earth Science Frontiers, 2017.

- Wu, Y., Jiang, Z., Zhao, J., Liu, X., Wei, W., Liu, Q., Li, Q., Zou, Z., and Zhang, L.: Crustal deformation before the 2008 Wenchuan M S 8.0 earthquake studied using GPS data, Journal of Geodynamics, 85, 11–23, 2015.
- Xiong, X., Barnet, C. D., Zhuang, Q., Machida, T., Sweeney, C., and Patra, P. K.: Mid-upper tropospheric methane in the high Northern Hemisphere: Spaceborne observations by AIRS, aircraft measurements, and model simulations, Journal of Geophysical Research Atmospheres,
- 30 115, -, 2010.

25

Xiong, X., Barnet, C., Maddy, E., Sweeney, C., Liu, X., Zhou, L., and Goldberg, M.: Characterization and validation of methane products from the Atmospheric Infrared Sounder (AIRS), Journal of Geophysical Research Biogeosciences, 113, 0–1, 2015.

Yue, Z. Q.: Cause and mechanism of highly compressed and dense methane gas mass for Wenchuan earthquake and associated rockavalanches and surface co-seismic ruptures, Earth Science Frontiers, 20, 15–20, 2013.

35 Zhang, P. Z., Xu, X. W., Wen, X. Z., and Ran, Y. K.: Slip rates and recurrence intervals of the Longmen Shan active fault zone, and tectonic implications for the mechanism of the May 12 Wenchuan earthquake, 2008, Sichuan, China, Chinese Journal of Geophysics, 51, 1066–1073, 2008.

- Zhang, X., Shen, X., and Miao, Y.: Electromagnetic Anomalies around Wenchuan Earthquake and Their Relationship with Earthquake Preparation, International Journal of Geophysics, 2011, (2011-8-7), 2011, 693–701, 2011a.
- Zhang, X. Y., Bai, W. G., Zhang, P., and Wang, W. H.: Spatiotemporal variations in mid-upper tropospheric methane over China from satellite observations, Chinese Science Bulletin, 56, 3321–3327, 2011b.
- 5 Zhang, Y. S., Guo, X., Zhong, M. J., Shen, W. R., Li, W., and He, B.: Wenchuan earthquake: Brightness temperature changes from satellite infrared information, Chinese Science Bulletin, 55, 1917–1924, 2010.
 - Zheng, G., Xu, S., Liang, S., Shi, P., and Zhao, J.: Gas emission from the Qingzhu River after the 2008 Wenchuan Earthquake, Southwest China, Chemical Geology, 339, 187–193, 2013.

Zhu, F., Wu, Y., Lin, J., and Zhou, Y.: Temporal and spatial characteristics of VTEC anomalies before Wenchuan Ms8.0 earthquake, Geodesy

^{10 &}amp; Geodynamics, 1, 23–28, 2010.