

# ***Interactive comment on “Numerical Simulations of the 2004 Indian Ocean Tsunami Deposits Thicknesses and Emplacements” by Syamsidik et al.***

**Syamsidik et al.**

syamsidik@tdmrc.org

Received and published: 8 March 2019

COMMENTS FROM REFEREE #2: This study investigated deposits by the Indian Ocean tsunami in 2004 through the field measurement and reproduced it by using numerical simulations, which yields very interesting and valuable findings. However, the significance of this research should be more emphasized in the introduction. In the present manuscript, only the applicability of numerical models seem to be the main subject. The authors should show idea on how to make use of the results of this research in disaster prevention and reduction, such as estimating the magnitude of past tsunamis from sediments and evaluating the energy that the tsunami transports

[Printer-friendly version](#)

[Discussion paper](#)



earth and sand. Although the authors stated that "this study estimated the energy required to transport sediments via a tsunami wave", the answer to this question is not clearly shown in the manuscript.

## RESPONSE:

Dear Referee #2,

We sincerely appreciate your input to our paper. In recent years, interests on using tsunami deposits information to estimate source of the tsunami and its magnitude have been in significant increase. Our study offers the use of numerical modeling prior to field investigation. This would provide a complimentary knowledge to locate potential locations of the tsunami deposits. In some studies, such as Jankaew et al. (2008) and Monecke et al. (2008), their important tsunami deposit findings were located after several attempts and identification of geological setting of the locations. Significance of our research is the use of the 2004 Indian Ocean tsunami with two different times of investigations, i.e one in 2005 (Jaffe et al., 2006 ) and in 2015 by authors. Understanding the energy of the tsunami from sediment transport process could be useful to validate energy generated by the tsunami. As in the 2004 Indian Ocean tsunami case, where monitoring equipments were rarely deployed in the affected area, information deduced from the tsunami deposit could better explain number of waves, wave heights, and wave velocity. In section 4.4, we discuss tsunami wave's shear stress simulated using the 2DH Delft3D model. The shear stress in sediment transport formulae represents the energy produced by the waves/currents during the process. We agree to include more explanation in the introduction part to elaborate the use of the study for disaster prevention and mitigation. Previous studies on tsunami deposits have inspired tsunami mitigation efforts in US and in Indonesia (Dunbar et al., 2008; Rubin et al., 2017). Data on tsunami deposits would provide more scientific evidence to past tsunamis for a significant period of records such as in the case of coastal cave in Aceh that preserved a very long record of past tsunamis in the region (Rubin et al., 2017). A set of tsunami deposits data could help us to estimate the recurrence period of the

[Printer-friendly version](#)[Discussion paper](#)

tsunami from the subduction zone (Fujiwara, 2007; Minoura et al., 2001). Finding more tsunami deposits would strengthen validation process and ease some ways to estimate its sources through inverse mechanism (Buckley et al., 2012; Moore et al., 2011). A similar explanation will be added to the Introduction part of our revised manuscript.

## References

Buckley, M.L., Wei, Y., Jaffe, B.E., Watt, S.G.: Inverse modeling of velocities and inferred cause of overwash that emplaced inland fields of boulders at Anegada, British Virgin Islands. *Natural Hazards* 63(1), 133-149. DOI: <https://doi.org/10.1007/s11069-011-9725-8>, 2012.

Dunbar P.K., Stroker, K.J., Brocko, V.R., Varner, J.D., McLean, S.J., Taylor, L.A., Eakins, B.W., Carignan, K.S., Warnken, R.R.: Long-Term Tsunami Data Archive Supports Tsunami Forecast, Warning, Research, and Mitigation. In: Cummins P.R., Satake K., Kong L.S.L. (eds) *Tsunami Science Four Years after the 2004 Indian Ocean Tsunami*. Pageoph Topical Volumes. Birkhäuser Basel, 2008.

Fujiwara, O.: Major contribution of Tsunami deposit studies to Quaternary Research. *The Quaternary Research* 46(3), 293-302. DOI: <https://doi.org/10.4116/jaqua.46.293>, 2007.

Jankaew, K., Atwater, B.F., Sawai, Y., Choowong, M., Charoentitirat, T., Martin M.E., and Prendergast A.: Medieval forewarning of the 2004 Indian Ocean tsunami in Thailand. *Nature* 455, 1228-1231, 2008. Minoura, K., Imamura, F., Sugawara, D., Kono, Y. and Iwashita, T.: The 869 Jogan tsunami deposit and recurrence interval of large-scale tsunami on the Pacific coast of northeast Japan. *J. Nat. Disast. Sci.* 23, 83–88, 2001.

Monecke, K., Finger, W., Klarer, D., Kongko, W., McAdoo, B.G., Moore A.L., and Sudrajat S.U.: A 1,000-year sediment record of tsunami recurrence in Northern Sumatra. *Nature* 455, 1232-1234, 2008.

Moore, A., Goff, J., McAdoo, B.G., Fritz, H.M., Gusman, A., Kalligeris, N., Kalsum, K.,

[Printer-friendly version](#)[Discussion paper](#)

Susanto, A., Suteja, D., Synolakis, C.E.: Sedimentary Deposits from the 17 July 2006 Western Java Tsunami, Indonesia: Use of Grain Size Analyses to Assess Tsunami Flow Depth, Speed, and Traction Carpet Characteristics. *Pure and Applied Geophysics* 168(11), 1951-1961. DOI: <https://doi.org/10.1007/s00024-011-0280-8>, 2011.

Rubin, C.M, Horton, B.P, Sieh, K., Pilarczyk, J.E., Daly, P., Ismail N., and Parnell A.C.: Highly variable recurrence of tsunamis in the 7,400 years before the 2004 Indian Ocean tsunami. *Nature Communications* 8, 16019, 2017.

---

Interactive comment on *Nat. Hazards Earth Syst. Sci. Discuss.*, <https://doi.org/10.5194/nhess-2018-348>, 2018.

[Printer-friendly version](#)

[Discussion paper](#)

