## **Precipitation extremes in a EURO-CORDEX 0.11° ensemble at hourly resolution: Supplementary material**

Peter Berg<sup>1</sup>, Ole B. Christensen<sup>2</sup>, Katharina Klehmet<sup>1</sup>, Geert Lenderink<sup>3</sup>, Jonas Olsson<sup>1</sup>, Claas Teichmann<sup>4</sup>, and Wei Yang<sup>1</sup>

<sup>1</sup>Swedish Meteorological and Hydrological Institute, Folkborgsvägen 17, 610 76 Norrköping, Sweden <sup>2</sup>Danish Meteorological Institute, Lyngbyvej 100, 2100 Copenhagen, Denmark

<sup>3</sup>KNMI Royal Netherlands Meteorological Institute, Utrechtseweg 297, 3731 GA De Bilt, the Netherlands <sup>4</sup>Climate Service Center Germany (GERICS), Helmholtz-Zentrum Geesthacht, Fischertwiete 1, 20095 Hamburg, Germany

**Correspondence:** Peter Berg (peter.berg@smhi.se)



**Figure 1.** Shape parameter (middle column) for each model for the historical period 1971-2000 at **1 h duration**. The 95% confidence interval values are shown for the (left) low estimate and (right) high estimate.



**Figure 2.** Scale parameter (middle column) for each model for the historical period 1971-2000 at **1 h duration**. The 95% confidence interval values are shown for the (left) low estimate and (right) high estimate.



**Figure 3.** Shape parameter (middle column) for each model for the historical period 1971-2000 at **12 h duration**. The 95% confidence interval values are shown for the (left) low estimate and (right) high estimate.



**Figure 4.** Scale parameter (middle column) for each model for the historical period 1971-2000 at **12 h duration**. The 95% confidence interval values are shown for the (left) low estimate and (right) high estimate.



**Figure 5.** Scatterplot of the relative change in 10-year **1 h** depths against summertime mean temperature change between future and historical time periods, for different sub-regions, emission scenarios, and time periods according to the legend. Each panel show the result for different RCM-GCM combinations. Linear fits to all data are presented in each panel, along with slope and intercept coefficients as well as the R-square value of the fit. CC-rate changes of 7%/K are shown as gray lines in the plots.