

Dear Editor,

Thank you very much for the acknowledgement to our effort and the result reflected in the manuscript. We are going to go carefully through the aspects indicated by the reviewers to continue improving the manuscript, with a special focus on the figures.

Best regards,

Diana Contreras

Reviewer 1

Thank you very much for your observations. You kindly spent time again delving into our manuscript, and we are grateful. We have used a colour code to answer your questions. Please find your comments already solved in grey, and the respective answers in black. The corresponding paragraphs in the paper are in dark blue.

Comments

Using the two leading databases on scientific journal publications (Scopus/Elsevier and SCI/Clarivate Analytics), the authors performed an assessment of articles published between 2010 and 2020 addressing the social and economic dimensions of vulnerability. A number of 27 publications were finally chosen for a detailed analysis on data sources, spatial variables, indicators, methods, indexes and tools for the spatial assessment of socio-economic vulnerability (SEV) related to geohazards. In the revised version, the paper reads much clearer and is more concise so that the potential readers can follow the string of argumentation as well as (by the included Tables) the relevant indicators discussed.

Overall, the manuscript is therefore suitable for publication in the target journal. My comments on the first version of the manuscript have been thoroughly acknowledged. The authors may wish to additionally specify the following:

- Page 1, line 28: my recommendation here is to refer to the original UNDRR source and not to Aubrecht et al. (2013) – this statement can be found in nearly all publications related to the topic, and I strongly recommend to cite the original source here.

Thank you very much for your recommendation. Please find the change accordingly:

‘(...) Vulnerability is defined by United Nations (UN) as ‘the conditions determined by physical, social, economic and environmental factors or processes which increase the susceptibility of an individual, a community, assets or systems to the impact of hazards’ (UN, 2016) (...)’.

- Page 5, line 15: I would recommend to insert “Austria” after Salzburg for those readers not familiar with European cities.

Thank you very much for your recommendation. Please find the change accordingly:

‘(...) In the case study area of Salzburg (Austria), an expert-based approach was chosen, and several experts were asked to allocate weights according to the contribution of each variable to the vulnerability of floods (Contreras and Kienberger, 2011) (...)’.

- Page 6, lines 8ff: I would prefer to use “endogenic” and “exogenic” geohazards instead of “internal” and “external”, to be in line with the geoscience literature (also page 7, line 21).

Thank you very much for enlightening us with regards to the appropriate terms to use; we appreciate this insight. Please find the changes accordingly:

‘(...) Geohazards can be **endogenic** such as earthquakes, tsunamis, and volcanic eruptions and **exogenic** such as landslides, soil erosion, and land degradation (...)’.

‘(...) The terms selected for the search query refer to vulnerability in the socio-economic dimension, the spatial variables listed by Meentemeyer (1989), Béné (2009), Contreras et al. (2013) and Buzai and Villerías Alarcón (2018) and the aforementioned **endogenic** and **exogenic** geohazards (...)’.

‘(...) The lack of articles that tackle **exogenic** geohazards can be explained by the fact that we excluded from the search query words such as “climate change” OR “ecological” OR “drought”, which are indirectly related to these phenomena (...)’.

- Moreover, the argument that “...we chose these hazards because they produced high losses... in Chile” seems a bit strange in the context of this manuscript, and can simply be deleted. If it is the choice of the authors to focus on the hazard types indicated, from my point of view, it is not necessary to state the reason. For the overall review it is not important whether or not certain hazard types chosen are relevant in Chile, it simply detracts the reading flow.

Thank you very much for your comment. As we stated, the reason to choose geohazards phenomena came as a request from a previous reviewer. Therefore, we agree to delete the reference to Chile, but we must keep the remainder of the text that explains the reason to choose geohazards phenomena.

‘(...) We particularly focus on these phenomena for two reasons: first, geohazards are the natural phenomena that have produced the highest quantity of losses in recent years in the urban environments, **particularly earthquakes**, and, second, because geohazards are the phenomena addressed by the institutions involved in the present research (...)’.

- Besides, I still have a certain doubt about leaving out other geohazards of exogenic type which – in particular in the US literature – are often subsumed under the term “landslides”, such as the “debris flow”-type hazard group, or the entire group of (gravitational) mass movements – please see e.g. the recent publication of Mazzorana et al. (2019, page 1092) explicitly focusing on Chile. Nevertheless, given the statements made in these lines it is ok for me just to name the hazard groups that were analysed, which in turn explains why other hazard types also relevant (for Chile) are missing.

Thank you very much for your comment. We also expected to have more literature references related to landslides in our systematic review, but after running the query using the terms defined in Table 1 (please see below), we managed only to obtain 5 references, which is 17% of the references identified for this systematic review. We deleted the reference to Chile following your advice.

Topic	(social vulnerability* OR societal vulnerability* OR socioeconomic vulnerability* OR socio-economic vulnerability* OR economic vulnerability*)
	AND
Topic	(area* OR distance* OR range* OR distance* OR direction* OR spatial geometries* OR patterns* OR spatial connectivity* OR isolation* OR diffusion* OR spatial association* OR scale* OR accessibility* OR network* OR cluster*)
	AND
Topic	(earthquakes* OR tsunamis* OR volcanic eruptions* OR landslides* OR soil erosion* OR land degradation)
	NOT
Topic	(climate change* OR ecological* OR drought* OR resilience* OR debris* OR epidemiological* OR substance* OR behavioral* OR evacuation* OR recovery* OR pollution* OR leptospirosis* OR violence* OR illness* OR disease* OR heat* OR

crisis* OR Conflict* OR deaths* OR obesity* OR criminal* OR chemical* OR symptoms* OR syndrome* OR food insecurity* OR air pollution* OR stress* OR diabetes* OR depressive* OR alcohol* OR cancer* OR drugs* OR palm oil* OR tobacco* OR smoke* OR storm* OR psychometric* OR cocaine* OR toxic* OR palliative* OR therapy* OR HIV* OR dengue* OR ecosystem* OR rheumatoid arthritis* OR nutritional* OR malaria* OR resources* OR sexual activity* OR sexual health*).

Table 1. Terms included and excluded to select relevant literature references

Literature references related to landslides identified in our systematic review using the terms listed in Table 1. Please see below:

- Chen, Y. (2016). Conceptual Framework for the Development of an Indicator System for the Assessment of Regional Land Subsidence Disaster Vulnerability. *Sustainability*, 8(8), 757.
- Ley-García, J., Denegri de Dios, F. M., & Ortega Villa, L. M. (2015). Spatial dimension of urban hazardscape perception: The case of Mexicali, Mexico. *International Journal of Disaster Risk Reduction*, 14, 487-495. doi:<https://doi.org/10.1016/j.ijdrr.2015.09.012>
- Toke, N. A., Boone, C. G., & Arrowsmith, J. R. (2014). Fault zone regulation, seismic hazard, and social vulnerability in Los Angeles, California: Hazard or urban amenity? *Earth's Future*, 2(9), 440-457. doi:10.1002/2014ef000241
- Zeng, J., Zhu, Z. Y., Zhang, J. L., Ouyang, T. P., Qiu, S. F., Zou, Y., & Zeng, T. (2012). Social vulnerability assessment of natural hazards on county-scale using high spatial resolution satellite imagery: a case study in the Luogang district of Guangzhou, South China. *Environmental Earth Sciences*, 65(1), 173-182. doi:10.1007/s12665-011-1079-8
- Ebert, A., Kerle, N., & Stein, A. (2009). Urban social vulnerability assessment with physical proxies and spatial metrics derived from air- and spaceborne imagery and GIS data. *Natural Hazards*, 48(2), 275-294. doi:10.1007/s11069-008-9264-0

Thanks for the recommendation of the reference:

- Mazzorana, B., Picco, L., Rainato, R., Iroumé, A., Ruiz-Villanueva, V., Rojas, C., Valdebenito, G., Iribarren-Anacona, P., and Melnick, D.: Cascading processes in a changing environment: Disturbances on fluvial ecosystems in Chile and implications for hazard and risk management, *Science of the Total Environment*, 655, 1089-1103, 2019.

However, given that this reference does not address any of the geohazards included in this systematic review, we feel that we should not include it in this paper. However, we do appreciate the suggestion and will keep it in mind for future research studies.

References

- Alcorn, R., Panter, K. S., and Gorsevski, P. V.: A GIS-based volcanic hazard and risk assessment of eruptions sourced within Valles Caldera, New Mexico, *Journal of Volcanology and Geothermal Research*, 267, 1-14, 2013.
- Armaş, I., Toma-Danila, D., Ionescu, R., and Gavriş, A.: Vulnerability to Earthquake Hazard: Bucharest Case Study, Romania, *International Journal of Disaster Risk Science*, 8, 182-195, 2017.
- Béné, C.: Are Fishers Poor or Vulnerable? Assessing Economic Vulnerability in Small-Scale Fishing Communities, *The Journal of Development Studies*, 45, 911-933, 2009.
- Buzai, G. and Villerías Alarcón, I.: Análisis espacial cuantitativo de los determinantes sociales de la salud (DSS) en la cuenca del río Luján (provincia de Buenos Aires, Argentina), *Estudios Socioterritoriales*, 23, 2018.
- Contreras, D., Blaschke, T., Kienberger, S., and Zeil, P.: Spatial connectivity as a recovery process indicator: The L'Aquila earthquake, *Technological Forecasting and Social Change*, 80, 1782-1803, 2013.

Contreras, D. and Kienberger, S. (Eds.): Handbook of Vulnerability Assessment in Europe, European Commission DG Environment, 2011.

Deichmann, U.: Accessibility Indicators in GIS, United Nations Statistics Division, Department for Economic and Policy Analysis, New York, NY, USA, 1997.

Meentemeyer, V.: Geographical perspectives of space, time, and scale., Landscape Ecology 3, 163-173, 1989.

OECD: Handbook on Constructing Composite Indicators: Methodology and User Guide, OECD Publishing, 2008.

Shen, S., Cheng, C. X., Yang, J., and Yang, S. L.: Visualized analysis of developing trends and hot topics in natural disaster research, PLoS One, 13, 15, 2018.

Toke, N. A., Boone, C. G., and Arrowsmith, J. R.: Fault zone regulation, seismic hazard, and social vulnerability in Los Angeles, California: Hazard or urban amenity?, Earth Future, 2, 440-457, 2014.

Toké, N. A., Boone, C. G., and Arrowsmith, J. R.: Fault zone regulation, seismic hazard, and social vulnerability in Los Angeles, California: Hazard or urban amenity?, Earth's Future, 2, 440-457, 2014.

UN: Report of the open-ended intergovernmental expert working group on indicators and terminology relating to disaster risk reduction. UNDRR (Ed.), Preventionweb, 2016.

Zhou, Y., Li, N., Wu, W., Wu, J., and Shi, P.: Local Spatial and Temporal Factors Influencing Population and Societal Vulnerability to Natural Disasters, Risk Analysis, 34, 614-639, 2014.

Reviewer 2

Thank you very much for the acknowledgement to our effort to improve the manuscript, we highly appreciate your insights. You kindly spent time again delving into our manuscript, and we are grateful. Please find your comments already solved in grey and the respective answers in black. The corresponding paragraph in the paper is in dark blue.

Comments

Very good and detailed improvements, impressive energy, time and very detailed documentation of changes made. Congratulations for such efforts.

■ Make another technical check - some typos.

Thank you very much for this recommendation, we went again carefully through the manuscript and still, we asked for additional support of our English proof-reader to be sure that we eliminate all possible typos, grammar or punctuation mistakes on the manuscript.

- ‘society and the **economy** are only two of the dimensions of vulnerability (...)’.
- ‘(...) Data constraints **play** a key role in the results of the SEV assessment (...)’.
- ‘(...) The construction of an index implies selection of indicators, indicator normalization and weighting, and aggregation into an index (OECD, 2008) that must collectively represent aspects of **a** society’s ability to prepare for, deal with and recover from a disaster (...)’.
- ‘(...) We limited the query to articles published in academic journals because they typically are rigorous in the selection of their publications and therefore contain **a** complete **and accurate** description of methodologies and consistent results (...)’.
- ‘(...) To prioritize disaster-prone areas **that** are known as potential demand points (PDPs) given their vulnerability under large-scale earthquakes (...)’ .
- ‘(...) To apply an artificial neural network (ANN) and geographic information system (GIS) for estimating the social vulnerability to earthquakes in the Tabriz city, Iran (...)’

- ‘For the purpose of the systematic review, we found that the Clarivate Analytics database more accurately identified the references for this systematic review, and it is more user-friendly than other databases (...)’.
 - ‘(...) Housing quality and tenancy conditions [describe](#) the vulnerability of the population to become homeless after a disaster (Toke et al., 2014) (...)’.
 - ‘(...) Accessibility as [a](#) spatial indicator is defined as the ability [to](#) contact and [interact](#) with places of economic or social opportunities (Deichmann, 1997). (...)’
 - ‘(...) Alcorn et al. (2013) used an improved version of the same index but specifically adapted it to the variability in SEV in the case study area that was focused on census-designated places (CDPs) on a small scale. (...)’
 - ‘(...) Toké et al (2014), build upon the SoVI® to create their own SV indexes that incorporate the spatial dimension. According to the LA-SoVIC developed by Toket et al. (2014), SV is highly linked to the normalised difference vegetation index (NDVI) as a proxy for urban green space (...)’.
 - ‘(...) Armaş et al., (2017) applied a pairwise comparative method in the AHP implemented in the SMCE module of the Integrated Land and Water Information System ([IIWIS](#)) software (...)’.
 - ‘Based on the evidence, we can state that most of the spatial assessments of SEV in urban environments have been done for earthquakes and landslides and that Indonesia, China, Iran, and the USA lead the research in [the](#) spatial assessment of SEV related to geohazards in urban environments (...)’.
 - ‘The lack of data availability hinders the understanding of the concept of vulnerability (Zhou et al., 2014) and that is why VGI is essential today to obtain updated information in real-time at [the](#) local scale when other data sources are not available (...)’.
 - ‘(...) The spatial assessment of SEV in the areas where it is requested must depend not only on the financial resources for research but also on the availability of opensource software with the functionalities of spatial statistics, such as QGIS, GeoDa or [IIWIS](#) (...)’.
 - ‘(...) Likewise, spatial assessments of SEV must be considered before taking resettlement decisions for not creating again spatial conditions that [favour](#) the SEV (...)’.
- Make sure figures are provided in high resolution at the end.
Thank you very much for your observation. The high-resolution images were provided to the editor in separate files in the previous version. This time the high-resolution images (TIFF/300dpi) will be included in the text.
 - Figure 1. may be hard to read due to colour red and rather thin font.

Thank you very much for your observation. This time, we have included a high-resolution image (TIFF/300dpi) in the text; we also increased and changed the font from Arial to Times New Roman to improve the readability of the figure for the reader. All the text is in bold. The colour, rather than red, is orange (68R/68G/72B). We selected this colour for two reasons, on the one hand, it is an eye-catching colour; on the other hand, it is a colour in the corporate image of the Research Centre for Integrated Disaster Risk Management (CIGIDEN), one of the institutions involved in this research. In case the editor considers that it is not a proper colour, we are willing to change it to grey (255R/55G/0B). Please see the result below:




Figure 1. Methodology applied for the systematic literature review.

- Figure 2: Font size may be too small in final published version to read easily for terms under columns.

Thank you very much for your observation. The font size of the terms under the bars (geohazards) under the column was increased from 9pt to 11pt.




Figure 2. Number of literature references in the systematic review that addresses each geohazard

- Table 3: Check if this is intended that second column sometimes directly starts with author names. To me this is rather confusing.

Thanks for your question. The second column in Table 3 lists, in some cases, the specific name of the census, the kind of satellite images, the name of the disaster databases, the name of the land use/land cover (LULC) maps, and the name of population datasets used by the authors. This detail is not necessary in the case of the other data sources; that is why, in these cases, the second column starts directly with the author names. We explain this aspect in the manuscript as follows:

‘(...) The second column in Table 3 includes, in some cases specific details of the data source used by the authors (...)’.

- Figures 2&3 are nice, but I am a bit doubtful whether they actually do not raise more questions such as how much is overpresence of earthquakes due to selection method / search terms or due to occurrence. And the title of your paper is about urban contexts - so are these all truly (spatial) urban studies? If so, would maybe a list of cities reveal more details than the world map - like are certain countries always having 1-2 cities presented mainly, while in other countries, it is 3-4 cities. That could another interesting insight. The map as such rather prompts to think about the representativeness of just about 20 analysed papers. As does Figure 2. But I am also fine if you keep both figure 2 and 3 as they are.

Thanks for this comment. The literature references identified are based on a highly detailed search query to avoid bias. The query could be repeated any time, and the results will be always the same although additional publications from 2020 could appear in the result. Rather than go deeply in highlighting specific countries leading, and case study areas used in the research on spatial assessment of socio-economic vulnerability (SEV), we only wanted to include an additional finding supported by the research undertaken by Shen et al., (2018). We prefer to focus on the main research question of this manuscript: What is the state of the art in the spatial assessment of SEV to geohazards in urban environments? The state of the art is not necessarily linked to specific case study areas.

The total number of references reviewed were many more than 20. In the first version of the manuscript, based on a previous more general query not focused on geohazards, we identified 235 literature references, from which we found 84 relevant references, and 42 highly relevant references. Finally, 21 references were selected for review. From the previous search and given their relevance, six of these references were considered again in the current version of the manuscript. In the current version, we carefully reviewed all 29 references, but eventually we selected 18 and ruled 11 out for the reasons already explained in the manuscript. The query is extremely specific, which explains the reduced number of references identified, but the literature reviewed in this research included many more than the 24 selected at the end. Additionally, the result in the previous search regarding case study areas was similar, with the USA and China leading the research in the spatial assessment of SEV. As a result, we prefer to keep the Figures 2 and 3 as they are, and as you agree also with this decision, we made some changes in the format as you also requested. Please find below the new version of Figure 3. We added this explanation in the discussion section of the manuscript. Please read below:

‘(...) The literature references identified are based on a high detailed search query to avoid bias. The query could be repeated any time and the results will be always the same, with additional publications from 2020 that could appear in the result. However, the total number of literature references reviewed were much more than 24. Previously, based on a more general query not specifically focused on geohazards, we identified 235 literature references, from which we found 84 relevant references, 42 highly relevant references, and finally 21 references were selected to be reviewed. Eventually, given their relevance, we decided to keep six of these references identified previously using the first query. In the current version, we reviewed all 29 references but eventually, we selected 18 and discarded 11 for the reasons already explained in the results section. The case study areas of the selected papers confirm the findings from Shen et al., (2018) and also ours using the previous query, relating to the USA, China, and Iran as major contributors to disaster research together with Italy, Indonesia, Germany, Turkey, England, India, and Spain in the topics of ‘prediction model’, ‘social vulnerability’ and ‘landslide inventory map’. Nevertheless, the references that use Indonesia as a case study area are focused on earthquakes and volcanic eruptions, not necessarily on the tsunami hazard as was suggested by Shen et al., (2018) (...)’.




Figure 3. Number of study areas per country addressed in the references identified through the systematic literature review.

References

- Alcorn, R., Panter, K. S., and Gorsevski, P. V.: A GIS-based volcanic hazard and risk assessment of eruptions sourced within Valles Caldera, New Mexico, *Journal of Volcanology and Geothermal Research*, 267, 1-14, 2013.
- Armaş, I., Toma-Danila, D., Ionescu, R., and Gavriş, A.: Vulnerability to Earthquake Hazard: Bucharest Case Study, Romania, *International Journal of Disaster Risk Science*, 8, 182-195, 2017.
- Deichmann, U.: Accessibility Indicators in GIS, United Nations Statistics Division, Department for Economic and Policy Analysis, New York, NY, USA, 1997.
- OECD: Handbook on Constructing Composite Indicators: Methodology and User Guide, OECD Publishing, 2008.
- Shen, S., Cheng, C. X., Yang, J., and Yang, S. L.: Visualized analysis of developing trends and hot topics in natural disaster research, *PLoS One*, 13, 15, 2018.
- Toke, N. A., Boone, C. G., and Arrowsmith, J. R.: Fault zone regulation, seismic hazard, and social vulnerability in Los Angeles, California: Hazard or urban amenity?, *Earth Future*, 2, 440-457, 2014.
- Toké, N. A., Boone, C. G., and Arrowsmith, J. R.: Fault zone regulation, seismic hazard, and social vulnerability in Los Angeles, California: Hazard or urban amenity?, *Earth's Future*, 2, 440-457, 2014.
- Zhou, Y., Li, N., Wu, W., Wu, J., and Shi, P.: Local Spatial and Temporal Factors Influencing Population and Societal Vulnerability to Natural Disasters, *Risk Analysis*, 34, 614-639, 2014.

The spatial dimension in the assessment of urban socio-economic vulnerability related to geohazards, a systematic review

Diana Contreras^{1,2}, Alondra Chamorro¹, Sean Wilkinson²

¹Research Center for Integrated Disaster Risk Management (CIGIDEN), Pontifical Catholic University, Santiago, 7820436, Chile

²School of Engineering, Newcastle University, Newcastle upon Tyne, NE1 7RU, United Kingdom (UK)

Correspondence to: Diana Contreras (diana.contreras@cigiden.com – diana.contreras-mojica@newcastle.ac.uk)

Abstract

Society and economy are only two of the dimensions of vulnerability. This paper aims to elucidate the state of the art in data sources, spatial variables, indicators, methods, indexes and tools for the spatial assessment of socio-economic vulnerability (SEV) related to geohazards. This review was first conducted in December 2018 and re-run in March 2020 for the period between 2010 and 2020. The gross number of articles reviewed were 27, from which we identified 18 relevant references using a revised search query, and six relevant references identified using the initial query giving a total sample of 24 references. The most common source of data remains population census. The most recurrent spatial variable used for the assessment of SEV is households without basic services, while critical facilities are the most frequent spatial categories. Traditional methods have been combined with more innovative and complex methods to select and weight spatial indicators and develop indices. The Social Vulnerability Index (SoVI®) remains the benchmark for the assessment of SEV and a reference for its spatial assessment. Geographic information systems (GIS) is the most common tool for conducting a spatial assessment of SEV regarding geohazards. For future spatial assessments of SEV regarding geohazards, we recommend considering 3D spatial indexes at the microscale in the urban level involving the community in the assessments.

1 Introduction

Vulnerability is defined by United Nations (UN) as ‘the conditions determined by physical, social, economic and environmental factors or processes which increase the susceptibility of an individual, a community, assets or systems to the

Field Code Changed

Deleted: s

Deleted: the economic

Deleted: five

Deleted: 23

Deleted: high-resolution

Deleted: and

Deleted: the

Deleted: Office for Disaster Risk Reduction

Deleted: DRR

Deleted: characteristics

Deleted: and

impact of hazards' (UN, 2016). In the past, vulnerability was considered a composite factor having only two dimensions:

exposure to risk and susceptibility (Béné, 2009; Chambers, 1989). More recently, Birkmann (2013) considered three factors: exposure, susceptibility, and fragility and lack of resilience. The degree of vulnerability of a specific community is a human value judgement that highly influences management decisions (McLaughlin et al., 2002). In addition, the concept of social

vulnerability (SV) to environmental hazards involves demographic and socio-economic factors that affect community resilience (Zebardast, 2013), and this is considered a hot topic in current disaster research (Shen et al., 2018). The social and economic dimensions are only two dimensions of vulnerability to multiple stressors and shocks. These shocks include disasters due to the fragility and susceptibility of human well-being damaged by disruption to individuals (physical and mental health) and collective social systems (e.g., education, services, health) and their characteristics (e.g., age, ethnicity, disabilities)

(Birkmann et al., 2013). Social vulnerability refers to the inability of people, organisations, and societies to cope with negative impacts from different stressors to which they are exposed (Eidsvig et al., 2014; Kuhlicke et al., 2011; Myers et al., 2008; Qasim et al., 2018). Typically, this inability results from pre-existing conditions that reduce a society's ability to prepare and recover from disasters (Alcorn et al., 2013; Cutter and Finch, 2008; Eidsvig et al., 2014; Zebardast, 2013; Zhou et al., 2014).

Social vulnerability additionally identifies sensitive populations that are less prepared to respond, cope with, and recover from a disaster (Zebardast, 2013) such as low-income populations, women, pregnant women, children below 5 years, elderly above 65 years (Bereitschaft, 2017a; Zhou et al., 2014) and physically and or mentally challenged individuals (Contreras and Kienberger, 2012). Other vulnerable population groups are people with language, cultural and spatial barriers (Eidsvig et al., 2014) such as migrants (Yuan et al., 2019a), rural population, people without post-secondary education (Bereitschaft, 2017a; Cutter et al., 2003; Eidsvig et al., 2014), high-density population (Cutter et al., 2003; Eidsvig et al., 2014) and public transport

captives (Bereitschaft, 2017a).

The concept of SV is complex and dynamic, changing through the time and over space, and therefore not easily captured by a single variable (Cutter and Finch, 2008; Zebardast, 2013). It represents the multidimensionality of disasters by focusing attention on the totality of relationships in a given social situation, which, in combination with environmental forces, such as

geohazards, result in a disaster (Oliver-Smith, 2003). Social vulnerability attracts less attention by researchers because many

Deleted: circumstances of a community, system, or asset that n
it susceptible to the damaging effects of a hazard

Deleted: (Aubrecht et al., 2013)

Deleted: and

Formatted: Font: (Default) Times New Roman, 10 pt

Deleted:

challenges are implied in its quantification (Qasim et al., 2018). Power relationships that exclude certain individuals or groups from benefiting from disaster risk reduction (DRR) or post-disaster recovery efforts are examples of SV (Contreras et al., 2011). These power relationships manifest between individuals or socio-economic groups in the framework of institutions or culturally determined dialogues about stressors (Warmer et al., 2007).

Deleted:

The economic dimension of vulnerability is the predisposition for the loss of economic value from damage to physical assets (Birkmann et al., 2013) and/or business interruption (activities, services or delivery of products). The assessment of SV is orientated to cast light on the most susceptible groups of a population to be impacted by a disaster, in the spatial and temporal dimensions (Zhou et al., 2014). Another important aspect to consider is the relationship between social and economic dimensions because, according to Noy (2009), no evidence exists of a correlation between consequences of disasters, such as the number of fatalities or affected population, and GDP growth. Nevertheless, the same author indicates that the degree of damage due to a disaster will negatively influence GDP growth. Thus, Noy (2015) proposes to integrate the number of fatalities and injuries with financial damage due to a disaster using a model similar to the estimation of disability-adjusted life years (DALYs). His index accounts for the number of human years lost as a result of the damage. The spatial dimension of socio-economic vulnerability (SEV) recognises that people and groups of similar characteristics tend to occupy the same or similar areas, while the temporal dimension of SEV makes reference to people's degree of vulnerability that can change depending on age, life situation, and season (Wisner and Uitto, 2009). To include urban vulnerability assessment into a spatial plan requires strategic, technical, substantial, and procedural integration (Hizbaron et al., 2012). According to Ebert et al., (2009) a spatial indicator of SV is an SV indicator with a physical component. Housing structures and the built environment were previously included by Shuang-Ye, Brent, and Ann (2002) in a GIS-based study of SV. The link between transportation infrastructure and land use had been already studied by Clark et al. (1998). The physical conditions were considered indicative of the social ones by Rashed and Weeks (2003). Kienberger et al.,(2009) proposed a methodology for the spatial quantification of vulnerability and the identification of vulnerability units build upon the *geon* concept, which is a framework for the clustering of homogeneous spatial information. Khazai et al., (2013) developed a sector-specific vulnerability index (IVIs), which included transport dependency indicators made up by the spatial variables such as freight transport volume road and

freight transport volume railway; this index also included the spatial variable of customer proximity as part of the indicator demand dependency.

In the context of disaster risk management, and mainly for exposure and impact assessment, the accuracy and reliability of input data are two of the most important factors (Aubrecht et al., 2013). Data constraints play a key role in the results of the

5 SEV assessment, with the number of variables changing the assessment, and the inclusion of ~~additional~~ variables enhancing its precision and enabling the proper presentation of SV assessment (Gautam, 2017). Thus, the assessment of vulnerability must be based on indicators and proxy indexes (Qasim et al., 2018) that can guarantee objectivity and can provide quantitative metrics to compare different places (Cerchiello et al., 2018). Indicators and indexes are defined as single qualitative or indirect quantitative measures of a characteristic (Chen, 2016) or a real phenomenon (Fekete, 2009) resulting from systematically
10 observed facts (OECD, 2008). Indicators transform complex data into manageable units of information for performance, change, and achievement assessment (Grace and Edwin, 2009). Indicators also summarise technical information into indexes, simplifying comprehension (Simpson and Katirai, 2006). The most important factor for indicator selection is the availability of data. The lack of data can lead to reliance on variables that may not be the most accurate indicators of vulnerability (Zhou et al., 2014). Vulnerability indicators are complex measures of a part of what constitutes a community. Scientific literature has
15 identified groups of social and economic indicators, which combined with physical and land data, are useful for the vulnerability assessment of communities (King, 2001). The use of these indicators has primarily been applied to the assessment of adaptive capacity and vulnerability (Chen, 2016).

Indexes are built up with those indicators and later mapped to display the different categories of vulnerability in each
20 administrative zone, limiting the spatial dimension to this stage. The construction of an index implies selection of indicators, indicator normalization and weighting, and aggregation into an index (OECD, 2008) that must collectively represent aspects of ~~a society's ability to prepare for, deal with, and recover from a disaster~~ (Eidsvig et al., 2014). The most sensitive step for constructing an index is the weighting of indicators. This can be undertaken either using participatory approaches such as the analytic hierarchy process (AHP), the budget allocation process, statistical assessment like the principal component analysis
25 (PCA), or factor analysis (FA) (Eidsvig et al., 2014; OECD, 2008). Weighting individual indicators is a major challenge for

Deleted: s

Deleted: more

Deleted: the

Deleted: ,

constructing a composite indicator for vulnerability (Adger et al., 2004; Zebardast, 2013). The objectives of indicators weighting are first, to investigate any correlation among indicators to detect overlapping information and second, to select a suitable weighting and aggregation approach for the final index calculation. Different weightings show varied spatial vulnerability patterns (Papathoma-Kohle et al., 2019); however, independent of the method applied, after comparing 106 studies for index construction with respect to risk assessment, Beccari (2016) found that the most common approach used (41.5%) was the 'equal weights' method. Eventually, the accuracy of SV assessment lies on the accuracy of input data (Yuan et al., 2019a) and not on the weighting method. After being weighted, indicators can be aggregated using additive, multiplicative, or decision rule models (Eidsvig et al., 2014). The method of aggregation is one of the most pressing problems in developing composite vulnerability indices (Rygel et al., 2006).

Composite indicators have been commonly employed by researchers, planners, and disaster managers for vulnerability assessments (Yuan et al., 2019a). Cutter, Boruff and Shirley (2003) have constructed an index of SV called (SoVI®) for environmental hazards in the United States using a factor analytic approach computed in a summary score based on an additive model. In the framework of the Methods for the Improvement of Vulnerability Assessment in Europe (MOVE) project, variables were grouped into single (Vinchon et al., 2011) and composite indicators. In the case study area of Salzburg ([Austria](#)), an expert-based approach was chosen, and several experts were asked to allocate weights according to the contribution of each variable to the vulnerability of floods (Contreras and Kienberger, 2011). Other composite indicators useful for the vulnerability assessment are the Prevalent Vulnerability Index (Cardona, 2005), Environmental Sustainability Index (Esty et al., 2005), and Human Development Index (UNDP, 2010). All these indexes face challenges when assessing vulnerability indicators, such as: ranking socio-economic data on an interval scale, dealing with temporal aspects (day-night changes), choosing the most suitable data resolution to avoid the 'modifiable areas unit problem' (MAUP) (Openshaw, 1983), deciding how to allocate a meaningful value to socio-economic variables, and how these aspects together affect the vulnerability assessment of each case study areas (McLaughlin et al., 2002). The compilation of all of the SV indicators used through time was undertaken by Fatemi, Ardalan, Aguirre, Mansouri and Mohammadfam (2017); however, they neither included the spatial dimension in their systematic review nor focused exclusively on geohazard as in this research.

Quantitative measures to develop indicators can be spatially explicit and based on spatial variables, such as location, area, range, distance, direction, spatial geometries, and patterns (Unwin, 1996), spatial connectivity, mobility (Béné, 2009), isolation, diffusion, distribution, spatial association, spatial interaction, spatial evolution, spatial synthesis and scale of the affected area, and surroundings (Béné, 2009; Buzai and Villerías Alarcón, 2018; Contreras et al., 2013; Meentemeyer, 1989).

5 The geographic patterns in vulnerability can increase due to spatial interactions; while additional patterns within these components may be related to the nature of vulnerability stemming from a specific hazard (Amram et al., 2011). The main aim of this research is to elucidate the state of the art in data sources, spatial variables, indicators, methods, indexes, and tools for the assessment of the SEV related to geohazards in urban environments. Geohazards can be endogenic such as earthquakes, tsunamis, and volcanic eruptions and exogenic, such as landslides, soil erosion, and land degradation. We particularly focus on
10 these phenomena for two reasons: first, geohazards are the natural phenomena that have produced the highest quantity of losses in recent years in the urban environments, particularly earthquakes, and, second, because geohazards are the phenomena addressed by the institutions involved in the present research.

The Indian Ocean tsunami in 2004, as a result of its large impact area, reignited the research community's interest in spatial
15 vulnerability analyses, illuminating the problems faced by low-income population after disasters (Fekete, 2012). This approach was aligned with the Hyogo Framework for Action (UNISDR, 2007), and confirmed by Gautam (2017), who notes that after 2005 a focus on construction and mapping of the SV index intensified. Thus, the use of geographic information systems (GIS) to collect and process data related to hazards and vulnerability was found very suitable (Fekete, 2012). Major earthquakes that
20 occurred during the same period as this systematic review (2010-2020), e.g. Chile (2010), New Zealand (2010 and 2011), Nepal (2015), Mexico (2017), Albania (2019), and Croatia (2020) demonstrate the vulnerability of urban areas to seismic damages (Armaş et al., 2017).

This research reviews case study areas, data sources, spatial variables, indicators, methods, indexes, and tools used in the spatial assessment of SEV vulnerability by different authors in the period between 2010 and 2020. This systematic review
25 aims to evaluate the literature to identify patterns and trends, as well as research gaps, to recommend new research areas. This

Deleted: internal

Formatted: Font: 10 pt, Font color: Auto

Deleted: external

Deleted: in Chile

Deleted: in

article aspires to guide scientists who want to perform any spatial assessment of SEV vulnerability. Socio-economic vulnerability is dynamic and changes across spatial and temporal scales, depending on demographic, geographic, economic, and cultural factors. Hence, no one-size-fits-all approach exists to measure and reduce SV (Zhou et al., 2014). This paper is divided into six sections. The introduction is the first section and includes a literature review. The second section, on methods, elaborates on the criteria for selecting the articles that comprise the systematic review and the format of the presentation of results. The third section focuses on the results. The fourth section includes discussion of the results supported by literature, and, the fifth section contains conclusions with recommendations proposed in the sixth section.

2 Methods

A systematic review searches for, appraises, and synthesises research evidence (Grant and Booth, 2009). In the present research, the systematic review was conducted to elucidate the state of the art of data sources, spatial variables, indicators, methods, indexes and tools for the spatial assessment of the SEV related to geohazards, which we consider is covered in the period between 2010 and 2020. Thus, the main research question is: what is the state of the art in the spatial assessment of SEV to geohazards in urban environments?

This review was conducted in December 2018 and re-run during the revision process in March 2020. For this research, Clarivate Analytics and Scopus/Elsevier were the sources of selected literature given their functionalities to run the search query. We limited the query to articles published in academic journals because they typically are rigorous in the selection of their publications and therefore contain a complete and accurate description of methodologies and consistent results. The terms selected for the search query refer to vulnerability in the socio-economic dimension, the spatial variables listed by Meentemeyer (1989), Béné (2009), Contreras et al. (2013) and Buzai and Villerías Alarcón (2018) and the aforementioned endogenic and exogenic geohazards. Based on several screenings, to refine the search strategy, we opted to exclude terms that were not related to geohazards and were recurring in the titles, abstracts and keywords of the resulting references. The final set of terms included and excluded in the search query are listed in Table 1 and the scheme of the methodology applied is depicted in Figure 1.

Deleted: more

Deleted: more

Formatted: Not Highlight

Formatted: Not Highlight

Deleted: external

Deleted: internal

D	Q	SEARCH TERMS
Clarivate analytics	TOPIC	"social vulnerability" OR "economic vulnerability" OR "socioeconomic vulnerability" OR "socio-economic vulnerability"
		AND
	TOPIC	"area" OR "distance" OR "range" OR "distance" OR "direction" OR "spatial geometries" OR "patterns" OR "spatial connectivity" OR "isolation" OR "diffusion" OR "spatial association" OR "scale" OR "accessibility" OR "network" OR "cluster"
		AND
	TOPIC	"earthquakes" OR "tsunamis" OR "volcanic eruptions" OR "landslides" OR "soil erosion" OR "land degradation"
Scopus/Elsevier		NOT
	TOPIC	"climate change" OR "ecological" OR "drought" OR "resilience" OR "debris" OR "epidemiological" OR "substance" OR "behavioural" OR "evacuation" OR "recovery" OR "pollution" OR "leptospirosis" OR "violence" OR "illness" OR "disease" OR "heat" OR "crisis" OR "conflict" OR "deaths" OR "obesity" OR "criminal" OR "chemical" OR "symptoms" OR "syndrome" OR "food insecurity" OR "air pollution" OR "stress" OR "diabetes" OR "depressive" OR "alcohol" OR "cancer" OR "drugs" OR "palm oil" OR "tobacco" OR "smoke" OR "storm" OR "psychometric" OR "cocaine" OR "toxic" OR "palliative" OR "therapy" OR "HIV" OR "dengue" OR "ecosystem" OR "rheumatoid" OR "arthritis" OR "nutritional" OR "malaria" OR "resources" OR "sexual activity" OR "sexual health"
	Article title, abstract, keywords	(TITLE-ABS-KEY ("social vulnerability*" AND "economic vulnerability*") AND TITLE-ABS-KEY ("socioeconomic vulnerability*") AND TITLE-ABS-KEY ("area" OR "distance" OR "range" OR "distance" OR "direction" OR "spatial geometries" OR "patterns" OR "spatial connectivity" OR "isolation" OR "diffusion" OR "spatial association" OR "scale" OR "accessibility" OR "network" OR "cluster") AND TITLE-ABS-KEY ("earthquakes" OR "tsunamis" OR "volcanic eruptions" OR "landslides" OR "soil erosion" OR "land degradation") AND NOT TITLE-ABS-KEY ("climate change" OR "ecological" OR "drought" OR "resilience" OR "debris" OR "epidemiological" OR "substance" OR "behavioral" OR "evacuation" OR "recovery" OR "pollution" OR "leptospirosis" OR "violence" OR "illness" OR "disease"))) AND DOCTYPE (ar) AND PUBYEAR > 2009 AND PUBYEAR <2021

D: Database

Q: Query

Table 1. Terms included and excluded to identify relevant literature references.




Figure 1. Methodology applied for the systematic literature review.

Deleted:

Deleted:

The findings will be presented in the results section in tables related to selected references, data sources, spatial variables, indicators, methods, spatial indexes, and tools. Table 2 is structured in four columns, namely author, year, research objective, geohazard addressed, and country where the case study area of the paper is located. The authors are listed from the most recent reference to the oldest one. Tables 3, 4, 5, 6 and 7 are structured mainly in two columns: the first column lists data sources, spatial variables, indicators, methods and indexes respectively. The second column contains the authors and the year of their publications, in which the mentioned topics are addressed. Moreover, the references in these tables are also listed in reverse chronological order. The second column in Table 3 includes, in some cases, specific details of the data source used by the authors. Table 8 includes three columns, namely, method, software and authors.

3 Results

The gross number of articles identified using the search query were 29, having two matching references in Clarivate Analytics and Scopus/Elsevier: Kurnianto et al., (2019) and Eidsvig (2014). Thus, eventually, we identified 27 references. Despite the precise search query, 11 references were discarded due to reasons explained as follows. In chronological order, the first reference discarded was Papathoma-Kohle et al., (2019) because they use variables in the physical dimension, rather than socio-economic one. Two references from Yuan et al., (2019a, b) were identified by the search query as using the same method for the spatial assessment of SEV; so, we decided to select only one of them. Zhang and Huang (2018) address the topic of SV but not its spatial assessment, while Shen et al. (2018) focused on calculating the impact of disasters, rather than estimating SEV. The paper written by Goncalves, M., & Vizintim, M. F. B. (2017) was written in Portuguese, which none of the authors is proficient. Postiglione et al., (2016) promote a culture of seismic risk prevention, rather than to estimate SEV due to earthquakes. Alcántara-Ayala and Oliver-Smith (2014) present the activities undertaken by the ICL Latin -American network (ICL LAB) related to capacity building to reduce risk due to landslides, with no specific emphasis on SEV. Khazai et al., (2014), in their book chapter, concentrate on modelling shelter needs and health impacts caused by earthquakes. Vilches et al. (2014) evaluate the socio-environmental effects of the 27/10/2010 tsunami in Chile, considering the SEV among other aspects, but they do not make use of any spatial variable, indicator, or index, which is similar to the vulnerability assessment relating

Deleted: 7

Deleted: ,

Formatted: Check spelling and grammar

to a tsunami in the Town of Tirua (Chile) undertaken by Jaque Castillo et al.,(2013). Six references from the previous search query carried out in 2018, and not identified in the refined search query, were included in the list given their relevance due to the geohazards and spatial variables, indicators, and indexes that they address. The 24 references finally reviewed are listed in Table 2.

Deleted: Five

Deleted: 23

5

AUTHOR	YEAR	RESEARCH OBJECTIVE	HAZARD	COUNTRY
Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W.	2020	To introduce a model for spatial multi-hazards risk assessment applied to Dharan, Nepal.	Earthquakes, floods and landslides	Nepal
Kurnianto, F. A., Ikhsan, F. A., Apriyanto, B., & Nurdin, E. A. (2019)	2019	To assess the level of vulnerability for an earthquake disaster in Lembang district, an area in West Java that includes the Bandung basin.	Earthquakes	Indonesia
Muir, J. A., Cope, M. R., Angeningsih, L. R., Jackson, J. E., & Brown, R. B.	2019	To explore whether return migration, compared to other migration options, results in superior improvements to mental health in the context of disasters.	Volcanic eruptions	Indonesia
Rezaei-Malek, M., Torabi, S. A., & Tavakkoli-Moghaddam, R.	2019	To prioritize disaster-prone areas that are known as potential demand points (PDPs) given their vulnerability under large-scale earthquakes	Earthquakes	Iran
Yuan, H. H., Gao, X. L., & Qi, W. (2019)	2019	To provide high spatial-temporal resolution information on vulnerable populations and population vulnerability using dasymetric population mapping with vulnerability index	Earthquakes	China
Alizadeh, M., Alizadeh, E., Kotenae, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L.	2018	To apply an artificial neural network (ANN) and geographic information system (GIS) for estimating the social vulnerability to earthquakes in the Tabriz city, Iran.	Earthquakes	Iran

Deleted: .

Deleted: .

Deleted: which

Deleted: ¶

Deleted: .

AUTHOR	YEAR	RESEARCH OBJECTIVE	HAZARD	COUNTRY
Qasim, S., Qasim, M., Shrestha, R. P., & Khan, A. N.	2018	To define the socio-economic determinants of landslide risk perception in Murree hills of Pakistan.	Landslides	Pakistan
Ponce-Pacheco, A. B., & Novelo-Casanova, D. A.	2018	To estimate the levels of vulnerability and risk to floods, earthquakes and subsidence of Valle de Chalco Solidaridad (VCS) in Mexico.	Earthquakes, floods and subsidence	Mexico
Armaş, I., Toma-Danila, D., Ionescu, R., & Gavriş, A.	2017	To develop an overall vulnerability index to seismic hazard based on a spatial approach applied to Bucharest, Romania.	Earthquakes	Romania
Bereitschaft, B.	2017	To explore inequity in neighbourhood walkability at the micro-scale level related to social vulnerability in terms of imageability, enclosure, human scale, transparency, complexity, tidiness and safety in Pittsburgh Streetscapes	Not walkability	USA
Gautam, D.	2017	To investigate social vulnerability to natural hazards in Nepal at district level.	Droughts, earthquakes, epidemics, floods and landslides,	Nepal
Chen, Y.	2016	To develop a set of valid and reliable indicators to evaluate the regional land subsidence disaster vulnerability in the Xixi-Chengnan area, in China.	Landslides	China
Garcia, R. A. C., Oliveira, S. C., & Zezere, J. L.	2016	To apply dasymetric cartography to improving population spatial resolution and to assess the potentially exposed population over large areas to deep rotational landslides and compare the results with those obtained with basic census units as the data source	Landslides	Portugal
Maharani, Y. N., Lee, S., & Ki, S. J.	2016	To propose the use of Self-Organizing Maps (SOM) approach to conducting the social	Volcanic eruptions	Indonesia

Deleted: .

Deleted:

Deleted: .

Deleted: .

Deleted: .

AUTHOR	YEAR	RESEARCH OBJECTIVE	HAZARD	COUNTRY
		vulnerability assessment around the Merapi volcano		
Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P.	2015	To assess the social vulnerability of informal settlements in Iquique and Puerto Montt in Chile	Earthquakes, floods, landslides and Tsunami	Chile
Ley-García, J., Denegri de Dios, F. M., & Ortega Villa, L. M.	2015	The aim is to identify visibility, invisibility and amplification of hazardscape perception in the city of Mexicali, Baja California, Mexico	Earthquake Landslide Tsunami Volcano Cyclone Thunderstorm Heavy rainfall Flood hail Snow-freeze Strong wind Drought Cold wave Heat wave	Mexico
Eidsvig, U. M. K., McLean, A., Vangelsten, B. V., Kalsnes, B., Ciurean, R. L., Argyroudis, S., . . . Kaiser, G.	2014	To propose a methodology to estimate socio-economic vulnerability to landslides at the local to regional scale using an indicator-based model	Landslides	Andorra, France, Greece, Norway, and Romania
Toké, N. A., Boone, C. G., & Arrowsmith, J. R.	2014	To construct a relative social vulnerability index classification for Los Angeles to examine the social condition within regions of significant seismic hazard, including areas regulated as Alquist-Priolo (AP) Act earthquake fault zones	Earthquakes landslides and wildfires	USA
Walker, B. B., Taylor-Noonan, C., Tabbernor,	2014	To model geophysical processes and identification of socio-economically	Earthquakes	Canada

Deleted: .


Deleted: .

AUTHOR	YEAR	RESEARCH OBJECTIVE	HAZARD	COUNTRY
A., McKinnon, T. B., Bal, H., Bradley, D., . . . Clague, J. J.		disadvantaged populations in Victoria, British Columbia		
Alcorn, R., Panter, K. S., & Gorsevski, P. V.	2013	To evaluate the spatial impact of a possible future eruption using a GIS-based volcanic hazard tool and to assess the social and economic vulnerabilities of the area at risk	Volcanic eruption	USA
Aubrecht, C., Özceylan, D., Steinnocher, K., & Freire, S.	2013	To review available multi-level geospatial information and modelling approaches from local to global scales, that could serve practitioners and researchers in disaster-related zones	Tsunami, floods	Austria Portugal Turkey USA
Zebardast, E.	2013	To develop a model that combines hybrid factor analysis and analytic network process (F'ANP) for constructing a composite social vulnerability index (SOVI)	Earthquakes	Iran
Hizbaron, D. R., Baiquni, M., Sartohadi, J., & Rijanta, R.	2012	To assess urban vulnerability due to seismic hazard using a risk based spatial plan	Earthquakes	Indonesia
Zeng, J., Zhu, Z. Y., Zhang, J. L., Ouyang, T. P., Qiu, S. F., Zou, Y., & Zeng, T.	2012	To introduce a new method to assess social vulnerability for county-scale regions using population density, based on land use	Landslides	China


Table 2. Articles identified and selected by the systematic review.

The most recurrent geohazards addressed among the selected papers are earthquakes, followed by landslides, volcanic eruptions, tsunamis, and subsidence, detailed information about the number of literature references that tackle each hazard is depicted in Figure 2. None of the references deals with soil erosion, nor land degradation. Case study areas selected from this set of papers are frequently located in Indonesia, China, Iran, and the USA, detailed information about the number of literature references that has case study areas on these countries can be appreciated in Figure 3. From the set of selected papers, the most common sources of data are the population census, followed by satellite images, field observations, disaster databases, surveys,

aerial photographs, and land use and land cover (LULC) maps. Other authors used [high definition \(HD\) videos](#), [orthophotos](#), [photographs](#), landslide susceptibility maps, and volunteered geographic information (VGI). The complete set of data sources identified in this systematic review are listed in Table 3.



Deleted: orthophotos,



Deleted:

5 Figure 2. Number of literature references in the systematic review that addresses each geohazard.





Figure 3. Number of study areas per country addressed in the references identified through the systematic literature review.



Deleted:

Deleted: literature references in the systematic review that add case ...

DATA SOURCES	AUTHORS
Census data	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
Nepal census	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)
City office of Dharan	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
National Institute of Statistics and Geography	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)
Municipal Government of Valle de Chalco Solidaridad	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)
Secretariat of Social Development of Mexico	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)
CBS 2011 Census	Gautam, D. (2017)
Xishan and Huishan Statistical Yearbook 2008	Chen, Y. (2016)

DATA SOURCES	AUTHORS
Population and Housing Census 2010	Lin, W.-Y., & Hung, C.-T. (2016)
National Census 2011	Garcia, R. A. C., Oliveira, S. C., & Zezere, J. L. (2016)
Statistics of Sleman Regency https://slemankab.bps.go.id/	Maharani, Y. N., Lee, S., & Ki, S. J. (2016)
National census of population and VI of housing	Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015).
2000 U.S. Census Bureau	Toké, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)
Statistical Office of Baden-Wuerttemberg	Khazai, B., Merz, M., Schulz, C., & Borst, D. (2013)
Regional Planning Board	Hizbaron, D. R., Baiquni, M., Sartohadi, J., & Rijanta, R. (2012)
Statistical Bureau	Hizbaron, D. R., Baiquni, M., Sartohadi, J., & Rijanta, R. (2012)
Armaş, I., Toma-Danila, D., Ionescu, R., & Gavriş, A. (2017)	
Garcia, R. A. C., Oliveira, S. C., & Zezere, J. L. (2016)	
Walker, B. B., Taylor-Noonan, C., Tabbernor, A., McKinnon, T. B., Bal, H., Bradley, D., . . . Clague, J. J. (2014)	
Satellite images	
WorldView-3	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
ASTER-DEM	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
PERSIANN-CDR	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
Google Earth satellite images	Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015)
GDEM-ASTER	Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015)
LANDSAT	Toké, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)

Deleted: Satellite images

	DATA SOURCES	AUTHORS
	LANDSAT™	Aubrecht, C., Özceylan, D., Steinnocher, K., & Freire, S. (2013)
	SPOT	Zeng, J., Zhu, Z. Y., Zhang, J. L., Ouyang, T. P., Qiu, S. F., Zou, Y., & Zeng, T. (2012)
	IKONOS	Aubrecht, C., Özceylan, D., Steinnocher, K., & Freire, S. (2013)
	NDVI	Aubrecht, C., Özceylan, D., Steinnocher, K., & Freire, S. (2013)
Field Observations	Alizadeh, M., Alizadeh, E., Kotenae, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L. (2018)	
	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)	
	Garcia, R. A. C., Oliveira, S. C., & Zezere, J. L. (2016)	
	Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015)	
	Hizbaron, D. R., Baiquni, M., Sartohadi, J., & Rijanta, R. (2012)	
Disaster Databases	Indonesian Disaster Data Information (DIBI)	Maharani, Y. N., Lee, S., & Ki, S. J. (2016)
	http://dibi.bnpb.go.id/dibi/	
	Risk Atlas of the Municipality of Mexicali 2011	Ley-García, J., Denegri de Dios, F. M., & Ortega Villa, L. M. (2015)
	Desinventar Data Base	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)
Surveys	Muir, J. A., Cope, M. R., Angeningsih, L. R., Jackson, J. E., & Brown, R. B. (2019)	
	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)	
	Qasim, S., Qasim, M., Shrestha, R. P., & Khan, A. N. (2018).	
Aerial Photograph	Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015)	
	Toké, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)	
LULC maps	CORINE	Aubrecht, C., Özceylan, D., Steinnocher, K., & Freire, S. (2013)
	HR Soil sealing layer	Aubrecht, C., Özceylan, D., Steinnocher, K., & Freire, S. (2013)
Population datasets	GPW/GPWv4	Aubrecht, C., Özceylan, D., Steinnocher, K., & Freire, S. (2013)

DATA SOURCES	AUTHORS
GRUMP	Aubrecht, C., Özceylan, D., Steinnocher, K., & Freire, S. (2013)
HD video	Bereitschaft, B. (2017)
Orthophotos	Armaş, I., Toma-Danila, D., Ionescu, R., & Gavriş, A. (2017)
Photographs	Bereitschaft, B. (2017)
Landslide susceptibility map (pixel terrain unit)	Garcia, R. A. C., Oliveira, S. C., & Zezere, J. L. (2016)
VGI	Aubrecht, C., Özceylan, D., Steinnocher, K., & Freire, S. (2013)

Table 3. Data sources for the spatial assessment of socio-economic vulnerability assessments.

The most common spatial variables used for the spatial assessment of SEV between 2010 and 2020 are households without basic services (piped water connection, electricity, sewerage infrastructure, cell phone, or landline), location, critical facilities (fire stations, medical emergency services, medical facilities, and hospitals), distance from faults/causative faults, precarious housing (low quality and/or precarious external walls, roofing, and floors), the total area of occupied space in the residences, and the presence of schools. The complete set of spatial variables identified in this systematic review are listed in Table 4.

SPATIAL VARIABLES	AUTHORS
Households without piped water connection, electricity, sewerage infrastructure, cell phone or landline	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)
	Gautam, D. (2017)
	Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015)
	Zebardast, E. (2013)
Location	Kurnianto, F. A., Ikhsan, F. A., Apriyanto, B., & Nurdin, E. A. (2019)
	Muir, J. A., Cope, M. R., Angeningsih, L. R., Jackson, J. E., & Brown, R. B. (2019)
	Qasim, S., Qasim, M., Shrestha, R. P., & Khan, A. N. (2018)
	Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015)

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto, German (Austria), Not Highlight

Formatted: Font: (Default) Times New Roman, 10 pt, German (Austria)

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto, German (Austria), Not Highlight

Formatted: Font: (Default) Times New Roman, 10 pt, German (Austria)

Deleted: Orthophoto

Deleted: ¶
¶

SPATIAL VARIABLES	AUTHORS
Critical facilities (fire stations, hospitals, health services, medical emergency services, medical facilities, etc.)	Rezaei-Malek, M., Torabi, S. A., & Tavakkoli-Moghaddam, R. (2019)
	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)
	Eidsvig, U. M. K., McLean, A., Vangelsten, B. V., Kalsnes, B., Ciurean, R. L., Argyroudis, S., . . . Kaiser, G. (2014)
	Alcorn, R., Panter, K. S., & Gorsevski, P. V. (2013)
	Zeng, J., Zhu, Z. Y., Zhang, J. L., Ouyang, T. P., Qiu, S. F., Zou, Y., & Zeng, T. (2012)
Distance from faults/ causative faults	Rezaei-Malek, M., Torabi, S. A., & Tavakkoli-Moghaddam, R. (2019)
	Hizbaron, D. R., Baiquni, M., Sartohadi, J., & Rijanta, R. (2012)
Household with low quality and/or precarious external walls, roofing and floors	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
	Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015)
Total area of occupied space in the residences	Armaş, I., Toma-Danila, D., Ionescu, R., & Gavriş, A. (2017)
	Zeng, J., Zhu, Z. Y., Zhang, J. L., Ouyang, T. P., Qiu, S. F., Zou, Y., & Zeng, T. (2012)
Schools	Alcorn, R., Panter, K. S., & Gorsevski, P. V. (2013)
	Zeng, J., Zhu, Z. Y., Zhang, J. L., Ouyang, T. P., Qiu, S. F., Zou, Y., & Zeng, T. (2012)
Families occupying rented houses	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
Households per housing unit	Zebardast, E. (2013)
Households with >1 family	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
City blocks	Yuan, H. H., Gao, X. L., & Qi, W. (2019)
Displaced, moved home, in transition, moved on	Muir, J. A., Cope, M. R., Angeningsih, L. R., Jackson, J. E., & Brown, R. B. (2019)
Distance to volcanoes	Kurnianto, F. A., Ikhsan, F. A., Apriyanto, B., & Nurdin, E. A. (2019)
Availability of evacuation roads	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)
Active uses/occupied storefronts	Bereitschaft, B. (2017)
Building color & design variety	Bereitschaft, B. (2017)
Building height & setback	Bereitschaft, B. (2017)

Deleted: (

Formatted: Font: (Default) Times New Roman, 10 pt, Not Highlight

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Black

Deleted: Muir et al., 2019), J. A., Cope, M. R., Angeningsih, L. R., Jackson, J. E., & Brown, R. B. (2019)

Formatted: Font: (Default) Times New Roman, 10 pt,

Formatted: Font: (Default) +Headings (Times New Roman, 10 pt

Formatted: Font: (Default) +Headings (Times New Roman, 10 pt

Formatted: Font: (Default) +Headings (Times New Roman, 10 pt,

Formatted: Font: (Default) +Headings (Times New Roman, 10 pt,

Formatted: Font: (Default) +Headings (Times New Roman, 10 pt

Formatted: Font: (Default) +Headings (Times New Roman, 10 pt

Formatted: Font: (Default) +Headings (Times New Roman, 10 pt

SPATIAL VARIABLES	AUTHORS
Housing type	Eidsvig, U. M. K., McLean, A., Vangelsten, B. V., Kalsnes, B., Ciurean, R. L., Argyroudis, S., . . . Kaiser, G. (2014)
Percentage of households with public assistance	Toke, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)
Percent of workers with a long commute	Toke, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)
Travel barriers to the trauma centres	Walker, B. B., Taylor-Noonan, C., Tabbernor, A., McKinnon, T. B., Bal, H., Bradley, D., . . . Clague, J. J. (2014)
Travel distance to trauma centres	Walker, B. B., Taylor-Noonan, C., Tabbernor, A., McKinnon, T. B., Bal, H., Bradley, D., . . . Clague, J. J. (2014)
Travel time to trauma centres	Walker, B. B., Taylor-Noonan, C., Tabbernor, A., McKinnon, T. B., Bal, H., Bradley, D., . . . Clague, J. J. (2014)
Walking time to trauma centres	Walker, B. B., Taylor-Noonan, C., Tabbernor, A., McKinnon, T. B., Bal, H., Bradley, D., . . . Clague, J. J. (2014)
Land use	Alcorn, R., Panter, K. S., & Gorsevski, P. V. (2013)
Housing with bathroom	Zebardast, E. (2013)
Housing with kitchen	Zebardast, E. (2013)
Migration status	Muir, J. A., Cope, M. R., Angeningsih, L. R., Jackson, J. E., & Brown, R. B. (2019)
Road type	Alcorn, R., Panter, K. S., & Gorsevski, P. V. (2013)
Spatial distribution of cell phone subscribers	Aubrecht, C., Özceylan, D., Steinnocher, K., & Freire, S. (2013)
Distance to hospital	Zeng, J., Zhu, Z. Y., Zhang, J. L., Ouyang, T. P., Qiu, S. F., Zou, Y., & Zeng, T. (2012)
Distance to road network	Hizbaron, D. R., Baiquni, M., Sartohadi, J., & Rijanta, R. (2012)
Distance to trauma centres	Walker, B. B., Taylor-Noonan, C., Tabbernor, A., McKinnon, T. B., Bal, H., Bradley, D., . . . Clague, J. J. (2014)
Distribution of urban greenspace	Toké, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)

SPATIAL VARIABLES	AUTHORS
Industry land, Office land and commercial and residential land ,	Zeng, J., Zhu, Z. Y., Zhang, J. L., Ouyang, T. P., Qiu, S. F., Zou, Y., & Zeng, T. (2012)
Population dependent on the land for the primary source of income	Eidsvig, U. M. K., McLean, A., Vangelsten, B. V., Kalsnes, B., Ciurean, R. L., Argyroudis, S., . . . Kaiser, G. (2014).
Road network	Zeng, J., Zhu, Z. Y., Zhang, J. L., Ouyang, T. P., Qiu, S. F., Zou, Y., & Zeng, T. (2012)

Table 4. Spatial variables for socio-economic vulnerability assessments.

Population density, housing density, hospital beds per 1,000 people, and living space per person are the most frequent spatial indicators of SEV. Global Moran’s I and local indicators of spatial association (LISA), which are traditional indicators in the spatial assessment, were also identified in this systematic research. We also found indicators such the access to environmental amenities and medical facilities, ~~mobility,~~ employed/unemployed density, ~~and~~ literate people density among others. The complete set of spatial indicators identified in this systematic review are listed in Table 5.

SPATIAL INDICATORS	AUTHORS
Population density (women/men density)	Alizadeh, M., Alizadeh, E., Kotenae, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L. (2018)
	Kurnianto, F. A., Ikhsan, F. A., Apriyanto, B., & Nurdin, E. A. (2019)
	Yuan, H. H., Gao, X. L., & Qi, W. (2019)
	Alizadeh, M., Alizadeh, E., Kotenae, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L. (2018)
	Armaş, I., Toma-Danila, D., Ionescu, R., & Gavriş, A. (2017)
	Chen, Y. (2016)
	Maharani, Y. N., Lee, S., & Ki, S. J. (2016)
	Eidsvig, U. M. K., McLean, A., Vangelsten, B. V., Kalsnes, B., Ciurean, R. L., Argyroudis, S., . . . Kaiser, G. (2014)
	Toké, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)
	Hizbaron, D. R., Baiquni, M., Sartohadi, J., & Rijanta, R. (2012)

Deleted: .

Deleted: and park space

Deleted: basic census units (BCU) target zones, and population

Deleted: floating population,

SPATIAL INDICATORS	AUTHORS
	Zeng, J., Zhu, Z. Y., Zhang, J. L., Ouyang, T. P., Qiu, S. F., Zou, Y., & Zeng, T. (2012)
Housing density	Alizadeh, M., Alizadeh, E., Kotenae, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L. (2018)
	Armaş, I., Toma-Danila, D., Ionescu, R., & Gavriş, A. (2017)
	Toké, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)
Hospital beds per 1,000 people	Eidsvig, U. M. K., McLean, A., Vangelsten, B. V., Kalsnes, B., Ciurean, R. L., Argyroudis, S., . . . Kaiser, G. (2014)
	Zebardast, E. (2013)
<u>Mobility</u>	<u>Yuan, H. H., Gao, X. L., & Qi, W. (2019)</u>
	<u>Bereitschaft, B. (2017)</u>
Living space pp	Armaş, I., Toma-Danila, D., Ionescu, R., & Gavriş, A. (2017)
	Zebardast, E. (2013)
Degree of population agglomeration	Yuan, H. H., Gao, X. L., & Qi, W. (2019)
Floating population	Yuan, H. H., Gao, X. L., & Qi, W. (2019)
Spatial distribution	Yuan, H. H., Gao, X. L., & Qi, W. (2019)
Employed/ Unemployed density	Alizadeh, M., Alizadeh, E., Kotenae, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L. (2018).
Household overcrowding	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)
Literate people density	Alizadeh, M., Alizadeh, E., Kotenae, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L. (2018)
<u>Businesses density</u>	<u>Bereitschaft, B. (2017)</u>
<u>Complexity</u>	<u>Bereitschaft, B. (2017)</u>
<u>Enclosure</u>	<u>Bereitschaft, B. (2017)</u>
<u>Human scale</u>	<u>Bereitschaft, B. (2017)</u>
<u>Imageability</u>	<u>Bereitschaft, B. (2017)</u>
<u>Safety & sensations</u>	<u>Bereitschaft, B. (2017)</u>
<u>Tidiness</u>	<u>Bereitschaft, B. (2017)</u>
<u>Traffic density</u>	<u>Bereitschaft, B. (2017)</u>
<u>Transparency</u>	<u>Bereitschaft, B. (2017)</u>
BCU target zones	Garcia, R. A. C., Oliveira, S. C., & Zezere, J. L. (2016)

Formatted: Font: (Default) Times New Roman, 10 pt, Sp (Spain), Not Highlight

Formatted: Font: (Default) Times New Roman, 10 pt, Sp (Spain), Not Highlight

Formatted: Font: (Default) Times New Roman, 10 pt, Sp (Spain), Not Highlight

Formatted: Font: (Default) Times New Roman, 10 pt, Sp (Spain), Not Highlight

Formatted: Font: (Default) Times New Roman, 10 pt, Sp (Spain), Not Highlight

Formatted: Font: (Default) Times New Roman, 10 pt, Sp (Spain), Not Highlight

Formatted: Font: (Default) Times New Roman, 10 pt, Sp (Spain), Not Highlight

Formatted: Font: (Default) Times New Roman, 10 pt, Sp (Spain), Not Highlight

SPATIAL INDICATORS	AUTHORS
BCU population	Garcia, R. A. C., Oliveira, S. C., & Zezere, J. L. (2016)
Density of agricultural/industrial production	Chen, Y. (2016)
Farming density	Chen, Y. (2016)
GDP density	Chen, Y. (2016)
Investment density of fixed assets	Chen, Y. (2016)
Global Moran's I	Ley-García, J., Denegri de Dios, F. M., & Ortega Villa, L. M. (2015)
LISA	Ley-García, J., Denegri de Dios, F. M., & Ortega Villa, L. M. (2015)
Access to environmental amenities (Park space, open spaces and walkable neighborhoods)	Toké, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)
Access to medical facilities	Walker, B. B., Taylor-Noonan, C., Tabbernor, A., McKinnon, T. B., Bal, H., Bradley, D., . . . Clague, J. J. (2014)
Infrastructure dependance	Toke, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)
Walkability	Toké, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)

Table 5. Spatial indicators for socio-economic vulnerability assessments.

Results extracted from the literature indicate that that the most common methods in the last 10 years for the reduction of variables is principal component analysis (PCA) and for indicators weighting is Analytic Hierarchy Process (AHP). The use of artificial neural networks (ANN) has been gaining ground in the last 10 years as a method for the spatial assessment of SEV. Other methods include dasymetric population mapping, factor analysis (FA), ordinal logistic regression (OLR), spatial multi-criteria evaluation (SMCE), and analytic network process (ANP). We also found hybrid methods that combine FA and ANP known as F'ANP, and others that combine fuzzy numbers with ANP, DEMATEL and PROMETHEE II (F-ADP). Other methods were simpler, such as an overlay analysis. The complete set of methods used by authors and identified in this systematic review is listed in Table 6.

METHODS		AUTHORS
PCA		Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
		Armaş, I., Toma-Danila, D., Ionescu, R., & Gavriş, A. (2017)
		Maharani, Y. N., Lee, S., & Ki, S. J. (2016)
		Toké, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)
		Alcorn, R., Panter, K. S., & Gorsevski, P. V. (2013)
AHP		Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
		Alizadeh, M., Alizadeh, E., Kotenaee, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L. (2018)
		Armaş, I., Toma-Danila, D., Ionescu, R., & Gavriş, A. (2017)
		Walker, B. B., Taylor-Noonan, C., Tabbernor, A., McKinnon, T. B., Bal, H., Bradley, D., . . . Clague, J. J. (2014)
ANN		Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
		Alizadeh, M., Alizadeh, E., Kotenaee, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L. (2018)
		Maharani, Y. N., Lee, S., & Ki, S. J. (2016)
Dasymeric mapping	population	Yuan, H. H., Gao, X. L., & Qi, W. (2019)
		Garcia, R. A. C., Oliveira, S. C., & Zezere, J. L. (2016)
FA		Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015)
		Zebardast, E. (2013)
MCE		Walker, B. B., Taylor-Noonan, C., Tabbernor, A., McKinnon, T. B., Bal, H., Bradley, D., . . . Clague, J. J. (2014)
		Alcorn, R., Panter, K. S., & Gorsevski, P. V. (2013)
SMCE		Armaş, I., Toma-Danila, D., Ionescu, R., & Gavriş, A. (2017)
		Hizbaron, D. R., Baiquni, M., Sartohadi, J., & Rijanta, R. (2012)
F-ADP		Rezaei-Malek, M., Torabi, S. A., & Tavakkoli-Moghaddam, R. (2019)
OLR		Muir, J. A., Cope, M. R., Angeningsih, L. R., Jackson, J. E., & Brown, R. B. (2019)
Binary Logistic regression		Qasim, S., Qasim, M., Shrestha, R. P., & Khan, A. N. (2018)
Logical analysis method		Chen, Y. (2016)
Distance-based analysis	network	Walker, B. B., Taylor-Noonan, C., Tabbernor, A., McKinnon, T. B., Bal, H., Bradley, D., . . . Clague, J. J. (2014)
Overlay analysis		Toké, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)

METHODS	AUTHORS
F*ANP	Zebardast, E. (2013)

Table 6. Methods applied to the spatial assessment of socio-economic vulnerability.

The Social Vulnerability Index (SoVI®) remains the benchmark for the assessment of SEV and a reference for its spatial assessment. [Nevertheless, indices such as Walk Scores® \(Bereitschaft, 2017a\) offer a proxy for the spatial assessment of SEV in a microscale urban level \(street level\) in 3 dimensions \(3D\).](#) The complete set of spatial indexes used by authors and identified in this systematic review is listed in Table 7.

SPATIAL INDEXES	AUTHORS
SoVI®	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
	Alcorn, R., Panter, K. S., & Gorsevski, P. V. (2013)
	Zebardast, E. (2013)
Population vulnerability Indexing	Yuan, H. H., Gao, X. L., & Qi, W. (2019)
Walk Scores®	Bereitschaft, B. (2017)
LA-SoVIC	Toké, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)

Table 7. Spatial indexes for socio-economic vulnerability assessments.

The tools to carry out the spatial assessment of SEV were selected according to the identified spatial variable and indicators, the method used, and the indexes used, adapted, or developed. The most frequent tool for the spatial assessment of SEV is GIS, followed by statistical analyses undertaken in the statistical package for the social sciences (SPSS), remote sensing (RS) using the environment for visualizing images (ENVI), programming languages, and interactive databases such as the retrieval of data for small Areas by microcomputer (REDATAM)(CELADE, 2015). The complete list of tools used by the authors selected is found in Table 8.

METHOD	SOFTWARE	AUTHORS
GIS	ArcGIS	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020).
		Yuan, H. H., Gao, X. L., & Qi, W. (2019)

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt, Sp (Spain)

METHOD	SOFTWARE	AUTHORS
		Alizadeh, M., Alizadeh, E., Kotenae, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L. (2018)
		Gautam, D. (2017)
		Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015)
	IDRISI	Alizadeh, M., Alizadeh, E., Kotenae, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L. (2018)
	ILWIS	Armaş, I., Toma-Danila, D., Ionescu, R., & Gavriş, A. (2017)
	GeoDa Version 16.6	Ley-García, J., Denegri de Dios, F. M., & Ortega Villa, L. M. (2015)
	Not specified	Ponce-Pacheco, A. B., & Novelo-Casanova, D. A. (2018)
		Eidsvig, U. M. K., McLean, A., Vangelsten, B. V., Kalsnes, B., Ciurean, R. L., Argyroudis, S., . . . Kaiser, G. (2014)
		Toke, N. A., Boone, C. G., & Arrowsmith, J. R. (2014)
		Walker, B. B., Taylor-Noonan, C., Tabernor, A., McKinnon, T., Bal, H., Bradley, D., . . . Clague, J. J. (2014)
		Alcorn, R., Panter, K. S., & Gorsevski, P. V. (2013)
		Hizbaron, D. R., Baiquni, M., Sartohadi, J., & Rijanta, R. (2012)
Statistical Analysis	SPSS 22.0	Aksha, S. K., Resler, L. M., Juran, L., & Carstensen, L. W. (2020)
	SPSS 16.0	Qasim, S., Qasim, M., Shrestha, R. P., & Khan, A. N. (2018)
	SPPS	Maharani, Y. N., Lee, S., & Ki, S. J. (2016)
RS	ENVI	Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015)
		Alizadeh, M., Alizadeh, E., Kotenae, S. A., Shahabi, H., Pour, A. B., Panahi, M., . . . Saro, L. (2018)
Programming language	MATLAB	Maharani, Y. N., Lee, S., & Ki, S. J. (2016)
Database	Redatam V5.0	Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., & Sarmiento, J. P. (2015)

Table 8. Tools for socio-economic vulnerability assessments.

4 Discussion

For the purpose of the systematic review, we found that the Clarivate Analytics database more accurately identified the references for this systematic review, and it is more user-friendly than other databases. The lack of articles that tackle exogenic geohazards can be explained by the fact that we excluded from the search query words such as “climate change” OR “ecological” OR “drought”, which are indirectly related to these phenomena. Nevertheless, considering that these geohazards usually takes place in rural, rather than urban environments, they are not relevant for this research.

The literature references identified are based on a high detailed search query to avoid bias. The query could be repeated any time and the results will be always the same, maybe additional publications from 2020 could appear on the result. However, the total number of literature references reviewed were much more than 24. Previously, based on a more general query not specifically focused on geohazards, we identified 235 literature references, from which we found 84 relevant references, 42 highly relevant references and finally 21 references were selected to be reviewed at that moment. Eventually given their relevance, we decided to keep six of these references identified previously using the first query. In the current version, we reviewed all 29 references but eventually, we selected 18 and discarded 11 for the reasons already explained in the results section. The case study areas of the selected papers confirm the findings from Shen et al., (2018) and also ours using the previous query, relating to the USA, China, and Iran as major contributors to disaster research together with Italy, Indonesia, Germany, Turkey, England, India, and Spain in the topics of ‘prediction model’, ‘social vulnerability’ and ‘landslide inventory map’. Nevertheless, the references that use Indonesia as a case study area are focused on earthquakes and volcanic eruptions, not necessarily on the tsunami hazard as was suggested by Shen et al., (2018). The reason to lead the research in those topics would be based on their degree of hazard considering that the USA, China and Indonesia are located along the Pacific Ring of Fire.

The research concentrated on the local level uses primary data collected via field observations, questionnaire surveys, or focus groups with representative members of the community to assess vulnerability (Birkmann, 2006; Khazai et al., 2017; Sarkar and Vogt, 2015), while for global or regional scales, primary data is derived from satellite images, aerial Photograph, LULC,

Deleted: ,

Deleted: external

Deleted: ¶

Deleted: ¶

landslide susceptibility maps, orthophotos, or VGI. Secondary data is obtained from the population census, disaster databases, and population datasets. For applications on the regional, national, international, or worldwide scale, coarse-scale raster data on population patterns are appropriate, but for city or local scales, representation of higher spatial resolution is requested, such as fine-scale population grids which finally go to individual building level (Aubrecht et al., 2013). Census data usually presents national data at the municipal level. Census and land databases are highly demanded by planners and disaster managers. However, there are several problems associated with using large community databases, such as scale, data decay, relevance (King, 2001), and time-constraints. Current data can easily change with the building of a new road or new houses (McLaughlin et al., 2002), and in the case of nomadic and/or geographically isolated groups, these data sets are rarely available (Béné, 2009) but they are necessary. Censuses are usually updated on an average of ten years, depending on the country, and some of the data could be altered by political biases. The surveys require significant resources, and the thematic scope is usually very narrow. These disadvantages can explain the strong demand for population data, independent of administrative areas, making it sometimes necessary to extract data from raster representations or using dasymetric mapping (Aubrecht et al., 2013; Garcia et al., 2016; Yuan et al., 2019a). Currently, data in 3D can be also extracted from VGI, which is an alternative source of real-time information based on the concept of citizens as sensors (Cervone and Hultquist, 2018).

Satellite images are useful to collect data from global to local scales. Rapid mapping concepts are mainly applied in structural post-disaster damage assessment, relaying on earth observation data from different sensors, sometimes provided by the International Charter Space and Major Disasters (2020) (Aubrecht et al., 2013). Lidar data are a good option for the city scale. The use of satellite images as data sources in the spatial assessment of SEV has been increasing in the last ten years, which can be explained because they offer quick, updated, and reliable data, making the satellite images currently the most effective source. One of the issues with using maps, air photos, or orthophotos as a resource is that they are not frequently updated.

The spatial variables found through this systematic review are similar to the variables identified by Meentemeyer (1989), Béné, (2009) Contreras et al. (2013), Buzai and Villerías Alarcón (2018). Based on the concept of spatial indicator of SV formulated by Ebert et al., (2009), we consider the lack of basic services as a spatial variable of SEV because all these networks are

distributed in a specific spatial area. The lack of life-supporting infrastructure and/or infrastructure necessary for the functioning of the society such as piped water, electricity networks, sewerage infrastructure, telecommunications and road networks hampers emergency management and therefore the recovery process (Eidsvig et al., 2014). Housing quality and tenancy conditions describe the vulnerability of the population to become homeless after a disaster (Toke et al., 2014). Housing type is an economic indicator of the economic status of individuals, communities, and nations. Thus, a house with low quality or precarious external walls located in a landslide-prone zone is usually associated with socially vulnerable communities having a negative influence on the quality of life. However, the typology of vulnerable houses depends also on the sort of landslide (Eidsvig et al., 2014). There are similar spatial variables used to produce an indicator of housing overcrowding (Ponce-Pacheco and Novelo-Casanova, 2018) such as households per housing unit (Zebardast, 2013) and households with >1 family (Aksha et al., 2020). We argue that besides spatial variables, we must also consider spatial categories in which critical and the other urban facilities must be included. These facilities are not only providers of services but are also sources of employment (Contreras et al., 2017); therefore, their presence or absence, access to, distance, travel time (Toke et al., 2014), and/or barriers (Walker et al., 2014) to reaching them highly influence the degree of spatial SEV of a community. Bereitschaft, (2017a) proposes innovative spatial variables of SEV at microscale urban level in 3D such as historic buildings, parks, place signs/identifiers, contiguous street wall, limited sightlines, street furniture, street vendors, first-floor windows, active uses/occupied storefronts, pedestrian activity, business type variety, crosswalks & pedestrian infrastructure, sidewalk condition and storefront/building condition. We also identify other spatial variables that are different to more traditional ones such as distance from faults (Hizbaron et al., 2012; Rezaei-Malek et al., 2019) and volcanoes (Kurnianto et al., 2019), land use (Alcorn et al., 2013), city blocks (Yuan et al., 2019a) and displacement (Muir et al., 2019) among others.

20

Based on the evidence found by this research, we agree with Zeng et al. (2012) that the most frequent spatial indicator in the assessment of SEV related to geohazards is population density, and it has the highest sensitivity coefficient (Yuan et al., 2019a). According to Kumianto et al., (2019), high population density is the factor that contributes most to the high SV, and it is usually linked to high population growth, which increases the SEV given the rise in the exposure of population and business. The reason, according to Gu et al. (2018), is that population density reveals the human resources of a neighbourhood and the

25

Deleted: s

Deleted: Households

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Deleted:

relief resources that could be required during a disaster. This is a key factor in large case study areas where different kinds of occupation can take place (urban, rural); therefore, important differences in population density are expected to be found. Disadvantaged population tends to live in denser neighbourhoods with more crowded parks and other recreational facilities (Sister et al., 2009; Toke et al., 2014; Wolch et al., 2005) and low levels of walkability (Bereitschaft, 2017a) that exacerbate the vulnerability making an evacuation difficult (Cutter et al., 2003) after an earthquake, tsunami, volcanic eruption, or landslide. It is also more difficult in such areas to find spaces to install temporary shelters near their households or areas for providing care after an emergency (Cutter et al., 2003). The density of the built environment is especially important in the case of seismic events (Toke et al., 2014). Innovative spatial indicators such as employed density, unemployed density, and literacy people density were proposed by Alizadeh et al. (2018). The importance of such fine-scale data and the temporal variations (daytime and night-time) for accurately estimating SV was highlighted by Yuan et al., (2019a), proposing the indicator: ‘floating population’. The consideration of the spatial and temporal dimension in the estimation of population exposure is a fundamental aspect of accurate catastrophe loss modelling, a key element for the integration of risk analysis and emergency management (Aubrecht et al., 2010), and therefore for the reduction of the SEV (Alizadeh et al., 2018). Chen (2016) proposes more spatial indicators in the economic rather than the social dimension. Ley-García et al. (2015), global Moran’s I and LISA enable the identification of dependence between attributes and localisations. As a result, these indicators are useful to determine whether the spatial distribution of elements influences the behaviour of a particular variable. The summary measure of autocorrelation in the territory is undertaken with global Moran’s I, while the autocorrelation of the spatial units included in the territory is measured using LISA. Cutter and Finch (2008) also previously utilised global Moran’s I and LISA to identify local variability and cluster similarity of low and SV. Besides the SoVI® and FA, Zhou et al. (2014) utilise exploratory spatial data analysis (ESDA) to identify the spatio-temporal patterns of SV based on the constructed SoVI® for each county in China. These authors used global and local Moran’s I or LISA as ESDA to determine the spatial autocorrelation among counties and identify the similarity and/or dissimilarity in the clustering of SV.

Accessibility as a spatial indicator is defined as the ability to contact and interact with places of economic or social opportunities (Deichmann, 1997). Goodall (1987) notes that accessibility is the ease to reach a location from another location, and this concept is also related to opportunities for attention (Aubrecht et al., 2013) in the case of, for example, hospitals and/or

Deleted:

Deleted: that

Deleted: ir

Deleted:

Deleted: for

Deleted: ion

Deleted: again

Deleted: the

Deleted: contact, interaction, and

trauma centres, accessibility is reduced by distance (Hizbaron et al., 2012; Zeng et al., 2012) increasing SEV level of the communities located far from these healthcare facilities. Besides the common spatial variables, indicators and indexes in 2D, there are also spatial indicators and indexes that include a 3D component, such as imageability, enclosure, human scale, transparency, complexity, safety & sensations and tidiness, (Bereitschaft, 2017a) satisfaction with the neighbourhood (Barata et al., 2011), and residential condition (de la Torre and de Riccitelli, 2017) that could be applied to the spatial assessment of SEV. Authors such as Yuan et al (2020) and Muir et al (2019) consider the spatial indicators of mobility and migration respectively in the framework of geohazards, being migration a topic mainly addressed by authors in the climate change community e.g. Nakayama et al.,(2019), Naugle et al.,(2019), van der Geest et al., (2020), Ayeb-Karlsson et al.,(2020) and others.

Deleted: Walk Score®, neighbourhood walkability (Bereitschaft 2017a, b), ...

Deleted: Locke,(2009)

Deleted: so forth

This systematic review identified the versatility of ANN, which can be either used to extract monthly rainfall data (Aksha et al., 2020), for deriving social vulnerability maps (SVM) (Alizadeh et al., 2018) or to train the self-organized map (SOM) algorithm cluster method (Maharani et al., 2016). The use of dasymetric population mapping not limited to administrative boundaries, even going to block-level to increase the spatial resolution of the population exposure analysis (Garcia et al., 2016) and additionally by including the temporal dimension with its day-night variability, enables improving the accuracy of the spatial assessments of SEV (Yuan et al., 2019a). Factor analysis (FA) is used by Castro et al. (2015) to establish the level of SEV and by Zebardast (2013) to extract primary dimensions and variables of SEV. Alcorn (2013) applied MCE to assess economic vulnerability using four significant factors: population, infrastructure, land use, and economic production. SMCE is applied by Armaş et al., (2017) to integrate social, education, housing, and social dependence vulnerability dimensions and by Hizbaron (2012) to develop deterministic SV scenarios. Zebardast (2013) enters the variables of SEV into a network model in an analytic network process (ANP) to rank the importance of each variable to complete the F'ANP method. This method is focused on developing a composite social vulnerability index (SOVI). Binary logistic regression was the statistical method applied by Qasim et al. (2018) to identify the determinants of landslide risk perception, location being one of them. Walker et al., (2014) present a multi-criteria evaluation (MCE) model that incorporates access to healthcare facilities using GIS to identify and rank residential areas in Victoria, British Columbia. The integration of the concept of uncertainty into ANP using fuzzy

numbers (F-ANP) is combined by Rezaei-Malek et al., (2019) with fuzzy DEMATEL (F-DEMATEL) to deal with the interdependency among a set of criteria and fuzzy PROMETHEE II (F- PROMETHEE II) to control the criteria weights, the complete method is denominated fuzzy ANP DEMATEL PROMETHEE II (F-ADP). Ordinal logistic regression (OLR) is used by Muir et al., (2019) to predict the mental health condition of people displaced by series of volcanic eruptions in Merapi, Indonesia, according to their migration status (displaced, moved home, in transition, and moved on), which implies a spatial component. Geological experience and logical analysis method were used by Chen (2016) to select indicators. Toke et al.,(2014) undertake an overlay analysis to identify the census block groups that intersect zones with an extreme ground shaking hazard.

Aksha et al., (2020) utilized the SoVI® to map the vulnerability levels in the study site with a multi-hazard map to produce a total risk map. Alcorn et al. (2013) used an improved version of the same index, but specifically adapted it to the variability in SEV in the case study area that was focused on census-designated places (CDPs) on a small scale. The population vulnerability indexing developed by Yuan et al., (2019a) considered most of the indicators available in the literature already identified by the SoVI®, but they adapted their index to the Chinese society, where according to the authors, race and ethnicity are not relevant indicators and rural-to-urban migrants are floating population with unequal access to public services and therefore a vulnerable population. Bereitschaft (2017a) explores the exiting inequities in the walkability of urban environments among neighbourhoods with low and high SEV using the Walk Scores®. This index could be used as a proxy spatial index of SEV in 3D at microscale urban level. The author found that neighbourhoods with high SV had fewer windows and less transparent storefronts, less continuous street walls, less well-maintained infrastructure, fewer business and generally less complexity than in neighbourhoods with low SV. Toké et al (2014), build upon the SoVI® to create their own SV indexes that incorporate the spatial dimension. According to the LA-SoVIC developed by Toket et al. (2014), SV is highly linked to the normalised difference vegetation index (NDVI) as a proxy for urban green space. Green areas are usually located in areas with lower SEV (Stow et al., 2007), and have also been recognised for their health benefits (Bedimo-Rung et al., 2005). Physical characteristics of green areas, such as attractive scenery, motivates people to stay and visit an area (Kurnianto et al., 2019), resulting in increased social control and reduced SEV.

Deleted: ,

It has been always difficult to quantify SV; hence, it is absent from post-disaster cost/loss estimation reports (Schmidtlein et al., 2008; Zhou et al., 2014). The use of spatial variables, indicators, and indexes will bridge the gap of integrating physical vulnerability and SV to achieve a holistic risk assessment. Davidson (1997) provides the first attempt to create an integrated risk assessment framework. Later, Carreño, Cardona, & Barbat, (2007) developed a risk index obtained by multiplying the physical risk index by an impact factor, which is, in fact, an aggravating coefficient consisting of socio-economic variables; nevertheless, in applying this method, the outcome will be similar to the assessment of physical vulnerability, without showing the contribution of SV to the assessment of integrated risk. Schmidtlein, Shafer, Berry and Cutter (2011) tested the link between SV and earthquake losses. The authors found that physical parameters related to hazard, such as distance from the epicentre and peak ground acceleration, were more significant in predicting impacts than SV. Nevertheless, the same authors established that SV is a significant predictor of earthquake losses when accounting for wealth (dollar losses per average income as the dependent variable). The previous finding reveals that those areas with higher levels of SV experience a greater relative impact than areas with lower degrees of SV.

Geospatial information systems are broadly utilised by several authors to collect and, process data, and map the SEV. GIS has been enabling researchers to have either large study regions, or equivalently, data sets at much finer spatial resolution (Unwin, 1996), for example, a comprehensive overview of the use of accessibility indicators in GIS was already provided by Deichmann (1997). Each author uses different versions of ArcGIS, which is the most widespread software used in GIS. The IDRISI software is utilised by Alizadeh et al. (2018) to generate a Social Vulnerability Map (SVM). Armaş et al., (2017) applied a pairwise comparative method in the AHP implemented in the SMCE module of the Integrated Land and Water Information System (ILWIS) software. GeoDa, an open-source software, focused on methods for spatial data, and has been used by authors who address the topic of spatial association (Gu et al., 2018; Ley-García et al., 2015). The aforementioned is an RS and GIS software, on which the robustness of the results from Armaş et al. (2017) was also tested, with a sensitivity analysis performed in the DEFINITE toolbox implemented in ILWIS. The MATLAB computation environment was used by Maharani et al. (2016) to develop the SOM toolbox. Sherly et al. (2015) also use MATLAB to perform multivariate data analyses, such as PCA and Data Envelopment Analysis (DEA). REDATAM used as a source of data by Castro et al. (2015), is an interactive hierarchical database that contains microdata and/or aggregate socio-economic information from any geographical division at a national

Deleted: Ilwis

Deleted: ,

level. This database combines data from the census, surveys and other sources, resulting in a very comprehensive and useful source of spatial and not-spatial variables for the SEV.

5 Conclusions

5 Based on the evidence, we can state that most of the spatial assessments of SEV in urban environments have been done for earthquakes and landslides and that Indonesia, China, Iran, and the USA lead the research in the spatial assessment of SEV related to geohazards in urban environments. The scale of the spatial level of assessment – namely global, continental, subcontinental, national, regional, provincial, municipal, or local – determines the type of data to be collected and the assessment approaches. Although there have been advances, census data continues to be the most frequent source of data for
10 the SEV assessments; however, in the case of spatial assessment, satellite images are now the main data source, facilitating the inclusion of the spatial component in SEV assessments. The spatial assessment of SEV allows visualising and communicating social phenomena and components that influence the degree of vulnerability that are not visible with other methods. The lack of data availability hinders the understanding of the concept of vulnerability (Zhou et al., 2014) and that is why VGI is essential today to obtain updated information in real-time at the local scale, when other data sources are not
15 available.

Traditional spatial variables and indicators continue to be used by authors, but combined with new variables, categories, and indicators, including the temporal dimension (day-night), and assessing at the local level, can increase the accuracy of spatial assessments of SEV and reduce uncertainty on their assessment. Each method for the spatial assessment of SV is selected according to the research aim, case study area, scale to cover, reliability of data sources, spatial variables and indicators
20 available; geohazard to address, the scope of the research, and the level of funding. Methods such as ANN are gaining ground in the assessment of SEV. Other methods such as dasymetric population mapping enable more accurate SEV assessment. Factor analysis continues to be a useful tool to define the level of SEV based on primary dimensions and variables. Multi-criteria evaluation method offers a robust decision-making technique based on flexible choice and combination in criteria (Alcorn et al., 2013). SMCE incorporates the spatial component to the MCE to integrate spatial and non-spatial data to generate
25 maps with multiple scenarios (Hizbaron et al., 2012). Classic methods such as FA are combined with more innovative ones

Deleted: ,

such as ANP and fuzzy numbers to generate hybrid methods such as F'ANP. These new methods encourage the development of more complex hybrid methods such as F-ADP that increase the accuracy and reduce the uncertainty levels in the spatial SEV assessments. Ordinal logistic regression and binary logistic regression are useful methods to identify spatial variables as determinants of SEV. The spatial component can be also be added by simply overlapping the areas with high SEV with hazard zones using GIS. Most authors have built upon the SoVI® developed by Cutter et al. (2003) to quantify SEV or to create their own SEV indexes, demonstrating that it remains the benchmark for the assessment of SEV and a reference for its spatial assessment. however there are new alternatives for the spatial assessment of SEV in 3D at microscale level such as Walk Scores® (Bereitschaft, 2017a).

Geographic Information Systems, statistical analysis, RS, programming languages, and interactive databases are the tools currently used by the scientists for the assessment of SEV vulnerability. The spatial assessment of SEV in the areas where it is requested must depend not only on the financial resources for research but also on the availability of opensource software with the functionalities of spatial statistics, such as QGIS, GeoDa or IIWIS. Authors combine traditional and new data sources, spatial variables and indicators, methods, indexes and tools including the temporal dimension, increasing the resolution to the local level with the aim to increase the accuracy and reduce the uncertainty of spatial assessments of SEV related to geohazard in urban environments.

6 Recommendations

The development of a global spatial index of SEV is an urgent task, with the aim of making informed decisions about priority in funding prevention and mitigation actions related to geohazards in urban environments. In the meantime, the priority for these types of assessments must be allocated to developing countries with high population density such as Bangladesh, Haiti, Philippines, Puerto Rico, el Salvador and Pakistan. More spatial assessment of SEV due to volcanic eruptions and tsunamis in urban environments are needed, but also due to soil erosion and land degradation in the rural zones. Furthermore, the priority must be to allocate funding for countries with high SEV to enable the update of their census information, as this is the most frequent source of secondary data for any SEV assessment. It is also important to encourage the population to share information

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, 10 pt

Deleted: the lowest GDPs

Deleted: and increased levels of SV (Zhou et al., 2014)

through social media (SM) about the vulnerable conditions in which they live, putting in practice the concept of citizen as a sensor (Cervone and Hultquist, 2018).

An assessment of SEV is a condition for the effective development of emergency management capabilities and to reduce the overall time for social recovery after an earthquake (Aubrecht et al., 2013; Garcia et al., 2016). Likewise, spatial assessments of SEV must be considered before taking resettlement decisions for not creating again spatial conditions that favour the SEV.

Authors such as Turvey (2007), Walker et al. (2014), Zhou et al. (2014) and Gautam (2017) highlight the need for place-specific, sub-provincial-level, neighbourhood-scale, or local level vulnerability indexes, due to geographic variations in population composition and social structures (Bell N et al., 2007). The macro-scale socio-economic assessment identifies general patterns but fails to capture the detail of the heterogeneity at the micro-scale. Thus, assessment at the provincial, county or state level can result in lost information (Zhou et al., 2014) or requires tackling issues such as ecological fallacy or MAUP (McLaughlin et al., 2002; Openshaw, 1983; Pacione, 2005). In the spatial assessment of SEV, it is necessary to go beyond the administrative boundaries or cartographic variables, with methods such as the dasymetric population mapping (Garcia et al., 2016; Yuan et al., 2019a), square mesh (Renard, 2017), pockets (Lin and Hung, 2016), or *geon* (Kienberger et al., 2009). We found interesting spatial indicators of SEV, such as population density based on land use (Zeng et al., 2012), which we consider more accurate than population density estimated at an area unit. This indicator can better integrate, using RS, the spatial dimension of the exposure and susceptibility of the population in the assessment of the SEV of a case study area. To improve the accuracy and reduce the uncertainty in spatial assessments of SEV must always be the aim. The presence of urban facilities must be included in the assessment of SV. Walker et al. (2014) suggest developing a weighted 'local resource' index for assessing systemic vulnerability since, for example, the absence of sports facilities is associated by Iguacel et al. (2018), Vandermeersch, Vos, & Scheerder (2015), and Aguilar-Palacio, Gil-Lacruz and Gil-Lacruz (2013) with high levels of SV. In the spatial assessment of SEV, it is also necessary to consider the influence of the spatial component represented by physical space in the degree of vulnerability of a specific area, such as the relationship between slums and a low degree of wellness and health (Buzai and Villerías Alarcón, 2018).

Deleted: s

It is necessary to take advantage of the versatility of methods such as ANN based on machine learning to make progress in the spatial assessment of SEV and SMCE in order to map multiple scenarios to inform urban communities and to integrate them in the decision making processes. Communities respond differently to vulnerability maps depending on the purpose behind the maps or the cultural background of the community. On the one hand, some communities reject being mapped as ‘victims’, but on the other hand, some request being identified as highly vulnerable to gain access to funding opportunities for activities of risk management (Fekete, 2012). The Walk Score® index developed by Bereitschaft (2017a) although originally orientated to measure only neighbourhood walkability (Bereitschaft, 2017b), ~~can be used a proxy index of spatial SEV in 3D at microscale urban level~~. The advantage over the SoVI® is that while the SoVI® can be spatialised, Walk Score® is a 3D high resolution spatial index *per se*. The use of the local scale for the assessment of SV will be more useful for the planning of resilient actions (Lee, 2014; Maharani et al., 2016) than would be vulnerability assessment at a regional scale, which is more orientated to the collection of pathologies in the social dimension. It is necessary to more closely examine so-called ‘proxy indicators’ to measure ~~spatial SEV~~ at micro-local scales or intra-city levels (Gu et al., 2018). The right management of the spatial component by a community can reduce its economic vulnerability. Groß (2017) presented the case of ski-lift entrepreneurs in Vorarlberg (Austria) who reduced the probability of business interruption by accelerating the uphill and downhill flows of people through manipulating snow and topography. Regarding tools, it is necessary to take full advantage of the functionalities of opensource software such as and QGIS and ILWIS to make the spatial assessment of SEV to the reach of all the scientific communities around the world.

Deleted: could

Deleted: be adapted as a tool to either complement or validate spatial index for assessing

Deleted: the local level

Acknowledgements

The authors thank ~~the National Agency of Research and Development (ANID by its acronym in Spanish) in Chile~~, which has funded the Research Center for Integrated Disaster Risk Management (CIGIDEN), ANID/FONDAP/15110017 and the Fondecyt Project 1181754/FONDECYT/ANID. Authors also acknowledge ~~funding received from~~ the Engineering and Physical Sciences Research Council (EPSRC) [Grant No. EP/P025951/1] ~~in UK to conclude this research~~. We want to extend our most thanks to Dr Carolina Martinez for the ~~literature~~ references suggested. We also appreciate the feedback received from Dr Magdalena Vicuña, Dr Cristina Vizconti, Dr Luis Maldonado and Marta Contreras, BCE. for their feedback during the review process. We also would like to thank the anonymous reviewers and Editors Prof. Dr Thomas Glade and Dr Heidi Kreibich for their contributions to the improvement of this research outcome and Dr Anne Bliss for her support with the English proofreading.

Deleted:

Deleted: .

Deleted: .

Deleted: .

Funding

Research Center for Integrated Disaster Risk Management (CIGIDEN), ANID/FONDAP/15110017
Fondecyt Project 1181754/FONDECYT/ANID

5 Engineering and Physical Sciences Research Council (EPSRC) [Grant No. EP/P025951/1]

References

<https://disasterscharter.org/web/guest/home;jsessionid=AA34233A7B32785905A53A64AB27541A.jvm1>, last access: The 13th March 2020 2020.

10 Adger, W. N., Brooks, N., Bentham, G., Agnew, M., and Eriksen, S.: New indicators of vulnerability and adaptive capacity, Tyndall Centre for Climate Research Technical Report 7, 2004.

Aguilar-Palacio, I., Gil-Lacruz, M., and Gil-Lacruz, A. I.: Salud, deporte y vulnerabilidad socioeconómica en una comunidad urbana, *Atención Primaria*, 45, 107-114, 2013.

15 Aksha, S. K., Resler, L. M., Juran, L., and Carstensen, L. W.: A geospatial analysis of multi-hazard risk in Dharan, Nepal, *Geomat. Nat. Hazards Risk*, 11, 88-111, 2020.

Alcántara-Ayala, I. and Oliver-Smith, A.: ICL Latin-American Network: on the road to landslide reduction capacity building, *Landslides*, 11, 315-318, 2014.

Alcorn, R., Panter, K. S., and Gorsevski, P. V.: A GIS-based volcanic hazard and risk assessment of eruptions sourced within Valles Caldera, New Mexico, *Journal of Volcanology and Geothermal Research*, 267, 1-14, 2013.

20 Alizadeh, M., Alizadeh, E., Kotenace, S. A., Shahabi, H., Pour, A. B., Panahi, M., Bin Ahmad, B., and Saro, L.: Social Vulnerability Assessment Using Artificial Neural Network (ANN) Model for Earthquake Hazard in Tabriz City, Iran, *Sustainability*, 10, 23, 2018.

Amram, O., Schuurman, N., and Hameed, S.: Mass casualty modelling: a spatial tool to support triage decision making, *Int J Health Geogr* 10, 2011.

Armaş, I., Toma-Danila, D., Ionescu, R., and Gavriş, A.: Vulnerability to Earthquake Hazard: Bucharest Case Study, Romania, *International Journal of Disaster Risk Science*, 8, 182-195, 2017.

25 Aubrecht, C., Köstl, M., and Steinnocher, K.: Population exposure and impact assessment: benefits of modeling urban land use in very high spatial and thematic detail. In: *Computational vision and medical image processing: recent trends. computational methods in applied sciences*, JMRS, T. and RM, N. J. (Eds.), Springer, Berlin 2010.

Aubrecht, C., Özceylan, D., Steinnocher, K., and Freire, S.: Multi-level geospatial modeling of human exposure patterns and vulnerability indicators, *Nat. Hazards*, 68, 147-163, 2013.

30 Ayeb-Karlsson, S., Kniveton, D., and Cannon, T.: Trapped in the prison of the mind: Notions of climate-induced (im)mobility decision-making and wellbeing from an urban informal settlement in Bangladesh, *Palgrave Communications*, 6, 62, 2020.

Barata, R. B., Sampaio de Almeida Ribeiro, M. C., and Cassanti, A. C.: Social vulnerability and health status: a household survey in the central area of a Brazilian metropolis, *CADERNOS DE SAUDE PUBLICA*, 27, S164-S175, 2011.

35 Beccari, B.: A Comparative Analysis of Disaster Risk, Vulnerability and Resilience Composite Indicators, *PLoS Curr*, 8, ecurrents.dis.453df025e434b682e9737f95070f95079b95970, 2016.

Bedimo-Rung, A. L., Mowen, A. J., and Cohen, D. A.: The significance of parks to physical activity and public health: A conceptual model, *American Journal of Preventive Medicine*, 28, 159-168, 2005.

Bell N, Schuurman N, Oliver L, and MV, H.: Towards the construction of place-specific measures of deprivation: a case study from the Vancouver metropolitan area. , *Can Geogr* 51, 444-461, 2007.

40 Béné, C.: Are Fishers Poor or Vulnerable? Assessing Economic Vulnerability in Small-Scale Fishing Communities, *The Journal of Development Studies*, 45, 911-933, 2009.

Bereitschaft, B.: Equity in Microscale Urban Design and Walkability: A Photographic Survey of Six Pittsburgh Streetscapes, *Sustainability*, 9, 1233, 2017a.

45 Bereitschaft, B.: Equity in neighbourhood walkability? A comparative analysis of three large U.S. cities, *Local Environment*, 22, 859-879, 2017b.

Birkmann, J.: Indicators and criteria for measuring vulnerability: theoretical bases and requirements. In: *Measuring Vulnerability to Natural Hazards. Towards Disaster Resilient Societies*, Birkmann, J. (Ed.), United Nations University Press, Tokyo, 2006.

Birkmann, J., Cardona, O. D., Carreño, M. L., Barbat, A. H., Pelling, M., Schneiderbauer, S., Kienberger, S., Keiler, M., Alexander, D., Zeil, P., and Welle, T.: Framing vulnerability, risk and societal responses: the MOVE framework, *Nat. Hazards*, 67, 193-211, 2013.

Formatted: English (United Kingdom)

Field Code Changed

Formatted: English (United Kingdom)

Formatted: Spanish (Spain)

Formatted: English (United Kingdom)

Formatted: Spanish (Spain)

- Buzai, G. and Villerías Alarcón, I.: Análisis espacial cuantitativo de los determinantes sociales de la salud (DSS) en la cuenca del río Luján (provincia de Buenos Aires, Argentina), *Estudios Socioterritoriales*, 23, 2018.
- Cardona, O. D.: Indicators of Disaster Risk and Risk Management: Program for Latin America and the Caribbean: Summary Report, Inter-American Development Bank, Washington, DC, USA, 2005.
- 5 Carreño, L., Cardona, O. D., and Barbat, A. H.: Urban Seismic Risk Evaluation: A Holistic Approach, *Natural Hazards*, 40, 137–172, 2007.
- Castro, C. P., Ibarra, I., Lukas, M., Ortiz, J., and Sarmiento, J. P.: Disaster risk construction in the progressive consolidation of informal settlements: Iquique and Puerto Montt (Chile) case studies, *International Journal of Disaster Risk Reduction*, 13, 109–127, 2015.
- CELADE: REtrieval of DATa for small Areas by Microcomputer (Redatam). 2015.
- 10 Cerchiello, V., Ceresa, P., Monteiro, R., and Komendantova, N.: Assessment of social vulnerability to seismic hazard in Nablus, Palestine, *International Journal of Disaster Risk Reduction*, 28, 491–506, 2018.
- Cervone, G. and Hultquist, C.: Citizen as indispensable sensors during disasters, 2018.
- Chambers, R.: Vulnerability, coping and policy, *IDS Bulletin*, 20, 1–7, 1989.
- Chen, Y.: Conceptual Framework for the Development of an Indicator System for the Assessment of Regional Land Subsidence Disaster Vulnerability, *Sustainability*, 8, 757, 2016.
- 15 Clark, G. E., Moser, S. C., Ratick, S. J., Dow, K., Meyer, W. B., Emani, S., Jin, W., Kasperson, J. X., Kasperson, R. E., and Schwarz, H. E.: Assessing the Vulnerability of Coastal Communities to Extreme Storms: The Case of Revere, MA., USA, *Mitigation and Adaptation Strategies for Global Change*, 3, 59–82, 1998.
- Contreras, D., Blaschke, T., and Hodgson, M. E.: Lack of spatial resilience in a recovery process: Case L'Aquila, Italy, *Technological Forecasting and Social Change*, 121, 76–88, 2017.
- 20 Contreras, D., Blaschke, T., Kienberger, S., and Zeil, P.: Spatial connectivity as a recovery process indicator: The L'Aquila earthquake, *Technological Forecasting and Social Change*, 80, 1782–1803, 2013.
- Contreras, D. and Kienberger, S.: GIS in the vulnerability assessment and recovery process in a community with elderly and disable people after a disaster In: *Rebuilding Sustainable Communities with vulnerable populations after the cameras have gone: a worldwide study* Awotona, A. (Ed.), Cambridge Scholar Publishing, United Kingdom, Cambridge, 2012.
- 25 Contreras, D. and Kienberger, S. (Eds.): *Handbook of Vulnerability Assessment in Europe*, European Commission DG Environment, 2011.
- Contreras, D., Thomas, B., Stefan, K., and Peter, Z.: Spatial Vulnerability Indicators: "Measuring" Recovery Processes after Earthquakes, *The 8th International conference on Information systems for crisis response and management*, Lisbon, Portugal, 2011.
- Cutter, S. L., Boruff, B. J., and Shirley, W. L.: Social vulnerability to environmental hazards, *Social Science Quarterly*, 84, 242–261, 2003.
- 30 Cutter, S. L. and Finch, C.: Temporal and spatial changes in social vulnerability to natural hazards, *Proceedings of the National Academy of Sciences*, 105, 2301–2306, 2008.
- Davidson, R.: A multidisciplinary urban earthquake disaster risk index, *Earthquake Spectra*, 13, 211–223, 1997.
- de la Torre, L. and de Riccietelli, M.: 'NiNis': Youth in Argentina who Neither Work nor Study. *A Social Integration Deficit*, *REVISTA ESPAÑOLA DE INVESTIGACIONES SOCIOLOGICAS*, doi: 10.5477/cis/reis.158.97, 2017, 97–115, 2017.
- Deichmann, U.: Accessibility Indicators in GIS, United Nations Statistics Division, Department for Economic and Policy Analysis, New York, NY, USA, 1997.
- 35 Ebert, A., Kerle, N., and Stein, A.: Urban social vulnerability assessment with physical proxies and spatial metrics derived from air- and spaceborne imagery and GIS data, *Nat. Hazards*, 48, 275–294, 2009.
- Eidsvig, U. M. K., McLean, A., Vangelsten, B. V., Kalsnes, B., Ciurean, R. L., Argyroudis, S., Winter, M. G., Mavrouli, O. C., Fotopoulou, S., Pitilakis, K., Baills, A., Malet, J. P., and Kaiser, G.: Assessment of socioeconomic vulnerability to landslides using an indicator-based approach: methodology and case studies, *Bull. Eng. Geol. Environ.*, 73, 307–324, 2014.
- 40 Esty, D., Levy, M., Srebotnjak, T., and De Sherbin, A.: *Environmental Sustainability Index*, Yale Center for Environmental Law and Policy New Haven, CT, USA, 2005.
- Fatemi, F., Ardalan, A., Aguirre, B., Mansouri, N., and Mohammadfam, I.: Social vulnerability indicators in disasters: Findings from a systematic review, *International Journal of Disaster Risk Reduction*, 22, 219–227, 2017.
- 45 Fekete, A.: Spatial disaster vulnerability and risk assessments: challenges in their quality and acceptance, *Nat. Hazards*, 61, 1161–1178, 2012.
- Fekete, A.: Validation of a social vulnerability index in context to river-floods in Germany, *Nat. Hazards Earth Syst. Sci.*, 9, 393–403, 2009.
- Garcia, R. A. C., Oliveira, S. C., and Zezere, J. L.: Assessing population exposure for landslide risk analysis using dasymetric cartography, *Natural Hazards and Earth System Sciences*, 16, 2769–2782, 2016.
- 50 Gautam, D.: Assessment of social vulnerability to natural hazards in Nepal, *Natural Hazards and Earth System Sciences*, 17, 2313–2320, 2017.
- Goncalves, M. and Vizintim, M. F. B.: Geographical features of the state of Parana in the face of natural disasters, *Confins*, 33, 25, 2017.
- Goodall, B.: *The Penguin dictionary of human geography*, Penguin Books, Harmondsworth, Middlesex, England; New York, N.Y., U.S.A., 1987.
- 55 Grace, K. L. L. and Edwin, H. W. C.: Indicators for evaluating environmental performance of the Hong Kong urban renewal projects, *Facilities*, 27, 515–530, 2019.

Formatted: English (United Kingdom)

Formatted: Spanish (Spain)

Formatted: English (United Kingdom)

- Grant, M. and Booth, A.: A typology of reviews: an analysis of 14 reviews types and associated methodologies, doi: 10.1111/j.1471-1842.2009.00848.x, 2009. 2009.
- Groß, R.: Uphill and Downhill Histories. How Winter Tourism Transformed Alpine Regions in Vorarlberg, Austria – 1930 to 1970. In: Zeitschrift für Tourismuswissenschaft, 1, 2017.
- 5 Gu, H., Du, S., Liao, B., Wen, J., Wang, C., Chen, R., and Chen, B.: A hierarchical pattern of urban social vulnerability in Shanghai, China and its implications for risk management, *Sustainable Cities and Society*, 41, 170-179, 2018.
- Hizbaron, D. R., Baiquni, M., Sartohadi, J., and Rijanta, R.: Urban Vulnerability in Bantul District, Indonesia-Towards Safer and Sustainable Development, *Sustainability*, 4, 2022-2037, 2012.
- 10 Iguacel, I., Fernández-Alvira, J. M., Bammann, K., Chadjigeorgiou, C., De Henaauw, S., Heidinger-Felső, R., Lissner, L., Michels, N., Page, A., Reisch, L. A., Russo, P., Sprengeler, O., Veidebaum, T., Börmhorst, C., Moreno, L. A., and consortium, O. b. o. t. I.: Social vulnerability as a predictor of physical activity and screen time in European children, *International Journal of Public Health*, 63, 283-295, 2018.
- Jaque Castillo, E., Contreras, A., Ríos, R., and Quezada Flory, J.: Assessment of Tsunami vulnerability in Central Chile. Factor for the local management of risk, *Revista Geografica Venezolana*, 54, 47-65, 2013.
- Khazai, B., Anhorn, J., Burton, C. G., Valcarcel, J., and Contreras, D.: Resilience Performance Scorecard (RPS) - Summary 2017.
- 15 Khazai, B., Daniell, J. E., Düzgün, Ş., Kunz-Plapp, T., and Wenzel, F.: Framework for systemic socio-economic vulnerability and loss assessment. In: *Geotechnical, Geological and Earthquake Engineering*, 2014.
- Khazai, B., Merz, M., Schulz, C., and Borst, D.: An integrated indicator framework for spatial assessment of industrial and social vulnerability to indirect disaster losses, *Nat. Hazards*, 67, 145-167, 2013.
- Kienberger, S., Lang, S., and Zeil, P.: Spatial vulnerability units – expert-based spatial modelling of socio-economic vulnerability in the Salzach catchment, Austria, *Nat. Hazards Earth Syst. Sci.*, 9, 767-778, 2009.
- 20 King, D.: Uses and limitations of socioeconomic indicators of community vulnerability to natural hazards: Data and disasters in northern Australia, *Nat. Hazards*, 24, 147-156, 2001.
- Kuhlicke, C., Scolobig, A., Tapsell, S., Steinführer, A., and De Marchi, B.: Contextualizing social vulnerability: findings from case studies across Europe, *Nat. Hazards*, 58, 789-810, 2011.
- 25 Kurnianto, F. A., Ikhsan, F. A., Apriyanto, B., and Nurdin, E. A.: Earthquake vulnerability disaster in the Lembang district of West Bandung Regency, Indonesia, *Earthq. Sci.*, 32, 40-46, 2019.
- Lee, Y.-J.: Social vulnerability indicators as a sustainable planning tool, *Environmental Impact Assessment Review*, 44, 31-42, 2014.
- Ley-García, J., Denegri de Dios, F. M., and Ortega Villa, L. M.: Spatial dimension of urban hazardscape perception: The case of Mexicali, Mexico, *International Journal of Disaster Risk Reduction*, 14, 487-495, 2015.
- 30 Lin, W.-Y. and Hung, C.-T.: Applying spatial clustering analysis to a township-level social vulnerability assessment in Taiwan, *Geomatics, Natural Hazards and Risk*, 7, 1659-1676, 2016.
- Maharani, Y. N., Lee, S., and Ki, S. J.: Social vulnerability at a local level around the Merapi volcano, *International Journal of Disaster Risk Reduction*, 20, 63-77, 2016.
- McLaughlin, S., McKenna, J., and Cooper, J. A. G.: Socio-economic data in coastal vulnerability indices: constraints and opportunities, *J. Coast. Res.*, 2002, 487-497, 2002.
- 35 Meentemeyer, V.: Geographical perspectives of space, time, and scale., *Landscape Ecology* 3, 163-173, 1989.
- Muir, J. A., Cope, M. R., Angeningsih, L. R., Jackson, J. E., and Brown, R. B.: Migration and Mental Health in the Aftermath of Disaster: Evidence from Mt. Merapi, Indonesia, *Int. J. Environ. Res. Public Health*, 16, 19, 2019.
- Myers, C. A., Slack, T., and Singelmann, J.: Social Vulnerability and Migration in the Wake of Disaster: The Case of Hurricanes Katrina and Rita, *Population and Environment*, 29, 271-291, 2008.
- 40 Nakayama, M., Drinkall, S., and Sasaki, D.: Climate Change, Migration, and Vulnerability: Overview of the Special Issue, *J. Disaster Res.*, 14, 1246-1253, 2019.
- Naugle, A. B., Backus, G. A., Tidwell, V. C., Kistin-Keller, E., and Villa, D. L.: A Regional Model of Climate Change and Human Migration, *Int. J. Syst. Dyn. Appl.*, 8, 1-22, 2019.
- 45 Noy, I.: Comparing the direct human impact of natural disasters for two cases in 2011: The Christchurch earthquake and the Bangkok flood, *International Journal of Disaster Risk Reduction*, 13, 61-65, 2015.
- Noy, I.: The macroeconomic consequences of disasters, *Journal of Development Economics*, 88, 221-231, 2009.
- OECD: Handbook on Constructing Composite Indicators: Methodology and User Guide, OECD Publishing, 2008.
- Oliver-Smith, A.: Theorizing Vulnerability in a Globalized World: A political Ecological Perspective In: *Mapping Vulnerability: Disasters, Development and People*, Bankoff, Frerik, and Hilhorst (Eds.), Earthscan, London, 2003.
- 50 Openshaw, S.: The Modifiable Areal Unit Problem, Geo Books, Norwich, 1983.
- Pacione, M.: Urban geography: a global perspective, Routledge, London etc., 2005.
- Papathoma-Kohle, M., Cristofari, G., Wenk, M., and Fuchs, S.: The importance of indicator weights for vulnerability indices and implications for decision making in disaster management, *International Journal of Disaster Risk Reduction*, 36, 12, 2019.
- 55 Ponce-Pacheco, A. B. and Novelo-Casanova, D. A.: Vulnerability and risk in Valle de Chalco solidaridad, Estado de Mexico, Mexico. Case study: El Triunfo, Avandaro and San Isidro, *Investigaciones Geograficas*, doi: 10.14350/ig.59675, 2018. 2018.

Formatted: Spanish (Spain)

- Postiglione, I., Masi, A., Mucciarelli, M., Lizza, C., Camassi, R., Bernabei, V., Piacentini, V., Chiauzzi, L., Brugagnoni, B., Cardoni, A., Calcara, A., Di Ludovico, M., Giannelli, M., Rita, R., La Pietra, M., Bernardini, F., Nostro, C., Pignone, M., and Peruzza, L.: The Italian communication campaign "I Do Not Take Risks - Earthquake", *Boll. Geofis. Teor. Appl.*, 57, 147-160, 2016.
- Qasim, S., Qasim, M., Shrestha, R. P., and Khan, A. N.: Socio-economic determinants of landslide risk perception in Murree hills of Pakistan, *AIMS Environ. Sci.*, 5, 305-314, 2018.
- 5 Rashed, T. and Weeks, J.: Assessing vulnerability to earthquake hazards through spatial multicriteria analysis of urban areas, *International Journal of Geographical Information Science*, 17, 547-576, 2003.
- Renard, F.: Flood risk management centred on clusters of territorial vulnerability, *Geomatics, Natural Hazards and Risk*, 8, 525-543, 2017.
- Rezaei-Malek, M., Torabi, S. A., and Tavakkoli-Moghaddam, R.: Prioritizing disaster-prone areas for large-scale earthquakes' preparedness: Methodology and application, *Socio-Econ. Plan. Sci.*, 67, 9-25, 2019.
- 10 Rygel, L., O'sullivan, D., and Yarnal, B.: A Method for Constructing a Social Vulnerability Index: An Application to Hurricane Storm Surges in a Developed Country, *Mitigation and Adaptation Strategies for Global Change*, 11, 741-764, 2006.
- Sarkar, R. and Vogt, J.: Drinking water vulnerability in rural coastal areas of Bangladesh during and after natural extreme events, *International Journal of Disaster Risk Reduction*, 14, 411-423, 2015.
- 15 Schmidtlein, M. C., Deutsch, R. C., Piegorsch, W. W., and Cutter, S. L.: A Sensitivity Analysis of the Social Vulnerability Index, *Risk Analysis*, 28, 1099-1114, 2008.
- Schmidtlein, M. C., Shafer, J. M., Berry, M., and Cutter, S. L.: Modeled earthquake losses and social vulnerability in Charleston, South Carolina, *Applied Geography*, 31, 269-281, 2011.
- Shen, S., Cheng, C. X., Yang, J., and Yang, S. L.: Visualized analysis of developing trends and hot topics in natural disaster research, *PLoS One*, 13, 15, 2018.
- 20 Sherly, M. A., Karmakar, S., Parthasarathy, D., Chan, T., and Rau, C.: Disaster Vulnerability Mapping for a Densely Populated Coastal Urban Area: An Application to Mumbai, India, *Annals of the Association of American Geographers*, 105, 1198-1220, 2015.
- Shuang-Ye, W., Brent, Y., and Ann, F.: Vulnerability of coastal communities to sea-level rise: a case study of Cape May County, New Jersey, USA, *Climate Research*, 22, 255-270, 2002.
- 25 Simpson, D. and Katirai, M.: Indicator Issues and Proposed Framework for a Disaster Preparedness Index (DPI), University of Louisville, Louisville, KY, USA, 2006.
- Sister, C., Wolch, J., and Wilson, J.: Got green? addressing environmental justice in park provision, *GeoJournal* \$V 75, 2009. 229-248, 2009.
- Stow, D., Lopez, A., Lippitt, C., Hinton, S., and Weeks, J.: Object-based classification of residential land use within Accra, Ghana based on QuickBird satellite data, *International Journal of Remote Sensing*, 28, 5167-5173, 2007.
- 30 Toke, N. A., Boone, C. G., and Arrowsmith, J. R.: Fault zone regulation, seismic hazard, and social vulnerability in Los Angeles, California: Hazard or urban amenity?, *Earth Future*, 2, 440-457, 2014.
- Toké, N. A., Boone, C. G., and Arrowsmith, J. R.: Fault zone regulation, seismic hazard, and social vulnerability in Los Angeles, California: Hazard or urban amenity?, *Earth's Future*, 2, 440-457, 2014.
- Turvey, R.: Vulnerability Assessment of Developing Countries: The Case of Small-island Developing States, *Development Policy Review*, 25, 243-264, 2007.
- 35 UN: Report of the open-ended intergovernmental expert working group on indicators and terminology relating to disaster risk reduction. UNDRR (Ed.), Preventionweb, 2016.
- UNDP: The Real Wealth of Nations: Pathways to Human Development, UNDP, New York, NY, USA, 2010.
- UNISDR: Hyogo Framework for Action 2005-2015: Building the Resilience of Nations and Communities to Disasters, 2007.
- 40 Unwin, D. J.: GIS, spatial analysis and spatial statistics, *Progress in Human Geography*, 20, 540-551, 1996.
- van der Geest, K., Burkett, M., Fitzpatrick, J., Stege, M., and Wheeler, B.: Climate change, ecosystem services and migration in the Marshall Islands: are they related?, *Climatic Change*, doi: 10.1007/s10584-019-02648-7, 2020. 2020.
- Vandermeersch, H., Vos, S., and Scheerder, J.: Who's joining the club? Participation of socially vulnerable children and adolescents in club-organised sports, *Sport, Education and Society*, 20, 941-958, 2015.
- 45 Vilches, O. R., Carrillo, K. S., Reyes, C. M., and Castillo, E. J.: Post -catastrophe social -environmental effects in vulnerable coastal areas affected by the tsunami of 02/27/2010 in Chile, *Interciencia*, 39, 383-390, 2014.
- Vinchon, C., Alexander, D., Barbat, A., Cardona, O., Carreño, M., Contreras, D., Decker, B., Eidsvig, U., Kienberger, S., Papathoma-köhle, M., Miniati, R., Pelling, M., Pratzler-Wanczura, S., Schneiderbauer, S., Ulbrich, T., Vidar, B., Welle, T., Angignard, M., Carvalho, S., Garcin, M., Marulanda, M.-C., Morabito, M., Pedoth, L., Pelling, M., Tedim, F., and Zehra, R.: Assessing vulnerability to natural hazards in Europe: From Principles to Practices. A manual on concept, methodology and tools, European Commission DG Environment, 2011.
- 50 Walker, B. B., Taylor-Noonan, C., Tabbernor, A., McKinnon, T. B., Bal, H., Bradley, D., Schuurman, N., and Clague, J. J.: A multi-criteria evaluation model of earthquake vulnerability in Victoria, British Columbia, *Nat. Hazards*, 74, 1209-1222, 2014.
- Warner, K., Kuhllicke, C., Vries, D. d., Sakdapolrak, P., Wutich, A., Real, B., Briones, F., and Verjee, F.: Perspectives on Social Vulnerability, *UNU-EHS, Bonn, Germany* 06/2007, 132 pp., 2007.
- 55 Wisner, B. and Uitto, J.: Life on the edge: urban social vulnerability and decentralized, citizen-based disaster risk reduction in four large cities of the Pacific Rim. In: *Facing global environmental change*, al, B. H. e. (Ed.), Springer, Berlin, 2009.

- Wolch, J., Wilson, J. P., and Fehrenbach, J.: Parks and Park Funding in Los Angeles: An Equity-Mapping Analysis, *Urban Geography*, 26, 4-35, 2005.
- Yuan, H. H., Gao, X. L., and Qi, W.: Fine-Scale Spatiotemporal Analysis of Population Vulnerability to Earthquake Disasters: Theoretical Models and Application to Cities, *Sustainability*, 11, 19, 2019a.
- 5 Yuan, H. H., Gao, X. L., and Qi, W.: Modeling the fine-scale spatiotemporal pattern of earthquake casualties in cities: Application to Haidian District, Beijing, *International Journal of Disaster Risk Reduction*, 34, 412-422, 2019b.
- Zebardast, E.: Constructing a social vulnerability index to earthquake hazards using a hybrid factor analysis and analytic network process (FANP) model, *Nat. Hazards*, 65, 1331-1359, 2013.
- 10 Zeng, J., Zhu, Z. Y., Zhang, J. L., Ouyang, T. P., Qiu, S. F., Zou, Y., and Zeng, T.: Social vulnerability assessment of natural hazards on county-scale using high spatial resolution satellite imagery: a case study in the Luogang district of Guangzhou, South China, *Environmental Earth Sciences*, 65, 173-182, 2012.
- Zhang, N. and Huang, H.: Assessment of world disaster severity processed by Gaussian blur based on large historical data: casualties as an evaluating indicator, *Nat. Hazards*, 92, 173-187, 2018.
- 15 Zhou, Y., Li, N., Wu, W., Wu, J., and Shi, P.: Local Spatial and Temporal Factors Influencing Population and Societal Vulnerability to Natural Disasters, *Risk Analysis*, 34, 614-639, 2014.