

# Responses to the Editor's and Reviewers' Comments and Suggestions

**Journal:** Natural Hazards and Earth System Sciences (NHESS)

**Manuscript number:** nhess-2019-212

**Manuscript title:** Integrated sea storm management strategy: the 29 October 2018 event in the Adriatic Sea

We would like to thank the Editor and Reviewers for their valuable comments and effort to improve the manuscript. We have responded to all comments as can be seen in the following list. We believe that with these revisions the manuscript has been improved and we hope that it is now ready for publication.

The original Editor's and Reviewers' comments and suggestions are shown in regular typeface, while our responses are shown in italics. The line and figures numbers we use refer to the revised document.

## Response to the Editor

- E1** a) I agree that you may not include in this study a detailed evaluation of the individual model performances (see reviewer 2), but I suggest that you insert a paragraph describing how to deal with the different accuracies of the models in a future operational framework.

*Response: Following the Editor's suggestion, we included a paragraph in the "Summary and concluding discussion" (Page 14, lines 15-22) describing the future development to account for the different accuracies of the models in the multi-model ensemble. The sentence reads: "It is not straightforward what averaging weights should be used for the multi-model ensemble forecast and therefore we used equally weighted members, despite the forecasts which are more precise than others should have more importance in the TMES (Salighehdar et al., 2017; Schevenhoven and Selten, 2017). Here we applied a simple average of the forecasts at every timestamp to compute the ensemble mean, but more sophisticated methods based on weighting function determined by comparison of the single model results with near real-time observations can be implemented in future (Di Liberto et al., 2011; Salighehdar et al., 2017). Taking advantage of the near real-time observations acquired by the aggregated monitoring network, the root mean square error of the individual forecast will be evaluated and stored for long-term statistics." See also the response to comment R2.4.*

- E2** Further, I note that there are two issues that are relevant for early warning systems and are likely important component in an operational framework. They are briefly mentioned in the "Summary and Conclusions", but you do not mention the former studies already available in the literature, specifically for Venice:

- b) the inclusion of a data assimilation procedure (e.g. Lionello et al., 2006)
- c) the probabilistic approach provided by a single model ensemble prediction. Note that papers showing this specifically for Venice are already available (e.g. Mel and Lionello, 2014a,b, 2016).

I suggest you paper your text is updated including these (and other) references.

*Response: We thank the editor for bringing these useful citations to our attention. We cited the study of Lionello et al. (2006) at Page 15 (line 25) when discussing the importance of data assimilation for storm surge forecasting in the Adriatic Sea. At Page 14 (lines 6-13), we extended the discussion of the probabilistic forecast by including the following sentence regarding single model ensemble prediction: “The awareness of the prediction uncertainties and errors has led many operational and research flood forecasting systems around the world to move toward numerical forecasts based on a probabilistic concept: the ensemble technique (Cloke and Pappenberger, 2009). In this contest, a probabilistic forecasting system could be based on perturbation of initial conditions, forcing and parameters of a single model (Flowerdew et al., 2010; Bernier and Thompson, 2015; Salighehdar et al., 2017). Such approach has been already applied to the Adriatic Sea for improving storm surge forecast and providing a realistic estimate of the prediction uncertainty (Mel and Lionello, 2014a,b, 2016; Bajo et al., 2019).”*

## Response to Reviewer #1

**R1.1** This manus describes a collaborative effort between agencies in Italy, Croatia and Slovenia to develop an integrated system for storm handling. A Transnational Multimodel Ensemble System (TMES) is used to handle the input from a number of operational model systems. The severe storm of 29 October 2018 is used as a convincing example of the usefulness of the new system. I have one comment: Why are no products from the Copernicus Marine Environment Monitoring System applied and part of the TMES? They are readily available and could provide an even more robust ensemble system.

*Response: We thank the reviewer for the positive comment. The developed Transnational Multimodel Ensemble System (TMES) integrates also some Copernicus Marine Environment Monitoring System (CMEMS) products. In details, the Mediterranean Forecasting System (managed by CMCC) and the MED-Waves (managed by HCMR) results are used in sea level height and the wave multi-model ensembles, respectively (see Tables 1 and 2 in the manuscript). In the revised manuscript (Page 7, line 6), we enhanced the description of the operational forecasting systems used in the TMES and clarified that some products are retrieved from CMEMS.*

## Response to Reviewer#2

**R2.1** General comments: I have just finished reviewing the manuscript entitles “Integrated sea storm management strategy: the 29 October 2018 event in the Adriatic Sea” by Ferrarin et al. Overall the article provides a good description of system that combines information from different forecasting models in order to provide an ensemble prediction. I believe that the article is of interest for the journal. It has a high technical character that is justifiable by the nature of the topic. However, I believe that in order for the manuscript to be accepted some significant changes need to be addressed. There are some fundamental misuse of the risk terminology that need further clearance this is principally in the introduction but also thought the document (see detailed comments). Introduction needs improvement especially the first part with the incorporation of objectives of the study. Some interesting information of the state of the art is placed in conclusion but

I think is more fitted for the introduction. A more detail comparison of the individual models and the model ensemble with observed data would be very interesting.

*Response: We appreciate the comments and we improved the manuscript following all reviewer's suggestions. We corrected the risk terminology over the whole document (see the responses to comments R2.7, R2.10 and R2.13). Some information we wrote in the "Conclusions" at first, had been repositioned within the "Introduction" and "Material and methods" (see also the response to comment R2.6). The aim of the study is presented at the end of the introduction section.*

*This manuscript aims at presenting the structure for sharing knowledge, data and forecasts in order to improve the prevention and protection measures to sea storm emergencies. Therefore, despite the multi-model ensemble system is a fundamental component of the developed systems, we decided not to include in this study a detailed evaluation of the individual models performance. The comparison of the models and the ensemble with observed data will be the subject of a future work which will consider the analysis of a more complete dataset of sea storm events (see also the response to comment R2.4).*

**R2.2** Forecasting System: Given technical character and it is describing a complex method of forecast ensemble I believe that Tables 2 and 3 can be expanded to include details indicating the forecast characteristics (e.g. forecast window, update time) would be very interesting for some readers. Also the meteorological models resolution are interesting, some data are already given in the text but I believe that a thorough description could be useful. I also believe that Table 1 and 2 are not very interesting and good be eliminated or moved to an appendix.

*Response: Following the reviewer's suggestion we included the forecast range and the meteorological models resolution in Table 3 and 4 (now Table 1 and 2). The tables providing the monitoring stations details have been moved to the appendix.*

**R2.3** Forecasting System: It is not clear if the IWS system is providing or receiving detail EWS. It would be useful to present the EWS in Figure 2. I believe that Figure 2 needs to be improved to provide a more detail presentation of the system. What about the other areas of the Adriatic that are not covered by the EWSs. Is a hazard map produced and how what kind of topographic information is used?

*Response: As described in section 2.3 and 3.3, IWS had been designed to provide multi-model forecast products to existing early warning systems, developed in areas where a deep knowledge of the coastal dynamics and high-resolution datasets (topography and bathymetry) are available. These concept were clarified in the manuscript (Page 5 line 7, Page 8 lines 20-21). Moreover, following the reviewer's suggestion we improved Figure 2.*

*The multi-model results have been used to provide a basin-wide overview of the physical processes acting in coastal areas and responsible for storm related hazards. TMES products are combined with the coastal characteristics (coast material and slope) provided by the MCD database for computing the total water level (TWL). For the coastal segments characterized by sandy beaches, TWL was computed combining the sea level height, wave setup and wave runup according to the Stockdon's formula ( $R_2$ , the 2% exceedance level of runup maxima; Stockdon et al., 2006). The resulting maps (Fig. 7) provide a basin-wide overview of the physical processes acting in coastal areas and responsible for storm related hazards. The forecasted TWLs are made available to the IWS users and can be combined with a digital elevation model (DEM) of the coast for estimating inundation intensity and extend. See also the responses to comments R2.5 and R2.14.*

**R2.4 Storm Predictability:** The section is well structured and with some interesting figures, however a more detailed analysis and statistical representation of the individual model and the model ensemble should be presented. An interesting question especial for regional assemblies is the spatial performance of the different model. Although Fig 5 present a good synoptic view of the ensemble performance a more detailed look (Fig6) reveals that there are two models that substantially under predict the sea level height for the final part of the storm. Why is this happening, are there any performance criteria for the models to enter the ensemble?

*Response: The multi-model ensemble forecasting model was created with the aim of combining together the existing available operational systems without providing a critical review of the individual model performance. The storm event of 29<sup>th</sup> October 2018 is here taken as a pilot study for applying and testing the developed approach. At the same time, the analysis of the model results provide an example of the scatter of the individual forecasts (ensemble standard deviation) to point the attention on the uncertainty of the sea condition prediction. To our opinion, the awareness of these uncertainties and prediction errors is crucial and should be accounted for in managing coastal risks related to sea storms.*

*Actually, there is no performance criteria for the models to enter the ensemble, but, as discussed at Page 14 (lines 15-22), weighting function determined by comparison of the single model results with observations will be implemented in future in the TMES. Therefore, the forecasts which are more precise than others will have more importance in the multi-model ensemble (see also the response to comment E1.*

**R2.5 Storm Hazard and Impact assessment on the coast:** Is the model ensemble always underestimates the events? Do you have other examples that also indicate the MEAN+STD is a better estimate for the events? Related with the hazard the authors mention that the calculated on each coastal assessment unit. Some more information on how many units were identified in the area and how they are distributed it would be interesting for the reader. It is unclear if the Stockton model is used in all areas. If yes this is contradicting the previous comments of the authors. Finally, the detail description of the storm in the study area gives valuable information however a directed comparison of the hazard intensity and extend predicted by the ensemble and the one observed is missing. Such comparison is important in order to identify the advantages of such a model.

*Response: We thank the reviewer for the useful comment, which helped us to clarify the approach adopted in this study. In this manuscript, we present the results of the TMES application to only one storm event and therefore we cannot indicate that the MAX (MEAN+STD) sea condition scenario is always the better estimation. It is however true that, generally, the wind condition over the Adriatic Sea are underestimated by many meteorological models (Cavaleri and Bertotti, 2004; Cavaleri et al., 2019). For these reasons, the most severe sea condition scenario can be considered for the investigated area as a conservative estimation of the peak storm conditions to be used for coastal risk management. As stated in the response to comment R2.1, the comparison of the models and the ensemble with observed data related to a more complete dataset of sea storm events will be the subject of a future work.*

*We clarified that, in order to assess the perception of the physical processes acting in coastal areas and responsible for storm related hazards, the coast is subdivided into segments of variable length in function of morphology, human settlements and administrative boundaries. The coastal assessment units were selected according to the Mediterranean Coastal Database (MCD) developed by Wolff et al. (2018). We let the reader to refer to*

above-cited article for more details about how the coastal units were defined. The MCD segments have an average length of 4.5 km. However, as discussed at Page 14 lines 29-35, the MCD segments are sometimes too coarse to represent complex morphologies, especially in confined coastal systems (lagoons) and along the eastern rocky coast.

The Stockdon's formula was improperly applied to all coastal segments. As specified in the response to comment R2.14, we corrected the manuscript specifying that the Stockdon's formula is applied only to the coastal segments characterised by sandy beaches (Page 8, lines 15-17).

We do agree with the reviewer that a more detailed comparison between predicted and observed hazard would add a significant contribution to the presented results. Unfortunately, detailed coastal observations of the hazard intensity and extend are not available. For the Emilia-Romagna and Slovenia coasts, flooding and erosion were reported by the local authorities and therefore the comparison between predicted and observed is qualitative and not quantitative. In the case of the City of Venice, the situation is similar, even if a more detailed information of the extend of the flooding is available as function of the sea level (see Figure 11). The flooding maps reported in the manuscript were obtained by the municipality imposing the sea level height observed/predicted at Punta della Salute (at intervals of 10 cm) to a centimetre accurate digital terrain model of the city.

**R2.6** Summary and Concluding discussion: The start of this section gives valuable detail information of the system that they should be placed in section 2.3.

*Response: Following the reviewer's suggestion we moved part of the first paragraph of the "Summary and Concluding discussion" section to the "Material and methods" section.*

**R2.7** Page 2 - Line 1: "Sea storm ... directly impact on the citizens quality of life". This statement is not exactly true in my opinion. Sea storms are a natural phenomenon and they do not affect the quality of life. Possible risks associated with sea storms can have this effect. The second paragraph provides a description of the process based models and it is more suited for the introduction.

*Response: We reformulated the first paragraph on the introduction as follow (Page 2, lines 2-9): "Sea storms represent the main threat in coastal areas. In fact, they can cause a range of potential hazards, such as coastal erosion and inundation, as well as damages to infrastructure and to the important cultural heritage exposed to these phenomena (Chauvillon et al., 2017; Reimann et al., 2018; Vousdoukas et al., 2018). Along the coast, extreme storms can also significantly affect businesses activities, such as aquaculture, fisheries, tourism and beach facilities. The potential future ...".*

**R2.8** Page 2 - Line 14: The reference of Roland et al., 2009 is not appropriate for wave setup maybe an older reference would be more appropriate (e.g. Longuet-Higgins, M.S., Stewart, R.W., 1963. A note on wave set-up. Journal of Marine Research 21, 4-10.).

*Response: Following the reviewer's suggestion we replaced Roland et al. (2009) with the more appropriate Longuet-Higgins and Steward (1963).*

**R2.9** Page 2 - Line 14: "they travel up and down the beach". Are you referring to swash processes?

*Response: Yes, we refer to swash processes. We modified the sentence (Page 2, line 14) as follow: "... they travel up and down the beach (swash processes)".*

**R2.10** Page 2 - Line 16-19: "Coastal flooding, erosion, impacts on ecosystems, damages to infrastructures and productive activities can worsen if combined with the absence of adequate

early warning systems, coordinated strategies, intervention procedures, coastal management and planning with significant related economic costs”. This sentence is mixing hazards and consequences with primary measures and management strategies. This can result in a confusion of terms that is undesirable. For example coastal flooding and erosion are not related with EWS. The absence of an EWS can result in increased damages if proper disaster risk reduction measures (DRRs) are not implemented. I suggest to restructure the sentence.

*Response: We concur with the reviewer that the mentioned sentence was not properly formulated. We modified it as follows (Page 2, lines 30-32): “Coastal flooding of urban areas, beach erosion, damages to infrastructures and productive activities can worsen if combined with the absence of an adequate sea storm management strategy with significant related economic costs.”*

**R2.11** Page 3 and throughout the document: The terms “Bora” and “Sirocco” are local wind names it is better to use italic font style or directly use the English name.

*Response: Following the reviewer’s suggestion we used italic font style for the terms Bora and Sirocco.*

**R2.12** Figure 2: What is the difference between the black thin lines the blue arrow and the dash line? Why the TMES exchanges information only with the Recourse layer (maybe a better explanation of what the TMES is doing could be useful). Only the resources layer is delimited I think it would be nice to show all 6 layers limits.

*Response: According to the reviewer’s suggestion we improved Figure 2 (see also response to comment R2.5). At Page 5 (lines 21 and 26), we clarified that the results from existing operational forecasting systems are stored into the Resource Layer and made available to the multi-model ensemble system. TMES outputs are also stored in the Resource Layer.*

**R2.13** Page 7 - Line 6-7: “hazard maps ... to identify vulnerable areas” there is a mix of the two terms that is common but not a good practice. I suggest following UNISDR Terminology where hazard is related with the physical aspects of the storm the coastal area and vulnerability with the socioeconomic aspects.

*Response: We concur with the reviewer that the hazard and vulnerability terms were not properly used. We modified the manuscript in the Introduction, section 2.3 and section 3.3 to clarify the terminology adopted in this study.*

**R2.14** Page 7 - Line 25: “It must be pointed out that the widely used Stockdon’s...” The Stockdon formula is not applicable in rocky and gravel beaches. The problem is not the underestimation of the runup is the use of an inappropriate formula that results in underestimation of the runup.

*Response: The Stockdon’s formula was improperly applied to all coastal segments.  $R_2$  results presented in section 2.5 (Fig. 7) are now reported only for the coastal segments characterized by sandy beaches (see also response to comment R2.5).*

**R2.15** Figure 3: Substitute the “C” by “L” for low pressure.

*Response: Done.*

**R2.16** Page 12 - Line 11-13: There is a large number of local names that are not shown in the figure and is difficult to follow by the reader. Please add a more detail figure.

*Response: We removed the local names from the manuscript and simplified the sentence, that now reads as (Page 13, lines 3-4): “As a consequence, the sea flooded several coastal locations, where the firemen set up anti-flooding barrages.”*

**R2.17** Page 14 - Line 27: “adaptation capacity” the dune and berm characteristics of a beach are not the adaptation capacity. The term adaptation is related with the ability of the system to overcome long term changes in forcing factors. Beach and berm characteristics can be combined with physical parameters (e.g. wave height water level) to calculate process based indexes that can serve as hazard intensity and extend parameters. A review of such indexes can be found in Ferreira, Ó., Plomaritis, T.A., Costas, S., 2017. Process based indicators to assess storm induced coastal hazards. *Earth-Science Reviews* 173, 159-167.

*Response: We concur with the reviewer that there was a mistake in the mentioned sentence. We modified the sentence as follow (Page 14, lines 35, Page 15, lines 1-2): “... comparing the magnitude of the impact (wave run-up for inundation and beach/shoreline retreat for erosion) with the morphological characteristics of the system (dune/berm height for inundation and beach width for erosion).”*

*We thank the reviewer for bringing this useful citation to our attention. We cited the study of Ferreira et al. (2017) in the Introduction (Page 2, line 19) and the discussion (Page 14, line 34).*

## References

- Bajo, M., Medugorac, I., Umgiesser, G., and Orlić, M.: Storm surge and seiche modelling in the Adriatic Sea and the impact of data assimilation, *Quart. J. Roy. Meteor. Soc.*, 145, 2070–2084, <https://doi.org/10.1002/qj.3544>, 2019.
- Bernier, N. B. and Thompson, K. R.: Deterministic and ensemble storm surge prediction for Atlantic Canada with lead times of hours to ten days, *Ocean Model.*, 86, 114 – 127, <https://doi.org/10.1016/j.ocemod.2014.12.002>, 2015.
- Cavaleri, L. and Bertotti, L.: Accuracy of the modelled wind and wave fields in enclosed seas, *Tellus A*, 56, 167–175, <https://doi.org/10.3402/tellusa.v56i2.14398>, 2004.
- Cavaleri, L., Bajo, M., Barbariol, F., Bastianini, M., Benetazzo, A., Bertotti, L., Chiggiato, J., Davolio, S., Ferrarin, C., Magnusson, L., Papa, A., Pezzutto, P., Pomaro, A., and Umgiesser, G.: The October 29, 2018 storm in Northern Italy - an exceptional event and its modeling, *Prog. Oceanogr.*, 178, 102178, <https://doi.org/10.1016/j.pocean.2019.102178>, 2019.
- Chaumillon, E., Bertin, X., Fortunato, A., Bajo, M., Schneider, J.-L., Dezileau, L., Walsh, J. P., Michelot, A., Chauveau, E., Créach, A., Hénaff, A., Sauzeau, T., Waeles, B., Gervais, B., Jan, G., Baumann, J., Breilh, J.-F., and Pedreros, R.: Storm-induced marine flooding: lessons from a multidisciplinary approach, *Earth-Sci. Rev.*, 165, 151 – 184, <https://doi.org/10.1016/j.earscirev.2016.12.005>, 2017.
- Cloke, H. L. and Pappenberger, F.: Ensemble flood forecasting: A review, *J. Hydrol.*, 375, 613 – 626, <https://doi.org/10.1016/j.jhydrol.2009.06.005>, 2009.
- Di Liberto, T., Colle, B. A., Georgas, N., Blumberg, A. F., and Taylor, A. A.: Verification of a Multimodel Storm Surge Ensemble around New York City and Long Island for the Cool Season, *Wea. Forecasting*, 26, 922–939, <https://doi.org/10.1175/waf-d-10-05055.1>, 2011.
- Ferreira, O., Plomaritis, T. A., and Costas, S.: Process-based indicators to assess storm induced coastal hazards, *Earth-Sci. Rev.*, 173, 159 – 167, <https://doi.org/10.1016/j.earscirev.2017.07.010>, 2017.

- Flowerdew, J., Horsburgh, K., Wilson, C., and Mylne, K.: Development and evaluation of an ensemble forecasting system for coastal storm surges, *Q.J.R. Meteorol. Soc.*, 136, 1444–1456, <https://doi.org/10.1002/qj.648>, 2010.
- Lionello, P., Sanna, A., Elvini, E., and Mufato, R.: A data assimilation procedure for operational prediction of storm surge in the northern Adriatic Sea, *Cont. Shelf Res.*, 26, 539–553, <https://doi.org/10.1016/j.csr.2006.01.003>, 2006.
- Longuet-Higgins, M. S. and Steward, R. W.: A note on wave set-up, *J. Mar. Res.*, 21, 4–10, 1963.
- Mel, R. and Lionello, P.: Verification of an ensemble prediction system for storm surge forecast in the Adriatic Sea, *Ocean Dyn.*, 64, 1803–1814, <https://doi.org/10.1007/s10236-014-0782-x>, 2014a.
- Mel, R. and Lionello, P.: Storm surge ensemble prediction for the city of Venice, *Wea. Forecasting*, 29, 1044–1057, <https://doi.org/10.1175/WAF-D-13-00117.1>, 2014b.
- Mel, R. and Lionello, P.: Probabilistic dressing of a storm surge prediction in the Adriatic Sea, *Adv. Meteorol.*, 3764519, <https://doi.org/10.1155/2016/3764519>, 2016.
- Reimann, L., Vafeidis, A. T., Brown, S., Hinkel, J., and Tol, R. S. J.: Mediterranean UNESCO World Heritage at risk from coastal flooding and erosion due to sea-level rise, *Nat. Commun.*, 9, 4161, <https://doi.org/10.1038/s41467-018-06645-9>, 2018.
- Roland, A., Cucco, A., Ferrarin, C., Hsu, T.-W., Liau, J.-M., Ou, S.-H., Umgiesser, G., and Zanke, U.: On the development and verification of a 2d coupled wave-current model on unstructured meshes, *J. Mar. Syst.*, 78, Supplement, S244–S254, <https://doi.org/10.1016/j.jmarsys.2009.01.026>, 2009.
- Salighehdar, A., Ye, Z., Liu, M., Florescu, I., and Blumberg, A. F.: Ensemble-Based Storm Surge Forecasting Models, *Wea. Forecasting*, 32, 1921–1936, <https://doi.org/10.1175/waf-d-17-0017.1>, 2017.
- Schevenhoven, F. J. and Selten, F. M.: An efficient training scheme for supermodels, *Earth Syst. Dynam.*, 8, 429–438, <https://doi.org/10.5194/esd-8-429-2017>, 2017.
- Stockdon, H. F., Holman, R. A., Howd, P. A., and Sallenger, A. H.: Empirical parameterization of setup, swash, and runup, *Coast. Eng.*, 53, 573–588, <https://doi.org/10.1016/j.coastaleng.2005.12.005>, 2006.
- Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Bianchi, A., Dottori, F., and Feyen, L.: Climatic and socioeconomic controls of future coastal flood risk in Europe, *Nat. Clim. Chang.*, 8, 776–780, <https://doi.org/10.1038/s41558-018-0260-4>, 2018.
- Wolff, C., Vafeidis, A. T., Muis, S., Lincke, D., Satta, A., Lionello, P., Jimenez, J. A., Conte, D., and Hinkel, J.: A Mediterranean coastal database for assessing the impacts of sea-level rise and associated hazards, *Sci. Data*, 5, 180 044, <https://doi.org/10.1038/sdata.2018.44>, 2018.

# Integrated sea storm management strategy: the 29 October 2018 event in the Adriatic Sea

Christian Ferrarin<sup>1</sup>, Andrea Valentini<sup>2</sup>, Martin Vodopivec<sup>3</sup>, Dijana Klaric<sup>4</sup>, Giovanni Massaro<sup>5</sup>, Marco Bajo<sup>1</sup>, Francesca De Pascalis<sup>1</sup>, Amedeo Fadini<sup>1</sup>, Michol Ghezzi<sup>1</sup>, Stefano Menegon<sup>1</sup>, Lidia Bressan<sup>2</sup>, Silvia Unguendoli<sup>2</sup>, Anja Fettich<sup>3</sup>, Jure Jerman<sup>3</sup>, Matjaž Ličer<sup>6</sup>, Lidija Fustar<sup>4</sup>, Alvisè Papa<sup>5</sup>, and Enrico Carraro<sup>7</sup>

<sup>1</sup>CNR - National Research Council of Italy, ISMAR - Marine Sciences Institute, Venice, Italy

<sup>2</sup>Arpaè-SIMC - Agency for Prevention, Environment and Energy of Emilia-Romagna, Hydro-Meteo-Climate Service, Bologna, Italy

<sup>3</sup>Slovenian Environment Agency, Ljubljana, Slovenia

<sup>4</sup>Meteorological and Hydrological Service of Croatia, Zagreb, Croatia

<sup>5</sup>Tide Forecast and Early Warning Center, City of Venice, Venice, Italy

<sup>6</sup>National Institute of Biology, Ljubljana, Slovenia

<sup>7</sup>City of Venice - European Policies, Venice, Italy

**Correspondence:** Christian Ferrarin (c.ferrarin@ismar.cnr.it)

**Abstract.** Addressing coastal risks related to sea storms requires an integrative approach which combines monitoring stations, forecasting models, early warning systems and coastal management and planning. Such great effort is sometimes possible only through transnational cooperation, which becomes thus vital to face effectively and promptly these marine events which are responsible for several damages impacting on the environment and citizens' life. Here we present a shared and interoperable system to allow a better exchange and elaboration of information related to sea storms among countries. The proposed Integrated Web System (IWS) is a combination of a common data system for sharing ocean observations and forecasts, a multi-model ensemble system, a geoportal and interactive geo-visualization tools to make results available to the general public. Multi-model ensemble mean and spread for sea level height and wave characteristics are used to describe three different sea condition scenarios. IWS is designed to provide sea state information required for issuing coastal risk alerts over the analysed region, as well as for being easily integrated into existing local early warning systems. This study describes the application of the developed system to the exceptional storm event of 29<sup>th</sup> of October 2018, that caused severe flooding and damages to coastal infrastructures in the Adriatic Sea. The forecasted ensemble products were successfully compared with in situ observations. The hazards estimated by integrating IWS results in existing early warning systems were confirmed by documented storm impacts along the coast of Slovenia, Emilia-Romagna and the City of Venice. For the investigated event, the most severe simulated scenario resulted to provide a realistic and conservative estimation of the peak storm conditions to be used in coastal risk management.

# 1 Introduction

Sea storms represent the main threat in coastal areas. In fact, they ~~directly impact on the citizens' quality of life (especially in urban areas where part of the inhabited areas is seldom covered by water), they create damages to~~ can cause a range of potential hazards, such as coastal erosion and inundation, as well as damages to infrastructure and to the important cultural heritage exposed to these phenomena ~~, and they affect businesses too~~ ((Chaumillon et al., 2017; Reimann et al., 2018; Vousdoukas et al., 2018a). Along the coast, extreme storms can also significantly affect businesses activities, such as aquaculture, fisheries, tourism ; beach facilities) and environment at large (coastal erosion, floods) (Chaumillon et al., 2017; Reimann et al., 2018; Vousdoukas et al., 2018a) and beach facilities. The potential future effects of global climate change emphasise the need for strategies based on an anticipatory approach particularly in coastal areas at immediate and high risk (Hinkel et al., 2014; Vousdoukas et al., 2018b). This is particularly true for coastal wetlands if enough additional accommodation space will not be created through careful nature-based adaptation solutions to coastal management (Schuerch et al., 2018).

Coastal flooding is induced by extreme sea levels, determined by the increase in sea level caused by strong winds and low atmospheric pressure (storm surge), often in combination with high tides (Muis et al., 2016). Under such extreme meteorological conditions, the coast could be also vulnerable by stormy waves with potential damages to infrastructures and erosion. Moreover, when waves reach the coast they interact with the bathymetry and drive an additional increase in water levels through wave setup (Roland et al., 2009) (Longuet-Higgins and Steward, 1963) and they travel up and down the beach before being reflected seaward (swash processes). The maximum vertical excursion of wave uprush on a beach or structure above the still water level is called the wave runup (Sorensen, 1997).

~~Coastal flooding, erosion, impacts on ecosystems,~~ The water levels along the coast can be estimated by numerical models and combined with a digital elevation model (DEM) for forecasting inundation intensity and extend. Several methodologies have been developed and applied at the basin and local scales for estimating hazard maps for coastal flooding (Hinkel et al., 2014; Vousdoukas et al., 2018a). It must be taken into account that meteorological and ocean models provide just an approximation of reality, despite their continuous development and improvements. Moreover, the interactions between atmospheric, oceanic and coastal processes are not fully understood, resulting in large uncertainties in the predictions of coastal flooding, in particular, under extreme conditions (Baart et al., 2011; Zou et al., 2013). This is mainly due to the chaotic nature of the atmosphere and the complexity of the air-sea interactions across scales over several orders of magnitude (Schevenhoven and Selten, 2017). Small errors in the initial conditions of a numerical weather prediction model grow rapidly and affect predictability; forecasted atmospheric conditions are then affected by errors (Molteni et al., 2001). However, as stated by Flowerdew et al. (2010), atmospheric forcing is not the only source of uncertainty in storm surge forecasting. Many other sources of uncertainty, as the model numerics, resolution, parametrization, boundary conditions and initial sea state, contribute non-linearly to the final forecast uncertainty.

Coastal flooding of urban areas, beach erosion, damages to infrastructures and productive activities can worsen if combined with the absence of ~~adequate early warning systems, coordinated strategies, intervention procedures, coastal management and planning an adequate sea storm management strategy~~ with significant related economic costs (Hinkel et al., 2014; Prah et al., 2018). The difficulty of reacting promptly to extreme events is also connected to the lack of shared data and know-how.

Recognizing the importance of information sharing for disaster risk reduction, human safety and well-being, the World Meteorological Organization (WMO, <https://public.wmo.int/>) promotes the standardization and exchange of observations since 1873. Similarly, the Permanent Service for Mean Sea Level (PSMSL, <http://www.psmsl.org/>) and the Global Sea Level Observing System (GLOSS, <http://www.ioc-sealevelmonitoring.org/>) are responsible for the collection, publication, analysis and interpretation of sea level data from the global network of tide gauges. In the same direction, at European level, the Copernicus Marine Environmental Monitoring Service (CMEMS, <http://marine.copernicus.eu/>), the European Marine Observation and Data Network (EMODnet, <http://www.emodnet.eu/>) and the European Global Ocean Observing System (<http://eurogoos.eu/>) aim at sharing information from both satellite and in situ observations, as well as state-of-the-art analyses and daily forecasts, which offer an unprecedented capability to observe, understand and anticipate marine environment events.

Despite such international effort on data sharing, as weather, climate and ocean know no national boundaries, the insufficient level of cooperation among neighbouring countries is often a cause of ineffective actions at local level and missed opportunities to collaborate with other actors to increase overall preparedness to sea storms (Chaumillon et al., 2017).

The problem of sea storms is particularly relevant for the Adriatic Sea, where extreme sea levels are higher than in other parts of the Mediterranean basin (Marcos et al., 2009) and several coastal cultural World Heritage sites (<http://whc.unesco.org/>) at risk from coastal flooding and erosion are located (Prahl et al., 2018; Reimann et al., 2018). This study presents the management approach for sea storm hazard initiated as part of the I-STORMS (Integrated Sea sTORM Management Strategies) project for the coastline of the Adriatic-Ionian macro-region (<https://istorms.adrioninterreg.eu/>). This manuscript describes a joint strategy to safeguard the coast from sea storm emergencies by sharing knowledge, data and forecasts among involved countries and improving their capacities in terms of early warning and management procedures. This study focuses on the recent exceptional storm event of 29<sup>th</sup> of October 2018, which is here taken as a pilot study for applying and testing the developed approach.

## 1.1 Study area

The Adriatic and Ionian seas are part of the Mediterranean Sea positioned between the eastern coastline of Italy, countries of the Balkan Peninsula (from Slovenia, south through Croatia, Bosnia-Herzegovina, Montenegro and to Albania) and Greece. The Adriatic Sea is an 800-km-long, 150-km-wide elongated semi-enclosed basin interacting with the Ionian Sea through the Otranto Strait in the southern part (Fig. 1). The shallow northern Adriatic Sea is the Mediterranean sub-basin where storm surges reach higher values (Marcos et al., 2009), mainly triggered by strong south-easterly moist and warm wind, called ~~Sirocco~~ *Sirocco*. For this reason, in this area storm surges and waves have been deeply investigated in the past (Orlić et al., 1992; Bajo and Umgiesser, 2010; Cavaleri et al., 2010; Lionello et al., 2012; Medugorac et al., 2015; Ferrarin et al., 2017; Pomaro et al., 2017; Vilibić et al., 2017; Bajo et al., 2019; Ferrarin et al., 2019). Tidal dynamics are particularly evident in the northern Adriatic Sea, where the most energetic tidal constituents, the semi-diurnal  $M_2$  and the diurnal  $K_1$ , reach amplitudes of 27 and 18 cm, respectively (Ferrarin et al., 2017).

The weather in the Adriatic area is strongly influenced by local orography and small-scale processes (Pasarić et al., 2009). The use of high-resolution meteorological models is essential to capture the temporal and spatial inhomogeneity of north-easterly ~~Bora~~ *Bora* winds, characterised by topographically controlled high-speed wind jets along the eastern shore (Signell

et al., 2005; Davolio et al., 2015). The same holds for ~~Siroeeo~~Sirocco: global and regional numerical models have been shown to consistently underestimate its speed due to the fact that orography, and hence the channelling of the air flow, is not well represented at typical model resolution (Cavaleri and Bertotti, 2004). Long term analyses of general wind conditions over the Adriatic basin further indicate a trend of reduction of the intensity of wind events - mostly due to ~~Bora-Bora~~ (Pirazzoli and Tomasin, 2003) and a general increase in terms of frequency, mostly associated to the increasing storminess of Scirocco (Pomaro et al., 2017).

The eastern and western coasts of the Adriatic Sea greatly differ in appearance and are therefore differently impacted by sea storms. The western coast is largely sedimentary with mild sloping and sandy beaches, while the eastern coast is composed of many islands and headlands and is generally high and rocky. Due to its alluvial origin, natural subsidence occurs in the northwestern Adriatic Sea because of compaction of fine-grained deposits (Carbognin and Tosi, 2002), that is worsened by the human exploitation of underground water and gas in some areas. Several shallow coastal transitional water bodies are present along the Italian coastline, the main of which are the Marano-Grado Lagoon, the Venice Lagoon, the system of lagoons of the Po Delta, the Lesina Lagoon and the Varano Lagoon (Umgiesser et al., 2014).

Extreme sea levels cause the flooding of several coastal cities on both sides of the Adriatic Sea (Lionello et al., 2012; Medugorac et al., 2015), especially when the storm is associated with spring tides (Bajo et al., 2017). Part of the western coast is below sea level, and therefore it is also very vulnerable to such hazards (Lionello et al., 2012). These coastal zones are also strongly impacted by north-easterly storms with severe morphological impacts on natural sectors and damage to structures along urbanised zones (Armaroli et al., 2012; Harley et al., 2016). Conversely, recurrent meteotsunami events occur on the eastern side of the Adriatic Sea, and particularly the Croatian coast and islands, causing flooding and damage in some harbours (Orlić, 2015). According to Rizzi et al. (2017) and Satta et al. (2017), the northern Adriatic coastline, due to its low elevation, will be one of the regions in the Mediterranean area most exposed - in terms of coastal risk for flooding and erosion - to future climate change.

## 2 Material and methods

In order to address the territorial challenges related to sea storms effect on the coastal areas, we developed a shared and interoperable system (Integrated Web System - IWS) to allow a better exchange of information at a basin scale. Therefore, available resources can be accessed simultaneously in an aggregated and standard way. IWS was designed to specifically store, visualize and share the following category of geospatial and informative contents:

- a. historical and real-time (or near real-time) time series of observations from fixed-point sensor networks;
- b. outputs from existing operational forecast models;
- c. localization and description of coastal sea storm events that have damaged the environment, social-cultural and economic assets;

- d. bi-dimensional geospatial layers to provide georeferenced representations of the study area. Such layers are organized in thematic categories (e.g. terrestrial and maritime boundaries, ports, shorelines, morphology and bathymetry, cultural heritage, coastal defence work);
  - e. datasets, model outputs and time-series metadata to improve discoverability and proper re-use of the shared resources.
- 5 All information on coastal disaster due to sea storm events (historical and more recent) are organized and mapped in geospatial layers which constitute the Sea Storms Atlas. That series can be used to draw the map of risk characterisation of the coast with the aim of identifying the most vulnerable areas and supporting the planning of coastal area use and development (Depellegrin et al., 2017).

The IWS ~~implementation is based on Free and Open Source Software and the~~ architecture design follows a resource-centred and service-oriented approach as described in Yang et al. (2007) and Longueville (2010). Following the so-called Service-Oriented Geoportal Architecture, the IWS includes three main layers:

- the resource layer corresponds to the physical storage of the structured information in databases or files;
- the access layer includes all code and software designed to provide access to the resources in the appropriate format;
- the Graphical User Interface (GUI) is the client-side component of the Geoportal architecture; the role of GUIs is not limited to the rendering of a given set of resources but also includes the aggregation of relevant resources through lightweight and loosely coupled JavaScript code. In other words, the GUI is not only a presentation layer but also creates a mash-up of relevant resources.

IWS overall architecture is described in Fig. 2. Furthermore, the schema highlights the user typologies served by the IWS and the interactions/connections with the ~~partner's nodes~~ data sources and with external portals and early warning systems.

20 IWS is structured into six main components:

1. the ~~Resource Layer for storing the datasets, metadata, resources and all the necessary information. It consists of a combination of different storage solutions in order to support the several and heterogeneous data models and formats shared, and all the information needed to achieve a fully operational infrastructure (e.g. metadata, catalogue information, user accounts and profiles);~~
- 25 2. ~~the Data importer~~ for data ingestion, harmonization ~~,~~ and preparation and deposit the datasets in the storage facilities of the Resource Layer. For this purpose, we implemented the use of data servers (e.g. THREDDS, Hyrax) with the advantages that such web systems are open-source and they already implement services like DAP (Data Access Protocol), WCS (Web Coverage Service), WMS (Web Map Service), SOS (Sensor Observation Service);
3. the ~~Transnational Multi-model Ensemble System (TMES) Resource Layer for collecting and storing the datasets, metadata,~~ model forecasts, resources and all the necessary information. It consists of a combination of different storage solutions in order to support the several and heterogeneous data models and formats shared, and all the information needed to achieve a fully operational infrastructure (e.g. metadata, catalogue information, user accounts and profiles);

4. the *Transnational Multi-model Ensemble System (TMES)* for combining the results from existing operational **forecast forecasting** systems (described in section 2.2). TMES outputs are also stored in the *Resource Layer*;
5. the *Task Manager middleware* for orchestrating the communication with IWS components (e.g. Data importer, TMES) in order to launch the process (e.g. download the data from the partners' node), monitor the execution status, and concatenate multiple tasks in a single processing pipeline. The Task Manager middleware supports a time-based job scheduler, synchronous-asynchronous task queue system and a message broker system;
6. the *Common Data Sharing System (CDSS)* (Access Layer) for publishing the API and the web services to interact (e.g. search, visualize, download, manage) with the informative resources through standardized interfaces (e.g OGC-Web service, web API);
7. the *Geoportal* (Graphical User Interface) for implementing the **end-user-end-user** interfaces and tools to search, visualize, explore and analyse informative resources. The Map Viewer and Composer is an interactive and dedicated GUI for creating, managing and sharing multi-layered maps and for navigating and querying them.

IWS implementation follows a full-fledged Free and Open Source Software (FOSS) approach, in order to foster transparency, transferability and durability of the system and to be in accord with open source software strategy of the European Commission (European Commission, 2016). IWS provides spatial data infrastructure functionalities for accessing geospatial layers and forecast model outputs through OGC (Open Geospatial Consortium, <http://www.opengeospatial.org/>) interoperable services. Such approach is widely accepted and implemented at European (INSPIRE directive, European Commission, 2007; EuroGEOSS initiative, Vaccari et al., 2012) and global level (GEOSS, Global Earth Observation System of Systems) to facilitate intergovernmental and inter-agency data exchange and harmonization (Maguire and Longley, 2005). Incorporating THREDDS data server, IWS provides access to stored resources also through OPeNDAP and NetCDF standard services and formats. These standards are all products of the scientific communities in oceanography, meteorology and climate sciences and are designed to specifically meet their needs (Hankin et al., 2010), providing coherent access to a large collection of real-time and archived datasets from a variety of environmental data sources at a number of distributed server site (Unidata, 2019).

## 2.1 The monitoring networks

A joint asset which could be exploited through fruitful cooperation is the presence in the whole Adriatic-Ionian coastal territories of large networks of sensors and stations. In the Adriatic region, we mapped 35 tide gauges (9 inside the Venice Lagoon) and 9 wave stations, with the highest concentration in the northern Adriatic Sea. The location of all reported monitoring stations is illustrated in Fig. 1, and their general characteristics are summarized in Table [??A1](#) and Table [??A2](#), for sea level and wave respectively. The stations' lists are not exhaustive since there are other monitoring stations active in the area, the data of which were not available at the time of writing this document.

In several cases, the stations are also equipped with sensors for monitoring meteorological (wind speed and direction, sea surface pressure, air temperature, relative humidity and precipitation) or oceanographic parameters (seawater temperature, salinity, current speed and direction).

## 2.2 The forecasting systems

5 A multi-model ensemble was developed to combine the outcomes of existing ocean and wave forecasting systems, helping in improving the forecast accuracy and reliability on one hand and by adding indications on the forecast uncertainty on the other hand. The error of multi-model ensemble products should be on average lowest compared to those of the ensemble members (Golbeck et al., 2015). According to Di Liberto et al. (2011), operational forecast benefits from the combination of different ocean models by considering different physical parameterization, numerical schemes, model resolution and forcing.

10 Several operational ocean forecasting models are currently available for the Adriatic-Ionian region. Here we combined 17 forecasting systems, with 10 predicting sea level height (either storm surge or total water level) and 9 predicting wave characteristics. The general characteristics of the forecasting systems are summarized in Table 1 and Table 2, respectively for sea level and wave. [MED-Currents and MED-Waves forecasts were retrieved from from CMEMS \(http://marine.copernicus.eu/\)](http://marine.copernicus.eu/). We would like to point out that there are other operational systems active in the area (e.g. the pan-European Storm Surge  
15 Forecasting System, Fernández-Montblanc et al., 2019), the results of which were not available at the time of writing this document.

The different operational models are forced at the surface boundary by several meteorological models (ECMWF, BOLAM, MOLOCH, COSMO, WFR and ALADIN) with horizontal resolution ranging from ~~16 to 1.4~~ 55 to 1.25 km. The length of the ocean forecast is mostly related to the length of the meteorological forecast and varies from ~~1.5~~ 3 to 10 days. There is a  
20 large variability in the model's set-up in terms of spatial resolution, temporal frequency, spatial domain (Mediterranean Sea, Adriatic Sea, northern Adriatic Sea), grid arrangement (e.g. structured or unstructured) and data format (NetCDF, GRIB). ~~Only two~~ Three of the considered systems (Kassandra, [MED-Currents](#) and Adriac) account for the current-wave coupling and two forecasting systems perform data assimilation of tide gauge observations in the operational chain (SIMMb and SIMMe).

TMES is implemented as an internal processing engine which interacts directly with the Resource Layer to access the  
25 datasets (e.g. time series and forecasts) and to deposit the processing results (e.g. ensemble model result, report, statistics). Such outputs are available to the end-users and external portal through the Common Data Sharing System and the Geoportals web interfaces.

All numerical model results are interpolated, through a distance-weighted average remapping of the nearest neighbours, on a common regular lat-lon grid covering the Adriatic-Ionian macro-region with a resolution of 0.02 deg. For coastal flooding  
30 hazard purpose, the total sea level height must be forecasted. Therefore, the astronomical tidal level values obtained by a specific SHYFEM application over the Mediterranean Sea (Ferrarin et al., 2018) are added to the residual sea level simulated by the operational systems not accounting for the tide (SHYMED, ISSOS, SIMMb, SIMMe and MFS). The so obtained sea level height simulated by the different models are all referred to the geoid. The spread among the operational simulations is expected to represent a measure of the uncertainty of prediction and should be linked to the forecast error ~~;~~ so that cases

with the largest spread are those with the highest uncertainty and where a large error of the ensemble mean (and also of the deterministic forecast) is more likely (Mel and Lionello, 2014a). TMES produces results in terms of the ensemble mean and standard deviation, accounted for a measure of the forecast uncertainty (Flowerdew et al., 2010).

### 2.3 Hazard Storm impact assessment and early warning systems

5 The vulnerability to sea storms of a particular segment of coast depends on a wide number of variables, not only related to the magnitude of the storm but including the land characteristics and the social and economic activities that distinguish that area. In order to draw ~~the hazard a~~ map showing the region coastal areas affected by stormy conditions along the Adriatic region,  
~~with the aim of identifying most vulnerable areas to a forecasted storm event,~~ Adriatic-Ionian region, the coast is subdivided  
into segments of variable length in function of morphology, human settlements and administrative boundaries. The coastal  
10 assessment units were selected according to the Mediterranean Coastal Database (MCD) developed by Wolff et al. (2018).  
The MCD segments have an average length of 4.5 km. For each of these units, the database provides information on the characteristics of the natural and socio-economic subsystems, such as vertical land movement, coastal slope, coastal material and number of people exposed to sea-level rise and to extreme sea levels. ~~Furthermore, the level of vulnerability to sea storm events are complemented with the results of consultation among different public and private socio-economic actors having their~~  
15 ~~main activity on the coast or at the marine level which is affected from the sea storms or bearing responsibilities for informing the citizens.~~

At each location, three sea condition scenarios are computed considering mean and standard deviation of predicted sea level and wave ensembles:

- MIN: Ens. Mean – Ens. St.Dev
- 20 – MEAN: Ens. Mean
- MAX: Ens. Mean + Ens. St.Dev.

~~Several methodologies have been developed and applied at the basin and local scales for estimating hazard maps for coastal flooding (Hinkel et al., 2014; Vousdoukas et al., 2016; Wolff et al., 2016; Rizzi et al., 2017; Armaroli and Duo, 2018).~~ Over the whole Adriatic-Ionian coastal region, the ~~hazard assessment to sea storm is computed considering nearshore forecasts~~  
25 ~~provided by the TMES were combined with the coastal characteristics (coast material and slope) provided by the MCD database for computing~~ the total water level (TWL)~~obtained~~. For the coastal segments characterized by sandy beaches, TWL was computed combining the sea level height, wave setup and wave runup according to the Stockdon's formula ( $R_2$ , the 2% exceedance level of runup maxima; Stockdon et al., 2006), ~~using nearshore forecasts provided by the TMES for each coastal unit. It must be pointed out that the widely used Stockdon's formula – developed for sandy beaches – could underestimate and~~  
30 ~~overestimate wave runup on gravel beaches (Poate et al., 2016).~~ For gravel beaches and rocky cliffs (Dodet et al., 2018) other methodologies should be used for estimating wave runup (Poate et al., 2016; Dodet et al., 2018), but they could not be applied in this study due to the lack of the required detailed coastal information (sediment grain size, type of rocks, permeability of the structure).

It is well known that the estimation of the total water level is strongly influenced by the local coast typology and morphology and the MCD segments are sometimes too coarse to represent complex morphologies, especially in confined coastal systems (lagoons) and along the eastern rocky coast. Therefore, in order to provide more reliable and resolute hazard assessment at a very fine coastal scale, the IWS has been designed to ~~be easily integrated into~~ [provide multi-model forecast products to](#) existing early warning systems, developed in areas where a deep knowledge of the coastal dynamics and high-resolution datasets (topography and bathymetry) are available. In this study, we present three existing local forecasting and early warning systems operative in the Adriatic Sea (Slovenia, Emilia-Romagna region and the City of Venice) to which IWS provides the information required for issuing coastal risk alerts.

### 2.3.1 Slovenia

TMES forecasts can be used directly by regional authorities for assessing the hazard of a particular segment of the coast to the storm event according to predefined thresholds. As an example, we report here the IWS based hazard estimates for the Slovenian coast, which is predominantly rocky and steep (flysch cliffs), and therefore well protected from flooding during storm surges. Important exceptions are the salt pans (Sečovlje and Strunjan) and urban areas such as Piran, Koper and Izola where lower parts get flooded up to 17 times per year (data for the 1963-2003 period; Kolega, 2006), with consequent damage to private properties and cultural heritage. The Slovenian Environment Agency issues a warning when the predicted sea level at Koper exceeds the yellow alert level which is set at 300 cm (above local datum). This is the value that marks the flooding of the lowest coastal urban areas. Orange and red alert levels are set to 330 and 350 cm, respectively.

### 2.3.2 Emilia-Romagna

In addition to the evaluation of thresholds for identifying critical storm conditions at sea (Armaroli et al., 2012), since December 2012, the Emilia-Romagna region (northern Italy) daily provides three-day forecast of coastal storm hazard at eight key sites along the coast, where several past sea storms have induced significant morphological change and damages. The Emilia-Romagna coastline is particularly vulnerable to sea storms due to its low-lying nature and high coastal urbanization (Armaroli and Duo, 2018). During major storm events, the water levels often exceed those of the dune crest and building foundations (Harley et al., 2016). The existing coastal Early Warning System (Harley et al., 2016) is based on the 1D cross-shore implementation of the XBeach morphodynamic model (Roelvink et al., 2009), a 2DH (depth-averaged) cross-shore process-based model that solves intra-wave flow and surface elevation variations for waves in intermediate and shallow water depths. The XBeach model is used to forecast wave runup and total water level during storm events. For each key site, IWS provides to the XBeach model the sea level and wave characteristics for the three above-mentioned sea condition scenarios. Hence, the developed methodology allows converting the forecast uncertainty on nearshore sea conditions into a coastal flooding hazard range of predictions. Coastal hazard is here estimated in terms of two storm impact indicators:

- Safe Corridor Width (SCW), a measure of the amount of dry beach available between the dune foot and waterline for safe passage by beach users;

- Building Waterline Distance (BWD), a measure of the amount of dry beach available between the seaward edge of a building and the model-derived waterline.

### 2.3.3 City of Venice

The City of Venice is located in the centre of a shallow lagoon and is composed of more than a hundred islands linked by bridges. The elevation of these islands is extremely low, subjecting them to flooding during storm tides (resulting from the combination of storm surge and the astronomical tide), which in turn threatens the unique cultural heritage of this city and affects its everyday life, causing among all: difficulties in transport, the practicability of roads and internal channels, emergency procedure response, commercial activities. In the city of Venice, a bulletin of forecasted sea level up to 3 days is emitted three times per day (at 9 am, 1 pm, 5 pm) by the Tide Forecast and Early Warning Center (CPSM). The forecast is based on a combination of statistical and deterministic models, as well as an evaluation of the synoptic meteorological conditions (<https://www.comune.venezia.it/it/content/centro-previsioni-e-segnalazioni-maree>).

Since Venice is protected from the sea by barrier islands (separated by three inlets), storm waves do not affect significantly - through setup and runup - the sea level height inside the lagoon (Roland et al., 2009). While propagating from the sea to the lagoon through the inlets, the tidal signal is deformed, either damped or amplified, according to a relationship between local flow resistance and inertia and the characteristics of the incoming open sea signal (Ferrarin et al., 2015). For those reasons, sea level height forecasts are used instead of TWL predictions in the operational system. To propagate the sea level from the inlets to the inner lagoon, nearshore TMES values of sea level height - for each of the above-mentioned three sea condition scenarios - are referred to the local sea level reference datum (Punta della Salute) and used as open sea boundary conditions in the SHYFEM finite element hydrodynamic model of the Venice Lagoon (Bajo and Umgiesser, 2010; Cavaleri et al., 2019). Such model adequately reproduces the complex geometry and bathymetry of the Venice Lagoon using an unstructured numerical mesh composed of triangular elements of variable form and size (down to a few meters in the tidal channels). Flooding maps of the city floor are produced by imposing the sea level height observed/predicted at Punta della Salute (at intervals of 10 cm) to a centimetre accurate digital terrain model of the city (<http://www.ramses.it/>).

The Municipality plan of procedures in case of high and low tide (City of Venice, 2002) defines the actions the several stakeholders (civil protection, public security and rescue forces, transport companies, public services) adopt in case of risk for flooding, with respect to the specific forecasted sea levels. Depending on the forecasted sea level, particular categories of stakeholders are informed by CPSM and elevated wooden walkways are installed in areas of the city that are prone to flooding. The communication channels for the warning includes a website, messages (SMS, social network), e-mails, phone calls, acoustic signals, direct information (door to door). Moreover, an operating room with forecasters is functioning 24 hours a day at CPSM during the most severe storm tide event.

### 3 The 29 October 2018 event

#### 3.1 Storm description

On 29 October 2018, an exceptional storm hit the central and northern part of Italy with very strong south-easterly winds (called ~~Sirocco~~-Sirocco over the Adriatic Sea. The basic meteorological situation of the 2018 storm was similar to the 1966 and 1979 ones, although with a weaker pressure gradient over the Adriatic area (Cavaleri et al., 2019). The weather condition was governed by a semi-stationary upper level trough which was positioned over West Mediterranean on 28<sup>th</sup> of October and was only slowly moving north-eastward on 29<sup>th</sup> and 30<sup>th</sup> (Fig. 3). The upper level southerly flow on the East side of the trough was very intense due to strong pressure gradients throughout the whole period of the event. The occurrence of the upper level trough resulted in a formation of a very intense low level low-pressure system over the Alps and Central Mediterranean which was the most prominent surface feature of the event.

The air mass over the Italian peninsula and Adriatic was very unstable on 28<sup>th</sup> and 29<sup>th</sup> of October due to meridional flow which was bringing moist and warm air from North Africa and Central Mediterranean. In this sense, it was a typical Autumn situation when the amount of precipitation can be extreme, especially on the windward side of orographic barriers. The flow at the surface was further intensified by extreme convection over the Apennines and the Alps. The amount of precipitation in northern Italy and wind storms over the Alps and northern Adriatic were rather extreme and not often observed in such intensity.

It is worth mentioning that ~~Sirocco~~-Sirocco wind started already on 26<sup>th</sup> at the most of Adriatic and lasted for almost four days without interruption with the strongest wind in North Adriatic on 29<sup>th</sup> afternoon, just before the passage of the cold front. Most of this time, the ~~Sirocco~~-Sirocco was well developed over the entire Adriatic basin and even further south in the Ionian Sea. This meant that the fetch was exceptionally long for the locations in the northern Adriatic Sea.

Consequently, sea level raised in the northern end of the Adriatic Sea - the area most affected by flooding - reaching peak values around 13 UTC in Venice, Koper and Rovinj (Fig. 4a). Exceptional sea levels were reached also along the Emilia-Romagna region with values higher than 1 m for about 5 hours, as measured at Porto Garibaldi. It has to be noted that these maximum values were not registered during the storm peak (happened around 19 UTC in this location) due to an out-of-phase with the astronomical tide. A secondary maximum was recorded in Koper and Rovinj just after the cold front passed and when the wind and waves were decreasing but the tide was rising. Along the central and southern Croatian coast, sea level resulted to be marginally affected by storm surge, even if weak meteotsunamis were observed during the frontal passage late in the evening on the 29<sup>th</sup> October.

The very long wind fetch allowed waves to develop over the whole basin. Available wave monitoring stations recorded values of significant wave height (the average height of the highest one-third of all waves measured; SWH) up to 6 m at the Piattaforma Acqua Alta (PTF), 15 km offshore the Venetian littoral, and up to 4.7 m (8 m of maximum wave height) near the city of Rovinj (Fig. 4b). Along the north-western Italian coastline, due to its mild slope, wave breaking occurs and the measured SWH reaches values of about 2 m during the storm peak (Nausica and Senigallia monitoring stations). On the Trieste Gulf, the highest waves occurred 6 hours later (Zarja wave buoy), probably due to the eastward shift of the wind induced by the passage

of the cold front. In the south-eastern Adriatic Sea, high wind and wave values were recorded even before the cold front on 28<sup>th</sup> October. The highest waves recorded in Dubrovnik reached values of about 5 and 9 m for significant and maximum wave height, respectively. Rough sea conditions (SWH > 2.5 m) lasted for 57 hours while the very rough sea state (SWH > 4 m) was recorded for 9.5 hours. According to long-term time series of observations, the 29 October 2018 event reached the records of the second highest sea state ever measured all along the Adriatic coast.

### 3.2 Storm predictability

Here we present the results of the forecasting system at hourly time step and for the day of the event only. However, as described by Cavaleri et al. (2019), up to five (six for the surge) days earlier there were indications of a severe event. Fig. 5 shows the TMES results in terms of the ensemble mean and standard deviation for both the sea level height (panels a and b) and the significant wave height (panels c and d). Storm surge during the 29 October event affected mostly the northern Adriatic Sea (Fig. 5a), while severe sea condition occurred over most of the Adriatic Sea with the higher waves impacting the Croatian coast (Fig. 5c). The ensemble operational system provides also the estimation of the uncertainty associated with the forecast of this event. Uncertainty is generally higher where the sea level and the waves reach the highest values (Fig. 5b and d). The ensemble standard deviation reached maximum values of about 30 cm for the sea level and 1.5 m for the significant wave height.

Model forecasts could be extracted at any location in the domain to provide a clear representation of sea storm evolution. As an example, we reported in Fig. 6 the values extracted at PTF, Rovinj and Dubrovnik (see Fig. 1 for their location). The comparison with the observations highlights the good performance of the ensemble methodology in reproducing such a strong event. The ensemble mean time-series are smoother than the observations and slightly underestimate the maximum sea level in the northern Adriatic Sea (Figs 5a and 6b), as well as the peak wave height at 20 UTC (5 m of forecasted significant wave height with respect to almost 6 m of observed at PTF; Fig. 6c). However, the observed values are - almost always - within the TMES spread (i.e. the standard deviation of the ensemble members). It is worth noting that the forecast uncertainty increases with the forecast lead time and with the severity of the storm, the maximum of which was reached in the northern Adriatic Sea between 19 and 20 UTC. In the southern Adriatic Sea (Fig. 6d), the ensemble mean well reproduces the observed significant wave height, which remained between 3 and 5 m for the whole day. For this specific location the spread of the ensemble assumed values between 0.7 and 1.1 m on 29<sup>th</sup> October.

### 3.3 Storm ~~hazard and impact~~ and hazard assessment on the coast

In order to ~~assess the storm hazard and impact at a basin scale~~ provide the perception of the physical processes acting along the Adriatic-Ionian coastal areas and responsible for storm related hazards, the results of the multi-model ensemble - in terms of sea level and wave conditions - were processed for each coastal assessment unit ~~of the investigated area~~. Considering the general underestimation of the ensemble means during the peak of the storm, we will ~~focus our storm hazard~~ mostly focus our basin-wide storm analysis on the ~~MEAN and MAX~~ sea condition ~~scenarios. The scenario~~.

The 2% exceedance level of wave runup maxima ( $R_2$ ) computed according the Stockdon's formula (Stockdon et al., 2006) and the total water levels forecasted for the 29 October 2018 event (at 19 UTC) are reported in Fig. 7 for ~~scenarios MEAN and~~

~~MAX the MAX scenario.~~ As for sea level height results (Fig. 5a), the maximum values of TWL are found in the North Adriatic along the Veneto and Friuli Venezia Giulia coasts. Indeed, during the 29 October storm, several coastal lowlands in the northern Adriatic were flooded. At these locations, characterised by gently sloping sandy beaches, the estimated ~~2% exceedance level of wave runup maxima ( $R_2$ ) reached values of 0.7 m in the MAX sea condition scenario~~  $R_2$  reached maximum values of 0.8 m,  
5 accounting therefore for about 50% of the total water level. ~~TWL differences between the MEAN and MAX scenarios reached there the maximum values of about 0.4 m, that is higher than the standard deviation of the multi-model ensemble for the sea level height.~~

The combination of the sea level height and the wave setup/runup generated high values of the total water level (TWL > ~~1.5 m, with  $R_2$  > 1 m~~) also ~~along the Istria peninsula, south of Dubrovnik and close to Ancona~~ at some locations on the along  
10 the eastern Adriatic coast and the Marche region. Along the Istrian-Croatian coast, the extremely high waves and the high sea levels caused inundation and widespread damages to the coastal infrastructure (Opatija-Istria and Zadar). Moreover, because of the rough sea conditions, there was a disruption of the maritime traffic during the 27-30 October and the ferry service cancelled almost all the scheduled sailings on 29<sup>th</sup> October, so most of the Croatian islands were cut off from the mainland for more than a day. TWL differences between the MEAN and MAX scenarios (not shown) reached there the maximum values of about  
15 0.4 m, that is higher than the standard deviation of the multi-model ensemble for the sea level height.

As stated in section 2.3, previous studies demonstrated that the wave runup estimation increases with the ~~slope of the structure~~ beach slope. Therefore, the high wave runup values found at some coastal segments (e.g. along the Marche and Abruzzo regions) are due to the severe wave conditions, but also to the fact that they are characterized by steep ~~rocky~~ coast (slope > 0.15). On such reflective conditions, the use of an alongshore-averaged beach slope in practical applications of the  
20 runup parameterization may result in large runup error (Stockdon et al., 2006). ~~Moreover, according to Dodet et al. (2018), wave runup could be overestimated at locations with rocky cliffs (e.g. the coast south of Ancona, the Croatian coast and islands), which act on the wave transformation by increasing dissipation and/or shifting offshore the breaking point.~~

In the following paragraphs we describe the results of the application of the multi-model ensemble to the existing early warning systems and investigate into details the storm ~~hazard and impact~~ impact and hazard at the three selected locations.

25 Due to its northward orientation, the Slovenian coast is relatively well protected from the waves generated by southerly winds, as in the case of the 29 October 2018 storm. Indeed, over there and for this event the wave contribution to the total water level is negligible. According to the 10-min observation data presented in Fig. 8, the sea level in Koper reached peak values well above the orange alert level (343 cm at 12:50 UTC and 341 cm at 23:20 UTC) and lasted for almost 10 hours above the yellow alert level. As a consequence, the sea flooded ~~Punta, Prešernovo nabrežje (Prešeren Seafront), Cankarjevo nabrežje (Cankar Seafront), the red pier and Tartinijev trg (Tartini square) in Piran; Veliki trg (Grand Square), Sončno nabrežje (Sunny Seafront) and parts of Dantejeva ulica (Dante street) in Izola.~~ During the storm, several coastal locations, where the firemen set up anti-flooding barrages ~~at many locations~~. As shown in Fig. 8, the MEAN scenario predicted the exceeding of the yellow flooding alert level but underestimated the observed peak values. A more realistic - even if slightly overestimated - prediction of the flooding hazard for the Slovenian coast is provided by the MAX scenario.

Along Emilia Romagna region, several coastal sites were affected by flooding and erosion during the 29 October 2018 sea storm, due to the combination of high sea level and energetic wave conditions. The documented coastal impacts are reported in Fig. 9b and include erosion of the beach and of the winter dunes, coastal flooding and damage to coastal infrastructures and defence structures. Damages and impacts were notified especially for the northern part of the region, while along the southern coastal area between ~~Cesenatico~~ Cesenao and Riccione real impact data are not available. The hazard index computed for this event using the XBeach model forced with the three (MEAN, MIN and MAX) conditions, reveals that the most severe sea condition scenario (MAX scenario) provides an exceeding of the predefined alert thresholds indicating a high level of coastal risk. An example of the Safe Corridor Width (described in section 2.3.2) calculated for a single cross-shore section, located at Lido di Classe, is reported in Fig. 10. The predicted coastal hazard (Fig. 9a) shows that the most critical scenario is in good agreement with the documented coastal impacts, displayed in the right panel. For this event, by comparing hazard forecasts and observations, the use of IWS provides a good prediction (MAX scenario) of coastal impacts for the major part of the Emilia-Romagna coastal area. Moreover, considering the distance between the MIN and the MAX conditions, IWS provides useful information about the range of the possible impacts.

On the 29 October 2018, the City of Venice was inundated by the exceptional sea storm. At 13:40 UTC the sea level reached the peak value of 156 cm at Punta della Salute (fourth historical level of flooding in Venice since 1872), which put three-quarters of the pedestrian public area of the historic town under water. Sea level reaching 120 cm (above local datum), at which point flooding covers 28% of Venice, lasted for about 14 hours. The SHYFEM application to the Venice Lagoon, forced by the open sea TMES results, forecasted sea level peak values of 142 and 161 cm for the MEAN and MAX scenarios, respectively. Fig. 11 shows the corresponding flooding map of the City of Venice according to the predicted peak values (rounded at 140 and 160 cm). The 59% and 77% of the pedestrian surface are flooded for the two scenarios, respectively. In the first case, the public navigation transport is strongly limited and only links to the islands are guaranteed; besides, most of the elevated walkways are impracticable. In the second case, the public navigation transport is no more guaranteed, as well as all of the elevated walkways. Moreover, many internal channels are no longer navigable due to insufficient vertical space under some bridges and hence the emergency rescue response may be strongly delayed. Since the observed peak was 156 cm, the MAX scenario provided a realistic prediction of the flooding hazard for the city of Venice.

#### 4 Summary and concluding discussion

To improve knowledge on sea storms events in order to progress the prevention and protection measures integrated into coastal defence plan and procedures, we developed a transnational integrated web system (IWS) for sharing [information](#), observations and forecasts across the Adriatic and Ionian seas. ~~IWS implementation follows a full-fledged Free and Open Source Software (FOSS) approach, in order to foster transparency, transferability and durability of the system and to be in accord with open source software strategy of the European Commission (European Commission, 2016). IWS provides spatial data infrastructure functionalities for accessing geospatial layers and forecast model outputs through OGC (Open Geospatial Consortium, ) interoperable services. Such approach is widely accepted and implemented at European (INSPIRE directive,~~

European Commission, 2007; EuroGEOSS initiative, Vaccari et al., 2012) and global level (GEOSS, Global Earth Observation System of Systems) to facilitate intergovernmental and interagency data exchange and harmonization (Maguire and Longley, 2005). Incorporating THREDDS data server, IWS provides access to stored resources also through OPeNDAP and NetCDF standard services and formats. These standards are all products of the scientific communities in oceanography, meteorology and climate sciences and are designed to specifically meet their needs (Hankin et al., 2010), providing coherent access to a large collection of real-time and archived datasets from a variety of environmental data sources at a number of distributed server sites (Unidata, 2019).

It must be taken into account that meteorological and ocean models provide just an approximation of reality, despite their continuous development and improvements. Moreover, the interactions between atmospheric, oceanic and coastal processes are not fully understood, resulting in large uncertainties in the predictions of coastal flooding, in particular, under extreme conditions (Baart et al., 2011; Zou et al., 2013). This is mainly due to the chaotic nature of the atmosphere and the complexity of the air-sea interactions across scales over several orders of magnitude (Schevenhoven and Selten, 2017). Small errors in the initial conditions of a numerical weather prediction model grow rapidly and affect predictability; forecasted atmospheric conditions are then affected by errors (Molteni et al., 2001). However, as stated by Flowerdew et al. (2010), atmospheric forcing is not the only source of uncertainty in storm surge forecasting. Many other sources of uncertainty, as the model numerics, resolution, parametrization, boundary conditions and initial sea state, contribute non-linearly to the final forecast uncertainty. The IWS allows to strengthen the forecasts with useful information of their degree of uncertainty and hence analyse the propagation of uncertainty towards the coastal forecasts. The awareness of ~~these uncertainties and prediction~~ the prediction uncertainties and errors has led many operational and research flood forecasting systems around the world to move toward numerical forecasts based on a probabilistic concept: the ensemble technique (Cloke and Pappenberger, 2009).

~~On that basis, the IWS allows to strengthen the forecasts with useful information of their degree of uncertainty and hence analyse the propagation of uncertainty towards the coastal forecasts, starting from the meteorological models~~In this contest, a probabilistic forecasting system could be based on perturbation of initial conditions, forcing and parameters of a single model (Flowerdew et al., 2010; Bernier and Thompson, 2015; Salighehdar et al., 2017). Such approach has been already applied to the Adriatic Sea for improving storm surge forecast and providing a realistic estimate of the prediction uncertainty (Mel and Lionello, 2014a). In order to improve sea storm predictions, ~~we here we instead~~ we here we instead implemented for the Adriatic-Ionian macro-region a ~~transnational~~ multi-model ensemble system which combines several existing oceanographic and wave forecasting systems. The magnitude of ensemble spread is a good indicator of how the forecast accuracy varies between different forecasting situations, indicating a decrease of reliability when the spread increases (World Meteorological Organization, 2012). It is not straightforward what averaging weights should be used for the multi-model ensemble forecast and therefore we used equally weighted members, despite the forecasts which are more precise than others should have more importance in the TMES (Salighehdar et al., 2017; Schevenhoven and Selten, 2017). Here we applied a simple average of the forecasts at every timestamp to compute the ensemble mean, but more sophisticated methods based on weighting function determined by comparison of the single model results with near real-time observations can be implemented in future (Di Liberto et al., 2011; Salighehdar et al., 2017). Taking

advantage of the near real-time observations acquired by the aggregated monitoring network, the root mean square error of the individual forecast will be evaluated and stored for long-term statistics.

Nearshore TMES sea levels and wave characteristics can be directly used in an early warning procedure on the basis of predefined thresholds for morphological change and for coastal erosion/flooding (e.g., Armaroli et al., 2012). TMES predictions are also used to compute the alongshore total water level time-series. ~~FWL can be used to quantify the vulnerability of the coast to extreme inundation and erosion events, but the~~ The estimated run-up values need to be considered with care due to the uncertainty associated to the geomorphological characteristics of the coastal segment units (beach material and slope in particular). Indeed, Bosom and Jiménez (2011) and De Leo et al. (2019) found large variability in hazards intensity and vulnerability along limited coast sectors (20 to 50 km long), even with homogeneous offshore wave conditions. Therefore, the choice of the coastal segment database and its resolution has a substantial effect on the accuracy of the hazard model. The MCD dataset has some limits in reproducing detailed coastal morphologies (i.e. northern Adriatic lagoons and Croatian islands) as well as storm defence structures as breakwaters and seawalls. However, the developed IWS has been designed to be flexible in integrating better defined coastal segment units. If detailed beach geomorphological information is available, ~~the approach of Bosom and Jiménez (2011)~~ process-based hazard indicators (Ferreira et al., 2017) could be used for assessing the potential of a coastal system to be harmed by the impact of a storm (inundation or erosion), comparing the magnitude of the impact (~~wave run-up total water level~~ for inundation and beach/shoreline retreat for erosion) with the ~~adaptation capacity morphological characteristics~~ of the system (dune/berm height for inundation and beach width for erosion).

The developed system has been tested against observations acquired during a very severe storm that affected the Adriatic Sea on 29 October 2018. TMES forecasts resulted to be in agreement - even if slightly underestimated during the storm peak - with the observed sea level height and significant wave height. The predicted ensemble mean and standard deviation were combined for creating three different sea condition scenarios all along the Adriatic and Ionian coastline, allowing to evaluate a range of coastal hazard forecast. Moreover, nearshore forecasts were successfully integrated into existing early warning systems for estimating storm hazard at three locations (Slovenia, Emilia-Romagna region, City of Venice). Through this system coupling, the predicted sea conditions were translated into local storm impact indicators and very detailed flooding maps. The underestimation of predicted sea levels and waves during the 29 October storm peak is probably a consequence of a general underestimation of the wind forecasts produced by the operational meteorological models. Cavaleri and Bertotti (2004) provided clear evidence of the wind speed problem over the Adriatic Sea. In particular, for the sea storm of the 29 October 2018, Cavaleri et al. (2019) compared the ECMWF model wind with scatterometer data and estimated a 1.11 average enhancement factor.

For the reasons stated above and considering the results presented in this study, the most severe sea condition scenario (MAX = ensemble mean + ensemble standard deviation) can be considered for the investigated area as a ~~realistic and~~ conservative estimation of the peak storm conditions to be used for coastal risk management. Another possible application of TMES outputs could be to use all possible combinations of ensemble mean and standard deviation for the sea level and wave characteristics, providing a large number of sea state combinations. In that way, it would be possible to calculate and estimate the frequency of

exceeding predefined thresholds for coastal hazard. This approach is closer to the idea of the probability of threshold exceeding and will be explored in future.

The aggregating approach for collecting and sharing observations is crucial for providing real-time information about the sea state - and its evolution - to be used by several countries for prompt emergency response and to increase the overall preparedness to sea storms. Moreover, by merging data from several stations and sensors, IWS is an important storage server for any data assimilation system. According to [Lionello et al. \(2006\)](#) and Bajo et al. (2019), the assimilation of tide-gauge data in the Adriatic Sea has a strong positive impact on the forecast performance, lasting several days, despite the underestimation in the atmospheric forcing. The forecast improvement is particularly relevant in the case of consecutive sea storms when storm surge levels are influenced by pre-existing oscillations of the basin (seiches) induced by previous events. It is worth mentioning that in the case of the Adriatic Sea - but there could be many other similar situations - the transnational cooperation is crucial for sharing observations acquired along the whole basin in order to provide real-time information on the sea state to be used in a data assimilation system.

Real-time observations and numerical model forecasts required to address environmental management problems such as sea storms are commonly excessively intricate for civil protection and stakeholders to use (Magaña et al., 2018). IWS is equipped with geoportals functionalities and interactive geo-visualization tools for simplifying search and access to geospatial data (including forecast model outputs) and monitoring networks time series. Such tools help and assist people who want to use IWS concepts, databases and results in their work and to support their activities. Moreover, a dedicated web site (<http://www.seastorms.eu/>), designed to foster the data dissemination according to the community-based paradigm and to the Open Data principles (<https://opendatacharter.net/>), will allow the public data, the forecast results and related statistics to be explored by non-experts over Internet through the use of shared maps, dashboards, graphics, tables and other interactive geo-visualization tools.

Concluding, to improve the capacity to react to sea storms, all relevant actors of the coastal area (public authorities, regional and national authorities in charge of Civil Protection, meteorological and forecast services, universities and research institutes, cruise ship enterprises, maritime business enterprises, marinas, aquaculture SMEs, stakeholders from touristic sector) should be involved - through the web and socials - in a transnational cooperation strategy to foster:

- knowledge and data exchange for providing real-time information about the basin-wide sea state through the use of standardized formats, protocols and services;
- coordination of all available ocean forecasting systems in a multi-model ensemble for enhancing the predictability of extreme events and for evaluating the uncertainty of the operational ocean products;
- integration of observations and numerical models through data assimilation techniques for improving the forecast accuracy;
- downscaling of open sea ensemble forecasts to be integrated in site specific early warning systems managed by local authorities;

– data and forecasts dissemination to all relevant coastal actors and citizens over Internet.

*Code and data availability.* The IWS model is available as an open-source code from <https://github.com/CNR-ISMAR/iws>. The SHYFEM hydrodynamic model is open source and freely available at <https://github.com/SHYFEM-model>. The data and forecast results used in this study can be accessed, depending on the specific provider’s data policy, upon request to the corresponding author. TMES operational forecasts are daily available at <http://www.seastorms.eu/>.

*Author contributions.* CF conceived the idea of the study with the support of AV, SM and MV. SM and AF designed the IWS structures. SM, AF, AV, LB, CF, MB, MV, ML and GM prepared the model results and developed the multi-model ensemble. AF, CF and SM developed the scripts for computing TWL on the MCD coastal segments. JJ described the meteorological situation of the 29 October 2018 storm. SU and AV elaborated the TMES results for computing the ER-EWS storm impact indicators for the Emilia-Romagna coast. GM, ES, AP, CF and MB applied the TMES results to the early warning system of the City of Venice. All authors discussed, reviewed and edited the different versions of the manuscript.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This work was supported by the I-STORMS project (Integrated Sea sTORM Management Strategies) funded by the European Union under the Interreg V-B Adriatic-Ionian programme with agreement n. 69 of 09-03-2018. The authors wish to thank: Isabella Marangoni, Denise Florean, Anna Zarotti, Alessia Porcu and Silvia Comiati from City of Venice - European Policies Division for promoting and supporting the project activities; Dr. Fabio Raicich from CNR-ISMAR for providing sea level data for the Trieste tide gauge; Dr. Pierluigi Penna from CNR-IRBIM for providing data for the Senigallia monitoring station; Dr. Luigi Cavaleri and Dr. Luciana Bertotti from CNR-ISMAR for sharing wave results of the Nettuno and Henetus systems; wave data of the Nettuno forecasting systems were kindly provided by Centro Nazionale di Meteorologia e Climatologia Aeronautica (CNMCA); Ing. Maurizio Ferla, Dr. Elisa Coraci and Dr. Roberto Inghilesi from ISPRA for providing sea level and wave forecasts of the SIMM systems; Janez Polajnar from Slovenian Environment Agency for the data on storm impacts to the Slovenian coast; Dr. Luisa Perini, Dr. Lorenzo Calabrese and Dr. Paolo Luciani from the Geological Seismic and Soil Survey of the Emilia-Romagna Region for providing the report of impacts occurred along the regional coast during the event; Dr. Alessandro Coluccelli and Dr. Aniello Russo from Università Politecnica delle Marche for the support on the AdriaROMS model; the Croatian Hydrographic Institute in Split for the tide-gauge records originating from Rovinj and Dubrovnik and for wave riders records originating from buoys near Rovinj and Dubrovnik/Sv-Andrija.

~~[urlstyle](#)~~

~~[Armaroli, C. and Duo, E.: Validation of the coastal storm risk assessment framework along the Emilia-Romagna coast, Coast. Eng., 134, 159–167, , 2018.](#)~~

- Armaroli, C., Ciavola, P., Perini, L., Calabrese, L., Lorito, S., Valentini, A., and Masina, M.: Critical storm thresholds for significant morphological changes and damage along the Emilia-Romagna coastline, Italy, *Geomorphology*, 143-144, 34–51, , 2012.
- 5 Baart, F., van Gelder, P. H. A. J. M., and van Koningsveld, M.: Confidence in real-time forecasting of morphological storm impacts, *J. Coast. Res.*, pp. 1835–1839, 2011.
- Bajo, M. and Umgiesser, G.: Storm surge forecast through a combination of dynamic and neural network models, *Ocean Model.*, 33, 1–9, , 2010.
- Bajo, M., De Biasio, F., Umgiesser, G., Vignudelli, S., and Zecchetto, S.: Impact of using scatterometer and altimeter data on storm surge forecasting, *Ocean Model.*, 113, 85–94, , 2017.
- 10 Bajo, M., Medugorac, I., Umgiesser, G., and Orlić, M.: Storm surge and seiche modelling in the Adriatic Sea and the impact of data assimilation, *Quart. J. Roy. Meteor. Soc.*, , 2019.
- Bertotti, L., Canestrelli, P., Cavaleri, L., Pastore, F., and Zampato, L.: The Henetus wave forecast system in the Adriatic Sea, *Nat. Hazards Earth Syst.*, 11, 2965–2979, , 2011.
- Bertotti, L., Cavaleri, L., Loffredo, L., and Torrisi, L.: Nettuno: Analysis of a Wind and Wave Forecast System for the 15 Mediterranean Sea, *Mon. Weather Rev.*, 141, 3130–3141, , 2013.
- Bosom, E. and Jiménez, J. A.: Probabilistic coastal vulnerability assessment to storms at regional scale—application to Catalan beaches (NW Mediterranean), *Nat. Hazards Earth Syst.*, 11, 475–484, , 2011.
- Bressan, L., Valentini, A., Paccagnella, T., Montani, A., Marsigli, C., and Tesini, M. S.: Sensitivity of sea-level forecasting to the horizontal resolution and sea surface forcing for different configurations of an oceanographic model of the Adriatic Sea, 20 *Adv. Sci. Res.*, 14, 77–84, , 2017.
- Carbognin, L. and Tosi, L.: Interaction between Climate Changes, Eustacy and Land Subsidence in the North Adriatic Region, Italy, *Mar. Ecol.*, 23, 38–50, , 2002.
- Cavaleri, L. and Bertotti, L.: Accuracy of the modelled wind and wave fields in enclosed seas, *Tellus A*, 56, 167–175, , 2004.
- 25 Cavaleri, L., Bertotti, L., Buizza, R., Buzzi, A., Masato, V., Umgiesser, G., and Zampieri, M.: Predictability of extreme meteo-oceanographic events in the Adriatic Sea, *Q. J. Roy. Meteor. Soc.*, 136, 400–413, , 2010.
- Cavaleri, L., Bajo, M., Barbariol, F., Bastianini, M., Benetazzo, A., Bertotti, L., Chiggiato, J., Davolio, S., Ferrarin, C., Magnusson, L., Papa, A., Pezzutto, P., Pomaro, A., and Umgiesser, G.: The October 29, 2018 storm in Northern Italy— an exceptional event and its modeling, Submitted to *Prog. Oceanogr.*, 2019.
- 30 Chaumillon, E., Bertin, X., Fortunato, A., Bajo, M., Schneider, J.-L., Dezileau, L., Walsh, J. P., Michelot, A., Chauveau, E., Créach, A., Hénaff, A., Sauzeau, T., Waeles, B., Gervais, B., Jan, G., Baumann, J., Breilh, J.-F., and Pedreros, R.: Storm-induced marine flooding: lessons from a multidisciplinary approach, *Earth-Sci. Rev.*, 165, 151–184, , 2017.
- City of Venice: Attachment to Municipality Deliberation n.129 22/07/2002: Piano integrato degli interventi in caso di alta marea e bassa marea, 2002.
- 35 Cloke, H. L. and Pappenberger, F.: Ensemble flood forecasting: A review, *J. Hydrol.*, 375, 613–626, , 2009.

- Davolio, S., Stocchi, P., Benetazzo, A., Bohm, E., Riminucci, F., Ravaioli, M., Li, X.-M., and Carniel, S.: Exceptional Bora outbreak in winter 2012: Validation and analysis of high-resolution atmospheric model simulations in the northern Adriatic area, *Dynam. Atmos. Ocean*, 71, 1–20, , 2015.
- De Leo, F., Besio, G., Zolezzi, G., and Bezzi, M.: Coastal vulnerability assessment: through regional to local downscaling of wave characteristics along the Bay of Lalzit (Albania), *Nat. Hazards Earth Syst.*, 19, 287–298, , 2019.
- Depellegrin, D., Menegon, S., Farella, G., Ghezzi, M., Gissi, E., Sarretta, A., Venier, C., and Barbanti, A.: Multi-objective spatial tools to inform maritime spatial planning in the Adriatic Sea, *Sci. Total Environ.*, 609, 1627–1639, , 2017.
- Di Liberto, T., Colle, B. A., Georgas, N., Blumberg, A. F., and Taylor, A. A.: Verification of a Multimodel Storm Surge Ensemble around New York City and Long Island for the Cool Season, *Wea. Forecasting*, 26, 922–939, , 2011.
- Dodet, G., Leckler, F., Sous, D., Ardhuin, F., Filipot, J., and Suanez, S.: Wave Runup Over Steep Rocky Cliffs, *J. Geophys. Res. Oceans*, 123, 7185–7205, , 2018.
- Dutour-Sikirić, M., Ivanković, D., Roland, A., Ivatek-Šahdan, S., and Tudor, M.: Operational Wave Modelling in the Adriatic Sea with the Wind Wave Model, *Pure and Applied Geophysics*, 175, 3801–3815, , 2018.
- EMODnet Bathymetry Consortium: EMODnet Digital Bathymetry (DTM), , 2018.
- European Commission: Commission of the European Communities, Directive 2007/2/EC of the European Parliament and of the Council of 14 March 2007 Establishing an Infrastructure for Spatial Information in the European Community (INSPIRE), , 2007.
- European Commission: Open Source Software Strategy 2014–2017, , 2016.
- Fernández-Montblanc, T., Vousdoukas, M. I., Ciavola, P., Voukouvalas, E., Mentaschi, L., Breyiannis, G., Feyen, L., and Salamon, P.: Towards robust pan-European storm surge forecasting, *Ocean Model.*, 133, 129–144, , 2019.
- Ferrarin, C., Roland, A., Bajo, M., Umgiesser, G., Cucco, A., Davolio, S., Buzzi, A., Malguzzi, P., and Drofa, O.: Tide-surge-wave modelling and forecasting in the Mediterranean Sea with focus on the Italian coast, *Ocean Model.*, 61, 38–48, , 2013.
- Ferrarin, C., Tomasin, A., Bajo, M., Petrizzo, A., and Umgiesser, G.: Tidal changes in a heavily modified coastal wetland, *Cont. Shelf Res.*, 101, 22–33, , 2015.
- Ferrarin, C., Maieu, F., and Umgiesser, G.: The effect of lagoons on Adriatic Sea tidal dynamics, *Ocean Model.*, 119, 57–71, , 2017.
- Ferrarin, C., Bellafiore, D., Sannino, G., Bajo, M., and Umgiesser, G.: Tidal dynamics in the inter-connected Mediterranean, Marmara, Black and Azov seas, *Prog. Oceanogr.*, 161, 102–115, , 2018.
- Ferrarin, C., Davolio, S., Bellafiore, D., Ghezzi, M., Maieu, F., Drofa, O., Umgiesser, G., Bajo, M., De Pascalis, F., Marguzzi, P., Zaggia, L., Lorenzetti, G., Manfè, G., and Mc Kiver, W.: Cross-scale operational oceanography in the Adriatic Sea, *J. Oper. Oceanogr.*, 12, 86–103, , 2019.
- Flowerdew, J., Horsburgh, K., Wilson, C., and Mylne, K.: Development and evaluation of an ensemble forecasting system for coastal storm surges, *Q.J.R. Meteorol. Soc.*, 136, 1444–1456, , 2010.
- Golbeck, I., Li, X., Janssen, F., Brüning, T., Nielsen, J. W., Huess, V., Söderkvist, J., Büchmann, B., Siiriä, S.-M., Vähä-Piikkiö, O., Hackett, B., Kristensen, N. M., Engedahl, H., Blockley, E., Sellar, A., Lagemaat, P., Ozer, J., Legrand, S., Ljungemyr, P.,

- and Axell, L.: Uncertainty estimation for operational ocean forecast products – a multi-model ensemble for the North Sea and the Baltic Sea, *Ocean Dyn.*, 65, 1603–1631, , 2015.
- Hankin, S., Blower, J., Carval, T., Casey, K., Donlon, C., Lauret, O., Loubrieu, T., Srinivasan, A., Trinanes, J., Godøy, Ø., Mendelsohn, R., Signell, R., de La Beaujardiere, J., Cornillon, P., Blanc, F., Rew, R., and Harlan, J.: NetCDF-CF-OPeNDAP: Standards for ocean data interoperability and object lessons for community data standards processes, in: *Proceedings of OceanObs'09: Sustained Ocean Observations and Information for Society*, edited by Hall, J., Harrison, D., and Stammer, D., vol. 2, ESA Publication, , 2010.
- Harley, M. D., Valentini, A., Armaroli, C., Perini, L., Calabrese, L., and Ciavola, P.: Can an early warning system help minimize the impacts of coastal storms? A case study of the 2012 Halloween storm, Northern Italy, *Nat. Hazards Earth Syst.*, 16, 209–222, , 2016.
- Hinkel, J., Lincke, D., Vafeidis, A. T., Perrette, M., Nicholls, R. J., Tol, R. S. J., Marzeion, B., Fettweis, X., Ionescu, C., and Levermann, A.: Coastal flood damage and adaptation costs under 21st century sea level rise, *Proc. Natl. Acad. Sci.*, 111, 3292–3297, , 2014.
- Kolega, N.: Slovenian coast sea floods risk, *Acta geographica Slovenica*, 46, 143–167, , 2006.
- Lionello, P., Cavaleri, L., Nissen, K., Pino, C., Raicich, F., and Ulbrich, U.: Severe marine storms in the Northern Adriatic: Characteristics and trends, *Phys. Chem. Earth., Parts A/B/C*, 40–41, 93–105, , 2012.
- Ličer, M., Smerkol, P., Fettich, A., Ravdas, M., Papapostolou, A., Mantziafou, A., Strajnar, B., Cedilnik, J., Jeromel, M., Jerman, J., Petan, S., Malaveive, V., and Sofianos, S.: Modeling the ocean and atmosphere during an extreme bora event in northern Adriatic using one-way and two-way atmosphere–ocean coupling, *Ocean Sci.*, 12, 71–86, , 2016.
- Longueville, B. D.: Community-based geoportals: The next generation? Concepts and methods for the geospatial Web 2.0, *Comput. Environ. Urban Syst.*, 34, 299–308, , 2010.
- Magaña, P., Bergillos, R. J., Del-Rosal-Salido, J., Reyes-Merlo, M. A., Díaz-Carrasco, P., and Ortega-Sánchez, M.: Integrating complex numerical approaches into a user-friendly application for the management of coastal environments, *Sci. Total Environ.*, 624, 979–990, , 2018.
- Maguire, D. and Longley, P.: The emergence of geoportals and their role in spatial data infrastructures, *Comput. Environ. Urban Syst.*, 29, 3–14, , 2005.
- Marcos, M., Tsimplis, M. N., and Shaw, A. G. P.: Sea level extremes in southern Europe, *J. Geophys. Res.*, 144, C01007, , 2009.
- Mariani, S., Casaioli, M., Coraci, E., and Malguzzi, P.: A new high-resolution BOLAM-MOLOCH suite for the SIMM forecasting system: assessment over two HyMeX intense observation periods, *Nat. Hazards Earth Syst.*, 15, 1–24, , 2015.
- Medugorac, I., Pasarić, M., and Orlić, M.: Severe flooding along the eastern Adriatic coast: the case of 1 December 2008, *Ocean Dyn.*, 65, 817–830, , 2015.
- Mel, R. and Lionello, P.: Verification of an ensemble prediction system for storm surge forecast in the Adriatic Sea, *Ocean Dyn.*, 64, 1803–1814, , 2014.

- Molteni, F., Buizza, R., Marsigli, C., Montani, A., Nerozzi, F., and Paccagnella, T.: A strategy for high-resolution-ensemble prediction. I: Definition of representative members and global-model experiments, *Quart. J. Roy. Meteor. Soc.*, 127, 2069–2094, ,2001.–
- 5 Muis, S., Verlaan, M., Winsemius, H. C., Aerts, J. C. J. H., and Ward, P. J.: Aglobal reanalysis of storm surges and extreme sea levels, *Nat. Commun.*, 7, 11969, , 2016.–
- Orlić, M.: The first attempt at cataloguing tsunami-like waves of meteorological origin in Croatian coastal waters, *Acta Adriat.*, 56, 83–96, 2015.–
- Orlić, M., Gačić, M., and La Violett, P. E.: The currents and circulation of the Adriatic Sea, *Oceanologica Acta*, 15, 109–124, 1992.–
- 10 Pasarić, Z., Belušić, D., and Chiggiato, J.: Orographic effects on meteorological fields over the Adriatic from different models, *J. Mar. Syst.*, 78, S90–S100, , 2009.–
- Pirazzoli, P. A. and Tomasin, A.: Recent near-surface wind changes in the central Mediterranean and Adriatic area, *Int. J. Climatol.*, 23, 963–973, , 2003.–
- 15 Poate, T. G., McCall, R. T., and Masselink, G.: A new parameterisation for runup on gravel beaches, *Coast. Eng.*, 117, 176–190, , 2016.–
- Pomaro, A., Cavaleri, L., and Lionello, P.: Climatology and trends of the Adriatic Sea wind waves: analysis of a 37-year long instrumental data set, *Int. J. Climatol.*, 37, 4237–4250, , 2017.–
- Prahl, B. F., Boettle, M., Costa, L., Kropp, J. P., and Rybski, D.: Damage and protection cost curves for coastal floods within the 600 largest European cities, *Sci. Data*, 5, 180034, , 2018.–
- 20 Reimann, L., Vafeidis, A. T., Brown, S., Hinkel, J., and Tol, R. S. J.: Mediterranean UNESCO World Heritage at risk from coastal flooding and erosion due to sea-level rise, *Nat. Commun.*, 9, 4161, , 2018.–
- Rizzi, J., Torresan, S., Zabeo, A., Critto, A., Tosoni, A., Tomasin, A., and Marcomini, A.: Assessing storm surge risk under future sea-level rise scenarios: a case study in the North Adriatic coast, *J. Coast. Conser.*, 21, 453–471, , 2017.–
- 25 Roelvink, D., Reniers, A., van Dongeren, A., van Thiel de Vries, J., McCall, R., and Lescinski, J.: Modelling storm impacts on beaches, dunes and barrier islands, *Coast. Eng.*, 56, 1133–1152, , 2009.–
- Roland, A., Cucco, A., Ferrarin, C., Hsu, T.-W., Liau, J.-M., Ou, S.-H., Umgiesser, G., and Zanke, U.: On the development and verification of a 2d coupled wave-current model on unstructured meshes, *J. Mar. Syst.*, 78, Supplement, S244–S254, , 2009.–
- Russo, A., Colucci, A., Carniel, S., Benetazzo, A., Valentini, A., Paccagnella, T., Ravaioli, M., and Bortoluzzi, G.: 30 Operational models hierarchy for short term marine predictions: The Adriatic Sea example, in: 2013 MTS/IEEE OCEANS –Bergen, IEEE, , 2013.–
- Salighehdar, A., Ye, Z., Liu, M., Floreseu, I., and Blumberg, A. F.: Ensemble-Based Storm Surge Forecasting Models, *Wea. Forecasting*, 32, 1921–1936, , 2017.–
- 35 Satta, A., Puddu, M., Venturini, S., and Giupponi, C.: Assessment of coastal risks to climate change related impacts at the regional scale: The case of the Mediterranean region, *Int. J. Disaster Risk Reduc.*, 24, 284–296, , 2017.–

- Schevenhoven, F. J. and Selten, F. M.: An efficient training scheme for supermodels, *Earth Syst. Dynam.*, **8**, 429–438, , 2017.
- Schuerch, M., Spencer, T., Temmerman, S., Kirwan, M. L., Wolff, C., Lincke, D., McOwen, C. J., Pickering, M. D., Reef, R., Vafeidis, A. T., Hinkel, J., Nicholls, R. J., and Brown, S.: Future response of global coastal wetlands to sea-level rise, *Nature*, **561**, 231–234, , 2018.
- 5 Signell, R. P., Carniel, S., Cavaleri, L., Chiggiato, J., Doyle, J. D., Pullen, J., and Selavo, M.: Assessment of wind quality for oceanographic modelling in semi-enclosed basins, *J. Marine Syst.*, **53**, 217–233, , 2005.
- Sorensen, R. M.: Coastal Engineering, in: *Basic Coastal Engineering*, Springer US, , 1997.
- Stockdon, H. F., Holman, R. A., Howd, P. A., and Sallenger, A. H.: Empirical parameterization of setup, swash, and runup, *Coast. Eng.*, **53**, 573–588, , 2006.
- 10 Tonani, M., Pinardi, N., Fratianni, C., Pistoia, J., Dobricic, S., Pensieri, S., de Alfonso, M., and Nittis, K.: Mediterranean Forecasting System: forecast and analysis assessment through skill scores, *Ocean Sci.*, **5**, 649–660, , 2009.
- Umgiesser, G., Ferrarin, C., Cucco, A., De Pascalis, F., Bellafiore, D., Ghezzi, M., and Bajo, M.: Comparative hydrodynamics of 10 Mediterranean lagoons by means of numerical modeling, *J. Geophys. Res. Oceans*, **119**, 2212–2226, , 2014.
- 15 Unidata: THREDDS Data Server (TDS), version 4.6.13 software, Boulder, CO: UCAR/Unidata, , 2019.
- Vaccari, L., Craglia, M., Fugazza, C., Nativi, S., and Santoro, M.: Integrative Research: The EuroGEOSS Experience, *IEEE J. Sel. Topics Appl. Earth Observ. Remote Sens.*, **5**, 1603–1611, , 2012.
- Valentini, A., Delli Passeri, L., Paccagnella, T., Patruno, P., Marsigli, C., Deserti, M., Chiggiato, J., and Tibaldi, S.: The Sea State forecast system of ARPA-SIM, *Boll. Geof. Teor. Appl.*, **48**, 333–349, 2007.
- 20 Vilibić, I., Šepić, J., Pasarić, M., and Orlić, M.: The Adriatic Sea: A Long-Standing Laboratory for Sea Level Studies, *Pure Appl. Geophys.*, , 2017.
- Vousdoukas, M. I., Voukouvalas, E., Mentaschi, L., Dottori, F., Giardino, A., Bouziotas, D., Bianchi, A., Salamon, P., and Feyen, L.: Developments in large-scale coastal flood hazard mapping, *Nat. Hazards Earth Syst.*, **16**, 1841–1853, , 2016.
- Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., and Feyen, L.: Extreme sea levels on the rise along Europe’s  
25 coasts, *Earth’s Future*, **5**, 304–323, , 2017.
- Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Bianchi, A., Dottori, F., and Feyen, L.: Climatic and socioeconomic controls of future coastal flood risk in Europe, *Nat. Clim. Chang.*, **8**, 776–780, , 2018.
- Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., Jevrejeva, S., Jackson, L. P., and Feyen, L.: Global probabilistic projections of extreme sea levels show intensification of coastal flood hazard, *Nat. Commun.*, **9**, 2360, , 2018.
- 30 Wolff, C., Vafeidis, A. T., Lincke, D., Marasmi, C., and Hinkel, J.: Effects of Scale and Input Data on Assessing the Future Impacts of Coastal Flooding: An Application of DIVA for the Emilia-Romagna Coast, *Front. Mar. Sci.*, **3**, 2016.
- Wolff, C., Vafeidis, A. T., Muis, S., Lincke, D., Satta, A., Lionello, P., Jimenez, J. A., Conte, D., and Hinkel, J.: A Mediterranean coastal database for assessing the impacts of sea-level rise and associated hazards, *Sci. Data*, **5**, 180044, , 2018.

~~World Meteorological Organization: Guidelines on Ensemble Prediction Systems and Forecasting, World Meteorological Organization: Weather, Climate, and Water, 1091, 2012.~~

~~Yang, P., Evans, J., Cole, M., Marley, S., Alameh, N., and Bambacus, M.: The Emerging Concepts and Applications of the Spatial-Web-Portal, Photogramm. Eng. Remote Sens., 73, 691–698, , 2007.~~

5 ~~Zacharioudaki, A., Korres, G., and Perivoliotis, L.: Wave climate of the Hellenic Seas obtained from a wave hindcast for the period 1960–2001, Ocean Dyn., 65, 795–816, , 2015.~~

~~Zou, Q.-P., Chen, Y., Cluckie, I., Hewston, R., Pan, S., Peng, Z., and Reeve, D.: Ensemble prediction of coastal flood risk arising from overtopping by linking meteorological, ocean, coastal and surf zone models, Quart. J. Roy. Meteor. Soc., 139, 298–313, , 2013.~~

## References

- Armaroli, C. and Duo, E.: Validation of the coastal storm risk assessment framework along the Emilia-Romagna coast, *Coast. Eng.*, 134, 159–167, <https://doi.org/10.1016/j.coastaleng.2017.08.014>, 2018.
- Armaroli, C., Ciavola, P., Perini, L., Calabrese, L., Lorito, S., Valentini, A., and Masina, M.: Critical storm thresholds for significant morphological changes and damage along the Emilia-Romagna coastline, Italy, *Geomorphology*, 143-144, 34–51, <https://doi.org/10.1016/j.geomorph.2011.09.006>, 2012.
- Baart, F., van Gelder, P. H. A. J. M., and van Koningsveld, M.: Confidence in real-time forecasting of morphological storm impacts, *J. Coast. Res.*, pp. 1835–1839, 2011.
- Bajo, M. and Umgiesser, G.: Storm surge forecast through a combination of dynamic and neural network models, *Ocean Model.*, 33, 1–9, <https://doi.org/10.1016/j.ocemod.2009.12.007>, 2010.
- Bajo, M., De Biasio, F., Umgiesser, G., Vignudelli, S., and Zecchetto, S.: Impact of using scatterometer and altimeter data on storm surge forecasting, *Ocean Model.*, 113, 85 – 94, <https://doi.org/10.1016/j.ocemod.2017.03.014>, 2017.
- Bajo, M., Medugorac, I., Umgiesser, G., and Orlić, M.: Storm surge and seiche modelling in the Adriatic Sea and the impact of data assimilation, *Quart. J. Roy. Meteor. Soc.*, 145, 2070–2084, <https://doi.org/10.1002/qj.3544>, 2019.
- Bernier, N. B. and Thompson, K. R.: Deterministic and ensemble storm surge prediction for Atlantic Canada with lead times of hours to ten days, *Ocean Model.*, 86, 114 – 127, <https://doi.org/10.1016/j.ocemod.2014.12.002>, 2015.
- Bertotti, L., Canestrelli, P., Cavaleri, L., Pastore, F., and Zampato, L.: The Henetus wave forecast system in the Adriatic Sea, *Nat. Hazards Earth Syst.*, 11, 2965–2979, <https://doi.org/10.5194/nhess-11-2965-2011>, 2011.
- Bertotti, L., Cavaleri, L., Loffredo, L., and Torrisi, L.: Nettuno: Analysis of a Wind and Wave Forecast System for the Mediterranean Sea, *Mon. Weather Rev.*, 141, 3130–3141, <https://doi.org/10.1175/mwr-d-12-00361.1>, 2013.
- Bosom, E. and Jiménez, J. A.: Probabilistic coastal vulnerability assessment to storms at regional scale - application to Catalan beaches (NW Mediterranean), *Nat. Hazards Earth Syst.*, 11, 475–484, <https://doi.org/10.5194/nhess-11-475-2011>, 2011.
- Bressan, L., Valentini, A., Paccagnella, T., Montani, A., Marsigli, C., and Tesini, M. S.: Sensitivity of sea-level forecasting to the horizontal resolution and sea surface forcing for different configurations of an oceanographic model of the Adriatic Sea, *Adv. Sci. Res.*, 14, 77–84, <https://doi.org/10.5194/asr-14-77-2017>, 2017.
- Carbognin, L. and Tosi, L.: Interaction between Climate Changes, Eustacy and Land Subsidence in the North Adriatic Region, Italy, *Mar. Ecol.*, 23, 38–50, <https://doi.org/10.1111/j.1439-0485.2002.tb00006.x>, 2002.
- Cavaleri, L. and Bertotti, L.: Accuracy of the modelled wind and wave fields in enclosed seas, *Tellus A*, 56, 167–175, <https://doi.org/10.3402/tellusa.v56i2.14398>, 2004.
- Cavaleri, L., Bertotti, L., Buizza, R., Buzzi, A., Masato, V., Umgiesser, G., and Zampieri, M.: Predictability of extreme meteo-oceanographic events in the Adriatic Sea, *Q. J. Roy. Meteor. Soc.*, 136, 400–413, <https://doi.org/10.1002/qj.567>, 2010.
- Cavaleri, L., Bajo, M., Barbariol, F., Bastianini, M., Benetazzo, A., Bertotti, L., Chiggiato, J., Davolio, S., Ferrarin, C., Magnusson, L., Papa, A., Pezzutto, P., Pomaro, A., and Umgiesser, G.: The October 29, 2018 storm in Northern Italy - an exceptional event and its modeling, *Prog. Oceanogr.*, 178, 102 178, <https://doi.org/10.1016/j.pocean.2019.102178>, 2019.
- Chaumillon, E., Bertin, X., Fortunato, A., Bajo, M., Schneider, J.-L., Dezileau, L., Walsh, J. P., Michelot, A., Chauveau, E., Créach, A., Hénaff, A., Sauzeau, T., Waeles, B., Gervais, B., Jan, G., Baumann, J., Breilh, J.-F., and Pedreros, R.: Storm-induced marine flooding: lessons from a multidisciplinary approach, *Earth-Sci. Rev.*, 165, 151 – 184, <https://doi.org/10.1016/j.earscirev.2016.12.005>, 2017.


- City of Venice: Attachment to Municipality Deliberation n.129 22/07/2002: Piano integrato degli interventi in caso di alta marea e bassa marea, 2002.
- Cloke, H. L. and Pappenberger, F.: Ensemble flood forecasting: A review, *J. Hydrol.*, 375, 613 – 626, <https://doi.org/10.1016/j.jhydrol.2009.06.005>, 2009.
- 5 Davolio, S., Stocchi, P., Benetazzo, A., Bohm, E., Riminucci, F., Ravaioli, M., Li, X.-M., and Carniel, S.: Exceptional Bora outbreak in winter 2012: Validation and analysis of high-resolution atmospheric model simulations in the northern Adriatic area, *Dynam. Atmos. Ocean*, 71, 1 – 20, <https://doi.org/10.1016/j.dynatmoce.2015.05.002>, 2015.
- De Leo, F., Besio, G., Zolezzi, G., and Bezzi, M.: Coastal vulnerability assessment: through regional to local downscaling of wave characteristics along the Bay of Lalzit (Albania), *Nat. Hazards Earth Syst.*, 19, 287–298, <https://doi.org/10.5194/nhess-19-287-2019>, 2019.
- 10 Depellegrin, D., Menegon, S., Farella, G., Ghezzi, M., Gissi, E., Sarretta, A., Venier, C., and Barbanti, A.: Multi-objective spatial tools to inform maritime spatial planning in the Adriatic Sea, *Sci. Total Environ.*, 609, 1627 – 1639, <https://doi.org/10.1016/j.scitotenv.2017.07.264>, 2017.
- Di Liberto, T., Colle, B. A., Georgas, N., Blumberg, A. F., and Taylor, A. A.: Verification of a Multimodel Storm Surge Ensemble around New York City and Long Island for the Cool Season, *Wea. Forecasting*, 26, 922–939, <https://doi.org/10.1175/waf-d-10-05055.1>, 2011.
- 15 Dodet, G., Leckler, F., Sous, D., Ardhuin, F., Filipot, J., and Suanez, S.: Wave Runup Over Steep Rocky Cliffs, *J. Geophys. Res. Oceans*, 123, 7185–7205, <https://doi.org/10.1029/2018jc013967>, 2018.
- Dutour Sikirić, M., Ivanković, D., Roland, A., Ivatek-Šahdan, S., and Tudor, M.: Operational Wave Modelling in the Adriatic Sea with the Wind Wave Model, *Pure and Applied Geophysics*, 175, 3801–3815, <https://doi.org/10.1007/s00024-018-1954-2>, 2018.
- EMODnet Bathymetry Consortium: EMODnet Digital Bathymetry (DTM), <https://doi.org/10.12770/18ff0d48-b203-4a65-94a9-5fd8b0ec35f6>, 2018.
- 20 European Commission: Commission of the European Communities, Directive 2007/2/EC of the European Parliament and of the Council of 14 March 2007 Establishing an Infrastructure for Spatial Information in the European Community (INSPIRE), <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=OJ>, 2007.
- European Commission: Open Source Software Strategy 2014-2017, [https://ec.europa.eu/info/departments/informatics/open-source-software-strategy\\_en](https://ec.europa.eu/info/departments/informatics/open-source-software-strategy_en), 2016.
- 25 Fernández-Montblanc, T., Vousedoukas, M. I., Ciavola, P., Voukouvalas, E., Mentaschi, L., Breyiannis, G., Feyen, L., and Salamon, P.: Towards robust pan-European storm surge forecasting, *Ocean Model.*, 133, 129 – 144, <https://doi.org/10.1016/j.ocemod.2018.12.001>, 2019.
- Ferrarin, C., Roland, A., Bajo, M., Umgiesser, G., Cucco, A., Davolio, S., Buzzi, A., Malguzzi, P., and Drofa, O.: Tide-surge-wave modelling and forecasting in the Mediterranean Sea with focus on the Italian coast, *Ocean Model.*, 61, 38–48, <https://doi.org/10.1016/j.ocemod.2012.10.003>, 2013.
- 30 Ferrarin, C., Tomasin, A., Bajo, M., Petrizzo, A., and Umgiesser, G.: Tidal changes in a heavily modified coastal wetland, *Cont. Shelf Res.*, 101, 22 – 33, <https://doi.org/10.1016/j.csr.2015.04.002>, 2015.
- Ferrarin, C., Maicu, F., and Umgiesser, G.: The effect of lagoons on Adriatic Sea tidal dynamics, *Ocean Model.*, 119, 57–71, <https://doi.org/10.1016/j.ocemod.2017.09.009>, 2017.
- 35 Ferrarin, C., Bellafiore, D., Sannino, G., Bajo, M., and Umgiesser, G.: Tidal dynamics in the inter-connected Mediterranean, Marmara, Black and Azov seas, *Prog. Oceanogr.*, 161, 102–115, <https://doi.org/10.1016/j.pocean.2018.02.006>, 2018.

- Ferrarin, C., Davolio, S., Bellafiore, D., Ghezzi, M., Maicu, F., Drofa, O., Umgiesser, G., Bajo, M., De Pascalis, F., Marguzzi, P., Zaggia, L., Lorenzetti, G., Manfè, G., and Mc Kiver, W.: Cross-scale operational oceanography in the Adriatic Sea, *J. Oper. Oceanogr.*, 12, 86–103, <https://doi.org/10.1080/1755876X.2019.1576275>, 2019.
- 5 Ferreira, O., Plomaritis, T. A., and Costas, S.: Process-based indicators to assess storm induced coastal hazards, *Earth-Sci. Rev.*, 173, 159 – 167, <https://doi.org/10.1016/j.earscirev.2017.07.010>, 2017.
- Flowerdew, J., Horsburgh, K., Wilson, C., and Mylne, K.: Development and evaluation of an ensemble forecasting system for coastal storm surges, *Q.J.R. Meteorol. Soc.*, 136, 1444–1456, <https://doi.org/10.1002/qj.648>, 2010.
- 10 Golbeck, I., Li, X., Janssen, F., Brüning, T., Nielsen, J. W., Huess, V., Söderkvist, J., Büchmann, B., Siiriä, S.-M., Vähä-Piikkiö, O., Hackett, B., Kristensen, N. M., Engedahl, H., Blockley, E., Sellar, A., Lagema, P., Ozer, J., Legrand, S., Ljungemyr, P., and Axell, L.: Uncertainty estimation for operational ocean forecast products - a multi-model ensemble for the North Sea and the Baltic Sea, *Ocean Dyn.*, 65, 1603–1631, <https://doi.org/10.1007/s10236-015-0897-8>, 2015.
- Hankin, S., Blower, J., Carval, T., Casey, K., Donlon, C., Lauret, O., Loubrieu, T., Srinivasan, A., Trinanes, J., Godøy, Ø., Mendelssohn, R., Signell, R., de La Beaujardiere, J., Cornillon, P., Blanc, F., Rew, R., and Harlan, J.: NetCDF-CF-OPeNDAP: Standards for ocean data interoperability and object lessons for community data standards processes, in: *Proceedings of OceanObs'09: Sustained Ocean Observations and Information for Society*, edited by Hall, J., Harrison, D., and Stammer, D., vol. 2, ESA Publication, <https://doi.org/10.5270/OceanObs09.cwp.41>, 2010.
- 15 Harley, M. D., Valentini, A., Armaroli, C., Perini, L., Calabrese, L., and Ciavola, P.: Can an early warning system help minimize the impacts of coastal storms? A case study of the 2012 Halloween storm, Northern Italy, *Nat. Hazards Earth Syst.*, 16, 209–222, <https://doi.org/10.5194/nhess-16-209-2016>, 2016.
- 20 Hinkel, J., Lincke, D., Vafeidis, A. T., Perrette, M., Nicholls, R. J., Tol, R. S. J., Marzeion, B., Fettweis, X., Ionescu, C., and Levermann, A.: Coastal flood damage and adaptation costs under 21st century sea-level rise, *Proc. Natl Acad. Sci.*, 111, 3292–3297, <https://doi.org/10.1073/pnas.1222469111>, 2014.
- Kolega, N.: Slovenian coast sea floods risk, *Acta geographica Slovenica*, 46, 143–167, <https://doi.org/10.3986/ags46201>, 2006.
- Lionello, P., Sanna, A., Elvini, E., and Mufato, R.: A data assimilation procedure for operational prediction of storm surge in the northern Adriatic Sea, *Cont. Shelf Res.*, 26, 539–553, <https://doi.org/10.1016/j.csr.2006.01.003>, 2006.
- 25 Lionello, P., Cavaleri, L., Nissen, K., Pino, C., Raicich, F., and Ulbrich, U.: Severe marine storms in the Northern Adriatic: Characteristics and trends, *Phys. Chem. Earth, Parts A/B/C*, 40-41, 93–105, <https://doi.org/10.1016/j.pce.2010.10.002>, 2012.
- Ličer, M., Smerkol, P., Fettich, A., Ravdas, M., Papapostolou, A., Mantziafou, A., Strajnar, B., Cedilnik, J., Jeromel, M., Jerman, J., Petan, S., Malavcivc, V., and Sofianos, S.: Modeling the ocean and atmosphere during an extreme bora event in northern Adriatic using one-way and two-way atmosphere-ocean coupling, *Ocean Sci.*, 12, 71–86, <https://doi.org/10.5194/os-12-71-2016>, 2016.
- 30 Longuet-Higgins, M. S. and Steward, R. W.: A note on wave set-up, *J. Mar. Res.*, 21, 4–10, 1963.
- Longueville, B. D.: Community-based geoportals: The next generation? Concepts and methods for the geospatial Web 2.0, *Comput. Environ. Urban Syst.*, 34, 299–308, <https://doi.org/10.1016/j.compenvurbsys.2010.04.004>, 2010.
- Magaña, P., Bergillos, R. J., Del-Rosal-Salido, J., Reyes-Merlo, M. A., Díaz-Carrasco, P., and Ortega-Sánchez, M.: Integrating complex numerical approaches into a user-friendly application for the management of coastal environments, *Sci. Total Environ.*, 624, 979–990, <https://doi.org/10.1016/j.scitotenv.2017.12.154>, 2018.
- 35 Maguire, D. and Longley, P.: The emergence of geoportals and their role in spatial data infrastructures, *Comput. Environ. Urban Syst.*, 29, 3–14, [https://doi.org/10.1016/s0198-9715\(04\)00045-6](https://doi.org/10.1016/s0198-9715(04)00045-6), 2005.

- Marcos, M., Tsimplis, M. N., and Shaw, A. G. P.: Sea level extremes in southern Europe, *J. Geophys. Res.*, 144, C01007, <https://doi.org/10.1029/2008JC004912>, 2009.
- Mariani, S., Casaioli, M., Coraci, E., and Malguzzi, P.: A new high-resolution BOLAM-MOLOCH suite for the SIMM forecasting system: assessment over two HyMeX intense observation periods, *Nat. Hazards Earth Syst.*, 15, 1–24, <https://doi.org/10.5194/nhess-15-1-2015>, 5 2015.
- Medugorac, I., Pasarić, M., and Orlić, M.: Severe flooding along the eastern Adriatic coast: the case of 1 December 2008, *Ocean Dyn.*, 65, 817–830, <https://doi.org/10.1007/s10236-015-0835-9>, 2015.
- Mel, R. and Lionello, P.: Verification of an ensemble prediction system for storm surge forecast in the Adriatic Sea, *Ocean Dyn.*, 64, 1803–1814, <https://doi.org/10.1007/s10236-014-0782-x>, 2014a.
- 10 Mel, R. and Lionello, P.: Storm surge ensemble prediction for the city of Venice, *Wea. Forecasting*, 29, 1044–1057, <https://doi.org/10.1175/WAF-D-13-00117.1>, 2014b.
- Mel, R. and Lionello, P.: Probabilistic dressing of a storm surge prediction in the Adriatic Sea, *Adv. Meteorol.*, 3764519, <https://doi.org/10.1155/2016/3764519>, 2016.
- Molteni, F., Buizza, R., Marsigli, C., Montani, A., Nerozzi, F., and Paccagnella, T.: A strategy for high-resolution ensemble prediction. I: Definition of representative members and global-model experiments, *Quart. J. Roy. Meteor. Soc.*, 127, 2069–2094, <https://doi.org/10.1002/qj.49712757612>, 2001.
- 15 Muis, S., Verlaan, M., Winsemius, H. C., Aerts, J. C. J. H., and Ward, P. J.: A global reanalysis of storm surges and extreme sea levels, *Nat. Commun.*, 7, 11 969, <https://doi.org/10.1038/ncomms11969>, 2016.
- Orlić, M.: The first attempt at cataloguing tsunami-like waves of meteorological origin in Croatian coastal waters, *Acta Adriat.*, 56, 83 – 96, 20 2015.
- Orlić, M., Gačić, M., and La Violett, P. E.: The currents and circulation of the Adriatic Sea, *Oceanologica Acta*, 15, 109–124, 1992.
- Pasarić, Z., Belušić, D., and Chiggiato, J.: Orographic effects on meteorological fields over the Adriatic from different models, *J. Mar. Syst.*, 78, S90 – S100, <https://doi.org/10.1016/j.jmarsys.2009.01.019>, 2009.
- Pirazzoli, P. A. and Tomasin, A.: Recent near-surface wind changes in the central Mediterranean and Adriatic area, *Int. J. Climatol.*, 23, 25 963–973, <https://doi.org/10.1002/joc.925>, 2003.
- Poate, T. G., McCall, R. T., and Masselink, G.: A new parameterisation for runup on gravel beaches, *Coast. Eng.*, 117, 176–190, <https://doi.org/10.1016/j.coastaleng.2016.08.003>, 2016.
- Pomaro, A., Cavaleri, L., and Lionello, P.: Climatology and trends of the Adriatic Sea wind waves: analysis of a 37-year long instrumental data set, *Int. J. Climatol.*, 37, 4237–4250, <https://doi.org/10.1002/joc.5066>, 2017.
- 30 Prael, B. F., Boettle, M., Costa, L., Kropp, J. P., and Rybski, D.: Damage and protection cost curves for coastal floods within the 600 largest European cities, *Sci. Data*, 5, 180 034, <https://doi.org/10.1038/sdata.2018.34>, 2018.
- Reimann, L., Vafeidis, A. T., Brown, S., Hinkel, J., and Tol, R. S. J.: Mediterranean UNESCO World Heritage at risk from coastal flooding and erosion due to sea-level rise, *Nat. Commun.*, 9, 4161, <https://doi.org/10.1038/s41467-018-06645-9>, 2018.
- Rizzi, J., Torresan, S., Zabeo, A., Critto, A., Tosoni, A., Tomasin, A., and Marcomini, A.: Assessing storm surge risk under future sea-level rise scenarios: a case study in the North Adriatic coast, *J. Coast. Conser.*, 21, 453–471, <https://doi.org/10.1007/s11852-017-0517-5>, 2017.
- 35 Roelvink, D., Reniers, A., van Dongeren, A., van Thiel de Vries, J., McCall, R., and Lescinski, J.: Modelling storm impacts on beaches, dunes and barrier islands, *Coast. Eng.*, 56, 1133–1152, <https://doi.org/10.1016/j.coastaleng.2009.08.006>, 2009.

- Roland, A., Cucco, A., Ferrarin, C., Hsu, T.-W., Liau, J.-M., Ou, S.-H., Umgiesser, G., and Zanke, U.: On the development and verification of a 2d coupled wave-current model on unstructured meshes, *J. Mar. Syst.*, 78, Supplement, S244–S254, <https://doi.org/10.1016/j.jmarsys.2009.01.026>, 2009.
- Russo, A., Coluccelli, A., Carniel, S., Benetazzo, A., Valentini, A., Paccagnella, T., Ravaioli, M., and Bortoluzzi, G.: Operational models hierarchy for short term marine predictions: The Adriatic Sea example, in: 2013 MTS/IEEE OCEANS - Bergen, IEEE, <https://doi.org/10.1109/oceans-bergen.2013.6608139>, 2013.
- Salighehdar, A., Ye, Z., Liu, M., Florescu, I., and Blumberg, A. F.: Ensemble-Based Storm Surge Forecasting Models, *Wea. Forecasting*, 32, 1921–1936, <https://doi.org/10.1175/waf-d-17-0017.1>, 2017.
- Satta, A., Puddu, M., Venturini, S., and Giupponi, C.: Assessment of coastal risks to climate change related impacts at the regional scale: The case of the Mediterranean region, *Int. J. Disaster Risk Reduc.*, 24, 284–296, <https://doi.org/10.1016/j.ijdrr.2017.06.018>, 2017.
- Schevenhoven, F. J. and Selten, F. M.: An efficient training scheme for supermodels, *Earth Syst. Dynam.*, 8, 429–438, <https://doi.org/10.5194/esd-8-429-2017>, 2017.
- Schuerch, M., Spencer, T., Temmerman, S., Kirwan, M. L., Wolff, C., Lincke, D., McOwen, C. J., Pickering, M. D., Reef, R., Vafeidis, A. T., Hinkel, J., Nicholls, R. J., and Brown, S.: Future response of global coastal wetlands to sea-level rise., *Nature*, 561, 231–234, <https://doi.org/10.1038/s41586-018-0476-5>, 2018.
- Signell, R. P., Carniel, S., Cavaleri, L., Chiggiato, J., Doyle, J. D., Pullen, J., and Sclavo, M.: Assessment of wind quality for oceanographic modelling in semi-enclosed basins, *J. Marine Syst.*, 53, 217 – 233, <https://doi.org/10.1016/j.jmarsys.2004.03.006>, 2005.
- Sorensen, R. M.: Coastal Engineering, in: Basic Coastal Engineering, Springer US, [https://doi.org/10.1007/978-1-4757-2665-7\\_1](https://doi.org/10.1007/978-1-4757-2665-7_1), 1997.
- Stockdon, H. F., Holman, R. A., Howd, P. A., and Sallenger, A. H.: Empirical parameterization of setup, swash, and runup, *Coast. Eng.*, 53, 573–588, <https://doi.org/10.1016/j.coastaleng.2005.12.005>, 2006.
- Tonani, M., Pinardi, N., Fratianni, C., Pistoia, J., Dobricic, S., Pensieri, S., de Alfonso, M., and Nittis, K.: Mediterranean Forecasting System: forecast and analysis assessment through skill scores, *Ocean Sci.*, 5, 649–660, <https://doi.org/10.5194/os-5-649-2009>, 2009.
- Umgiesser, G., Ferrarin, C., Cucco, A., De Pascalis, F., Bellafiore, D., Ghezzi, M., and Bajo, M.: Comparative hydrodynamics of 10 Mediterranean lagoons by means of numerical modeling, *J. Geophys. Res. Oceans*, 119, 2212–2226, <https://doi.org/10.1002/2013JC009512>, 2014.
- Unidata: THREDDS Data Server (TDS), version 4.6.13 [software], Boulder, CO: UCAR/Unidata, <https://doi.org/10.5065/D6N014KG>, 2019.
- Vaccari, L., Craglia, M., Fugazza, C., Nativi, S., and Santoro, M.: Integrative Research: The EuroGEOSS Experience, *IEEE J. Sel. Topics Appl. Earth Observ. Remote Sens.*, 5, 1603–1611, <https://doi.org/10.1109/jstars.2012.2190382>, 2012.
- Valentini, A., Delli Passeri, L., Paccagnella, T., Patrino, P., Marsigli, C., Deserti, M., Chiggiato, J., and Tibaldi, S.: The Sea State forecast system of ARPA-SIM, *Boll. Geof. Teor. Appl.*, 48, 333–349, 2007.
- Vilibić, I., Šepić, J., Pasarić, M., and Orlić, M.: The Adriatic Sea: A Long-Standing Laboratory for Sea Level Studies, *Pure Appl. Geophys.*, <https://doi.org/10.1007/s00024-017-1625-8>, 2017.
- Vousdoukas, M. I., Voukouvalas, E., Mentaschi, L., Dottori, F., Giardino, A., Bouziotas, D., Bianchi, A., Salamon, P., and Feyen, L.: Developments in large-scale coastal flood hazard mapping, *Nat. Hazards Earth Syst.*, 16, 1841–1853, <https://doi.org/10.5194/nhess-16-1841-2016>, 2016.
- Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., and Feyen, L.: Extreme sea levels on the rise along Europe’s coasts, *Earth’s Future*, 5, 304–323, <https://doi.org/10.1002/2016ef000505>, 2017.
- Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Bianchi, A., Dottori, F., and Feyen, L.: Climatic and socioeconomic controls of future coastal flood risk in Europe, *Nat. Clim. Chang.*, 8, 776–780, <https://doi.org/10.1038/s41558-018-0260-4>, 2018a.

- Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., Jevrejeva, S., Jackson, L. P., and Feyen, L.: Global probabilistic projections of extreme sea levels show intensification of coastal flood hazard, *Nat. Commun.*, 9, 2360, <https://doi.org/10.1038/s41467-018-04692-w>, 2018b.
- Wolff, C., Vafeidis, A. T., Lincke, D., Marasmi, C., and Hinkel, J.: Effects of Scale and Input Data on Assessing the Future Impacts of Coastal Flooding: An Application of DIVA for the Emilia-Romagna Coast, *Front. Mar. Sci.*, 3, 2016.
- Wolff, C., Vafeidis, A. T., Muis, S., Lincke, D., Satta, A., Lionello, P., Jimenez, J. A., Conte, D., and Hinkel, J.: A Mediterranean coastal database for assessing the impacts of sea-level rise and associated hazards, *Sci. Data*, 5, 180 044, <https://doi.org/10.1038/sdata.2018.44>, 2018.
- World Meteorological Organization: Guidelines on Ensemble Prediction Systems and Forecasting, World Meteorological Organization: Weather, Climate, and Water, 1091, 2012.
- Yang, P., Evans, J., Cole, M., Marley, S., Alameh, N., and Bambacus, M.: The Emerging Concepts and Applications of the Spatial Web Portal, *Photogramm. Eng. Remote Sens.*, 73, 691–698, <https://doi.org/10.14358/pers.73.6.691>, 2007.
- Zacharioudaki, A., Korres, G., and Perivoliotis, L.: Wave climate of the Hellenic Seas obtained from a wave hindcast for the period 1960–2001, *Ocean Dyn.*, 65, 795–816, <https://doi.org/10.1007/s10236-015-0840-z>, 2015.
- Zou, Q.-P., Chen, Y., Cluckie, I., Hewston, R., Pan, S., Peng, Z., and Reeve, D.: Ensemble prediction of coastal flood risk arising from overtopping by linking meteorological, ocean, coastal and surf zone models, *Quart. J. Roy. Meteor. Soc.*, 139, 298–313, <https://doi.org/10.1002/qj.2078>, 2013.



**Figure 1.** Bathymetry of the Adriatic Sea with monitoring stations for sea surface height (yellow dots) and waves (red stars). The 50-year extreme sea levels (ESL) from Vousdoukas et al. (2017) are also reported. Background: EMODnet bathymetry (EMODnet Bathymetry Consortium, 2018).




Figure 2. Schematic representation of the IWS architecture.







Figure 3. ECMWF 10m wind speed and mean sea level pressure fields over the Mediterranean Sea of 29 October 2018 at 18 UTC.




**Figure 4.** Observed sea level height (a) and significant wave heights (b) measured at different locations (see Fig. 1 for reference).




**Figure 5.** 29 October 2018 results of TMES in terms of the ensemble mean (a, c) and ensemble standard deviation (b, d) for sea level height at 13 UTC (a, b) and significant wave height at 19 UTC (c, d), respectively.




**Figure 6.** TMES sea level height extracted at PTF (a) and Rovinj (b), and significant wave height extracted at PTF (c) and Dubrovnik (d).




**Figure 7.** Forecasted total water 2% exceedance level along the Adriatic coastline for MEAN of runup maxima (a) and MAX total water level (b) for the MAX sea condition scenarios scenario at 19 UTC of October the 29<sup>th</sup>, 2018. Background: EMODnet bathymetry (EMODnet Bathymetry Consortium, 2018).




**Figure 8.** Observed and predicted (according to the three sea condition scenarios) sea level height at Koper (Slovenia). The yellow, orange and red lines indicate the adopted thresholds for flooding alerts.



**Figure 9.** Predicted coastal hazard (a) and the documented coastal impacts (b) along the coast of Emilia-Romagna, Italy. Background: image Google, ©2019 TerraMetrics.



**Figure 10.** Forecasted Safe Corridor Width Index for the beach profile of Classe06 (Lido di Classe, Emilia-Romagna, Italy). The dashed orange and red lines indicate respectively the medium and high thresholds for coastal alerts.



**Figure 11.** Flooding map of the City of Venice according to the predicted sea level height at Punta della Salute (black dot). The colours represent the flooded pedestrian area for sea levels of 140 cm (blue; MEAN scenario) and 160 cm (blue and red; MAX scenarios). Light blue indicates the canals.

**Table 1.** Details of the sea level forecasting systems used in the TMES. Key references are reported at the bottom of the table.

Managing authority - Country	System name	Domain	Forecast range	Horizontal res.	Core engine	Tide	Baroclinic	Meteo forcing
City of Venice - IT	SHYMED <sup>1</sup>	Mediterranean Sea	10 days	var. up to 200 m	SHYFEM	no	no	ECMWF 1/8 deg
National Research Council - IT	Kassandra <sup>2</sup>	Mediterranean Sea	4 days	var. up to 100 m	SHYFEM	yes	no	BOLAM 8.3 km, MOLOCH 1.25 km
National Research Council - IT	ISSOS	Mediterranean Sea	3 days	var. up to 200 m	SHYFEM	no	no	BOLAM 8.3 km
National Research Council - IT	Tiresias <sup>3</sup>	Adriatic Sea	3 days	var. up to 10 m	SHYFEM	yes	yes	MOLOCH 1.25 km
Ag. for Env. Protection and Energy ER - IT	AdriaROMS <sup>4</sup>	Adriatic Sea	3 days	2 km	ROMS	yes	yes	COSMO-5M 5 km
Ag. for Env. Protection and Energy ER - IT	Adriae <sup>5</sup>	Adriatic Sea	3 days	1 km	ROMS	yes	yes	COSMO-2I 2.2 km, 5M 5km
Inst. for Env. Protection and Research - IT	SIMMb <sup>6</sup>	Mediterranean Sea	3.5 days	var. up to 1 km	SHYFEM	no	no	BOLAM 11 km
Inst. for Env. Protection and Research - IT	SIMMe <sup>6</sup>	Mediterranean Sea	4 days	var. up to 1 km	SHYFEM	no	no	ECMWF 55 km
Slovenian Environment Agency - SL	SMMO <sup>7</sup>	Adriatic Sea	3 days	1/216 deg	NEMO	yes	yes	ALADIN 4.4 km
CMCC - IT	MED-Currents <sup>8*</sup>	Mediterranean Sea	10 days	1/24 deg	NEMO	no	yes	ECMWF 1/8 deg

<sup>1</sup>Bajo and Umgiesser (2010); <sup>2</sup>Ferrarin et al. (2013); <sup>3</sup>Ferrarin et al. (2019); <sup>4</sup>Russo et al. (2013); <sup>5</sup>Bressan et al. (2017); <sup>6</sup>Mariani et al. (2015); <sup>7</sup>Ličer et al. (2016); <sup>8</sup>Tonani et al. (2009); \*retrieved from CMEMS, <http://marine.copernicus.eu/>.

**Details of the wave monitoring stations (red stars in Fig. 1).**

**Table 2.** Details of the wave forecasting systems used in the TMES. Key references are reported at the bottom of the table.

Managing authority - Country	System name	Domain	Forecast range	Horizontal res.	Core engine	Meteo forcing
National Research Council - IT	Kassandra <sup>1</sup>	Mediterranean Sea	4 days	var. up to 100 m	WWMIII	BOLAM 8.3 km, MOLOCH 1.25 km
CNMCA / National Research Council - IT	Nettuno <sup>2</sup>	Mediterranean Sea	3 days	4.5 km	WAM	COSMO-ME 5km
National Research Council - IT	Henetus <sup>3</sup>	Adriatic Sea	5 days	1/12 deg	WAM	ECMWF 1/8 deg
Ag. for Env. Protection and Energy ER - IT	SWAN <sup>4</sup>	Med., Adriatic Sea	3 days	25 km, 8 km	SWAN	COSMO-5M 5 km
Ag. for Env. Protection and Energy ER - IT	Adriac	Adriatic Sea	3 days	1 km	SWAN	COSMO-2I 2.2 km, 5M 5km
Inst. for Env. Protection and Research - IT	SIMM <sup>5</sup>	Med., Adriatic Sea	3.5 days	1/30; 1/240 deg	WAM, SWAN	BOLAM 11 km
Slovenian Environment Agency - SL	SMMO	Central Med. Sea	3 days	1/60 deg	WAM	ALADIN 4.4 km
Met. and Hydrol. Serv. - HR	WWM <sup>6</sup>	Adriatic Sea	3 days	var. up to 10 m	WWM	ALADIN 8 km
HCMR - GR	MED-Waves <sup>7*</sup>	Mediterranean Sea	10 days	1/24 deg	WAM	ECMWF 1/8 deg

<sup>1</sup>Ferrarin et al. (2013); <sup>2</sup>Bertotti et al. (2013); <sup>3</sup>Bertotti et al. (2011); <sup>4</sup>Valentini et al. (2007); <sup>5</sup>Mariani et al. (2015); <sup>6</sup>Dutour Sikirić et al. (2018); <sup>7</sup>Zacharioudaki et al. (2015);

\*retrieved from CMEMS, <http://marine.copernicus.eu/>.

**Table A1.** Details of the sea level monitoring stations (yellow dots in Fig. 1).

Managing authority - Country	Station name	Longitude °E	Latitude °N
City of Venice - IT	Diga Sud Lido	12.43	45.42
City of Venice - IT	Diga Nord Malamocco	12.34	45.33
City of Venice - IT	Diga Sud Chioggia	12.31	45.23
City of Venice - IT	Punta della Salute CG	12.34	45.43
City of Venice - IT	Laguna Nord Saline	12.47	45.50
City of Venice - IT	Misericordia	12.34	45.45
City of Venice - IT	Burano	12.42	45.48
City of Venice - IT	Malamocco Porto	12.29	45.34
City of Venice - IT	Chioggia Porto	12.28	45.23
City of Venice - IT	Chioggia Vigo	12.28	45.22
City of Venice - IT	Fusina	12.26	45.41
City of Venice - IT	Punta Salute Giudecca	12.34	45.43
National Research Council - IT	PTF Acqua Alta	12.51	45.31
National Research Council - IT	Meda Abate	12.78	45.25
National Research Council - IT	Senigallia	13.20	43.75
Ag. for Env. Protection and Energy ER - IT	Porto Garibaldi	12.25	44.68
Ag. for Env. Protection and Energy ER - IT	Volano	12.27	44.80
Ag. for Env. Protection and Energy ER - IT	Faro	12.40	44.79
Inst. for Env. Protection and Research - IT	Trieste	13.76	45.65
Inst. for Env. Protection and Research - IT	Ancona	13.51	43.62
Inst. for Env. Protection and Research - IT	San Benedetto del T.	13.89	42.96
Inst. for Env. Protection and Research - IT	Vieste	16.18	41.89
Inst. for Env. Protection and Research - IT	Otranto	18.50	40.15
Inst. for Env. Protection and Research - IT	Crotone	17.14	39.08
Slovenian Environment Agency - SL	Koper	13.72	45.55
Hydrographic Institute - HR	Rovinj	13.63	45.08
Hydrographic Institute - HR	Dubrovnik	18.07	42.67
Institute of Oceanography and Fisheries - HR	Kaštelanskizaljev	16.39	43.51
Institute of Oceanography and Fisheries - HR	Vela Luka*	16.71	42.96
Institute of Oceanography and Fisheries - HR	Starigrad*	16.60	43.18
Institute of Oceanography and Fisheries - HR	Sobra*	17.60	42.74
Institute of GeoSciences - AL	Vlore Triport	19.39	40.51
Institute of GeoSciences - AL	Durres	19.45	41.30
Institute of GeoSciences - AL	Vlore	19.48	40.45
Institute of GeoSciences - AL	Sarande	20.00	39.87
Institute of GeoSciences - AL	Shengjin	19.59	41.81

\* Available through <http://www.ioc-sealevelmonitoring.org/>.

**Table A2.** [Details of the wave monitoring stations \(red stars in Fig. 1\).](#)

Managing authority - Country	Station name	Longitude °E	Latitude °N
City of Venice - IT	Misericordia	12.34	45.45
National Research Council - IT	Senigallia	13.20	43.75
National Research Council - IT	PTF Acqua Alta	12.51	45.31
Ag. for Env. Protection and Energy ER - IT	Nausicaa	12.48	44.22
National Institute of Biology - SL	Vida	13.55	45.55
Slovenian Environment Agency - SL	Zora	13.67	45.60
Slovenian Environment Agency - SL	Zarja	13.54	45.60
Hydrographic Institute - HR	Rovinj	13.52	45.07
Hydrographic Institute - HR	Dubrovnik	17.96	42.65