

**10<sup>th</sup> April, 2020.**

Resubmission of manuscript "Summer variation of the UTCI index and Heat Waves in Serbia", **nhess-2019-270**

Dear Editor,

Thank you very much for your kind consideration of this resubmitted version of our manuscript. Please find enclosed our revised manuscript, entitled "Analysis of UTCI index during heat waves in Serbia" by Milica Pecelj, Milica Lukić, Dejan Filipović, Branko Protić, Uroš Bogdanović, for publication consideration in the Natural Hazards and Earth System Sciences (NHES).

We value the comments received greatly, as they have pointed out a number of issues to be addressed. We would like to thank the editor and the reviewers for taking time in reading and suggesting modifications to the paper. We highly appreciate it, as the comments have been very useful to improve the paper.

We did significant modifications to the initial manuscript based on the suggestions of the reviewers. The modification has been done in all section in the manuscript:

- We have completely modified the Introduction, where we have emphasized and explained the concept of heat budget index more clearly. We have supplemented the introduction with a new text outlining the concept of heat budget indices and their importance for bioclimatic analysis. We modified the paragraphs followed by detailed recent studies (lines 47-58; lines 63-74; lines 77-94) and moved it into the Discussion section (lines 471-484; lines 438-454; lines 455-461).
- To provide a clearer introduction we have inserted a new subheading "Study area"
- The section "Materials and methods" is improved with two subheadings about detailed explanation of UTCI index and indices considered in the study.
- The previous chapter "Results and discussion" are divided into two different ones, "Results" with inserted three subheadings and "Discussion".

It should be emphasized that in order to improve the initial manuscript, it has been included one more co-author whose contribution was related to the interpretation of the results and the improvement of the figures. To address the comments 5 and 6 from the reviewer RC1 regarding Results and Figures, we conducted more detail analysis between very strong heat stress category with tmax and interpreted all results with improved figures. This was executed by Uroš Bogdanović. Therefore, in the revised manuscript we have decided to include him as a co-author of the paper. This inclusion was discussed with and has received consent from all the co-authors of the paper.

For ease to follow the changes in revised manuscript, we uploaded another revised manuscript, where the modified text is colored green and the new added text is colored blue. Hopefully this will be useful for the new review.

We hope that the editor will find the paper suitable for publication.

Thank you again for consideration.

Yours sincerely,

Milica Pecelj (On behalf of the authors of the manuscript)

Please find enclosed

1. Response to the reviewer 1 (RC1)
2. Response to the reviewer 2 (RC2)
3. Marked up manuscript version

18<sup>th</sup> February 2020

Response to peer reviewer comments

nhess-2019-270: **Summer variation of the UTCI index and Heat Waves in Serbia**

Milica M. Pecelj, Milica Z. Lukić, Dejan J. Filipović, Branko M. Protić

Dear Reviewer,

We would like to thank you for giving us the opportunity to submit a revised draft of our manuscript titled Summer variation of the UTCI index and Heat Waves in Serbia to Journal of Natural Hazard and Earth System Science. We appreciate the time and effort that you dedicated to providing valuable feedback on our manuscript.

We have been able to incorporate changes to reflect most of the suggestions. Since you emphasized major revisions to the manuscript, significant changes have been made to the paper. The revision, based on your guidance, includes changed interpretation of research in all chapters.

Here are the responses to yours comments and concerns, listed step by step. To make the changes in the manuscript easier to follow, we have highlighted the modified text in **green** and the new added text in **blue**.

#### **Comment 1: Title**

*To me the title could be: "Analysis of UTCI index during heat waves in Serbia"*

**Author response:** Thank you. We agree with this suggestion and have made changes, so the revised title reads exactly as suggested: **Analysis of UTCI index during heat waves in Serbia**

#### **Comment 2: General comment**

*This work could potential be interesting however improvement of the paper is need previous to publication in order to reach the expected international standards requested by the journal. First of all, according to the elements presented in the introduction, many similar studies have already been conducted in Serbia. Within the current study, a more recent meteorological data sets are used to compute the UTCI index values for the selected stations. Starting before 1998 would have been even more interesting to identify the trend of such extreme heat waves.*

**Author response:** Thank you for pointing this out. We agree with the reviewer's statement however, in the case of our study, it seems slightly out of scope because the procedure for obtaining daily hourly data from the national weather center is very slow, so it would be impossible to respect the time limit for correction. It would certainly be interesting to explore this aspect for future research. Since many studies have been conducted in Serbia concerning the analysis of heat waves in the last 50 years, we considered their results relevant as they indicate an increase in heat waves, especially since 2000. Starting from this fact, we used the data available only from 1999 to 2017 and for this period we did a bioclimatic identification of thermal discomfort. We changed a part in introduction to emphasize this point.

The revised text reads as follows on:

(lines 77-81) **Considering extreme temperature is one of the most significant climatic parameters in the universal context of climate change, analyses of heat waves in Serbia have been performed in several different approaches and all of them show a growing trend in max temperature (Drljača et al., 2009; Unkašević and Tošić, 2009a; Unkašević and Tošić, 2009b; Unkašević and Tošić, 2013; Malinović-Milićević et al., 2013) and heat waves (Unkašević and Tošić, 2011; Unkašević and Tošić, 2015 Milićević-Malinović et al., 2016) especially since 2000.**

**Keeping this fact in mind, the basic idea of this study is to conduct the analysis of human bioclimatic conditions in Serbia in the last 20 years, where according to the previous studies mentioned above, warming has been perceived and recorded, especially since 2000. Determining extreme weather event from the aspect of heat budget indices**

allows other meteorological parameters to be taken into account because high temperatures and humidity generate a heat load more rapidly, unless the wind and direct radiation are taken into account.

(lines 89-93) The conducted human bioclimatic evaluation of UTCI thermal stress in Serbia was considered to be of great importance due to the identification of biothermal heat hazard and the study of the evolution of such indices regarding climate change. This evaluation is aimed at addressing the following topics in this study:

- providing a comprehensive assessment of the human heat stress associated with UTCI index and
- comparing it to heat waves defined by meteorological parameters.

### **Comment 3: Abstract/Introduction**

*The abstract is concise and seems complete. However, the introduction needs to be improved. The first part introduces the UTCI index, then heat waves in Serbia are presented, followed by recent studies (line 75 to 94). This final part of the introduction is difficult to understand. It is better to rewrite it and introduce that there is a need to study the evolution of such indices regarding climate change, as trend has been identified during the period 1999-2007.*

**Author response:** Thank you for pointing this out. We agree with this comment. Therefore, in relation to your remarks, we have completely modified the introduction, where we have emphasized and explained the concept of heat budget index more clearly. We have supplemented the introduction with a new text outlining the concept of heat budget indices and their importance for bioclimatic analysis (lines 35-52).

The revised text reads as follows on:

(lines 32-38) Over the last century many models and indices have been developed for the assessment of human exposure to heat, ranging from simple physical instruments designed to imitate the human heat exchange with the environment, to complex thermophysiological models that simulate external and internal body heat transfer and allow detailed simulation of different work load, clothing, and climate scenarios (Havenith and Fiala, 2015).

(lines 39-52) Many thermal indices have been developed for the purpose of describing the complex conditions of heat exchange between the human body and its thermal environment (Fanger, 1970; Landsberg, 1972; Parsons, 2003). Those are two meteorological parameter indices – the ones used for cold environment conditions combining the air temperature and the wind velocity (Osczevski and Bluestein, 2005; Siple and Passel, 1945;), and the ones used for heat environment conditions combining the air temperature and humidity (Masterton and Richardson, 1979; Steadman, 1984; Yaglou and Minard, 1957). Considering their shortcomings, i.e. the fact that they are not universally applicable to all climatic regions, including different seasons, the dominance in the analysis of biothermal conditions and thermal stress over the last 30 years has been taken over by so-called heat budget indices (Blazejczyk, 1994; Hoppe, 1999; Jendritzky, 2012). The heat budget indices are based on the thermal exchange between humans and the environment. Methodologically, they use variables related to meteorological (air temperature, wind velocity, radiation, air humidity) and physiological processes (most commonly metabolic heat) and clothing insulation. The issue that should be emphasized during the analysis of the thermal stress is the influence that extreme temperatures have on the physiological parameters in humans. According to McGregor and Vanos (2018), generated heat load can undermine the human body's ability to hold its core temperature within the range of optimal physiological achievement.

Then, we modified the paragraph followed by detailed recent studies (lines 47-58; lines 63-74; lines 77-94) and moved it into the Discussion section (lines 471-484; lines 438-454; lines 455-461).

More specifically, lines 47-58 have been moved into the discussion lines 471-484, where a reference about heat wave related mortality in 2007 by Bogdanović et al. (2013) was added (lines 485-498).

The revised text reads as follows on:

(lines 471-484) Identified biothermal discomfort in July and August as the hottest summer months in Serbia, could provoke the health disorders, more frequent. Specific characteristics of HWs such as intensity and duration has been

found to have devastating impacts on human health and wellbeing. In certain cases, heat waves cause problems, especially to children, elderly people, chronic patients and workers. Analysis of the UTCI index correlated with mortality in Europe shows the deaths occur in the southern Europe (Italy, Portugal, Greece, Spain) for the categories of moderate and strong stress. There is an impact of heat load in increasing mortality, as conditions become more thermally stressful (Di Napoli et al., 2018). The intensity of heat stress in Poland was assessed by the UTCI index based on daily mortality and weather data for 1991–2000. There is increase in mortality for the days with strong and very strong heat stress, in relation to no thermal stress days, of 12% and 47%, respectively (Błażejczyk et al., 2017). Biothermal conditions and mortality rates show that during summer in southern Europe thresholds of Physiological Subjective Temperature (PST) increase when moving from west to east. The general finding is that population of central Europe is more sensitive for thermal stimuli in summer than citizens of the south of continent (Błażejczyk and McGregor, 2008). The statement about the PST index is in support with the comparison of UTCI index with other thermal indices, including PST index, which is very well correlated with UTCI index (Błażejczyk et al., 2012).

(lines 485-498) Similarly, Bogdanović et al. (2013) reported a significant short-term excess mortality on 16<sup>th</sup> July in Belgrade, when the maximum daily temperature exceeded 35°C, leading to 167 excess deaths (38% increase compared to the number of expected deaths) for nine consecutive days of heat, with a progressive return to almost normal mortality as the maximum temperature dropped below 35°C on 25<sup>th</sup> July.

In addition, lines 63-74 have been moved into the discussion lines 438-454, and there is an explanation of what a tropical day is (line 456-457) in the analysis by Unkasevic and Tosic (2009b).

The revised text reads as follows on:

(lines 438-454) This agrees with findings of Unkašević and Tošić (2011, 2015) that trend of heat waves in Serbia is increase, especially since 2000. In order to justify present research for the last 20 years, the earlier results related to the increase in heat waves will be discussed in more detail. In that regard, previous research of heat waves in Niš for 1949-2007, based on the autoregressive-moving-average model, show observed the warmest years during three periods, in 1951-1952, 1987-1998 and 2000-2007 (Unkašević and Tošić, 2009b). According to this research, the longest heat wave was recorded in 1952 in Niš with 21 days while during 2003 in was observed 29 consecutive tropical days. It should be considered that according to the Serbian National Weather Service, tropical days are days with a maximum temperature over 30°C. Further, based on Heat Wave Duration index (HWD) i.e. daily maximum values of air temperature, Drljača et al. (2009) determined the length and strength of heat waves in Niš during the summer season. The analysis showed Niš with the greater number of heat waves compared to Belgrade larger urban area. As stated in research, since the mid-1980s, heat waves have a higher frequency and occur on average every year. Prior to that period heat wave fluctuations have generally been reported in one of two years (Niš during summer) (Drljača et al., 2009). An increase of the heat wave duration, in addition to increase of the heat wave frequency of occurrence was detected by Unkašević and Tošić (2011) during the period 1999 – 2007, studying the characteristics of the heat waves in central Serbia (1949 – 2007).

Lastly, lines 77-94 have been moved into the discussion section (lines 455-461). The revised text reads as follows:

(lines 455-461) As regards other heat budget index of Heat Load (HL) in Serbia (July) for 2000-2010 is noted that in, Niš, Zlatibor and Novi Sad expired “extremely hot” in 2007 and 2000 (Milovanović et al., 2017). Referring to heat budget index Physiological Equivalent Temperature PET for the period 1949–2012, the highest number of heat waves in Novi Sad was observed in the last two decades and in the first decade of the investigated period (Basarin et al., 2016) while the number of the days above particular thresholds for 1961-2014, show increase along with a number of heat waves per year since 1981. The highest values of mean annual PET are mostly detected during the last 15 years. (Basarin et al., 2018).


To provide a clearer introduction we have inserted a new text. Namely, we added a heading “Study area” where we provided a relief map of Serbia, inside Europe, with the location of the three stations studied in the paper (lines 95-105). Basic information about the climate of Serbia has been added. Thus, we switched the first paragraph from the Materials and methods section (lines 100-115), which in the revised version takes up lines 106-120.

The revised text reads as follows:

(lines 95-105) Study area

The Republic of Serbia is located in south-eastern Europe, in the area of the southern Pannonian Plain and the central Balkans. Northern Serbia is mainly flat, while its central and southern areas consist of highlands and mountains (Gocić and Trajković, 2014), as the map of the relief characteristics shows (Figure 1). The territory of Serbia is characterised by temperate continental and mountain climate and the spatial distribution of climate parameters is determined by geographical location. The research involved studying three synoptic stations located in different geographical areas of Serbia: (i) Novi Sad, (ii) Niš and (iii) Mt. Zlatibor (Figure 1).

(lines 106-120) The first weather station (at an altitude of 86 m) is located in the territory of Novi Sad, which is the administrative urban center of Vojvodina province and South Bačka District (Figure 1<sub>1</sub>). The city is located in the southern part of the Pannonian Basin, on the Danube River bank near Mt. Fruška Gora, and the national park bearing the same name. Novi Sad has a temperate continental climate, summers are warm and winters are cold with a small amount of snow (Lazić et al., 2006). The second station is Niš (at an altitude of 202 m) and it is located in the Niš Fortress. This city is the administrative urban center of the Nišava District in southern Serbia and it is situated in the Nišava valley, located in the central part of the spacious geological depression called the Nišava Basin (Figure 1<sub>2</sub>). According to Köppen's climate classification, the Nišava valley belongs to the Cfwax type – the Danube type of moderately warm and humid climate characterized by hot summers (the highest precipitation is recorded at the beginning of the summer) and somewhat dry winters (Prokić, 2018). Compared to other cities in this valley (Dimitrovgrad, Pirot and Bela Palanka), Niš is the hottest one with an average annual temperature of 11.8°C (Prokić, 2018). The third station is Zlatibor, a mountain in western Serbia, which belongs to the mountain range of the Dinaric Alps (Figure 1<sub>3</sub>). Zlatibor weather station lies at an altitude of 1029 m a.s.l. In the area of Zlatibor plateau mountains meet air currents from the Adriatic Sea, which, as it can be assumed, creates a favorable climate and for this reason Zlatibor has already been confirmed as a climatic resort (Pecelj et al., 2017). The mountain climate of Zlatibor involves long and cold winters, short and fresh summers, and less pronounced spring and autumn.



**Figure 1.** Relief map of Serbia with the areas studied: 1-Novi Sad, 2-Niš, 3-Zlatibor and a map of geographical location of Serbia in Europe

Maps were created using QGIS 3.8 software on the basis of the official European Commission's (Eurostat) datasets, available at <https://ec.europa.eu/eurostat/web/gisco/geodata/reference-data/>, (Map ratio: 1:1900000; Map projection: WGS 84/UTM, Zone 34N, the official national coordinate system)

#### Comment 4: Materials and methods

*Materials and methods: The model used to compute the index is not explained and presented. The reader does not know how the index are computed and therefore cannot judge the methodology and evaluate the results. Therefore, within the material and method section, it is needed to add the equations of the model. Moreover, to me a validation of the model should be introduced in the section, this will give more weight to the results, moreover this will help the reader to understand the indices and also help the writer to introduce the result and limits to this method. To catch the reader attention, it is necessary to present climate in Serbia earlier in the paper. For example, adding a map of Serbia who presents the location of the three stations studied and the cities presents in the paper. Information concerning the weather in Serbia should also be added: most severe droughts, their consequences on the populations...*

**Author response:** We have, accordingly, a modified description of the model used to compute the index and we have made changes, clarifying the index analysis methodology. More specifically, we have divided the Materials and methods into two sections. In the beginning, an overview of meteorological and physiological components is given (lines 122-131), where we describe in more detail the metabolic rate as a key physiological parameter. The revised text reads as follows:

(lines 122-131) The present study implements the methodological approach of Universal Thermal Climate Index (UTCI) based on human heat budget model relying on the evaluation of human energy balance. As a thermal comfort indicator, the UTCI considers combined meteorological and physiological parameters describing thermal comfort through the evaluation of human energy balance. In terms of physiological conditions, the metabolic rate plays the most important role. Metabolic processes in the human body create heat that is constantly exchanged with the environment achieving a state of thermal equilibrium in the body for maintaining a constant body temperature. The amount of the heat that is produced and released depends on the following: physical activity, clothing, sex, age, body mass, diet, mental state, health, external conditions, acclimatization, etc. As a measure of physical activity, a unit of "MET" was defined, which corresponds to the release of human heat of  $58.2 \text{ W}\cdot\text{m}^{-2}$  for average body surface area ( $1.8 \text{ m}^2$ ), i.e. it is equal to the energy rate produced per unit surface area of an average person seated at rest (ANSI/ASHRAE Standard 55). According to ISO 8996 for standard applications, the metabolic heat energy is  $M = 135 \text{ W}\cdot\text{m}^{-2}$  i.e. 2.3 MET, for a person moving at a speed of  $1.1 \text{ m}\cdot\text{s}^{-1}$ .

The first section deals with the Universal Thermal Climate Index (UTCI), where we provide detailed definition, parameter inputs, and their limitations including stress classification (lines 135-142; lines 144-165). The revised text reads as follows:

(lines 133-140) The Universal Thermal Climate Index (UTCI)

For valid assessment of the outdoor thermal environment in the fields of public weather services, public health systems, urban planning, tourism and recreation and climate impact research, the Universal Thermal Climate Index (UTCI) has been developed based upon the most recent scientific progress in human thermo-physiology in biophysics and heat exchange theory. The UTCI is the result of an approach which was developed in the International Society of Biometeorology (ISB) Commission 6 and was later improved by COST Action 730 (Jendritzky et al., 2012). The design of UTCI is of great importance due to the fact that it is applicable to all seasons and climates together with all spatial and temporal scales.

(lines 144-165) In other words, this model simulates the same sweat production or skin wettedness in human body response as the actual environment condition (Błażejczyk et al., 2013; Błażejczyk et al., 2014). This is derived from the multi-dimensional dynamic response of a state-of-the-art multi-node thermo-physiological model of human heat transfer and thermoregulation (Fiala et al., 2012). The UTCI can be represented in general function as bellow:

$$\text{UTCI} = f(t, f, v, t_{\text{mrt}}) \quad (\text{Eq.1})$$

UTCI = f (air temperature, relative humidity, wind speed, mean radiant temperature)

For appropriate use, UTCI can only be approximated using a regression equation abbreviated from sample calculations performed by computing centers (Bröde et al., 2012; Jendritzky et al., 2012). It causes a narrow range of input parameters it can manage with. For the reference environment, the ISB Commission on the UTCI, was decided to use: (i) a wind speed ( $v$ ) of  $=0.5 \text{ m}\cdot\text{s}^{-1}$  at 10 m height (approximately  $=0.3 \text{ m}\cdot\text{s}^{-1}$  at 1.1 m), (ii) a mean radiant temperature ( $T_{\text{mrt}}$ ) equal to air temperature, (iii) vapour pressure ( $vp$ ) that represents a relative humidity ( $f$ ) of 50%; at high air temperatures ( $>29^\circ\text{C}$ ) the reference humidity was taken to be constant at 20 hPa (Błażejczyk et al., 2013). Physiological parameters, the metabolic rate ( $M$ ) and the thermal properties of clothing (clothing insulation, permeability) are taken as universal constants in the model due to the evaluation by means of the regression equation. This implies an outdoor activity of a person walking at the speed of  $4 \text{ km}\cdot\text{h}^{-1}$  ( $1.1 \text{ m}\cdot\text{s}^{-1}$ ), corresponding to heat production of  $135 \text{ W}\cdot\text{m}^{-2}$  (2.3 MET) of metabolic energy (Błażejczyk et al., 2013; Jendritzky et al., 2012) and the clothing insulation which is self-adapting according to the environmental conditions (Havenith et al., 2012). Clothing insulation, vapor resistance and the insulation of surface air layers, are strongly influenced by changes in wind speed and body movement and will therefore also influence physiological responses (Błażejczyk et al., 2012). Particular ranges of UTCI are categorised according to thermal stress (Table 1).

In order to save the lines, we have modified Table 1.

**Table 1.** UTCI thermal stress classification

UTCI (°C)	> 46	38 to 46	32 to 38	26 to 32	9 to 26	0 to 9	0 to -13	-13 to -27	-27 to -40	< -40
Stress	extreme heat stress	very strong heat stress	strong heat stress	moderate heat stress	no thermal stress	slight cold stress	moderate cold stress	strong cold stress	very strong cold stress	extreme cold stress
Abbr.	EHS	VSHS	SHS	MHS	NTS	SLCS	MCS	SCS	VSCS	ECS

Source: Błażejczyk et al. 2013

The second section explains the kind of analysis which was conducted in the study (lines 170-205). The revised text reads as follows:

(lines 167-205) Data and indices considered in the study

The meteorological data set from the period from 1998 to 2017 was recorded for two urban weather stations (Novi Sad and Niš) and one rural mountain station (Mt. Zlatibor). Mean daily and hourly meteorological parameters (07h, 14h, CET) of air temperature ( $t$ ), air humidity ( $f$ ), and wind speed ( $v$ ) from the above weather stations were considered for the calculation of particular UTCI thermal heat stress in the summer months (July, August and September). The meteorological data set used in the study was retrieved from the Meteorological Yearbook for the period from 1998 to 2017 (Republic Hydrometeorological Service of Serbia) while the UTCI index was calculated by applying the BioKlima 2.6 software package (available at <http://www.igipz.pan.pl/Bioklima-zgik.html>).

Although extreme temperatures are one of the most effective climatic parameters in the universal case of hot days and heat waves, the influence of other parameters, especially clothing insulation and metabolic rate, can justify the thermal discomfort and the other way round. In that regard, the summer (July, August and September) daily maximum and minimum temperatures ( $t_{\text{max}}$ ,  $t_{\text{min}}$ ), for the same stations, were analysed in terms of particular thresholds to identify the frequency of extremely high temperatures and heat waves in the period observed to compare them with the discomfort recorded by the UTCI index.

The first stage of the study presented mean daily ( $\text{UTCI}_{\text{avg}}$ ), morning ( $\text{UTCI}_{7\text{h}}$ ) and midday ( $\text{UTCI}_{14\text{h}}$ ) UTCI indices. They are presented by months (July, Aug, Sep) for each station to identify the general differences caused by the geographical origin of the stations under study. The quantification of human bioclimatic conditions in Serbia was designed for the UTCI indices on a daily basis of each defined stress categories. To obtain a better insight into the  $\text{UTCI}_{14\text{h}}$  index values during the summer months, averaged monthly  $\text{UTCI}_{14\text{h}}$  index was provided for each year to see how much the  $\text{UTCI}_{14\text{h}}$  index fluctuated in relation to the average monthly value.



For the second stage, midday UTCI thermal stress indices were identified (strong heat stress SHS, very strong heat stress VSHS, extreme heat stress EHS) and thermal indices based on the maximum and minimum temperatures (hot day HD and hot night HN). During the midday observation (14 CET), particularly prominent are strong heat stress (SHS), which refers to UTCI range of 32°C to 38°C and very strong heat stress (VSHS), which refers to UTCI range of 38°C to 46°C (Table 2). The occurrence of thermal stress days was presented for each month (July, August and September) in the last 20 years. Extreme heat stress (EHS), which refers to UTCI range above 46°C was presented separately when it occurred. Thermal indices, hot day (HD) and hot night (HN), were identified on the basis of the threshold of the maximum and minimum temperatures. Considering the influence of extreme temperatures, as stated by Collins et al. (2000) the following indices were marked in relation to the thresholds of maximum and minimum temperature: Hot days ( $HD=t_{max} \geq 35^{\circ}C$ ) and Hot nights ( $HN=t_{min} \geq 20^{\circ}C$ ) (Table 2).

For the last stage of the study, for the purpose of identifying thermal discomfort during the period observed, as an illustration of a threshold based on the duration of hot days (HD) and hot nights (HN) a hot day event is determined ( $HWE = 3$  HD days of  $t_{max} \geq 35^{\circ}C$ ). In addition, as an illustration of a threshold based on the duration of very strong heat stress (VSHS), a very strong heat stress event (VSHSE) is determined and it is caused by the occurrence of 5 consecutive VSHS days (Table 2).

As a result of comparing UTCI thermal stress to the selected thresholds of maximum and minimum temperatures, biothermal discomfort was identified during extremely high temperatures for the geographical area of Serbia. The recorded number of days with VSHS stress was compared to the number of days when the maximum air temperature was above 35°C.

For clearer insight into the methodology, Table 2 has been modified and follows the revised text.

**Table 2.** Definition of indices used in the study

Abbreviations	UTCI Indices	Definition
UTCI <sub>avg</sub>	Universal Thermal Climate Index	$UTCI_{avg} = f(t, f, v, t_{mrt})$
UTCI <sub>07h</sub>	Universal Thermal Climate Index <sub>07h</sub>	$UTCI_{07h} = f(t_{7h}, f_{7h}, v_{7h}, t_{mrt})$
UTCI <sub>14h</sub>	Universal Thermal Climate Index <sub>14h</sub>	$UTCI_{14h} = f(t_{14h}, f_{14h}, v_{14h}, t_{mrt})$
Abbreviations	UTCI heat stress indices	Definition
SHS	Strong Heat Stress	SHS= UTCI (32°C to 38°C)
VSHS	Very Strong Heat Stress	VSHS = UTCI (38°C to 46°C)
VSHSE	Very Strong Heat Stress Event	minimum 5 VSHS days in a row
EHS	Extreme Strong Stress	EHS= UTCI >46°C
Abbreviations	Thermal indices	Definition
HD	Hot days	$t_{max} \geq 35^{\circ}C$
HN	Hot nights	$t_{min} \geq 20^{\circ}C$
HWE	Heat waves event	minimum 3 HD days in a row

t-air temperature; f-relative humidity; v-wind speed; t<sub>mrt</sub>-mean radiant temperature; t<sub>max</sub>-maximum air temperature, t<sub>min</sub>-minimum air temperature

Information regarding the most severe droughts in Serbia is provided in the Results section (lines 233-236) and lines 241-243). The revised text reads as follows:

(lines 233-236) This is confirmed by Tošić and Unkasević (2014) in the study of dry periods in Serbia during the period between 1949 and 2011. It was found that the frequency of droughts in the southern part of Serbia was higher than in the other parts of the country. The most severe drought occurred in Niš and lasted from July 2006 to March 2008 (with a duration of 21 months and severity of 63.0).

(lines lines 241-243) Nevertheless, this does not rule out the occurrence of high temperatures and severe dry periods during the warm part of the year. This idea is based on the findings of Leščešen et al. (2018), when they analyzed

drought periods in Vojvodina for over 60 years (1956-2016). All the regions of Vojvodina that were investigated experienced at least one extreme drought event over the reference period, in particular in 2000, 2001-2002, and 2011-2012. Moreover, the same results were obtained in other similar studies (Gocic and Trajković, 2013; 2014; Milanović, et al., 2014).

#### **Comment 5: Results**

*The result section is a numeration of the results. This section is the most important of the paper, it is important to bring more analysis to the presented results. Results of the analysis also need to be more accurate and explained. They also need to be more synthetic. For example, as the paper objective is to rate occurrence of thermal stress a trend of the UTCI indices could be computed. Correlation between the different indices obtained could be computed the complete the analysis. Presenting a figure showing the intensity of UTCI index values may help to improve this section and help to reach substantial conclusions.*

**Author response:** We have divided the section Results and discussion in two separate sections. According to the 3 stages of the study, described in revised Methodology (Data and indices considered in the study), the results are presented in three sections: (i) UTCI indices ( $UTCI_{avg}$ ,  $UTCI_{7h}$ ,  $UTCI_{14h}$ ), (ii) UTCI heat stress indices (strong heat stress SHS, very strong heat stress VSHS, extreme heat stress EHS) and thermal indices (hot day HD, hot night HN), (ii) Heat waves (HWE and VSHSE).

In all three sections the results are explained in more detail along with the modified and new added figures. Alongside the trend of UTCI index, the correlation between UTCI and maximum air temperature, is provided, too.

The revised text reads as follows:


(lines 207-215) **Results**

On the basis of the conducted bioclimatic analysis and comparison of biothermal conditions given in the two urban meteorological stations Novi Sad, Niš and Mt. Zlatibor, which represents a rural and lower mountain area, some differences in weather conditions were perceived in the summer period, as was expected. The results are presented in three sections: (i) UTCI indices ( $UTCI_{avg}$ ,  $UTCI_{7h}$ ,  $UTCI_{14h}$ ), (ii) UTCI heat stress indices (strong heat stress SHS, very strong heat stress VSHS, extreme heat stress EHS) and thermal indices (hot day HD, hot night HN), (iii) Heat waves (HWE and VSHSE).

UTCI indices ( $UTCI_{avg}$ ,  $UTCI_{7h}$ ,  $UTCI_{14h}$ )

This section presents the results of the heat budget index UTCI calculated for mean daily data ( $UTCI_{avg}$ ) morning ( $UTCI_{7h}$ ) and midday ( $UTCI_{14h}$ ) for the period of 20 years.

(lines 222-227) These are the years when significant heat waves were recorded. The results depicted in Fig. 2 show the frequency (in percentages) of all stress categories for each index ( $UTCI_{avg}$ ,  $UTCI_{7h}$ ,  $UTCI_{14h}$ ) that occurred in July, August and September for the period that was observed.



**Figure 2.** Frequency of the days with different UTCI thermal stress categories for mean daily ( $UTCI_{avg}$ ), morning ( $UTCI_{7h}$ ) and midday ( $UTCI_{14h}$ ) index during the period from July to September, 1998-2017: a) Niš, b) Novi Sad, c) Zlatibor. X-axis: time (months), y-axis: frequency (number of days)

(lines 229-230) There are 5 days observed with “extreme heat stress” for midday  $UTCI_{14h}$ .


(lines 243-254) This is confirmed by Tošić and Unkasević (2014) in the study of dry periods in Serbia during the period between 1949 and 2011. It was found that the frequency of droughts in the southern part of Serbia was higher than in the other parts of the country. The most severe drought occurred in Niš and lasted from July 2006 to March 2008 (with a duration of 21 months and severity of 63.0).

(lines 239-282) Nevertheless, this does not rule out the occurrence of high temperatures and severe dry periods during the warm part of the year. This idea is based on the findings of Leščešen et al. (2018), when they analyzed drought periods in Vojvodina for over 60 years (1956-2016). All the regions of Vojvodina that were investigated experienced at least one extreme drought event over the reference period, in particular in 2000, 2001-2002, and 2011-2012. Moreover, the same results were obtained in other similar studies (Gocić and Trajković, 2013; 2014; Milanović, et al., 2014). On the mean daily level ( $UTCI_{avg}$ ) the most frequent category is “moderate heat stress” while on the hourly level  $UTCI_{14h}$  the most frequent category is “strong heat stress”. Days with “very strong heat stress” were observed in July and August (Fig. 2b) and there was one day with “extreme heat stress” in July 2007. Mt. Zlatibor has distinctive characteristics of the sub-mountain and mountain climate. Among these three stations, the bioclimatic conditions of Mt. Zlatibor are the most pleasant ones considering the dominant category of “no thermal stress” on the daily level of  $UTCI_{avg}$ . The thermal conditions with a morning  $UTCI_{7h}$  are similar to  $UTCI_{avg}$  while the  $UTCI_{14h}$  shows the prevailing categories of “moderate heat stress” and slightly less “strong heat stress” and “very strong heat stress”. It is important to emphasize the UTCI category of “slight cold stress” for all the indices ( $UTCI_{avg}$ ,  $UTCI_{7h}$ ,  $UTCI_{14h}$ ) was recorded several times during the three months of the period which was observed. The lowest value of the UTCI index was  $-2.84^{\circ}\text{C}$  and it occurred on September 5<sup>th</sup>, 2007 in Mt. Zlatibor as “moderate cold stress” (Fig. 2c).

Of all the calculated UTCI indices, mean daily data ( $UTCI_{avg}$ ), morning data ( $UTCI_{7h}$ ) and midday ( $UTCI_{14h}$ ) data, the midday  $UTCI_{14h}$  index shows the most extreme values for identifying heat waves. In this regard, for the further data analysis in this study, the  $UTCI_{14h}$  is used. To gain better insight into the  $UTCI_{14h}$  index values during the summer months, for the investigated period, averaged monthly  $UTCI_{14h}$  index for each year was provided for all three

stations. The mean monthly values are presented in Fig. 3, together with the trends in  $UTCI_{14h}$  and mean absolute deviation of  $UTCI_{14h}$  in order to see how much the  $UTCI_{14h}$  index fluctuated in relation to the mean monthly value. The results show a significant increase in the extreme values of the  $UTCI_{14h}$  index over the last ten years, presented mainly as “very strong heat stress category” (VSHS). A series of peaks can be observed in the years that have been marked as extremely warm. In addition, there is a growing trend of the  $UTCI_{14h}$  index in all the summer months and all the weather stations, i.e. a series of peaks with successively higher values during the investigated period (1998-2017). The highest growing trend can be seen in September in Nis and Mt. Zlatibor, while the lowest growing trend can be observed in August on Zlatibor.


As regards the severity of daily UTCI index, there were 5 days recorded in Niš when the  $UTCI_{14h}$  exceeded the limit value for “extreme heat stress” (EHS). This happened on July 5<sup>th</sup>, 2000 ( $UTCI_{14h}=47.08^{\circ}C$ ); July 24<sup>th</sup>, 2007 ( $UTCI_{14h}=48.26^{\circ}C$ ); August 24<sup>th</sup>, 2007 ( $UTCI_{14h}=46.29^{\circ}C$ ); August 5<sup>th</sup> and 6<sup>th</sup>, 2017 ( $UTCI_{14h}=46.75$  and  $UTCI_{14h}=46.76^{\circ}C$ ). The maximum value of UTCI in Novi Sad was recorded on July 24<sup>th</sup>, 2007 ( $UTCI=48.42^{\circ}C$ ). In the area of Mt. Zlatibor, the “extreme heat stress” has not been recorded in the last 20 years. The highest values of UTCI in Zlatibor occurred on July 22<sup>nd</sup> and July 24<sup>th</sup>, 2007 ( $UTCI_{14h}=38.62^{\circ}C$  and  $UTCI_{14h}=38.37^{\circ}C$ ). The year 2007 is rated as the most unfavourable one, particularly the date of July 24<sup>th</sup>, when the highest temperature ever was recorded in Serbia (Smederevska Palanka,  $t_{max}=44,9^{\circ}C$ , source: Republic Hydro-meteorological Service of Serbia, RHMSS).



**Figure 3.** The mean monthly  $UTCI_{14h}$  index during the period which is investigated (1998-2017). x-axis: time (years), y-axis: UTCI ( $^{\circ}C$ )

#### Thermal indices and UTCI heat stress indices

This section identifies midday  $UTCI_{14h}$  heat stress indices (strong heat stress SHS, very strong heat stress VSHS) and thermal indices marked in relation to the thresholds of the maximum and minimum temperatures in particular (hot day HD and hot night HN). Fig. 4 presents frequency (number of days) of hot days (HD), hot nights (HN), strong heat stress (SHS) and very strong heat stress (VSHS) for each month (July, August and September) in the last 20 years. The results of UTCI that were obtained show compatibility with previous analyses in relation to the occurrence of heat waves, especially since 2000 (Basarin et al., 2018; Unkašević and Tošić, 2009a; Unkašević and Tošić, 2009b; Unkašević and Tošić, 2013).



(lines 293-340) **Figure 4.** Number of hot days (HD), hot nights (HN), strong heat stress (SHS) and very strong heat stress (VSHS) in Niš, Novi Sad and Zlatibor (July, Aug, Sep) 1998-2017. x-axis: time (years), y-axis: number of days

For the period investigated (1998-2017), the total number of hot days recorded in July, August and September in Niš was 126, 133 and 21 HDs, respectively, while Novi Sad saw about a half of the said number of hot days, namely 52, 63 and 3 HDs, respectively. In Mt. Zlatibor 2 HD days were recorded in July.

As regards hot nights (HN), there is generally a lower intensity of HNs compared to HDs. Generally, during the observed period up to 4 HNs were recorded, although an increase in the HNs has been observed in the last ten years. For example, in Niš during 2012, 2015 and 2017 the number of HNs recorded in July was 12, 7 and 7 respectively and during 2000, 2013, 2015, and 2017 the number of HNs recorded in August was 8, 6, 5 and 5 respectively. Since 2010 up to 2 HNs have been recorded in September. In Novi Sad, during 2010, 2012, 2015 and 2017 there were 7, 5, 5 and 6 HNs respectively recorded in July, while in August during 2012 and 2017 there were 5 and 7 HNs respectively. In September, one HN was recorded during 2009 and 2011 (Figure 4). The total number of hot nights recorded in July, August and September for the investigated period in Niš was 60, 56 and 6 HNs, respectively, while in Novi Sad there was a significantly lower number of hot nights – 41, 38 and 2 HNs, respectively. In Mt. Zlatibor 9 HN days were recorded in July and 11 HN days were recorded in August for the period which was investigated.

Midday  $UTCI_{14h}$  “strong heat stress” (SHS) occurs most commonly in all three stations. For example, SHS is particularly important to Mt. Zlatibor, located at an altitude above 1000 m since there has been an increase in the number of days with SHS in Zlatibor over the last decade. The best indicator is the increase in SHS in September. Similarly, the indicator of biothermal discomfort, “very strong heat stress” (VSHS) has been occurring more frequently in the last decade, reaching a maximum of 17 days in July 2007 and 2012 and in August 2015 in Niš. The number of VSHS days in Novi Sad reached a maximum of 12 days in July 2015 and 14 days in August 2015 and 2017. The total number of VSHS days recorded in July, August and September for the investigated period in Niš was 167, 174 and 25 VSHS days, respectively, while in Novi Sad there was a significantly lower number of VSHS days – 83, 97 and 7 VSHS days, respectively. In Mt. Zlatibor 2 VSHS days were recorded in July and 3 VSHS days were recorded in August. The recorded numbers of VSHS days indicate slightly greater biothermal discomfort duration in relation to the number of HD days. The HD and VSHS indices are significantly correlated during the summer months as they are directly derived from the maximum air temperatures and  $UTCI_{14h}$  index (Table 3).  $UTCI_{14h}$  and  $t_{max}$  for the Niš and Novi Sad urban stations have a significantly high linear correlation.

Table 3. Linear correlation of  $UTCI_{14h}$  and  $t_{max}$

UTCI <sub>14h</sub> ~ t <sub>max</sub>	Niš	Novi Sad	Zlatibor
July	0.88	0.97	0.85
August	0.92	0.95	0.77
September	0.89	0.92	0.41

For Zlatibor the correlation is slightly lower, and in September it is weak compared to the urban stations, but neither HD nor VSHS indices occurred on Zlatibor. Similar results of the linear relationship between the air temperature and the UTCI index ( $r = 83$ ;  $r = 0.98$ ) were shown in the studies of Urban and Kysely (2014) for summer in the urban areas of South Bohemia in Czech Republic and in the studies of Nassiri et al. (2017) for Iran.

#### Heat waves (HWE and VSHSE)

After UTCI thermal stress was compared to the selected thresholds of maximum temperature, biothermal discomfort was identified during extremely high temperatures for the geographical area of Serbia. During the observed period, a heat wave event (HWE) was determined, caused by 3 consecutive HD days and very strong heat stress event (VSHSE), caused by the occurrence of 5 consecutive VSHS days.

Table 4 lists the identified heat waves that refer to UTCI<sub>14h</sub> VSHS index (VSHSE) and HD index (HWE). It has a very similar layout. However, shadowed differences can be observed, especially in the duration of the heat wave event. The results are presented for Niš and Novi Sad since there were no recorded heat waves in Mt. Zlatibor. More heat waves, both HWEs and VSHSEs, have been identified in the last ten years.

A heat wave event (HWE) of 10 days in a row (Niš, July from 15<sup>th</sup> to 24<sup>th</sup>, 2007) is the maximum number of consecutive days with such high temperatures when all the three stations are compared. In the same month, HWE of 3 days in a row (July 8<sup>th</sup> to 10<sup>th</sup>) and 4 days in a row (July 27<sup>th</sup> to 30<sup>th</sup>) were observed (Table 4, HWE). Altogether, the heat wave events (HWEs) occurred 3 times and they lasted 3, 10 and 4 days in July 2007, which is certainly an extreme for the observed period.

(lines 354-359) **Table 4.** Number of Heat waves events (HWE) and Very strong heat stress event (VSHSE) in Niš, Novi Sad - July, Aug and Sep, 1998-2017

HWE	July		Aug		Sep	VSHSE	July		Aug		Sep
	Niš	Novi Sad	Niš	Novi Sad	Niš		Niš	Novi Sad	Niš	Novi Sad	Niš
1998	4	/	5	/	/	1998	7	/	5	/	/
1999	/	/	3, 4	/	/	1999	/	/	5	/	/
2000	3	/	4, 6	6	/	2000	/	/	6	7	/
2001	3	/	3, 3	/	/	2001	/	/	/	/	/
2002	6	/	/	/	/	2002	/	/	/	/	/
2003	4	/	/	/	/	2003	/	/	/	/	/
2004	3, 4	3	/	/	/	2004	5, 5	/	/	/	/
2005	6	/	/	/	/	2005	6	/	/	/	/
2006	/	/	3	/	/	2006	/	/	/	/	/
2007	3, 10, 4	6	6	/	/	2007	10	7	/	/	/
2008	/	/	4	3	3	2008	/	/	5	/	/
2009	/	/	3	/	/	2009	/	/	/	/	/
2010	/	/	4	/	/	2010	/	/	5	/	/
2011	3	4	3	/	/	2011	/	/	/	/	/
2012	9	6	4, 6	3, 5	/	2012	12	6	5, 6	5, 5	/
2013	/	/	7	4	/	2013	/	/	8	8	/
2014	/	/	/	/	/	2014	/	/	/	/	/
2015	7	9	6, 4	4, 5	5, 3	2015	10	8	7	6	5
2016	3	/	/	/	/	2016	/	/	/	/	/
2017	4, 4	/	6, 4	6, 3, 3	/	2017	/	/	6, 5	6	/

Note: pink cells have more than one HWE and VSHSE duration defined by numbers of the consecutive days

Of the total number of hot days, it amounts to 21 HDs in September in Niš, only one HD was recorded in the first decade (1998-2007) of the investigated period, while the remaining HDs were recorded over the last decade (2008-2017), especially in 2011 and 2015. In the second decade of the investigated period the year 2015 stands out with two HWEs (September 1<sup>st</sup> to 5<sup>th</sup>, and 17<sup>th</sup> to 19<sup>th</sup>). In the same year the highest daily temperature was recorded in September (on September 18<sup>th</sup>,  $t_{\max}=37.5^{\circ}\text{C}$ ).

(lines 369-384) Bioclimatic conditions analyzed by means of the UTCI index show that the calculated data observed at 14 CET are related to the marked heat wave events (HWE). When  $\text{UTCI}_{14\text{h}}$  thermal stress category of “very strong heat stress” (VSHS) is compared to the selected heat wave events (HWE,) a sub-index is defined. It is called very strong heat stress event (VSHSE) and it is caused by the occurrence of 5 consecutive VSHS days. The VSHSE corresponds to heat wave event and provokes a severe biothermal discomfort, so it was used as an indicator of extremely unfavorable bioclimatic conditions (biothermal heat hazard). Along these lines, the VSHSE in Niš occurred during July in 1998, 2004, 2005, 2007, 2012 and 2015 where 7, 5 (on two occasions), 6, 10, 12 and 10 VSHS days in a row were recorded respectively (Table 4, VSHSE). In August, VSHSE was recorded on 5 VSHS days in a row in 1998, 1999, 2008, 2010, 2012 and 2017, while in 2000, 2012, 2013, 2015 and 2017 6, 6, 8, 7 and 6 VSHS days in a row were recorded respectively. It should be emphasized that there were two VSHSE events in 2012 and 2017, for each year 5 and 6 VSHS days in a row. September 2015 saw a VSHSE with 5 VSHS days in a row. According to Unkašević and Tošić (2009b) the highest temperatures in Niš ever were recorded during the summer of 2007 ( $44.2^{\circ}\text{C}$ ) and the summer of 2000 ( $42.5^{\circ}\text{C}$ ), covering the data period from 1948 to 2007. The VSHSE occurred in Novi Sad in July 2007 (7 VSHS days in a row), 2012 (6 VSHS days in a row) and 2015 (8 VSHS days in a row) and in August 2000 (7 VSHS days in a row), 2013 (8 VSHS days in a row), 2015 (6 VSHS days in a row) and 2017 (6 VSHS days in a row). In August 2012 VSHSE occurred twice and each time 5 VSHS days in a row were recorded. There was no sub-index VSHSE recorded in Zlatibor.

#### Comment 6: Figures

*The ordinate of the figure 2, 3 and 4 need to be change to “number of days” instead of “Days”. In general, figures and tables needs to be improved to make them easier to understand.*

**Author response:** We agree with this suggestion and we have improved the figures and tables. Figures 2,3 and 4 have been changed and merged to form one, which became Figure 4 in the revised manuscript. Furthermore, one new figure (Figure 3 in the revised version) has been added in the first section of Results to represent the trend of the  $\text{UTCI}_{14\text{h}}$  index. Table 3 has been supplemented with VSHSE index compared with HWE. In the revised manuscript it became Table 4. The revised figures and tables are provided in the previous author response (response for Comment 5: Results).

#### Additional clarifications

Considering that the revised version has separated the results and the discussions, we need to point out the changes in the discussion section. The revised text is provided as follows:

(line 387-419) **Discussion**

The assessment of human biothermal conditions during the summer made it possible to identify biothermal discomfort related to heat waves in Serbia. The results obtained in the study for the investigated period from 1998 to 2017, indicates the general increase in biothermal discomfort associated with heat waves defined by maximum air temperature above  $35^{\circ}\text{C}$  and UTCI “very strong heat stress” above  $38^{\circ}\text{C}$ .

The results obtained for morning, midday and average UTCI show significant occurrence of thermal heat stress during the summer months and increase in biothermal discomfort especially for  $\text{UTCI}_{14\text{h}}$ . This is confirmed by obtained distribution of the average monthly  $\text{UTCI}_{14\text{h}}$  showing a significant increase in the extreme values of the

UTCI<sub>14h</sub> index in the last ten years, presented mainly as “very strong heat stress category” (VSHS) together with a growing trend of the UTCI<sub>14h</sub> index in all the summer months (especially September) for all three investigated weather stations (primarily in Niš). According to Di Napoli et al. (2018), there are two thermal climates in Europe. One of them is associated with heat stress condition and it is predominant in the southern part of Europe (including the Balkans) when moderate and strong heat stress occurs at central day time hours, which reflects the general relationship between heat load and insolation.

The increase in extreme biothermal conditions is the most evident in the number of days with maximum air temperature thresholds above 35°C, defined in the study as hot days (HD) and in the number of days with UTCI index thresholds between 38°C and 46°C, defined in the study as “very strong heat stress” (VSHS). Thus, most HDs (18 days) occurred in 2007 and then in 2012 (16 days), which means that more than half of July in these years was characterised by extremely high temperatures. Similar results about increase in daily maximum air temperature were obtained by Unkašević and Tošić (2011), when the record values of the maximum temperatures were observed for almost the whole territory of Serbia in 2007. As reported by Papanastasiou et al. (2014), the summer of 2007 was the warmest summer in Athens in the last hundred years. Besides, Spinoni et al. (2015) in their study compiled a list of the most severe drought events which occurred in Europe from 1950 to 2012. They singled out the Balkan countries as extreme weather and drought-prone areas (particularly Serbia) with the longest drought registered in 2007, and the most severe drought event in 2011. Moreover, in 2012 the Balkan Peninsula and the south-eastern Europe faced the hottest summer and one of the worst droughts in nearly 40 years (Unkašević and Tošić, 2015). This is in relation with the investigation of heat waves (a sequence of at least 5 days with maximum temperature above the 95<sup>th</sup> annual percentile) in south-eastern Europe, where there was an increase in the number of hot days for the period from 1973 to 2010. The UTCI<sub>14h</sub> category of “very strong heat stress” (VSHS) correlates with the maximum air temperatures and there has been an increase in the number of such days over the last ten years. Particularly severe biothermal discomfort occurred in 2007 (18 VSHS days) and 2012 (17 VSHS days). This is in agreement with the statement of Di Napoli et al. (2018) about UTCI reference value from the period between 1979 and 2016, when the UTCI at 12:00 (UTC) was about 0.5°C colder in the period between 1980 and 1999, while it was 0.5°C warmer in the period between 2000 and 2009 and 1°C warmer in the period between 2010 and 2016.

(lines 437-443) In this sense, there is a difference between the definition of biothermal heat stress and potential heat stress, defined on the basis of the interpretation of maximum temperatures. Determining extreme weather event from the aspect of human heat budget indices allows other meteorological parameters to be taken into account. Apart from temperature and humidity, wind and direct solar radiation, have been considered together with physiological metabolic rate.

The identified heat waves based on the threshold of the UTCI index correspond to the identified heat waves based on the threshold of maximum air temperatures. Furthermore, there is a significant increase in heat waves (HWE and VSHSE) plus duration of such events over the last ten years in Niš and Novi Sad, especially in Niš.

(lines 438-461) This agrees with the findings of Unkašević and Tošić (2011, 2015) that there has been a growing trend of heat waves in Serbia, especially since 2000. In order to justify the present research for the last 20 years, the earlier results related to the increase in heat waves will be discussed in more detail. In that regard, previous research into heat waves in Niš for the period from 1949 and 2007, based on the autoregressive-moving-average model, observed the warmest years during three periods, 1951-1952, 1987-1998 and 2000-2007 (Unkašević and Tošić, 2009b). According to this research, the longest heat wave was recorded in Niš in 1952, with 21 days, while in 2003, 29 consecutive tropical days were observed. It should be taken into account that according to the Serbian National Weather Service, tropical days are days with a maximum temperature over 30°C. Furthermore, based on Heat Wave Duration index (HWD) i.e. daily maximum values of air temperature, Drljača et al. (2009) determined the duration and strength of heat waves in Niš during the summer season. The analysis showed Niš with a greater number of heat waves compared to the larger urban area of Belgrade. As stated in the research, since the mid-1980s, heat waves have had a higher frequency and on average they have occurred every year. Prior to that period heat wave fluctuations were generally reported in one of two years (Niš during the summer) (Drljača et al., 2009). While studying the characteristics of the heat waves in central Serbia (1949-2007), Unkašević and Tošić (2011) detected an increase in heat wave duration, in addition to an increase in heat wave frequency of occurrence during the period from 1999 to 2007. Besides, based on the HWDI index, heat waves in Novi Sad show a trend of increasing tropical



days ( $T_{\max} \geq 30.0^{\circ}\text{C}$ ) from 1960 to 2010, indicating years 1994, 1998, 2000, 2003, 2007, 2009 (Malinović-Milićević et al., 2013).

As regards the heat budget index of Heat Load (HL) in Serbia (July) for the period from 2000 to 2010, it was noticed that in, Niš, Zlatibor and Novi Sad the category of “extremely hot” occurred in 2000 and 2007 (Milovanović et al., 2017). As for the heat budget index Physiological Equivalent Temperature PET for the period from 1949 to 2012, the highest number of heat waves in Novi Sad was observed in the last two decades and in the first decade of the investigated period (Basarin et al., 2016), while the number of the days above particular thresholds for the period 1961-2014 shows an increase along with the number of heat waves per year since 1981. The highest values of mean annual PET were mostly detected during the last 15 years (Basarin et al., 2018).

(lines 468-495) The biothermal discomfort identified in July and August as the hottest summer months in Serbia, could provoke health disorders more frequently. Specific HWs characteristics such as intensity and duration have been found to have devastating effects on human health and wellbeing. In certain cases heat waves cause problems and children, elderly people, chronic patients and workers are particularly susceptible to them. The analysis of the UTCI index correlating with mortality in Europe shows that deaths occur in the southern Europe (Italy, Portugal, Greece, Spain) for the categories of moderate and strong stress. Heat load has an impact on increasing mortality as conditions become more thermally stressful (Di Napoli et al., 2018). The intensity of heat stress in Poland was assessed by means of the UTCI index on the basis of daily mortality and the weather data for the period from 1991 to 2000. There is an increase in mortality on the days with strong and very strong heat stress, in relation to no thermal stress days, of 12% and 47%, respectively (Błażejczyk et al., 2017). Biothermal conditions and mortality rates show that in southern Europe during the summer thresholds of Physiological Subjective Temperature (PST) increase when moving from the west to the east. The general finding is that the population of central Europe is more sensitive to thermal stimuli in the summer than the citizens in the south of the continent (Błażejczyk and McGregor, 2008). The statement about the PST index is in support of the comparison of UTCI index with other thermal indices, including PST index, which correlates very well with UTCI index (Błażejczyk et al., 2012). Human sensitivity to extreme heat weather can also be seen in the impact of heat waves on daily mortality in Belgrade during the summer, when the strong correlation between heat waves and daily mortality can be observed in July 2007 (Stanojević et al., 2014). Similarly, Bogdanović et al. (2013) reported a significant short-term excess mortality on 16<sup>th</sup> July in Belgrade, when the maximum daily temperature exceeded 35°C, leading to 167 excess deaths (38% increase compared to the number of expected deaths) for nine consecutive days of heat, with a progressive return to almost normal mortality as the maximum temperature dropped below 35°C on 25<sup>th</sup> July.

The application of the standardized bioclimatic heat budget index UTCI could help improve the understanding of biothermal conditions relative to heat wave. The daily  $\text{UTCI}_{14\text{h}}$  heat stress becomes more extreme in terms of severity and heat wave duration up to very strong heat stress (VSHS). The biothermal indices investigated for the three weather stations follow the trend of general warming. In the light of climatic changes and other negative factors resulting from this global phenomenon, it is becoming a true challenge to minimize their effects and improve living conditions in urban and rural areas (Stevović et al., 2017). However, the human heat budget indices are increasingly serving as a determinant of extreme weather condition (Di Napoli et al., 2018; Matzarakis et al., 2018; Theoharatos et al., 2010; Urban and Kyselý, 2014). UTCI index improves the understanding of extreme thermal stress, serving as one of the criteria for initiating heat alerts about extreme thermal stress in Serbia.

#### **Comment 7: Conclusion**

*In the introduction of the paper, the role of such indexes is introduced (weather alert, weather forecast). The conclusion should come back to the importance and role of this UTCI index values and the possible consequences of its increase on population.*

**Author response:** We have incorporated your suggestion throughout the conclusion section. The revised text is provided as follows:

(Lines 497-510) The present research of the human bioclimatic evaluation of UTCI thermal stress in Serbia is considered to be of great importance due to the identification of biothermal heat hazard and the study of the evolution of such indices regarding climate change. This evaluation aims at providing a comprehensive assessment of the human heat stress associated with UTCI index and heat waves defined by UTCI very strong heat stress and maximum air temperature.

The assessment of human biothermal conditions for the investigated period from 1998 to 2017 was provided for three synoptic stations – two urban (Niš and Novi Sad) and one station representing mountain areas of an altitude up to 1500 m (Zlatibor). All of them are located in different geographical areas in Serbia. The results obtained during the study indicate the significant increase in biothermal discomfort associated with heat waves defined by UTCI “very strong heat stress”. It was found that the most frequent heat stress categories are “strong heat stress” (SHS) and “very strong heat stress” (VSHS) and they have occurred in all three locations with the tendency of increase in the number of days in the last decade. The VSHS describes an alarming biothermal state and has occurred frequently in the last ten years, particularly in Niš and Novi Sad. The outcome of the study is UTCI<sub>14h</sub> sub-index “very strong heat stress event” (VSHSE) as an indicator of biothermal heat hazard. Undoubtedly, heat waves (HWs) are one of the natural hazards with increasing impact in urban areas related to higher population density.

(lines 514-519) Considering these facts, it can be deduced that heat waves are becoming more frequent, stronger and longer. Thus, frequent heat waves since 2007 certainly indicate biothermal heat hazard. The more lasting events of bioclimatic discomfort will indicate more stressful bioclimatic conditions for human health and wellbeing. However, the results of the study highlight the importance of UTCI index as a bioclimatic indicator of biothermal hazard in Serbia particularly if it can serve as one of the criteria for initiating heat warnings in Serbia.

#### **Comment 8: Reference**

*Reference: The following reference seems relevant and needs to be added: Bogdanović, D. and al. (2013). The impact of the July 2007 heat wave on daily mortality in Belgrade, Serbia. Central European journal of public health, 21(3), p. 140-145.*

**Author response:** We have added the suggested reference in the Discussion section (lines 492-495). The revised text is provided as follows:

(lines 483-486) Similarly, Bogdanović et al. (2013) reported a significant short-term excess mortality on 16<sup>th</sup> July in Belgrade, when the maximum daily temperature exceeded 35°C, leading to 167 excess deaths (38% increase compared to the number of expected deaths) for nine consecutive days of heat, with a progressive return to almost normal mortality as the maximum temperature dropped below 35°C on 25<sup>th</sup> July.

Other supplemented reference in the revised version is as follows:

- ANSI/ASHRAE Standard 55, Thermal Environmental Conditions for Human Occupancy, (ANSI/ASHRAE Approved), [www.techstreet.com/ashrae/standards](http://www.techstreet.com/ashrae/standards), 2017.
- Di Napoli, C., Pappenberger, F., and Cloke, H.L.: Verification of Heat Stress Thresholds for a Health-Based Heat-Wave Definition, *J. Appl. Meteorol. Climatol.*, 58(6), 177- 1194, <https://doi.org/10.1175/jamc-d-18-0246.1>, 2019.
- Easterling, D.R., Horton, B., Jones, P.D., Peterson, T.C., Thomas, R.K., David, E.P., Salinger, M.J., Razuvayev, V., Plummer, N., Jamason, P., and Folland, C.K.: Maximum and minimum temperature trends for the globe. *Sci.*, 277(5324), 364–367, <https://doi.org/10.1126/science.277.5324.364>, 1997.
- European Commission's (Eurostat) datasets: <https://ec.europa.eu/eurostat/web/gisco/geodata/reference-data/>, last access: 3<sup>rd</sup> February 2020.
- Fanger, P.O.: Thermal Comfort: analysis and applications in environmental engineering, Danish Technical Press, Copenhagen, Denmark, 1970.
- Gocić, M. and Trajković, S.: Analysis of precipitation and drought data in Serbia over the period 1980–2010, *J. Hydrol.*, 494, 32-42, <https://doi.org/10.1016/j.jhydrol.2013.04.044>, 2013.
- Gocić, M. and Trajković, S.: Spatiotemporal characteristics of drought in Serbia, *J. Hydrol.*, 510, 110-123, <https://doi.org/10.1016/j.jhydrol.2013.12.030>, 2014.

- Höppe, P.: The physiological equivalent temperature - a universal index for the biometeorological assessment of the thermal environment, *Int. J. Biometeorol.*, 43, 71–75, <https://doi.org/10.1007/s004840050118>, 1999.
- IPCC, Special Report of the Intergovernmental Panel on Climate Change, in: *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*, edited by Field, C. B., Barros, V., Stocker, T.F., Qin, D., Dokken, D.J., Ebi, K.L., Mastrandrea, M.D., Mach, K.J., Plattner, G.K., Allen, S.K., Tignor, M. and Midgley, P.M., Cambridge University Press, Cambridge, UK, 582, 2012.
- ISO 8996: Ergonomics of the thermal environment - Determination of metabolic rate, <https://www.iso.org/standard/34251.html>, 2004.
- Landsberg, H. E.: *The Assessment of Human Bioclimate: a limited review of physical parameters*, World Meteorological Organization WMO, 331, Geneva, 1972.
- Lee, V., Zermoglio, F., and Ebi, K.: Heat waves and human health - emerging evidence and experience to inform risk management in a warming world, Technical Report, United States Agency for International Development USAID, Washington 2019
- Leščešen, I., Dolijan, D., Pantelić, M., and Popov, S.: Drought assessment in Vojvodina (Serbia) using k-means cluster analysis, *J. Geogr. Inst. Jovan Cvijić SASA.*, 69(1), 17-27, <https://doi.org/10.2298/IJGI1901017L>, 2019.
- Matzarakis, A., Muthers, S., and Rutz, F.: Application and comparison of UTCI and PET in temperate climate conditions, *Finisterra*, 48(98), 21–31., <https://doi.org/10.18055/Finis6453>, 2014.
- Masterton, J. and Richardson, F.A.: Humidex, a method of quantifying human discomfort due to excessive heat and humidity, Atmospheric Environment Service Canada, Ontario, Downsview, 1979.
- McGregor, G. and Vanos, J.: Heat: A primer for public health researchers, *Public Health*, 161, 138–146, <https://doi.org/10.1016/j.puhe.2017.11.005>, 2018.
- Milewski, P.: 2013. Application of the UTCI to the local bioclimate of Poland's Ziemia Kłodzka region, *Geogr. Pol.*, 86(1), 47-54, <https://doi.org/10.7163/GPol.2013.6>, 2013.
- Milovanović, M., Gocić, M., and Trajković, S.: Analysis of meteorological and agricultural droughts in Serbia, *Architecture and Civil Engineering*, 12(3), 253-264, <https://doi.org/10.2298/FUACE1403253M>, 2014.
- Nassiri, P., Monazzam, M.R., Golbabaie, F., Dehghan, S.F., Rafieepour, A., Mortezaipoor, A., Asghari, M.P.: Application of Universal Thermal Climate Index (UTCI) for assessment of occupational heat stress in open-pit mines, *Industrial Health*, 55 (5), 437-443, <https://doi.org/10.2486/indhealth.2017-0018>, 2017.
- Nastos, P. and Matzarakis, A.: The effect of air temperature and human thermal indices on mortality in Athens, Greece, *Theor. Appl. Climatol.*, 108, 591–599, <https://doi.org/10.1007/s00704-011-0555-0>, 2012.
- Nemeth, A.: Changing thermal bioclimate in some Hungarian cities, *Acta Climatologica et Chorologica*, 44–45, 93–101, 2011.
- Oscewski, R.J. and Bluestein, M.: The new wind chill equivalent temperature chart, *Bull. Am. Meteorol. Soc.*, 86, 1453–1458, <https://doi.org/10.1175/BAMS-86-10-1453>, 2005.
- Parsons, K.C.: *Human thermal environments: the effects of hot, moderate, and cold environments on human health, comfort and performance*, Taylor & Francis Group, Abingdon, England, UK, 2003.
- Rahmstorf, S. and Coumou, D.: Increase of extreme events in a warming world, *P. Natl. Acad. Sci.*, 108(44), 17905–17909, <https://doi.org/10.1073/pnas.1101766108>, 2011.
- Siple, P. and Passel, C.F.: Measurements of dry atmospheric cooling in subfreezing temperatures, *Proc Am Philos Soc*, 89(1), 177–199, [www.jstor.org/stable/985324](http://www.jstor.org/stable/985324), 1945.
- Spinoni, J., Naumann, G., Vogt, J., and Barbosa, P.: The biggest drought events in Europe from 1950 to 2012, *J. Hydrol. Reg. Stud.*, 3, 509-524, <https://doi.org/10.1016/j.ejrh.2015.01.001>, 2015.
- Seneviratne, S.I. and Nicholls, N.: Changes in Climate extremes and their impacts on the natural physical environment, in: *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*, edited by Field, C. B., Barros, V., Stocker, T.F., Qin, D., Dokken, D.J., Ebi, K.L., Mastrandrea, M.D., Mach, K.J., Plattner, G.K., Allen, S.K., Tignor, M. and Midgley, P.M., Cambridge University Press, Cambridge, UK, 109–230, 2012.
- Steadman, R.G.: A Universal Scale of Apparent Temperature, *J. Appl. Meteor.*, 23, 1674-1687, [https://doi.org/10.1175/1520-0450\(1984\)023<1674:AUSOAT>2.0.CO;2](https://doi.org/10.1175/1520-0450(1984)023<1674:AUSOAT>2.0.CO;2), 1984.

- Tolika, K., Maheras, P., Pytharoulis, I., and Anagnostopoulou, C.: The anomalous low and high temperatures of 2012 over Greece - an explanation from a meteorological and climatological perspective, *Nat. Hazards Earth Syst. Sci.*, 14(3), 501–507, <https://doi.org/10.5194/nhess-14-501-2014>, 2014.
- Tomczyk, M.A. and Owczarek M.: Occurrence of strong and very strong heat stress in Poland and its circulation conditions, *Theor. Appl. Climatol.*, 139, 893-905, <https://doi.org/10.1007/s00704-019-02998-3>, 2020.
- Tošić, I. and Unkašević, M.: Analysis of wet and dry periods in Serbia, *Int. J. Climatol.*, 34, 1357-1368, <https://doi.org/10.1002/joc.3757>, 2014.
- Urban, A. and Kyselý J.: Comparison of UTCI with other thermal indices in the assessment of heat and cold effects on cardiovascular mortality in the Czech Republic, *Int. J. Environ. Res. Public Health*, 11(1), 952-967, <https://doi.org/10.3390/ijerph110100952>, 2014.
- Yaglou, C.P. and Minard, D.: Control of heat casualties at military training centers, *Am Med Assoc Arch Ind Health*, 16(4), 302–316, 1957.
- 

#### Additional clarifications

According to all changes that have made in the manuscript we have modified Abstract and in revised text is provided as follows:

(lines 9) The objective of this paper is to assess the bioclimatic conditions in Serbia so as to identify biothermal heat hazard.

(lines 13-16) In order to identify patterns of biothermal heat stress conditions, the thresholds of daily UTCI index ( $UTCI \geq 38^{\circ}C$  refers as very strong heat stress-VSHS) at 14 CET, are compared with the thresholds of daily maximum of air temperature ( $t_{max} \geq 35^{\circ}C$  refers as hot day-HD) which are further termed as heat wave events (VSHSE and HWE). The findings show that the UTCI heat stress category “very strong heat stress” at 14 CET indicates heat waves. (lines 18-21) Heat wave events (HWE) and very strong heat stress events (VSHSE) recorded in July 2007 (lasted 10 days each), 2012 (lasted 9 and 12 days) 2015 (lasted 7 and 10 days) were of the longest duration and refer as indicators of biothermal heat hazard. The daily  $UTCI_{14h}$  heat stress becomes more extreme in terms of severity and heat wave duration up to very strong heat stress (VSHS).

#### Comment 9: English

*English language must be checked, many spelling mistakes are presents.*

**Author response:** With regard to the above comment, all the spelling and grammatical errors pointed out by the reviewers have been corrected.

18<sup>th</sup> February 2020

Response to peer reviewer comments

nhess-2019-270: **Summer variation of the UTCI index and Heat Waves in Serbia**

Milica M. Pecelj, Milica Z. Lukić, Dejan J. Filipović, Branko M. Protić

Dear Reviewer,

We would like to thank you for giving us the opportunity to submit a revised draft of our manuscript titled Summer variation of the UTCI index and Heat Waves in Serbia to Journal of Natural Hazard and Earth System Science. We appreciate the time and effort that you dedicated to providing valuable feedback on our manuscript.

We have been able to incorporate changes to reflect most of the suggestions. Since you emphasized major revisions to the manuscript, significant changes have been made to the paper. The revision, based on your input, includes changed interpretation of research in all chapters.

Here are the responses to yours comments and concerns, listed step by step. To make the changes in the manuscript easier to follow, we have highlighted the modified text in **green** and the new added text in **blue**.

**Reviewer Comment:**

*The assessment of bioclimatic conditions in Serbia has been investigated in the present research. This work seems to be interesting and can be accepted for publication after major revisions. The content of the manuscript should be deeply improved in terms of scientific (results, interpretations and discussions) and technical approaches (methodological approach). The content is deemed to be not fully satisfactory to understand the main finding of the work. Introduction According to the introduction, many studies have been carried out in Serbia. It's important to underline the original contribution of the present research. This part should be improved and some part are difficult to understand. Methodological approach Any detailed explanation has been presented in this part. I have not understood the methodology and how the results have been calculated. Then, a more recent meteorological data sets are used for the computation of the UTCI index values for the selected stations. Starting in 1998 should not be enough to calculate the long-term trends. Starting before should be more interesting to identify the trend of such extreme heat waves. Results This part should be more explained. I cannot fully understand the results since the methodology is not detailed. More analysis should be added to explain the main finding of the paper. Parts of Discussions and Conclusions Both parts should be written differently. The work should be discussed with other previous works dealing with extremes in this zone and also other works using the same methodology (if this one is not new.. I don't know exactly since it's not clearly presented).*

**Author response:** Thank you for pointing this out. Since the major revisions to the manuscript have emphasized, significant changes were made to the manuscript. The revision, based on the review input, includes a number of following changes:

#### **Title**

We have modified the title and named in revised text as: [Analysis of UTCI index during heat waves in Serbia](#)

#### **Abstract**

*According to all changes that have made in the manuscript we have modified Abstract and in revised text is provided as follows:*

(lines 8-9) [The objective of this paper is to assess the bioclimatic conditions in Serbia so as to identify biothermal heat hazard.](#)

(lines 12-14) [In order to identify patterns of biothermal heat stress conditions, the thresholds of daily UTCI index, refers as very strong heat stress \(VSHS\), are compared with the thresholds of daily maximum of air temperature which are termed as heat wave events.](#)

(lines 16-18) [Heat wave events recorded \(HWE, VSHSE\) in July 2007 \(10 days\), 2012 \(9, 12 days\) 2015 \(7, 10 days\) were of the longest duration and refer as indicators of biothermal heat hazard.](#)

#### **Introduction**

In relation to your remarks, we have completely modified the introduction where have emphasized and explained the concept of heat budget index more clearly. We have supplemented the introduction with a new text outlining the concept of heat budget indices, their importance for bioclimatic analysis.

We agree with the reviewer statement about data set used in the study however, in the case of our study, it seems slightly out of scope because the procedure for obtaining daily hourly data from the national weather center is very slow, so it would be impossible to respect the time limit for correction. It would certainly have been interesting to explore this aspect for future research. Since many studies have been conducted in Serbia concerning the analysis of heat waves in the last 50 years, we considered their results relevant indicating an increase in heat waves especially since 2000. Starting from this fact, we used available data only from 1998-2017 and for that period we did a bioclimatic identification of thermal discomfort. We changed a part in introduction to emphasize this point.

Then, we modified the paragraph followed by detailed recent studies (lines 47-58; lines 63-74; lines 77-94) and moved it into the discussion section (lines 479-494; lines 448-462; lines 463-469). More specifically, lines 47-58 have been moved into the discussion lines 479-494, where a reference about heat wave related mortality in 2007 by Bogdanović et al. (2013) was added (lines 491-494). In addition, lines 63-74 have been moved into the discussion lines 448-462, and there is an explanation of what a tropical day is (line 455-456) in the analysis by Unkasevic and Tomic (2009b). Lastly, lines 77-94 have been moved into the discussion section (lines 466-472).

The revised text reads as follows on:

(lines 21-25) [Extreme weather events such as heat waves, floods, droughts or storms have shown an increased frequency in recent decades \(Brown et al., 2008; Easterling et al., 1997; IPCC, 2012; Rahmstorf and Coumou, 2011; Seneviratne and Nicholls, 2012\). Not only are heat and drought events of great importance in most Mediterranean](#)

climate regions, but also in most of southern and south-eastern Europe, because of the diverse and costly impact that they have on various economic sectors and on the environment (Peña-Gallardo et al., 2019).

(lines 27-29) They have both direct effects on human health, affecting the body's physiological responses and functions and indirect effects on human health, increasing challenges to food and water safety (Lee et al., 2019).

(lines 32-35) Over the last century many models and indices have been developed for the assessment of human exposure to heat, ranging from simple physical instruments designed to imitate the human heat exchange with the environment, to complex thermophysiological models that simulate external and internal body heat transfer and allow detailed simulation of different work load, clothing, and climate scenarios (Havenith and Fiala, 2015).

(lines 36-64) Many thermal indices have been developed for the purpose of describing the complex conditions of heat exchange between the human body and its thermal environment (Fanger, 1970; Landsberg, 1972; Parsons, 2003). Those are two meteorological parameter indices – the ones used for cold environment conditions combining the air temperature and the wind velocity (Osczevski and Bluestein, 2005; Siple and Passel, 1945;), and the ones used for heat environment conditions combining the air temperature and humidity (Masterton and Richardson, 1979; Steadman, 1984; Yaglou and Minard, 1957). Considering their shortcomings, i.e. the fact that they are not universally applicable to all climatic regions, including different seasons, the dominance in the analysis of biothermal conditions and thermal stress over the last 30 years has been taken over by so-called heat budget indices (Błażejczyk, 1994; Hoppe, 1999; Jendritzky, 2012). The heat budget indices are based on the thermal exchange between humans and the environment. Methodologically, they use variables related to meteorological (air temperature, wind velocity, radiation, air humidity) and physiological processes (most commonly metabolic heat) and clothing insulation. The issue that should be emphasized during the analysis of the thermal stress is the influence that extreme temperatures have on the physiological parameters in humans. According to McGregor and Vanos (2018), generated heat load can undermine the human body's ability to hold its core temperature within the range of optimal physiological achievement.

For the purpose of estimating the thermal effect of the environment on the human body, the total effects of all the thermal components are considered by the UTCI index. The UTCI is heat budget index considering both physiological and meteorological parameters describing the physiological heat stress that the human body experiences while achieving thermal equilibrium with the surrounding outdoor environment (Błażejczyk et al., 2013). Compared to other indices, UTCI is more sensitive to changes in all of the environment parameters, particularly air temperature, solar radiation, humidity, and wind speed (Błażejczyk et al., 2012). According to Jendritzky et al. (2012) the UTCI evaluates the outdoor thermal environment for biometeorological applications by simulating the dynamic physiological response with a model of human thermoregulation together with a modern clothing model. UTCI provides a multiple opportunity for investigation and has a wide application in weather forecast for outdoor activities, appropriate behavior and climate therapies, and extreme thermal stress alerts. These analyses can have application in tourism, health sector, and urban and regional planning (Jendritzky et al., 2012). In the past decade, a large number of publications in Europe concern analyses of bioclimatic condition in accordance with the UTCI index (Błażejczyk et al., 2014; Błażejczyk et al., 2017; Bleta et al., 2014; Di Napoli et al., 2018; Di Napoli et al., 2019; Matzarakis et al., 2014; Milewski, 2013; Nastos and Matzarakis 2012; Nemeth, 2011; Tomczyk and Owczarek, 2019; Urban and Kyselý, 2014).

(lines 72-83) There is a need to study the evolution of such indices regarding heat stress that affects the human thermoregulatory system as a result of heat exchange between the body and its thermal environment.

Considering extreme temperature is one of the most significant climatic parameters in the universal context of climate change, analyses of heat waves in Serbia have been performed in several different approaches and all of them show a growing trend in max temperature (Drljača et al., 2009; Unkašević and Tošić, 2009a; Unkašević and Tošić, 2009b; Unkašević and Tošić, 2013) and heat waves (Unkašević and Tošić, 2011; Unkašević and Tošić, 2015) especially since 2000.

Keeping this fact in mind, the basic idea of this study is to conduct the analysis of human bioclimatic conditions in Serbia in the last 20 years, where according to the previous studies mentioned above, warming has been perceived and recorded, especially since 2000. Determining extreme weather event from the aspect of heat budget indices allows other meteorological parameters to be taken into account because high temperatures and humidity generate a heat load more rapidly, unless the wind and direct radiation are taken into account.

(lines 86-90) The conducted human bioclimatic evaluation of UTCI thermal stress in Serbia was considered to be of great importance due to the identification of biothermal heat hazard and the study of the evolution of such indices regarding climate change. This evaluation is aimed at addressing the following topics in this study:


- providing a comprehensive assessment of the human heat stress associated with UTCI index and
- comparing it to heat waves defined by meteorological parameters.

To provide a clearer introduction we have inserted a new text. Namely, we added a heading “Study area” where we provided a relief map of Serbia, inside Europe, with the location of the three stations studied in the paper (lines 92-105). Basic information about the climate of Serbia has been added. Thus, we switched the first paragraph (lines 105-115) from the Materials and methods section, which in the revised version takes up lines 107-121.

The revised text reads as follows on:

(lines 92-121) *Study area*

The Republic of Serbia is located in south-eastern Europe, in the area of the southern Pannonian Plain and the central Balkans. Northern Serbia is mainly flat, while its central and southern areas consist of highlands and mountains (Gocić and Trajković, 2014), as the map of the relief characteristics shows (Figure 1). The territory of Serbia is characterised by temperate continental and mountain climate and the spatial distribution of climate parameters is determined by geographical location. The research involved studying three synoptic stations located in different geographical areas of Serbia: (i) Novi Sad, (ii) Niš and (iii) Mt. Zlatibor (Figure 1).



**Figure 1.** Relief map of Serbia with the areas studied: 1- Novi Sad, 2- Niš, 3- Zlatibor and a map of geographical location of Serbia in Europe. Maps were created using QGIS 3.8 software on the basis of the official European Commission's (Eurostat) datasets, available at <https://ec.europa.eu/eurostat/web/gisco/geodata/reference-data/>, (Map ratio: 1:1900000; Map projection: WGS 84/UTM, Zone 34N, the official national coordinate system)

The first weather station (at an altitude of 86 m) is located in the territory of Novi Sad, which is the administrative urban center of Vojvodina province and South Bačka District (Figure 1<sub>1</sub>). The city is located in the southern part of the Pannonian Basin, on the Danube River bank near Mt. Fruška Gora, and the national park bearing the same name.



Novi Sad has a temperate continental climate, summers are warm and winters are cold with a small amount of snow (Lazić et al., 2006). The second station is Niš (at an altitude of 202 m) and it is located in the Niš Fortress. This city is the administrative urban center of the Nišava District in southern Serbia and it is situated in the Nišava valley, located in the central part of the spacious geological depression called the Nišava Basin (Figure 1<sub>2</sub>). According to Köppen's climate classification, the Nišava valley belongs to the Cfwax type – the Danube type of moderately warm and humid climate characterized by hot summers (the highest precipitation is recorded at the beginning of the summer) and somewhat dry winters (Prokić, 2018). Compared to other cities in this valley (Dimitrovgrad, Pirot and Bela Palanka), Niš is the hottest one with an average annual temperature of 11.8°C (Prokić, 2018). The third station is Zlatibor, a mountain in western Serbia, which belongs to the mountain range of the Dinaric Alps (Figure 1<sub>3</sub>). Zlatibor weather station lies at an altitude of 1029 m a.s.l. In the area of Zlatibor plateau mountains meet air currents from the Adriatic Sea, which, as it can be assumed, creates a favorable climate and for this reason Zlatibor has already been confirmed as a climatic resort (Pecelj et al., 2017). The mountain climate of Zlatibor involves long and cold winters, short and fresh summers, and less pronounced spring and autumn.

## Materials and methods

We have, accordingly, a modified description of the model used to compute the index and we have made changes, clarifying the index analysis methodology. More specifically, we have divided the Materials and methods into two sections. In the beginning, an overview of meteorological and physiological components is given (lines 123-133), where we describe in more detail the metabolic rate as a key physiological parameter. The first section deals with the Universal Thermal Climate Index (UTCI), where we provide detailed definition, parameter inputs, and their limitations including stress classification (123-142; 147-169).

The revised text reads as follows on:

(lines 123-142) The present study implements the methodological approach of Universal Thermal Climate Index (UTCI) based on human heat budget model relying on the evaluation of human energy balance. As a thermal comfort indicator, the UTCI considers combined meteorological and physiological parameters describing thermal comfort through the evaluation of human energy balance. In terms of physiological conditions, the metabolic rate plays the most important role. Metabolic processes in the human body create heat that is constantly exchanged with the environment achieving a state of thermal equilibrium in the body for maintaining a constant body temperature. The amount of the heat that is produced and released depends on the following: physical activity, clothing, sex, age, body mass, diet, mental state, health, external conditions, acclimatization, etc. As a measure of physical activity, a unit of "MET" was defined, which corresponds to the release of human heat of  $58.2 \text{ W}\cdot\text{m}^{-2}$  for average body surface area ( $1.8 \text{ m}^2$ ), i.e. it is equal to the energy rate produced per unit surface area of an average person seated at rest (ANSI/ASHRAE Standard 55). According to ISO 8996 for standard applications, the metabolic heat energy is  $M = 135 \text{ W}\cdot\text{m}^{-2}$  i.e. 2.3 MET, for a person moving at a speed of  $1.1 \text{ m}\cdot\text{s}^{-1}$ .

### *The Universal Thermal Climate Index (UTCI)*

For valid assessment of the outdoor thermal environment in the fields of public weather services, public health systems, urban planning, tourism and recreation and climate impact research, the Universal Thermal Climate Index (UTCI) has been developed based upon the most recent scientific progress in human thermo-physiology in biophysics and heat exchange theory. The UTCI is the result of an approach which was developed in the International Society of Biometeorology (ISB) Commission 6 and was later improved by COST Action 730 (Jendritzky et al., 2012). The design of UTCI is of great importance due to the fact that it is applicable to all seasons and climates together with all spatial and temporal scales.

(lines 147-169) In other words, this model simulates the same sweat production or skin wettedness in human body response as the actual environment condition (Błażejczyk et al., 2013; Błażejczyk et al., 2014). This is derived from the multi-dimensional dynamic response of a state-of-the-art multi-node thermo-physiological model of human heat transfer and thermoregulation (Fiala et al., 2012). The UTCI can be represented in general function as bellow:

$$\text{UTCI} = f(t, f, v, t_{\text{mrt}}) \quad (\text{Eq.1})$$

UTCI = f (air temperature, relative humidity, wind speed, mean radiant temperature)

For appropriate use, UTCI can only be approximated using a regression equation abbreviated from sample calculations performed by computing centers (Bröde et al., 2012; Jendritzky et al., 2012). It causes a narrow range of input parameters it can manage with. The regression equation supports reference environment described as: (i) air temperature (t) in the range of -50.0°C to +50.0°C, (ii) relative humidity (f) from 0 % to 100 %, (iii) wind speed (v) at 10 m from 0.5 ms<sup>-1</sup> and up to 17.0 ms<sup>-1</sup>, (iiii) including a difference between mean radiant temperature (t<sub>mrt</sub>) and air temperature (t) of -30.0°C to +70.0°C. Physiological parameters, the metabolic rate (M) and the thermal properties of clothing (clothing insulation, permeability) are taken as universal constants in the model due to the evaluation by means of the regression equation. This implies an outdoor activity of a person walking at the speed of 4 km.h<sup>-1</sup> (1.1 m.s<sup>-1</sup>), corresponding to heat production of 135 W.m<sup>-2</sup> (2.3 MET) of metabolic energy (Błażejczyk et al., 2013; Jendritzky et al., 2012) and the clothing insulation which is self-adapting according to the environmental conditions (Havenith et al., 2012). Clothing insulation, vapor resistance and the insulation of surface air layers, are strongly influenced by changes in wind speed and body movement and will therefore also influence physiological responses (Błażejczyk et al., 2012). Particular ranges of UTCI are categorised according to thermal stress (Table 1).

**Table 1.** UTCI thermal stress classification

UTCI (°C)	> 46	38 to 46	32 to 38	26 to 32	9 to 26	0 to 9	0 to -13	-13 to -27	-27 to -40	< -40
Stress	extreme heat stress	very strong heat stress	strong heat stress	moderate heat stress	no thermal stress	slight cold stress	moderate cold stress	strong cold stress	very strong cold stress	extreme cold stress
Abbr.	EHS	VSHS	SHS	MHS	NTS	SLCS	MCS	SCS	VSCS	ECS

Source: Błażejczyk et al. 2013

The second section, named *Data and indices considered in the study*, explains what kind of analysis was done in the study. The revised text reads as follows on:

(lines 170-210) *Data and indices considered in the study*

The meteorological data set from the period from 1998 to 2017 was recorded for two urban weather stations (Novi Sad and Niš) and one rural mountain station (Mt. Zlatibor). Mean daily and hourly meteorological parameters (07h, 14h, CET) of air temperature (t), air humidity (f), and wind speed (v) from the above weather stations were considered for the calculation of particular UTCI thermal heat stress in the summer months (July, August and September). The meteorological data set used in the study was retrieved from the Meteorological Yearbook for the period from 1998 to 2017 (Republic Hydrometeorological Service of Serbia) while the UTCI index was calculated by applying the BioKlima 2.6 software package (available at <http://www.igipz.pan.pl/Bioklima-zgik.html>).

Although extreme temperatures are one of the most effective climatic parameters in the universal case of hot days and heat waves, the influence of other parameters, especially clothing insulation and metabolic rate, can justify the thermal discomfort and the other way round. In that regard, the summer (July, August and September) daily maximum and minimum temperatures (t<sub>max</sub>, t<sub>min</sub>), for the same stations, were analysed in terms of particular thresholds to identify the frequency of extremely high temperatures and heat waves in the period observed to compare them with the discomfort recorded by the UTCI index.

The first stage of the study presented mean daily (UTCI<sub>avg</sub>), morning (UTCI<sub>7h</sub>) and midday (UTCI<sub>14h</sub>) UTCI indices. They are presented by months (July, Aug, Sep) for each station to identify the general differences caused by the geographical origin of the stations under study. The quantification of human bioclimatic conditions in Serbia was designed for the UTCI indices on a daily basis of each defined stress categories. To obtain a better insight into the UTCI<sub>14h</sub> index values during the summer months, averaged monthly UTCI<sub>14h</sub> index was provided for each year to see how much the UTCI<sub>14h</sub> index fluctuated in relation to the average monthly value.

For the second stage, midday UTCI thermal stress indices were identified (strong heat stress SHS, very strong heat stress VSHS, extreme heat stress EHS) and thermal indices based on the maximum and minimum temperatures (hot day HD and hot night HN). During the midday observation (14 CET), particularly prominent are strong heat stress (SHS), which refers to UTCI range of 32°C to 38°C and very strong heat stress (VSHS), which refers to UTCI range of 38°C to 46°C (Table 2). The occurrence of thermal stress days was presented for each month (July, August and September) in the last 20 years. Extreme heat stress (EHS), which refers to UTCI range above 46°C was presented separately when it occurred. Thermal indices, hot day (HD) and hot night (HN), were identified on the basis of the threshold of the maximum and minimum temperatures. Considering the influence of extreme temperatures, as stated by Collins et al. (2000) the following indices were marked in relation to the thresholds of maximum and minimum temperature: Hot days ( $HD=t_{max} \geq 35^{\circ}C$ ) and Hot nights ( $HN=t_{min} \geq 20^{\circ}C$ ) (Table 2).

For the last stage of the study, for the purpose of identifying thermal discomfort during the period observed, as an illustration of a threshold based on the duration of hot days (HD) and hot nights (HN) a hot day event is determined ( $HWE = 3$  HD days of  $t_{max} \geq 35^{\circ}C$ ). In addition, as an illustration of a threshold based on the duration of very strong heat stress (VSHS), a very strong heat stress event (VSHSE) is determined and it is caused by the occurrence of 5 consecutive VSHS days (Table 2).

As a result of comparing UTCI thermal stress to the selected thresholds of maximum and minimum temperatures, biothermal discomfort was identified during extremely high temperatures for the geographical area of Serbia. The recorded number of days with VSHS stress was compared to the number of days when the maximum air temperature was above 35°C.

**Table 2.** Definition of indices used in the study

Abbreviations	UTCI Indices	Definition
UTCI <sub>avg</sub>	Universal Thermal Climate Index	$UTCI_{avg} = f(t, f, v, t_{mrt})$
UTCI <sub>07h</sub>	Universal Thermal Climate Index <sub>07h</sub>	$UTCI_{07h} = f(t_{7h}, f_{7h}, v_{7h}, t_{mrt})$
UTCI <sub>14h</sub>	Universal Thermal Climate Index <sub>14h</sub>	$UTCI_{14h} = f(t_{14h}, f_{14h}, v_{14h}, t_{mrt})$
Abbreviations	UTCI heat stress indices	Definition
SHS	Strong Heat Stress	SHS= UTCI (32°C to 38°C)
VSHS	Very Strong Heat Stress	VSHS = UTCI (38°C to 46°C)
VSHSE	Very Strong Heat Stress Event	minimum 5 VSHS days in a row
EHS	Extreme Strong Stress	EHS= UTCI >46°C
Abbreviations	Thermal indices	Definition
HD	Hot days	$t_{max} \geq 35^{\circ}C$
HN	Hot nights	$t_{min} \geq 20^{\circ}C$
HWE	Heat waves event	minimum 3 HD days in a row

t-air temperature; f-relative humidity; v-wind speed; tmrt-mean radiant temperature; tmax-maximum air temperature, tmin-minimum air temperature

## Results

We have divided the section Results and discussion in two separate sections. According to the 3 stages of the study, described in revised Methodology (Data and indices considered in the study), the results are presented in three sections: (i) UTCI indices (UTCI<sub>avg</sub>, UTCI<sub>7h</sub>, UTCI<sub>14h</sub>), (ii) UTCI heat stress indices (strong heat stress SHS, very strong heat stress VSHS, extreme heat stress EHS) and thermal indices (hot day HD, hot night HN), (ii) Heat waves (HWE

and VSHSE). In all three sections the results are explained in more detail along with the modified and new added figures. Alongside the trend of UTCI index, the correlation between UTCI and maximum air temperature, is provided, too.

We have improved the figures and tables. Figures 2,3 and 4 have been changed and merged to form one, which became Figure 4 in the revised manuscript. Furthermore, one new figure (Figure 3 in the revised version) has been added in the first section of Results to represent the trend of the UTCI<sub>14h</sub> index. Table 3 has been supplemented with VSHSE index compared with HWE. In the revised manuscript it became Table 4.


The revised text reads as follows:

(lines 213-222) On the basis of the conducted bioclimatic analysis and comparison of biothermal conditions given in the two urban meteorological stations Novi Sad, Niš and Mt. Zlatibor, which represents a rural and lower mountain area, some differences in weather conditions were perceived in the summer period, as was expected. The results are presented in three sections: (i) UTCI indices (UTCI<sub>avg</sub>, UTCI<sub>7h</sub>, UTCI<sub>14h</sub>), (ii) UTCI heat stress indices (strong heat stress SHS, very strong heat stress VSHS, extreme heat stress EHS) and thermal indices (hot day HD, hot night HN), (iii) Heat waves (HWE and VSHSE).

*UTCI indices (UTCI<sub>avg</sub>, UTCI<sub>7h</sub>, UTCI<sub>14h</sub>)*

This section presents the results of the heat budget index UTCI calculated for mean daily data (UTCI<sub>avg</sub>) morning (UTCI<sub>7h</sub>) and midday (UTCI<sub>14h</sub>) for the period of 20 years.

(lines 228-235) These are the years when significant heat waves were recorded. The results depicted in Fig. 2 show the frequency (in percentages) of all stress categories for each index (UTCI<sub>avg</sub>, UTCI<sub>7h</sub>, UTCI<sub>14h</sub>) that occurred in July, August and September for the period that was observed.




**Figure 2.** Frequency of the days with different UTCI thermal stress categories for mean daily (UTCI<sub>avg</sub>), morning (UTCI<sub>7h</sub>) and midday (UTCI<sub>14h</sub>) index during the period from July to September, 1998-2017: a) Niš, b) Novi Sad, c) Zlatibor. X-axis: time (months), y-axis: frequency (number of days)

(lines 241-244) This is confirmed by Tošić and Unkasević (2014) in the study of dry periods in Serbia during the period between 1949 and 2011. It was found that the frequency of droughts in the southern part of Serbia was higher than in the other parts of the country. The most severe drought occurred in Niš and lasted from July 2006 to March 2008 (with a duration of 21 months and severity of 63.0).

(lines 247-291) Nevertheless, this does not rule out the occurrence of high temperatures and severe dry periods during the warm part of the year. This idea is based on the findings of Leščešen et al. (2018), when they analyzed drought periods in Vojvodina for over 60 years (1956-2016). All the regions of Vojvodina that were investigated experienced at least one extreme drought event over the reference period, in particular in 2000, 2001-2002, and 2011-2012. Moreover, the same results were obtained in other similar studies (Gocic and Trajković, 2013; 2014; Milanović, et al., 2014). On the mean daily level ( $UTCI_{avg}$ ) the most frequent category is “moderate heat stress” while on the hourly level  $UTCI_{14h}$  the most frequent category is “strong heat stress”. Days with “very strong heat stress” were observed in July and August (Fig. 2b) and there was one day with “extreme heat stress” in July 2007. Mt. Zlatibor has distinctive characteristics of the sub-mountain and mountain climate. Among these three stations, the bioclimatic conditions of Mt. Zlatibor are the most pleasant ones considering the dominant category of “no thermal stress” on the daily level of  $UTCI_{avg}$ . The thermal conditions with a morning  $UTCI_{7h}$  are similar to  $UTCI_{avg}$  while the  $UTCI_{14h}$  shows the prevailing categories of “moderate heat stress” and slightly less “strong heat stress” and “very strong heat stress”. It is important to emphasize the  $UTCI$  category of “slight cold stress” for all the indices ( $UTCI_{avg}$ ,  $UTCI_{7h}$ ,  $UTCI_{14h}$ ) was recorded several times during the three months of the period which was observed. The lowest value of the  $UTCI$  index was  $-2.84^{\circ}C$  and it occurred on September 5<sup>th</sup>, 2007 in Mt. Zlatibor as “moderate cold stress” (Fig. 2c).

Of all the calculated  $UTCI$  indices, mean daily data ( $UTCI_{avg}$ ), morning data ( $UTCI_{7h}$ ) and midday ( $UTCI_{14h}$ ) data, the midday  $UTCI_{14h}$  index shows the most extreme values for identifying heat waves. In this regard, for the further data analysis in this study, the  $UTCI_{14h}$  is used. To gain better insight into the  $UTCI_{14h}$  index values during the summer months, for the investigated period, averaged monthly  $UTCI_{14h}$  index for each year was provided for all three stations. The mean monthly values are presented in Fig. 3, together with the trends in  $UTCI_{14h}$  and mean absolute deviation of  $UTCI_{14h}$  in order to see how much the  $UTCI_{14h}$  index fluctuated in relation to the mean monthly value. The results show a significant increase in the extreme values of the  $UTCI_{14h}$  index over the last ten years, presented mainly as “very strong heat stress category” (VSHS). A series of peaks can be observed in the years that have been marked as extremely warm. In addition, there is a growing trend of the  $UTCI_{14h}$  index in all the summer months and all the weather stations, i.e. a series of peaks with successively higher values during the investigated period (1998-2017). The highest growing trend can be seen in September in Nis and Mt. Zlatibor, while the lowest growing trend can be observed in August on Zlatibor.

As regards the severity of daily  $UTCI$  index, there were 5 days recorded in Niš when the  $UTCI_{14h}$  exceeded the limit value for “extreme heat stress” (EHS). This happened on July 5<sup>th</sup>, 2000 ( $UTCI_{14h}=47.08^{\circ}C$ ); July 24<sup>th</sup>, 2007 ( $UTCI_{14h}=48.26^{\circ}C$ ); August 24<sup>th</sup>, 2007 ( $UTCI_{14h}=46.29^{\circ}C$ ); August 5<sup>th</sup> and 6<sup>th</sup>, 2017 ( $UTCI_{14h}=46.75$  and  $UTCI_{14h}=46.76^{\circ}C$ ). The maximum value of  $UTCI$  in Novi Sad was recorded on July 24<sup>th</sup>, 2007 ( $UTCI=48.42^{\circ}C$ ). In the area of Mt. Zlatibor, the “extreme heat stress” has not been recorded in the last 20 years. The highest values of  $UTCI$  in Zlatibor occurred on July 22<sup>nd</sup> and July 24<sup>th</sup>, 2007 ( $UTCI_{14h}=38.62^{\circ}C$  and  $UTCI_{14h}=38.37^{\circ}C$ ). The year 2007 is rated as the most unfavourable one, particularly the date of July 24<sup>th</sup>, when the highest temperature ever was recorded in Serbia (Smederevska Palanka,  $t_{max}=44,9^{\circ}C$ , source: Republic Hydro-meteorological Service of Serbia, RHMSS).



**Figure 3.** The mean monthly  $UTCI_{14h}$  index during the period which is investigated (1998-2017). x-axis: time (years), y-axis:  $UTCI$  ( $^{\circ}C$ )  
*Thermal indices and UTCI heat stress indices*

This section identifies midday  $UTCI_{14h}$  heat stress indices (strong heat stress SHS, very strong heat stress VSHS) and thermal indices marked in relation to the thresholds of the maximum and minimum temperatures in particular (hot day HD and hot night HN). Fig. 4 presents frequency (number of days) of hot days (HD), hot nights (HN), strong heat stress (SHS) and very strong heat stress (VSHS) for each month (July, August and September) in the last 20 years. The results of  $UTCI$  that were obtained show compatibility with previous analyses in relation to the occurrence of heat waves, especially since 2000 (Basarin et al., 2018; Unkašević and Tošić, 2009a; Unkašević and Tošić, 2009b; Unkašević and Tošić, 2013).

(lines 302-350)



**Figure 4.** Number of hot days (HD), hot nights (HN), strong heat stress (SHS) and very strong heat stress (VSHS) in Niš, Novi Sad and Zlatibor (July, Aug, Sep) 1998-2017. x-axis: time (years), y-axis: number of days

For the period investigated (1998-2017), the total number of hot days recorded in July, August and September in Niš was 126, 133 and 21 HDs, respectively, while Novi Sad saw about a half of the said number of hot days, namely 52, 63 and 3 HDs, respectively. In Mt. Zlatibor 2 HD days were recorded in July.

As regards hot nights (HN), there is generally a lower intensity of HNs compared to HDs. Generally, during the observed period up to 4 HNs were recorded, although an increase in the HNs has been observed in the last ten years. For example, in Niš during 2012, 2015 and 2017 the number of HNs recorded in July was 12, 7 and 7 respectively and during 2000, 2013, 2015, and 2017 the number of HNs recorded in August was 8, 6, 5 and 5 respectively. Since 2010 up to 2 HNs have been recorded in September. In Novi Sad, during 2010, 2012, 2015 and 2017 there were 7, 5, 5 and 6 HNs respectively recorded in July, while in August during 2012 and 2017 there were 5 and 7 HNs respectively. In September, one HN was recorded during 2009 and 2011 (Figure 4). The total number of hot nights recorded in July, August and September for the investigated period in Niš was 60, 56 and 6 HNs, respectively, while in Novi Sad there was a significantly lower number of hot nights – 41, 38 and 2 HNs, respectively. In Mt. Zlatibor 9 HN days were recorded in July and 11 HN days were recorded in August for the period which was investigated.

Midday  $UTCI_{14h}$  “strong heat stress” (SHS) occurs most commonly in all three stations. For example, SHS is particularly important to Mt. Zlatibor, located at an altitude above 1000 m since there has been an increase in the number of days with SHS in Zlatibor over the last decade. The best indicator is the increase in SHS in September. Similarly, the indicator of biothermal discomfort, “very strong heat stress” (VSHS) has been occurring more frequently in the last decade, reaching a maximum of 17 days in July 2007 and 2012 and in August 2015 in Niš. The number of VSHS days in Novi Sad reached a maximum of 12 days in July 2015 and 14 days in August 2015 and 2017. The total number of VSHS days recorded in July, August and September for the investigated period in Niš was 167, 174 and 25 VSHS days, respectively, while in Novi Sad there was a significantly lower number of VSHS days – 83, 97 and 7 VSHS days, respectively. In Mt. Zlatibor 2 VSHS days were recorded in July and 3 VSHS days were recorded in August. The recorded numbers of VSHS days indicate slightly greater biothermal discomfort duration in relation to the number of HD days. The HD and VSHS indices are significantly correlated during the summer months as they are directly derived from the maximum air temperatures and  $UTCI_{14h}$  index (Table 3).  $UTCI_{14h}$  and  $t_{max}$  for the Niš and Novi Sad urban stations have a significantly high linear correlation.

Table 3. Linear correlation of  $UTCI_{14h}$  and  $t_{max}$

$UTCI_{14h} \sim t_{max}$	Niš	Novi Sad	Zlatibor
July	0.88	0.97	0.85
August	0.92	0.95	0.77
September	0.89	0.92	0.41

For Zlatibor the correlation is slightly lower, and in September it is weak compared to the urban stations, but neither HD nor VSHS indices occurred on Zlatibor. Similar results of the linear relationship between the air temperature and the UTCI index ( $r = 83$ ;  $r = 0.98$ ) were shown in the studies of Urban and Kysely (2014) for summer in the urban areas of South Bohemia in Czech Republic and in the studies of Nassiri et al. (2017) for Iran.

#### Heat waves (HWE and VSHSE)

After UTCI thermal stress was compared to the selected thresholds of maximum temperature, biothermal discomfort was identified during extremely high temperatures for the geographical area of Serbia. During the observed period, a heat wave event (HWE) was determined, caused by 3 consecutive HD days and very strong heat stress event (VSHSE), caused by the occurrence of 5 consecutive VSHS days.

Table 4 lists the identified heat waves that refer to  $UTCI_{14h}$  VSHS index (VSHSE) and HD index (HWE). It has a very similar layout. However, shadowed differences can be observed, especially in the duration of the heat wave event. The results are presented for Niš and Novi Sad since there were no recorded heat waves in Mt. Zlatibor. More heat waves, both HWEs and VSHSEs, have been identified in the last ten years.

A heat wave event (HWE) of 10 days in a row (Niš, July from 15<sup>th</sup> to 24<sup>th</sup>, 2007) is the maximum number of consecutive days with such high temperatures when all the three stations are compared. In the same month, HWE of 3 days in a row (July 8<sup>th</sup> to 10<sup>th</sup>) and 4 days in a row (July 27<sup>th</sup> to 30<sup>th</sup>) were observed (Table 4, HWE). Altogether, the heat wave

events (HWEs) occurred 3 times and they lasted 3, 10 and 4 days in July 2007, which is certainly an extreme for the observed period.

(lines 364-371) **Table 4.** Number of Heat waves events (HWE) and Very strong heat stress event (VSHSE) in Niš, Novi Sad - July, Aug and Sep, 1998-2017

HWE	July		Aug		Sep	VSHSE	July		Aug		Sep
	Niš	Novi Sad	Niš	Novi Sad	Niš		Niš	Novi Sad	Niš	Novi Sad	Niš
1998	4	/	5	/	/	1998	7	/	5	/	/
1999	/	/	3, 4	/	/	1999	/	/	5	/	/
2000	3	/	4, 6	6	/	2000	/	/	6	7	/
2001	3	/	3, 3	/	/	2001	/	/	/	/	/
2002	6	/	/	/	/	2002	/	/	/	/	/
2003	4	/	/	/	/	2003	/	/	/	/	/
2004	3, 4	3	/	/	/	2004	5,5	/	/	/	/
2005	6	/	/	/	/	2005	6	/	/	/	/
2006	/	/	3	/	/	2006	/	/	/	/	/
2007	3, 10, 4	6	6	/	/	2007	10	7	/	/	/
2008	/	/	4	3	3	2008	/	/	5	/	/
2009	/	/	3	/	/	2009	/	/	/	/	/
2010	/	/	4	/	/	2010	/	/	5	/	/
2011	3	4	3	/	/	2011	/	/	/	/	/
2012	9	6	4, 6	3, 5	/	2012	12	6	5, 6	5, 5	/
2013	/	/	7	4	/	2013	/	/	8	8	/
2014	/	/	/	/	/	2014	/	/	/	/	/
2015	7	9	6, 4	4, 5	5, 3	2015	10	8	7	6	5
2016	3	/	/	/	/	2016	/	/	/	/	/
2017	4, 4	/	6, 4	6, 3, 3	/	2017	/	/	6, 5	6	/

Note: pink cells have more than one HWE and VSHSE duration defined by numbers of the consecutive days

Of the total number of hot days, it amounts to 21 HDs in September in Niš, only one HD was recorded in the first decade (1998-2007) of the investigated period, while the remaining HDs were recorded over the last decade (2008-2017), especially in 2011 and 2015. In the second decade of the investigated period the year 2015 stands out with two HWEs (September 1<sup>st</sup> to 5<sup>th</sup>, and 17<sup>th</sup> to 19<sup>th</sup>). In the same year the highest daily temperature was recorded in September (on September 18<sup>th</sup>,  $t_{max}=37.5^{\circ}C$ ).

(lines 381-396) Bioclimatic conditions analyzed by means of the UTCI index show that the calculated data observed at 14 CET are related to the marked heat wave events (HWE). When  $UTCI_{14h}$  thermal stress category of “very strong heat stress” (VSHS) is compared to the selected heat wave events (HWE,) a sub-index is defined. It is called very strong heat stress event (VSHSE) and it is caused by the occurrence of 5 consecutive VSHS days. The VSHSE corresponds to heat wave event and provokes a severe biothermal discomfort, so it was used as an indicator of extremely unfavorable bioclimatic conditions (biothermal heat hazard). Along these lines, the VSHSE in Niš occurred during July in 1998, 2004, 2005, 2007, 2012 and 2015 where 7, 5 (on two occasions), 6, 10, 12 and 10 VSHS days in a row were recorded respectively (Table 4, VSHSE). In August, VSHSE was recorded on 5 VSHS days in a row in 1998, 1999, 2008, 2010, 2012 and 2017, while in 2000, 2012, 2013, 2015 and 2017 6, 6, 8, 7 and 6 VSHS days in a row were recorded respectively. It should be emphasized that there were two VSHSE events in 2012 and 2017, for each year 5 and 6 VSHS days in a row. September 2015 saw a VSHSE with 5 VSHS days in a row. According to Unkašević and Tošić (2009b) the highest temperatures in Niš ever were recorded during the summer of 2007 (44.2°C) and the summer of 2000 (42.5°C), covering the data period from 1948 to 2007. The VSHSE occurred in Novi Sad in July 2007 (7 VSHS days in a row), 2012 (6 VSHS days in a row) and 2015 (8 VSHS days in a row) and in August 2000 (7 VSHS days in a row), 2013 (8 VSHS days in a row), 2015 (6 VSHS days in a row) and 2017 (6 VSHS days in a row). In August



2012 VSHSE occurred twice and each time 5 VSHS days in a row were recorded. There was no sub-index VSHSE recorded in Zlatibor.

## Discussion

Considering that the revised version has separated the results and the discussions, we need to point out the changes in the discussion section. According to the mention changes regarding Introduction chapter, we modified the paragraph followed by detailed recent studies (lines 47-58; lines 63-74; lines 77-94) and moved it into the discussion section (lines 479-494; lines 448-462; lines 463-469). More specifically, lines 47-58 have been moved into the discussion lines 479-494. In addition, lines 63-74 have been moved into the discussion lines 448-462, and there is an explanation of what a tropical day is (line 456-457) in the analysis by Unkasevic and Tosic (2009b). Lastly, lines 77-94 have been moved into the discussion section (lines 463-469).

The revised text is provided as follows:

### Discussion

(lines 398-430) The assessment of human biothermal conditions during the summer made it possible to identify biothermal discomfort related to heat waves in Serbia. The results obtained in the study for the investigated period from 1998 to 2017, indicates the general increase in biothermal discomfort associated with heat waves defined by maximum air temperature above 35°C and UTCI “very strong heat stress” above 38°C.

The results obtained for morning, midday and average UTCI show significant occurrence of thermal heat stress during the summer months and increase in biothermal discomfort especially for UTCI<sub>14h</sub>. This is confirmed by obtained distribution of the average monthly UTCI<sub>14h</sub> showing a significant increase in the extreme values of the UTCI<sub>14h</sub> index in the last ten years, presented mainly as “very strong heat stress category” (VSHS) together with a growing trend of the UTCI<sub>14h</sub> index in all the summer months (especially September) for all three investigated weather stations (primarily in Niš). According to Di Napoli et al. (2018), there are two thermal climates in Europe. One of them is associated with heat stress condition and it is predominant in the southern part of Europe (including the Balkans) when moderate and strong heat stress occurs at central day time hours, which reflects the general relationship between heat load and insolation.

The increase in extreme biothermal conditions is the most evident in the number of days with maximum air temperature thresholds above 35°C, defined in the study as hot days (HD) and in the number of days with UTCI index thresholds between 38°C and 46°C, defined in the study as “very strong heat stress” (VSHS). Thus, most HDs (18 days) occurred in 2007 and then in 2012 (16 days), which means that more than half of July in these years was characterised by extremely high temperatures. Similar results about increase in daily maximum air temperature were obtained by Unkašević and Tošić (2011), when the record values of the maximum temperatures were observed for almost the whole territory of Serbia in 2007. As reported by Papanastasiou et al. (2014), the summer of 2007 was the warmest summer in Athens in the last hundred years. Besides, Spinoni et al. (2015) in their study compiled a list of the most severe drought events which occurred in Europe from 1950 to 2012. They singled out the Balkan countries as extreme weather and drought-prone areas (particularly Serbia) with the longest drought registered in 2007, and the most severe drought event in 2011. Moreover, in 2012 the Balkan Peninsula and the south-eastern Europe faced the hottest summer and one of the worst droughts in nearly 40 years (Unkašević and Tošić, 2015). This is in relation with the investigation of heat waves (a sequence of at least 5 days with maximum temperature above the 95<sup>th</sup> annual percentile) in south-eastern Europe, where there was an increase in the number of hot days for the period from 1973 to 2010. The UTCI<sub>14h</sub> category of “very strong heat stress” (VSHS) correlates with the maximum air temperatures and there has been an increase in the number of such days over the last ten years. Particularly severe biothermal discomfort occurred in 2007 (18 VSHS days) and 2012 (17 VSHS days). This is in agreement with the statement of Di Napoli et al. (2018) about UTCI reference value from the period between 1979 and 2016, when the UTCI at 12:00 (UTC) was about 0.5°C colder in the period between 1980 and 1999, while it was 0.5°C warmer in the period between 2000 and 2009 and 1°C warmer in the period between 2010 and 2016.

(lines 437-443) In this sense, there is a difference between the definition of biothermal heat stress and potential heat stress, defined on the basis of the interpretation of maximum temperatures. Determining extreme weather event from the aspect of human heat budget indices allows other meteorological parameters to be taken into account. Apart from temperature and humidity, wind and direct solar radiation, have been considered together with physiological metabolic rate.

The identified heat waves based on the threshold of the UTCI index correspond to the identified heat waves based on the threshold of maximum air temperatures. Furthermore, there is a significant increase in heat waves (HWE and VSHSE) plus duration of such events over the last ten years in Niš and Novi Sad, especially in Niš.

(lines 449-470) This agrees with the findings of Unkašević and Tošić (2011, 2015) that there has been a growing trend of heat waves in Serbia, especially since 2000. In order to justify the present research for the last 20 years, the earlier results related to the increase in heat waves will be discussed in more detail. In that regard, previous research into heat waves in Niš for the period from 1949 and 2007, based on the autoregressive-moving-average model, observed the warmest years during three periods, 1951-1952, 1987-1998 and 2000-2007 (Unkašević and Tošić, 2009b). According to this research, the longest heat wave was recorded in Niš in 1952, with 21 days, while in 2003, 29 consecutive tropical days were observed. It should be taken into account that according to the Serbian National Weather Service, tropical days are days with a maximum temperature over 30°C. Furthermore, based on Heat Wave Duration index (HWD) i.e. daily maximum values of air temperature, Drljača et al. (2009) determined the duration and strength of heat waves in Niš during the summer season. The analysis showed Niš with a greater number of heat waves compared to the larger urban area of Belgrade. As stated in the research, since the mid-1980s, heat waves have had a higher frequency and on average they have occurred every year. Prior to that period heat wave fluctuations were generally reported in one of two years (Niš during the summer) (Drljača et al., 2009). While studying the characteristics of the heat waves in central Serbia (1949-2007), Unkašević and Tošić (2011) detected an increase in heat wave duration, in addition to an increase in heat wave frequency of occurrence during the period from 1999 to 2007.

As regards the heat budget index of Heat Load (HL) in Serbia (July) for the period from 2000 to 2010, it was noticed that in, Niš, Zlatibor and Novi Sad the category of “extremely hot” occurred in 2000 and 2007 (Milovanović et al., 2017). As for the heat budget index Physiological Equivalent Temperature PET for the period from 1949 to 2012, the highest number of heat waves in Novi Sad was observed in the last two decades and in the first decade of the investigated period (Basarin et al., 2016), while the number of the days above particular thresholds for the period 1961-2014 shows an increase along with the number of heat waves per year since 1981. The highest values of mean annual PET were mostly detected during the last 15 years (Basarin et al., 2018).

(lines 477-505) The biothermal discomfort identified in July and August as the hottest summer months in Serbia, could provoke health disorders more frequently. Specific HWs characteristics such as intensity and duration have been found to have devastating effects on human health and wellbeing. In certain cases heat waves cause problems and children, elderly people, chronic patients and workers are particularly susceptible to them. The analysis of the UTCI index correlating with mortality in Europe shows that deaths occur in the southern Europe (Italy, Portugal, Greece, Spain) for the categories of moderate and strong stress. Heat load has an impact on increasing mortality as conditions become more thermally stressful (Di Napoli et al., 2018). The intensity of heat stress in Poland was assessed by means of the UTCI index on the basis of daily mortality and the weather data for the period from 1991 to 2000. There is an increase in mortality on the days with strong and very strong heat stress, in relation to no thermal stress days, of 12% and 47%, respectively (Błażejczyk et al., 2017). Biothermal conditions and mortality rates show that in southern Europe during the summer thresholds of Physiological Subjective Temperature (PST) increase when moving from the west to the east. The general finding is that the population of central Europe is more sensitive to thermal stimuli in the summer than the citizens in the south of the continent (Błażejczyk and McGregor, 2008). The statement about the PST index is in support of the comparison of UTCI index with other thermal indices, including PST index, which correlates very well with UTCI index (Błażejczyk et al., 2012). Human sensitivity to extreme heat weather can also be seen in the impact of heat waves on daily mortality in Belgrade during the summer, when the strong correlation between heat waves and daily mortality can be observed in July 2007 (Stanojević et al., 2014). Similarly, Bogdanović et al. (2013) reported a significant short-term excess mortality on 16<sup>th</sup> July in Belgrade, when the maximum daily temperature exceeded 35°C, leading to 167 excess deaths (38% increase compared to the number of expected deaths) for nine consecutive days of heat, with a progressive return to almost normal mortality as the maximum temperature dropped below 35°C on 25<sup>th</sup> July.

The application of the standardized bioclimatic heat budget index UTCI could help improve the understanding of biothermal conditions relative to heat wave. The daily UTCI<sub>14h</sub> heat stress becomes more extreme in terms of severity and heat wave duration up to very strong heat stress (VSHS). The biothermal indices investigated for the three weather stations follow the trend of general warming. In the light of climatic changes and other negative factors resulting from this global phenomenon, it is becoming a true challenge to minimize their effects and improve living conditions in urban and rural areas (Stevović et al., 2017). However, the human heat budget indices are increasingly serving as a determinant of extreme weather condition (Di Napoli et al., 2018; Matzarakis et al., 2018; Theoharatos et al., 2010; Urban and Kyselý, 2014). UTCI index improves the understanding of extreme thermal stress, serving as one of the criteria for initiating heat alerts about extreme thermal stress in Serbia.

## Conclusion

We have made improvement in conclusion and revised text is provided as follows:

(lines 507-519) The present research of the human bioclimatic evaluation of UTCI thermal stress in Serbia is considered to be of great importance due to the identification of biothermal heat hazard and the study of the evolution of such indices regarding climate change. This evaluation aims at providing a comprehensive assessment of the human heat stress associated with UTCI index and heat waves defined by UTCI very strong heat stress and maximum air temperature.

The assessment of human biothermal conditions for the investigated period from 1998 to 2017 was provided for three synoptic stations – two urban (Niš and Novi Sad) and one station representing mountain areas of an altitude up to 1500 m (Zlatibor). All of them are located in different geographical areas in Serbia. The results obtained during the study indicate the significant increase in biothermal discomfort associated with heat waves defined by UTCI “very strong heat stress”. It was found that the most frequent heat stress categories are “strong heat stress” (SHS) and “very strong heat stress” (VSHS) and they have occurred in all three locations with the tendency of increase in the number of days in the last decade. The VSHS describes an alarming biothermal state and has occurred frequently in the last ten years, particularly in Niš and Novi Sad. The outcome of the study is UTCI<sub>14h</sub> sub-index “very strong heat stress event” (VSHSE) as an indicator of biothermal heat hazard. Undoubtedly, heat waves (HWs) are one of the natural hazards with increasing impact in urban areas related to higher population density.

(lines 523-527) Considering these facts, it can be deduced that heat waves are becoming more frequent, stronger and longer. Thus, frequent heat waves since 2007 certainly indicate biothermal heat hazard. The more lasting events of bioclimatic discomfort will indicate more stressful bioclimatic conditions for human health and wellbeing. However, the results of the study highlight the importance of UTCI index as a bioclimatic indicator of biothermal hazard in Serbia particularly if it can serve as one of the criteria for initiating heat warnings in Serbia.

## Reference

Considering that the revised version has significant changes the supplemented reference in the revised version is added as follows:

- ANSI/ASHRAE Standard 55, Thermal Environmental Conditions for Human Occupancy, (ANSI/ASHRAE Approved), [www.techstreet.com/ashrae/standards](http://www.techstreet.com/ashrae/standards), 2017.
- Di Napoli, C., Pappenberger, F., and Cloke, H.L.: Verification of Heat Stress Thresholds for a Health-Based Heat-Wave Definition, *J. Appl. Meteorol. Climatol.*, 58(6), 177- 1194, <https://doi.org/10.1175/jamc-d-18-0246.1>, 2019.
- Easterling, D.R., Horton, B., Jones, P.D., Peterson, T.C., Thomas, R.K., David, E.P., Salinger, M.J., Razuvayev, V., Plummer, N., Jamason, P., and Folland, C.K.: Maximum and minimum temperature trends for the globe. *Sci.*, 277(5324), 364–367, <https://doi.org/10.1126/science.277.5324.364>, 1997.
- European Commission's (Eurostat) datasets: <https://ec.europa.eu/eurostat/web/gisco/geodata/reference-data/>, last access: 3<sup>rd</sup> February 2020.

- Fanger, P.O.: Thermal Comfort: analysis and applications in environmental engineering, Danish Technical Press, Copenhagen, Denmark, 1970.
- Gocić, M. and Trajković, S.: Analysis of precipitation and drought data in Serbia over the period 1980–2010, *J. Hydrol.*, 494, 32–42, <https://doi.org/10.1016/j.jhydrol.2013.04.044>, 2013.
- Gocić, M. and Trajković, S.: Spatiotemporal characteristics of drought in Serbia, *J. Hydrol.*, 510, 110–123, <https://doi.org/10.1016/j.jhydrol.2013.12.030>, 2014.
- Höppe, P.: The physiological equivalent temperature - a universal index for the biometeorological assessment of the thermal environment, *Int. J. Biometeorol.*, 43, 71–75, <https://doi.org/10.1007/s004840050118>, 1999.
- IPCC, *Special Report of the Intergovernmental Panel on Climate Change, in: Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*, edited by Field, C. B., Barros, V., Stocker, T.F., Qin, D., Dokken, D.J., Ebi, K.L., Mastrandrea, M.D., Mach, K.J., Plattner, G.K., Allen, S.K., Tignor, M. and Midgley, P.M., Cambridge University Press, Cambridge, UK, 582, 2012.
- ISO 8996: Ergonomics of the thermal environment - Determination of metabolic rate, <https://www.iso.org/standard/34251.html>, 2004.
- Landsberg, H. E.: The Assessment of Human Bioclimate: a limited review of physical parameters, World Meteorological Organization WMO, 331, Geneva, 1972.
- Lee, V., Zermoglio, F., and Ebi, K.: Heat waves and human health - emerging evidence and experience to inform risk management in a warming world, Technical Report, United States Agency for International Development USAID, Washington 2019
- Leščešen, I., Dolijan, D., Pantelić, M., and Popov, S.: Drought assessment in Vojvodina (Serbia) using k-means cluster analysis, *J. Geogr. Inst. Jovan Cvijić SASA.*, 69(1), 17–27, <https://doi.org/10.2298/IJGI1901017L>, 2019.
- Matzarakis, A., Muthers, S., and Rutz, F.: Application and comparison of UTCI and PET in temperate climate conditions, *Finisterra*, 48(98), 21–31., <https://doi.org/10.18055/Finis6453>, 2014.
- Masterton, J. and Richardson, F.A.: Humidex, a method of quantifying human discomfort due to excessive heat and humidity, Atmospheric Environment Service Canada, Ontario, Downsview, 1979.
- McGregor, G. and Vanos, J.: Heat: A primer for public health researchers, *Public Health*, 161, 138–146, <https://doi.org/10.1016/j.puhe.2017.11.005>, 2018.
- Milewski, P.: 2013. Application of the UTCI to the local bioclimate of Poland's Ziemia Kłodzka region, *Geogr. Pol.*, 86(1), 47–54, <https://doi.org/10.7163/GPol.2013.6.2013>.
- Milovanović, M., Gocić, M., and Trajković, S.: Analysis of meteorological and agricultural droughts in Serbia, *Architecture and Civil Engineering*, 12(3), 253–264, <https://doi.org/10.2298/FUACE1403253M>, 2014.
- Nassiri, P., Monazzam, M.R., Golbabaei, F., Dehghan, S.F., Rafieepour, A., Mortezaipoor, A., Asghari, M.P.: Application of Universal Thermal Climate Index (UTCI) for assessment of occupational heat stress in open-pit mines, *Industrial Health*, 55 (5), 437–443, <https://doi.org/10.2486/indhealth.2017-0018>, 2017.
- Nastos, P. and Matzarakis, A.: The effect of air temperature and human thermal indices on mortality in Athens, Greece, *Theor. Appl. Climatol.*, 108, 591–599, <https://doi.org/10.1007/s00704-011-0555-0>, 2012.
- Nemeth, A.: Changing thermal bioclimate in some Hungarian cities, *Acta Climatologica et Chorologica*, 44–45, 93–101, 2011.
- Osczevski, R.J. and Bluestein, M.: The new wind chill equivalent temperature chart, *Bull. Am. Meteorol. Soc.*, 86, 1453–1458, <https://doi.org/10.1175/BAMS-86-10-1453>, 2005.
- Parsons, K.C.: Human thermal environments: the effects of hot, moderate, and cold environments on human health, comfort and performance, Taylor & Francis Group, Abingdon, England, UK, 2003.
- Rahmstorf, S. and Coumou, D.: Increase of extreme events in a warming world, *P. Natl. Acad. Sci.*, 108(44), 17905–17909, <https://doi.org/10.1073/pnas.1101766108>, 2011.
- Siple, P. and Passel, C.F.: Measurements of dry atmospheric cooling in subfreezing temperatures, *Proc Am Philos Soc*, 89(1), 177–199, [www.jstor.org/stable/985324](http://www.jstor.org/stable/985324), 1945.
- Spinoni, J., Naumann, G., Vogt, J., and Barbosa, P.: The biggest drought events in Europe from 1950 to 2012, *J. Hydrol. Reg. Stud.*, 3, 509–524, <https://doi.org/10.1016/j.ejrh.2015.01.001>, 2015.
- Seneviratne, S.I. and Nicholls, N.: Changes in climate extremes and their impacts on the natural physical environment, in: *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*,

edited by Field, C. B., Barros, V., Stocker, T.F., Qin, D., Dokken, D.J., Ebi, K.L., Mastrandrea, M.D., Mach, K.J., Plattner, G.K., Allen, S.K., Tignor, M. and Midgley, P.M., Cambridge University Press, Cambridge, UK, 109–230, 2012.

- Steadman, R.G.: A Universal Scale of Apparent Temperature, *J. Appl. Meteor.*, **23**, 1674-1687, [https://doi.org/10.1175/1520-0450\(1984\)023<1674:AUSOAT>2.0.CO;2](https://doi.org/10.1175/1520-0450(1984)023<1674:AUSOAT>2.0.CO;2), 1984.
- Tolika, K., Maheras, P., Pytharoulis, I., and Anagnostopoulou, C.: The anomalous low and high temperatures of 2012 over Greece - an explanation from a meteorological and climatological perspective, *Nat. Hazards Earth Syst. Sci.*, 14(3), 501–507, <https://doi.org/10.5194/nhess-14-501-2014>, 2014.
- Tomczyk, M.A. and Owczarek M.: Occurrence of strong and very strong heat stress in Poland and its circulation conditions, *Theor. Appl. Climatol.*, 139, 893-905, <https://doi.org/10.1007/s00704-019-02998-3>, 2020.
- Tošić, I. and Unkašević, M.: Analysis of wet and dry periods in Serbia, *Int. J. Climatol.*, 34, 1357-1368, <https://doi.org/10.1002/joc.3757>, 2014.
- Urban, A. and Kyselý J.: Comparison of UTCI with other thermal indices in the assessment of heat and cold effects on cardiovascular mortality in the Czech Republic, *Int. J. Environ. Res. Public Health*, 11(1), 952-967, <https://doi.org/10.3390/ijerph110100952>, 2014.
- Yaglou, C.P. and Minard, D.: Control of heat casualties at military training centers, *Am Med Assoc Arch Ind Health*, 16(4), 302–316, 1957.

## English

With regard to English, all the spelling and grammatical errors have been corrected.

We look forward to hearing from you in due time regarding our submission and responding to any further questions and comments you may have.

# Analysis of UTCI index during heat waves in Serbia

Milica M. Pecelj<sup>123</sup>, Milica Z. Lukić<sup>4</sup>, Dejan J. Filipović<sup>4</sup>, Branko M. Protić<sup>4</sup>, Uroš M. Bogdanović<sup>5</sup>

<sup>1</sup>Geographical Institute Jovan Cvijić Serbian Academy of Science and Arts, Belgrade, Serbia

<sup>2</sup>Department of Geography, University of East Sarajevo, East Sarajevo, RS, Bosnia and Herzegovina

<sup>3</sup>South Ural State University, Institute of Sports, Tourism and Service, Chelyabinsk, Russia

<sup>4</sup>University of Belgrade – Faculty of Geography, Belgrade, Serbia

<sup>5</sup>University of Belgrade, Faculty of Organisational Sciences, Belgrade, Serbia

Correspondence to: Milica M. Pecelj (milicapecelj@gmail.com)

**Abstract.** The objective of this paper is to assess the bioclimatic conditions in Serbia so as to identify biothermal heat hazard. Special emphasis is placed on the bioclimatic index UTCI (Universal Thermal Climate Index), whose purpose is to evaluate the degree of thermal stress that the human body is exposed to. For this research, mean daily and hourly (7 and 14 CET) meteorological data from three weather stations (Mt. Zlatibor, Novi Sad, Niš) have been collected for the period from 1998 to 2017. In order to identify patterns of biothermal heat stress conditions, the thresholds of daily UTCI index ( $UTCI \geq 38^\circ\text{C}$  refers as very strong heat stress-VSHS) at 14 CET, are compared with the thresholds of daily maximum of air temperature ( $t_{\text{max}} \geq 35^\circ\text{C}$  refers as hot day-HD) which are further termed as heat wave events (VSHSE and HWE). The findings show that the UTCI heat stress category “very strong heat stress” at 14 CET indicates heat waves. The most extreme heat wave events occurred in 2007, 2012, 2015 and 2017. Moreover, three heat wave events (HWE) in Niš that occurred in July 2007 lasted 3, 10 and 4 days. Heat wave events (HWE) and very strong heat stress events (VSHSE) recorded in July 2007 (lasted 10 days each), 2012 (lasted 9 and 12 days) 2015 (lasted 7 and 10 days) were of the longest duration and refer as indicators of biothermal heat hazard. The daily  $UTCI_{14h}$  heat stress becomes more extreme in terms of severity and heat wave duration up to very strong heat stress (VSHS).

**Key words:** Bioclimatic conditions, Universal Thermal Climate Index (UTCI), heat waves, VSHS.

## 1. Introduction

Extreme weather events such as heat waves, floods, droughts or storms have shown an increased frequency in recent decades (Brown et al., 2008; Easterling et al., 1997; IPCC, 2012; Rahmstorf and Coumou, 2011; Seneviratne and Nicholls, 2012). Not only are heat and drought events of great importance in most Mediterranean climate regions, but also in most of southern and south-eastern Europe, because of the diverse and costly impact that they have on various economic sectors and on the environment (Peña-Gallardo et al., 2019). Extremely high temperatures, especially during the summer months, beg the question as to how heat could affect everyday life of humans. Heat waves as a hazard often have a negative effect, causing heat stress in the human body. They have both direct effects on human health, affecting the body’s physiological responses and functions and indirect effects on human health, increasing challenges to food and water safety (Lee et al., 2019). Bioclimatic condition provides a base for considering the effects of climatic conditions on humans and gives special importance to the social factors that mitigate or reinforce the consequences of environmental changes (Bleta et al., 2014). The impact of the weather and climate on humans is most commonly described as the biothermal condition of a certain area, presented by bioclimatic indices. Over the last century many models and indices have been developed for the assessment of human exposure to heat, ranging from simple physical instruments designed to imitate the human heat exchange with the environment, to complex thermophysiological models that simulate external and internal body heat transfer and allow detailed simulation of different work load, clothing, and climate scenarios (Havenith and Fiala, 2015). According to Epstein et al. (2006), there are over 100 heat stress indices that could describe extreme bioclimatic conditions in relation to humans. Many thermal indices have been developed for the purpose of describing the complex conditions of heat exchange between the human body and its thermal environment (Fanger,

1970; Landsberg, 1972; Parsons, 2003). Those are two meteorological parameter indices – the ones used for cold environment conditions combining the air temperature and the wind velocity (Osczevski and Bluestein, 2005; Siple and Passel, 1945;), and the ones used for heat environment conditions combining the air temperature and humidity (Masterton and Richardson, 1979; Steadman, 1984; Yaglou and Minard, 1957). Considering their shortcomings, i.e. the fact that they are not universally applicable to all climatic regions, including different seasons, the dominance in the analysis of biothermal conditions and thermal stress over the last 30 years has been taken over by so-called heat budget indices (Błażejczyk, 1994; Hoppe, 1999; Jendritzky, 2012). The heat budget indices are based on the thermal exchange between humans and the environment. Methodologically, they use variables related to meteorological (air temperature, wind velocity, radiation, air humidity) and physiological processes (most commonly metabolic heat) and clothing insulation. The issue that should be emphasized during the analysis of the thermal stress is the influence that extreme temperatures have on the physiological parameters in humans. According to McGregor and Vanos (2018), generated heat load can undermine the human body's ability to hold its core temperature within the range of optimal physiological achievement.

For the purpose of estimating the thermal effect of the environment on the human body, the total effects of all the thermal components are considered by the UTCI index. The UTCI is heat budget index considering both physiological and meteorological parameters describing the physiological heat stress that the human body experiences while achieving thermal equilibrium with the surrounding outdoor environment (Błażejczyk et al., 2013). Compared to other indices, UTCI is more sensitive to changes in all of the environment parameters, particularly air temperature, solar radiation, humidity, and wind speed (Błażejczyk et al., 2012). According to Jendritzky et al. (2012) the UTCI evaluates the outdoor thermal environment for biometeorological applications by simulating the dynamic physiological response with a model of human thermoregulation together with a modern clothing model. UTCI provides a multiple opportunity for investigation and has a wide application in weather forecast for outdoor activities, appropriate behavior and climate therapies, and extreme thermal stress alerts. These analyses can have application in tourism, health sector, and urban and regional planning (Jendritzky et al., 2012). In the past decade, a large number of publications in Europe concern analyses of bioclimatic condition in accordance with the UTCI index (Błażejczyk et al., 2014; Błażejczyk et al., 2017; Bleta et al., 2014; Di Napoli et al., 2018; Di Napoli et al., 2019; Matzarakis et al., 2014; Milewski, 2013; Nastos and Matzarakis 2012; Nemeth, 2011; Tomczyk and Owczarek, 2019; Urban and Kysely, 2014).

Although over recent years in Serbia articles have been published containing analysis of bioclimatic condition conducted by means of other heat budget indices (Basarin et al., 2016; Basarin et al., 2018; Lukić et al., 2019; Milovanović et al., 2017; Pecelj et al., 2017; Pecelj et al., 2018; Stojicević et al., 2016), detailed bioclimatic condition has not yet been thoroughly analysed by means of the UTCI index, especially in the case of heat waves. The heat wave is a prolonged period of unusually high air temperature that causes changes in everyday life and may cause health and wellbeing effects of heat stress. Even though a heat wave is understood as a meteorological event, its significance and influence could not be adequately presented without showing its clear impact on man (Robinson, 2001). There is a need to study the evolution of such indices regarding heat stress that affects the human thermoregulatory system as a result of heat exchange between the body and its thermal environment.

Considering extreme temperature is one of the most significant climatic parameters in the universal context of climate change, analyses of heat waves in Serbia have been performed in several different approaches and all of them show a growing trend in max temperature (Drljača et al., 2009; Unkašević and Tošić, 2009a; Unkašević and Tošić, 2009b; Unkašević and Tošić, 2013) and heat waves (Unkašević and Tošić, 2011; Malinović-Milićević et al., 2013; Unkašević and Tošić, 2015, Milićević-Malinović et al., 2016) especially since 2000.

Keeping this fact in mind, the basic idea of this study is to conduct the analysis of human bioclimatic conditions in Serbia in the last 20 years, where according to the previous studies mentioned above, warming has been perceived and recorded, especially since 2000. Determining extreme weather event from the aspect of heat budget indices allows other meteorological parameters to be taken into account because high temperatures and humidity generate a heat load more rapidly, unless the wind and direct radiation are taken into account. The analysis of bioclimatic conditions implies the determination of UTCI thermal stress during the summer months and different heat stress

that has occurred in the past 20 years in different geographical landscapes in Serbia. In addition to UTCI thermal stress, extreme temperature thresholds were analysed in order to observe the biothermal stress related to the heat wave. The conducted human bioclimatic evaluation of UTCI thermal stress in Serbia was considered to be of great importance due to the identification of biothermal heat hazard and the study of the evolution of such indices regarding climate change. This evaluation is aimed at addressing the following topics in this study:

- providing a comprehensive assessment of the human heat stress associated with UTCI index and
- comparing it to heat waves defined by meteorological parameters.

#### a. Study area

The Republic of Serbia is located in south-eastern Europe, in the area of the southern Pannonian Plain and the central Balkans. Northern Serbia is mainly flat, while its central and southern areas consist of highlands and mountains (Gocić and Trajković, 2014), as the map of the relief characteristics shows (Figure 1). The territory of Serbia is characterised by temperate continental and mountain climate and the spatial distribution of climate parameters is determined by geographical location. The research involved studying three synoptic stations located in different geographical areas of Serbia: (i) Novi Sad, (ii) Niš and (iii) Mt. Zlatibor (Figure 1).

#### Figure 1.

The first weather station (at an altitude of 86 m) is located in the territory of Novi Sad, which is the administrative urban center of Vojvodina province and South Bačka District (Figure 1<sub>1</sub>). The city is located in the southern part of the Pannonian Basin, on the Danube River bank near Mt. Fruška Gora, and the national park bearing the same name. Novi Sad has a temperate continental climate, summers are warm and winters are cold with a small amount of snow (Lazić et al., 2006). The second station is Niš (at an altitude of 202 m) and it is located in the Niš Fortress. This city is the administrative urban center of the Nišava District in southern Serbia and it is situated in the Nišava valley, located in the central part of the spacious geological depression called the Nišava Basin (Figure 1<sub>2</sub>). According to Köppen's climate classification, the Nišava valley belongs to the Cfwx type – the Danube type of moderately warm and humid climate characterized by hot summers (the highest precipitation is recorded at the beginning of the summer) and somewhat dry winters (Prokić, 2018). Compared to other cities in this valley (Dimitrovgrad, Pirot and Bela Palanka), Niš is the hottest one with an average annual temperature of 11.8°C (Prokić, 2018). The third station is Zlatibor, a mountain in western Serbia, which belongs to the mountain range of the Dinaric Alps (Figure 1<sub>3</sub>). Zlatibor weather station lies at an altitude of 1029 m a.s.l. In the area of Zlatibor plateau mountains meet air currents from the Adriatic Sea, which, as it can be assumed, creates a favorable climate and for this reason Zlatibor has already been confirmed as a climatic resort (Pecelj et al., 2017). The mountain climate of Zlatibor involves long and cold winters, short and fresh summers, and less pronounced spring and autumn.

## 2. Materials and methods

The present study implements the methodological approach of Universal Thermal Climate Index (UTCI). As a thermal comfort indicator, the UTCI considers combined meteorological and physiological parameters describing thermal comfort through the evaluation of human energy balance. In terms of physiological conditions, the metabolic rate plays an important role. Metabolic processes in the human body create heat that is constantly exchanged with the environment achieving a state of thermal equilibrium in the body for maintaining a constant body temperature. The amount of the heat that is produced and released depends on the following: physical activity, clothing, sex, age, body mass, diet, mental state, health, external conditions, acclimatization, etc. As a measure of physical activity, a unit of "MET" was defined, which corresponds to the release of human heat of 58.2 W.m<sup>-2</sup> for average body surface area (1.8 m<sup>2</sup>), i.e. it is equal to the energy rate produced per unit surface area of an average person seated at rest



(ANSI/ASHRAE Standard 55). According to ISO 8996 for standard applications, the metabolic heat energy is  $M = 135 \text{ W}\cdot\text{m}^{-2}$  i.e. 2.3 MET, for a person moving at a speed of  $1.1 \text{ m}\cdot\text{s}^{-1}$ .

#### a. The Universal Thermal Climate Index (UTCI)

For valid assessment of the outdoor thermal environment in the fields of public weather services, public health systems, urban planning, tourism and recreation and climate impact research, the Universal Thermal Climate Index (UTCI) has been developed based upon the most recent scientific progress in human thermo-physiology in biophysics and heat exchange theory. The UTCI is the result of an approach which was developed in the International Society of Biometeorology (ISB) Commission 6 and was later improved by COST Action 730 (Jendritzky et al., 2012). The design of UTCI is of great importance due to the fact that it is applicable to all seasons and climates together with all spatial and temporal scales.

Human reaction was simulated by the UTCI-Fiala multi-node model of human thermoregulation, which was integrated with an adaptive clothing model. For any combination of meteorological parameters (Eq. 1), based on the conception of an equivalent temperature, the UTCI is the air temperature of the reference environment which, according to the model, produces an equivalent dynamic physiological response under a set of reference conditions (Bröde et al., 2012). In other words, this model simulates the same sweat production or skin wettedness in human body response as the actual environment condition (Błażejczyk et al., 2013; Błażejczyk et al., 2014). This is derived from the multi-dimensional dynamic response of a state-of-the-art multi-node thermo-physiological model of human heat transfer and thermoregulation (Fiala et al., 2012). The UTCI can be represented in general function as below:

$$\text{UTCI} = f(t, f, v, t_{\text{mrt}}) \quad (\text{Eq.1})$$

UTCI = f (air temperature, relative humidity, wind speed, mean radiant temperature)

For appropriate use, UTCI can only be approximated using a regression equation abbreviated from sample calculations performed by computing centers (Bröde et al., 2012; Jendritzky et al., 2012). It causes a narrow range of input parameters it can manage with. For the reference environment, the ISB Commission on the UTCI, was decided to use: (i) a wind speed ( $v$ ) of  $0.5 \text{ m}\cdot\text{s}^{-1}$  at 10 m height (approximately  $0.3 \text{ m}\cdot\text{s}^{-1}$  at 1.1 m), (ii) a mean radiant temperature ( $T_{\text{mrt}}$ ) equal to air temperature, (iii) vapour pressure ( $vp$ ) that represents a relative humidity ( $f$ ) of 50%; at high air temperatures ( $>29^\circ\text{C}$ ) the reference humidity was taken to be constant at 20 hPa (Błażejczyk et al., 2013). Physiological parameters, the metabolic rate ( $M$ ) and the thermal properties of clothing (clothing insulation, permeability) are taken as universal constants in the model due to the evaluation by means of the regression equation. This implies an outdoor activity of a person walking at the speed of  $4 \text{ km}\cdot\text{h}^{-1}$  ( $1.1 \text{ m}\cdot\text{s}^{-1}$ ), corresponding to heat production of  $135 \text{ W}\cdot\text{m}^{-2}$  (2.3 MET) of metabolic energy (Błażejczyk et al., 2013; Jendritzky et al., 2012) and the clothing insulation which is self-adapting according to the environmental conditions (Havenith et al., 2012). Clothing insulation, vapor resistance and the insulation of surface air layers, are strongly influenced by changes in wind speed and body movement and will therefore also influence physiological responses (Błażejczyk et al., 2012). Particular ranges of UTCI are categorised according to thermal stress (Table 1).

Table 1.

#### b. Data and indices considered in the study

The meteorological data set from the period from 1998 to 2017 was recorded for two urban weather stations (Novi Sad and Niš) and one rural mountain station (Zlatibor). Mean daily and hourly meteorological parameters (07h, 14h, CET) of air temperature ( $t$ ), air humidity ( $f$ ), and wind speed ( $v$ ) from the above weather stations were considered for the calculation of particular UTCI thermal heat stress in the summer months (July, August and September). The

meteorological data set used in the study was retrieved from the Meteorological Yearbook for the period from 1998 to 2017 (Republic Hydrometeorological Service of Serbia) while the UTCI index was calculated by applying the BioKlima 2.6 software package (available at <http://www.igipz.pan.pl/Bioklima-zgik.html>).

Although extreme temperatures are one of the most effective climatic parameters in the universal case of hot days and heat waves, the influence of other parameters, especially clothing insulation and metabolic rate, can justify the thermal discomfort and the other way round. In that regard, the summer (July, August and September) daily maximum and minimum temperatures ( $t_{max}$ ,  $t_{min}$ ), for the same stations, were analysed in terms of particular thresholds to identify the frequency of extremely high temperatures and heat waves in the period observed to compare them with the discomfort recorded by the UTCI index.

The first stage of the study presented mean daily ( $UTCI_{avg}$ ), morning ( $UTCI_{7h}$ ) and midday ( $UTCI_{14h}$ ) UTCI indices. They are presented by months (July, Aug, Sep) for each station to identify the general differences caused by the geographical origin of the stations under study. The quantification of human bioclimatic conditions in Serbia was designed for the UTCI indices on a daily basis of each defined stress categories. To obtain a better insight into the UTCI index values during the summer months, averaged monthly  $UTCI_{14h}$  index was provided for each year to see how much the  $UTCI_{14h}$  index fluctuated in relation to the average monthly value.

For the second stage, midday  $UTCI_{14h}$  thermal stress indices were identified (strong heat stress SHS, very strong heat stress VSHS, extreme heat stress EHS) and thermal indices based on the maximum and minimum temperatures (hot day HD and hot night HN). During the midday observation (14 CET), particularly prominent are strong heat stress (SHS), which refers to UTCI range of 32°C to 38°C and very strong heat stress (VSHS), which refers to UTCI range of 38°C to 46°C (Table 2). The occurrence of thermal stress days was presented for each month (July, August and September) in the last 20 years. Extreme heat stress (EHS), which refers to UTCI range above 46°C was presented separately when it occurred.

Thermal indices, hot day (HD) and hot night (HN), were identified on the basis of the threshold of the maximum and minimum temperatures. Considering the influence of extreme temperatures, as stated by Collins et al. (2000) the following indices were marked in relation to the thresholds of maximum and minimum temperature: Hot days ( $HD=t_{max} \geq 35^{\circ}C$ ) and Hot nights ( $HN=t_{min} \geq 20^{\circ}C$ ) (Table 2). It should be taken into account that according to the Serbian National Weather Service, a daily maximum temperature equal to or greater than 30°C is taken as the threshold for what is called tropical days and is mainly used for the analysis of heat waves.

For the last stage of the study, with the purpose of identifying thermal discomfort during the period observed, as an illustration of a threshold based on the duration of hot days (HD) and hot nights (HN) a hot day event is determined ( $HWE = 3$  HD days of  $t_{max} \geq 35^{\circ}C$ ). In addition, as an illustration of a threshold based on the duration of very strong heat stress (VSHS), a very strong heat stress event (VSHSE) is determined and it is caused by the occurrence of 5 consecutive VSHS days (Table 2).

As a result of comparing UTCI thermal stress to the selected thresholds of maximum and minimum temperatures, biothermal discomfort was identified during extremely high temperatures for the geographical area of Serbia. The recorded number of days with VSHS stress was compared to the number of days when the maximum air temperature was above 35°C.

## Table 2.

### 3. Results

On the basis of the conducted bioclimatic analysis and comparison of biothermal conditions given in the two urban meteorological stations Novi Sad, Niš and Mt. Zlatibor, which represents a rural and lower mountain area, some differences in weather conditions were perceived in the summer period, as was expected. The results are presented in three sections: (i) UTCI indices ( $UTCI_{avg}$ ,  $UTCI_{7h}$ ,  $UTCI_{14h}$ ), (ii) UTCI heat stress indices (strong heat stress SHS, very strong heat stress VSHS, extreme heat stress EHS) and thermal indices (hot day HD, hot night HN), (iii) Heat waves (HWE and VSHSE).

#### a. UTCI indices ( $UTCI_{avg}$ , $UTCI_{7h}$ , $UTCI_{14h}$ )

This section presents the results of the heat budget index UTCI calculated for mean daily data ( $UTCI_{avg}$ ) morning ( $UTCI_{7h}$ ) and midday ( $UTCI_{14h}$ ) for the period of 20 years. In particular, the physiological stress in the category of “strong heat stress” occurred quite frequently in 1998, 2000, 2004, 2007, 2011, 2012, 2015 and 2017 in Niš (July and August) and slightly less frequently in Novi Sad, where “strong heat stress” occurred more frequently in July and August of 2000, 2002, 2006, 2007, 2009, 2013, 2015, 2016 and 2017. A much smaller amount of “strong heat stress” was observed in Mt. Zlatibor. There were periods with the categories of “moderate” and “strong heat stress” evenly distributed in the months of July and August over the period that was investigated. It is important to emphasize that the category of “strong heat stress” most frequently occurred in Mt. Zlatibor in 2000, 2004, 2007, 2012, 2015 and 2017, which generally coincided with the other two stations. These are the years when significant heat waves were recorded. The results depicted in Fig. 2 show the frequency (in percentages) of all stress categories for each index ( $UTCI_{avg}$ ,  $UTCI_{7h}$ ,  $UTCI_{14h}$ ) that occurred in July, August and September for the period that was observed.

#### Figure 2.

Bioclimatic conditions in Niš are certainly the most unpleasant ones, given the domination of “strong heat stress” and not so rare “very strong heat stress” categories during the period of 20 years. There are 5 days observed with “extreme heat stress” for midday  $UTCI_{14h}$ . The most pleasant bioclimatic conditions can be observed in the morning hours ( $UTCI_{7h}$ ), when the dominant categories are “moderate heat stress” (July and August) and “no thermal stress” in September (Fig. 2a). Niš is located in the south of Serbia and belongs to the areas that are most endangered by drought and extreme high temperatures in Serbia. This is confirmed by Tošić and Unkasević (2014) in the study of dry periods in Serbia during the period between 1949 and 2011. It was found that the frequency of droughts in the southern part of Serbia was higher than in the other parts of the country. The most severe drought occurred in Niš and lasted from July 2006 to March 2008 (with a duration of 21 months and severity of 63.0).

Novi Sad is located in the northern part of Serbia, on the slopes of Mt. Fruška Gora and is characterized by a temperate continental and continental climate. Therefore, bioclimatic conditions in Novi Sad are more pleasant than those in Niš. Nevertheless, this does not rule out the occurrence of high temperatures and severe dry periods during the warm part of the year. This idea is based on the findings of Leščešen et al. (2018), when they analyzed drought periods in Vojvodina for over 60 years (1956-2016). All the regions of Vojvodina that were investigated experienced at least one extreme drought event over the reference period, in particular in 2000, 2001-2002, and 2011-2012. Moreover, the same results were obtained in other similar studies (Gocic and Trajković, 2013; 2014; Milanović, et al., 2014). On the mean daily level ( $UTCI_{avg}$ ) the most frequent category is “moderate heat stress” while on the hourly level  $UTCI_{14h}$  the most frequent category is “strong heat stress”. Days with “very strong heat stress” were observed in July and August (Fig. 2b) and there was one day with “extreme heat stress” in July 2007. Mt. Zlatibor has distinctive characteristics of the sub-mountain and mountain climate. Among these three stations, the bioclimatic conditions of Mt. Zlatibor are the most pleasant ones considering the dominant category of “no thermal stress” on the daily level of  $UTCI_{avg}$ . The thermal conditions with a morning  $UTCI_{7h}$  are similar to  $UTCI_{avg}$  while the  $UTCI_{14h}$  shows the prevailing categories of “moderate heat stress” and slightly less “strong heat stress” and “very strong heat stress”. It is important to emphasize the UTCI category of “slight cold stress” for all the indices ( $UTCI_{avg}$ ,  $UTCI_{7h}$ ,  $UTCI_{14h}$ ) was recorded several times during the three months of the period which was observed. The lowest value of the UTCI index was  $-2.84^{\circ}\text{C}$  and it occurred on September 5<sup>th</sup>, 2007 in Mt. Zlatibor as “moderate cold stress” (Fig. 2c).

Of all the calculated UTCI indices, mean daily data ( $UTCI_{avg}$ ), morning data ( $UTCI_{7h}$ ) and midday ( $UTCI_{14h}$ ) data, the midday  $UTCI_{14h}$  index shows the most extreme values for identifying heat waves. In this regard, for the further data analysis in this study, the  $UTCI_{14h}$  is used. To gain better insight into the  $UTCI_{14h}$  index values during the summer months, for the investigated period, averaged monthly  $UTCI_{14h}$  index for each year was provided for all three stations. The mean monthly values are presented in Fig. 3, together with the trends in  $UTCI_{14h}$  and mean absolute deviation of  $UTCI_{14h}$  in order to see how much the  $UTCI_{14h}$  index fluctuated in relation to the mean monthly value.

The results show a significant increase in the extreme values of the  $UTCI_{14h}$  index over the last ten years, presented mainly as “very strong heat stress category” (VSHS). A series of peaks can be observed in the years that have been marked as extremely warm. In addition, there is a growing trend of the  $UTCI_{14h}$  index in all the summer months and all the weather stations, i.e. a series of peaks with successively higher values during the investigated period (1998-2017). The highest growing trend can be seen in September in Nis and Mt. Zlatibor, while the lowest growing trend can be observed in August on Zlatibor.

As regards the severity of daily  $UTCI$  index, there were 5 days recorded in Niš when the  $UTCI_{14h}$  exceeded the limit value for “extreme heat stress” (EHS). This happened on July 5<sup>th</sup>, 2000 ( $UTCI_{14h}=47.08^{\circ}C$ ); July 24<sup>th</sup>, 2007 ( $UTCI_{14h}=48.26^{\circ}C$ ); August 24<sup>th</sup>, 2007 ( $UTCI_{14h}=46.29^{\circ}C$ ); August 5<sup>th</sup> and 6<sup>th</sup>, 2017 ( $UTCI_{14h}=46.75$  and  $UTCI_{14h}=46.76^{\circ}C$ ). The maximum value of  $UTCI$  in Novi Sad was recorded on July 24<sup>th</sup>, 2007 ( $UTCI=48.42^{\circ}C$ ). In the area of Mt. Zlatibor, the “extreme heat stress” has not been recorded in the last 20 years. The highest values of  $UTCI$  in Zlatibor occurred on July 22<sup>nd</sup> and July 24<sup>th</sup>, 2007 ( $UTCI_{14h}=38.62^{\circ}C$  and  $UTCI_{14h}=38.37^{\circ}C$ ). The year 2007 is rated as the most unfavourable one, particularly the date of July 24<sup>th</sup>, when the highest temperature ever was recorded in Serbia (Smederevska Palanka,  $t_{max}=44,9^{\circ}C$ , source: Republic Hydro-meteorological Service of Serbia, RHMSS).

### Figure 3.

#### b. Thermal indices and $UTCI$ heat stress indices

This section identifies midday  $UTCI_{14h}$  heat stress indices (strong heat stress SHS, very strong heat stress VSHS) and thermal indices marked in relation to the thresholds of the maximum and minimum temperatures in particular (hot day HD and hot night HN). Fig. 4 presents frequency (number of days) of hot days (HD), hot nights (HN), strong heat stress (SHS) and very strong heat stress (VSHS) for each month (July, August and September) in the last 20 years. The results of  $UTCI$  that were obtained show compatibility with previous analyses in relation to the occurrence of heat waves, especially since 2000 (Basarin et al., 2018; Unkašević and Tošić, 2009a; Unkašević and Tošić, 2009b; Unkašević and Tošić, 2013).

In accordance with the thresholds of maximum temperatures in July, there were several years in Niš with 10 or more HDs in July, when the temperature was above  $35^{\circ}C$ , as shown in Fig. 4. This can be observed in July of 2000, 2007, 2012, 2015, and 2017, and in total, for each year it amounts to 10, 18, 16, 11 and 11 HD days, respectively. As regards the thresholds of maximum temperature in Novi Sad, there were 10 or more HD days with temperatures above  $35^{\circ}C$ , more specifically 10 and 11 HD days during July of 2012 and 2015, respectively. In Mt. Zlatibor only two HD days were recorded in July 2007. In August, this can be observed in 2000, 2007, 2012, 2013, 2015, and 2017 and in total it amounts to 12, 12, 10, 10, 13 and 13 HDs days respectively for each year in Niš. The highest number of days with a temperature over  $35^{\circ}C$  degrees was recorded in 2000, 2012, 2015 and 2017 and in total for each year it amounts to 9, 8, 11 and 13, respectively. In Mt. Zlatibor there are no HD days recorded in August (Figure 4).

### Figure 4.

For the period investigated (1998-2017), the total number of hot days recorded in July, August and September in Niš was 126, 133 and 21 HDs, respectively, while Novi Sad saw about a half of the said number of hot days, namely 52, 63 and 3 HDs, respectively. In Mt. Zlatibor 2 HD days were recorded in July.

As regards hot nights (HN), there is generally a lower intensity of HNs compared to HDs. Generally, during the observed period up to 4 HNs were recorded, although an increase in the HNs has been observed in the last ten years. For example, in Niš during 2012, 2015 and 2017 the number of HNs recorded in July was 12, 7 and 7 respectively and during 2000, 2013, 2015, and 2017 the number of HNs recorded in August was 8, 6, 5 and 5 respectively. Since 2010 up to 2 HNs have been recorded in September. In Novi Sad, during 2010, 2012, 2015 and 2017 there were 7, 5, 5 and 6 HNs respectively recorded in July, while in August during 2012 and 2017 there were 5 and 7 HNs respectively. In September, one HN was recorded during 2009 and 2011 (Figure 4). The total number of hot nights recorded in July,

August and September for the investigated period in Niš was 60, 56 and 6 HNs, respectively, while in Novi Sad there was a significantly lower number of hot nights – 41, 38 and 2 HNs, respectively. In Mt. Zlatibor 9 HN days were recorded in July and 11 HN days were recorded in August for the period which was investigated.

Midday  $UTCI_{14h}$  “strong heat stress” (SHS) occurs most commonly in all three stations. For example, SHS is particularly important to Mt. Zlatibor, located at an altitude above 1000 m since there has been an increase in the number of days with SHS in Zlatibor over the last decade. The best indicator is the increase in SHS in September. Similarly, the indicator of biothermal discomfort, “very strong heat stress” (VSHS) has been occurring more frequently in the last decade, reaching a maximum of 17 days in July 2007 and 2012 and in August 2015 in Niš. The number of VSHS days in Novi Sad reached a maximum of 12 days in July 2015 and 14 days in August 2015 and 2017. The total number of VSHS days recorded in July, August and September for the investigated period in Niš was 167, 174 and 25 VSHS days, respectively, while in Novi Sad there was a significantly lower number of VSHS days – 83, 97 and 7 VSHS days, respectively. In Mt. Zlatibor 2 VSHS days were recorded in July and 3 VSHS days were recorded in August. The recorded numbers of VSHS days indicate slightly greater biothermal discomfort duration in relation to the number of HD days. The HD and VSHS indices are significantly correlated during the summer months as they are directly derived from the maximum air temperatures and  $UTCI_{14h}$  index (Table 3).  $UTCI_{14h}$  and  $t_{max}$  for the Niš and Novi Sad urban stations have a significantly high linear correlation.

Table 3.

For Zlatibor the correlation is slightly lower in July and August, while in September is weak compared to the urban stations, but neither HD nor VSHS indices occurred on Zlatibor. Similar results of the linear relationship between the air temperature and the UTCI index ( $r = 0.83$ ;  $r = 0.98$ ) were shown in the studies of Urban and Kysely (2014) for summer in the urban areas of South Bohemia in Czech Republic and in the studies of Nassiri et al. (2017) for Iran.

### c. Heat waves (HWE and VSHSE)

After UTCI thermal stress was compared to the selected thresholds of maximum temperature, biothermal discomfort was identified during extremely high temperatures for the geographical area of Serbia. During the observed period, a heat wave event (HWE) was determined, caused by 3 consecutive HD days and very strong heat stress event (VSHSE), caused by the occurrence of 5 consecutive VSHS days.

Table 4 lists the identified heat waves that refer to  $UTCI_{14h}$  VSHS index (VSHSE) and HD index (HWE). It has a very similar layout. However, shadowed differences can be observed, especially in the duration of the heat wave event. The results are presented for Niš and Novi Sad since there were no recorded heat waves in Mt. Zlatibor. More heat waves, both HWEs and VSHSEs, have been identified in the last ten years.

A heat wave event (HWE) of 10 days in a row (Niš, July from 15<sup>th</sup> to 24<sup>th</sup>, 2007) is the maximum number of consecutive days with such high temperatures when all the three stations are compared. In the same month, HWE of 3 days in a row (July 8<sup>th</sup> to 10<sup>th</sup>) and 4 days in a row (July 27<sup>th</sup> to 30<sup>th</sup>) were observed (Table 4, HWE). Altogether, the heat wave events (HWEs) occurred 3 times and they lasted 3, 10 and 4 days in July 2007, which is certainly an extreme for the observed period. The heat wave events (HWE) which lasted 3 or 4 days in a row occurred during 9 years in the observed period of 20 years. From that source, heat wave events (HWE) occurred twice and lasted 3 to 4 days during July in 2004 and 2017 (Table 4). It is necessary to point out that heat waves in July 2002, 2005, 2007, 2012 and 2015 lasted more than five HD days in a row with  $t_{max}$  above 35°C, amounting to 6, 6, 10, 9 and 7 HD days respectively. During August heat wave events (HWE) with 3 to 6 hot days in a row occurred twice in 1999, 2000, 2001, 2012, 2015 and 2017. Furthermore, in August, series of heat wave events with 5 hot days in a row or more were observed in 1998, 2000, 2007, 2012, 2013, 2015 and 2017 and it amounts to 5, 6, 6, 6, 7 and 6 hot days in a row, respectively. As regards the thresholds in Novi Sad, there were two heat waves events (HWE) with 3 and 4 hot days in a row which occurred in August of 2012 and 2015. In 2017 3 heat wave events (HWE) occurred with 6, 3, and 3 hot days in a row. It is certainly important that two HWEs occurred during September 2015 and lasted 3 and 5 HD days in a row with  $t_{max}$  above 35°C (Table 4). This justifies the finding of Tomczyk (2016) that the heat waves began to occur in

September in the last decade. As shown in Fig. 4, the number of HDs in September, during the period of 20 years increased significantly after 2007, especially in Niš.

#### Table 4.

Of the total number of hot days, it amounts to 21 HDs in September in Niš, only one HD was recorded in the first decade (1998-2007) of the investigated period, while the remaining HDs were recorded over the last decade (2008-2017), especially in 2011 and 2015. In the second decade of the investigated period the year 2015 stands out with two HWEs (September 1<sup>st</sup> to 5<sup>th</sup>, and 17<sup>th</sup> to 19<sup>th</sup>). In the same year the highest daily temperature was recorded in September (on September 18<sup>th</sup>,  $t_{max}=37.5^{\circ}C$ ). A similar situation occurred in the area of Novi Sad. During the first 10 years, not a single HD was recorded in September while after 2007, 5 such days were recorded. For Niš and Novi Sad year 2015 stands out and the hottest September days were September 17<sup>th</sup> and 18<sup>th</sup> with  $t_{max}=36.7^{\circ}C$  and  $36.4^{\circ}C$ .

For both weather stations in the second decade, together with the increase in the number of HDs, there was also an increase in the number of HNs. In September, the daily air temperature increases, so days with temperatures around  $30^{\circ}C$  and higher are more frequent. In this weather station year 2015 stands out with the same date of the hottest day in September (September 18<sup>th</sup>,  $t_{max}=33.2^{\circ}C$ ). This correlates with the heat waves analysis in Athens, where period of heat wave lasted from mid-June to the beginning of September (Papanastasiou et al., 2014). In Mt. Zlatibor area there was no significant phenomenon of extreme temperatures and therefore no HWEs were recorded.

Bioclimatic conditions analyzed by means of the UTCI index show that the calculated data observed at 14 CET are related to the marked heat wave events (HWE). When  $UTCI_{14h}$  thermal stress category of "very strong heat stress" (VSHS) is compared to the selected heat wave events (HWE,) a sub-index is defined. It is called very strong heat stress event (VSHSE) and it is caused by the occurrence of 5 consecutive VSHS days. The VSHSE corresponds to heat wave event and provokes a severe biothermal discomfort, so it was used as an indicator of extremely unfavorable bioclimatic conditions (biothermal heat hazard). Along these lines, the VSHSE in Niš occurred during July in 1998, 2004, 2005, 2007, 2012 and 2015 where 7, 5 (on two occasions), 6, 10, 12 and 10 VSHS days in a row were recorded respectively (Table 4, VSHSE). In August, VSHSE was recorded on 5 VSHS days in a row in 1998, 1999, 2008, 2010, 2012 and 2017, while in 2000, 2012, 2013, 2015 and 2017 6, 6, 8, 7 and 6 VSHS days in a row were recorded respectively. It should be emphasized that there were two VSHSE events in 2012 and 2017, for each year 5 and 6 VSHS days in a row. September 2015 saw a VSHSE with 5 VSHS days in a row. According to Unkašević and Tošić (2009b) the highest temperatures in Niš ever were recorded during the summer of 2007 ( $44.2^{\circ}C$ ) and the summer of 2000 ( $42.5^{\circ}C$ ), covering the data period from 1948 to 2007. The VSHSE occurred in Novi Sad in July 2007 (7 VSHS days in a row), 2012 (6 VSHS days in a row) and 2015 (8 VSHS days in a row) and in August 2000 (7 VSHS days in a row), 2013 (8 VSHS days in a row), 2015 (6 VSHS days in a row) and 2017 (6 VSHS days in a row). In August 2012 VSHSE occurred twice and each time 5 VSHS days in a row were recorded. There was no sub-index VSHSE recorded in Zlatibor.

#### 4. Discussion

The assessment of human biothermal conditions during the summer made it possible to identify biothermal discomfort related to heat waves in Serbia. The results obtained in the study for the investigated period from 1998 to 2017, indicates the general increase in biothermal discomfort associated with heat waves defined by maximum air temperature above  $35^{\circ}C$  and UTCI "very strong heat stress" above  $38^{\circ}C$ .

The results obtained for morning, midday and average UTCI show significant occurrence of thermal heat stress during the summer months and increase in biothermal discomfort especially for  $UTCI_{14h}$ . This is confirmed by obtained distribution of the average monthly  $UTCI_{14h}$  showing a significant increase in the extreme values of the

UTCI<sub>14h</sub> index in the last ten years, presented mainly as “very strong heat stress category” (VSHS) together with a growing trend of the UTCI<sub>14h</sub> index in all the summer months (especially September) for all three investigated weather stations (primarily in Niš). According to Di Napoli et al. (2018), there are two thermal climates in Europe. One of them is associated with heat stress condition and it is predominant in the southern part of Europe (including the Balkans) when moderate and strong heat stress occurs at central day time hours, which reflects the general relationship between heat load and insolation.

The increase in extreme biothermal conditions is the most evident in the number of days with maximum air temperature thresholds above 35°C, defined in the study as hot days (HD) and in the number of days with UTCI index thresholds between 38°C and 46°C, defined in the study as “very strong heat stress” (VSHS). Thus, most HDs (18 days) occurred in 2007 and then in 2012 (16 days), which means that more than half of July in these years was characterised by extremely high temperatures. Similar results about increase in daily maximum air temperature were obtained by Unkašević and Tošić (2011), when the record values of the maximum temperatures were observed for almost the whole territory of Serbia in 2007. As reported by Papanastasiou et al. (2014), the summer of 2007 was the warmest summer in Athens in the last hundred years. Besides, Spinoni et al. (2015) in their study compiled a list of the most severe drought events which occurred in Europe from 1950 to 2012. They singled out the Balkan countries as extreme weather and drought-prone areas (particularly Serbia) with the longest drought registered in 2007, and the most severe drought event in 2011. Moreover, in 2012 the Balkan Peninsula and the south-eastern Europe faced the hottest summer and one of the worst droughts in nearly 40 years (Unkašević and Tošić, 2015). This is in relation with the investigation of heat waves (a sequence of at least 5 days with maximum temperature above the 95<sup>th</sup> annual percentile) in south-eastern Europe, where there was an increase in the number of hot days for the period from 1973 to 2010. The UTCI<sub>14h</sub> category of “very strong heat stress” (VSHS) correlates with the maximum air temperatures and there has been an increase in the number of such days over the last ten years. Particularly severe biothermal discomfort occurred in 2007 (18 VSHS days) and 2012 (17 VSHS days). This is in agreement with the statement of Di Napoli et al. (2018) about UTCI reference value from the period between 1979 and 2016, when the UTCI at 12:00 (UTC) was about 0.5°C colder in the period between 1980 and 1999, while it was 0.5°C warmer in the period between 2000 and 2009 and 1°C warmer in the period between 2010 and 2016. Of particular importance is the increase in HD, HN, SHS and VSHS days in September since 2008. The resultant biothermal condition as an outcome of the heat budget index has some advantages. For instance, the situation in Novi Sad on July 30<sup>th</sup> 2007, when the maximum temperature was higher than the calculated UTIC index ( $T_{max}=26.9^{\circ}C$ ;  $UTCI_{14h}=15.7^{\circ}C$ , no thermal stress) does not necessarily indicate biothermal heat stress if the wind speed is high ( $6.7 \text{ m}\cdot\text{s}^{-1}$ ) or the relative humidity is lower (52%). Otherwise, the situation on August 26<sup>th</sup> 2002, when the maximum air temperature was lower than the calculated UTIC index ( $T_{max} = 24.5^{\circ}C$ ;  $UTCI_{14h} = 30.6^{\circ}C$ ) indicates that heat stress may still occur if the wind speed is low ( $0.8 \text{ m}\cdot\text{s}^{-1}$ ) or the relative humidity is higher (57%).

In this sense, there is a difference between the definition of biothermal heat stress and potential heat stress, defined on the basis of the interpretation of maximum temperatures. Determining extreme weather event from the aspect of human heat budget indices allows other meteorological parameters to be taken into account. Apart from temperature and humidity, wind and direct solar radiation, have been considered together with metabolic rate and clothing insulation.

The identified heat waves based on the threshold of the UTCI index correspond to the identified heat waves based on the threshold of maximum air temperatures. Furthermore, there is a significant increase in heat waves (HWE and VSHSE) plus duration of such events over the last ten years in Niš and Novi Sad, especially in Niš. The urban areas have a high risk of heat absorption of buildings and asphalt, which results in the formation of high temperatures during the night (Giannopoulou et al., 2014). Rural and mountainous areas rich in forests and greenery are less exposed to these phenomena, like Mt. Zlatibor, but during the hottest summer months, days and nights with high temperatures can occur (Fig. 2c). It is important to point out that two or even three heat waves occurred in some years. The analysis highlights 2007, 2012, 2015 and 2017 as the years with the most heat waves registered as heat wave events (HWE), with special emphasis on Niš, where the longest HWE lasted for 10 days in July, 2007 with UTCI index over 38°C and maximum temperature over 35°C (Table 4). This agrees with the findings of Unkašević and Tošić (2011, 2015) that there has been a growing trend of heat waves in Serbia, especially since 2000. In order to justify the present research for the last 20 years, the earlier results related to the increase in heat waves will be discussed in more detail. In that regard, previous research into heat waves in Niš for the period from 1949 and 2007, based on the autoregressive-

moving-average model, observed the warmest years during three periods, 1951-1952, 1987-1998 and 2000-2007 (Unkašević and Tošić, 2009b). According to this research, the longest heat wave was recorded in Niš in 1952, with 21 days, while in 2003, 29 consecutive tropical days were observed. It should be taken into account that according to the Serbian National Weather Service, tropical days are days with a maximum temperature over 30°C. Furthermore, based on Heat Wave Duration Index (HWDI) i.e. daily maximum values of air temperature, Drljača et al. (2009) determined the duration and strength of heat waves in Niš during the summer season. The analysis showed Niš with a greater number of heat waves compared to the larger urban area of Belgrade. As stated in the research, since the mid-1980s, heat waves have had a higher frequency and on average they have occurred every year. Prior to that period heat wave fluctuations were generally reported in one of two years (Niš during the summer) (Drljača et al., 2009). While studying the characteristics of the heat waves in central Serbia (1949-2007), Unkašević and Tošić (2011) detected an increase in heat wave duration, in addition to an increase in heat wave frequency of occurrence during the period from 1999 to 2007. Besides, based on the HWDI index, heat waves in Novi Sad show a trend of increasing tropical days ( $T_{\max} \geq 30.0^{\circ}\text{C}$ ) from 1960 to 2010, indicating years 1994, 1998, 2000, 2003, 2007, 2009 (Malinović-Milićević et al., 2013).

As regards the heat budget index of Heat Load (HL) in Serbia (July) for the period from 2000 to 2010, it was noticed that in, Niš, Zlatibor and Novi Sad the category of “extremely hot” occurred in 2000 and 2007 (Milovanović et al., 2017). As for the heat budget index Physiological Equivalent Temperature PET for the period from 1949 to 2012, the highest number of heat waves in Novi Sad was observed in the last two decades and in the first decade of the investigated period (Basarin et al., 2016), while the number of the days above particular thresholds for the period 1961-2014 shows an increase along with the number of heat waves per year since 1981. The highest values of mean annual PET were mostly detected during the last 15 years (Basarin et al., 2018).

On the basis of the investigation of heat waves (a sequence of at least 5 days with maximum temperature above the 95th annual percentile) in south-eastern Europe for 21 stations for the period from 1973 to 2010, Tomczyk (2016) concluded that there is an increase in the number of hot days. In south-eastern Europe, heat waves mostly occurred from May to September and the highest number of HWs was recorded in July and August. As stated, in the last decades, heat waves have begun to occur in May and September (Tomczyk, 2016). In this regard, 2 heat wave events (HWE) were highlighted and they lasted 5 and 3 days, as well as a very strong heat wave event (VSHSE) which lasted 5 days in September 2015.

The biothermal discomfort identified in July and August as the hottest summer months in Serbia, could provoke health disorders more frequently. Specific HWs characteristics such as intensity and duration have been found to have devastating effects on human health and wellbeing. In certain cases heat waves cause problems and children, elderly people, chronic patients and workers are particularly susceptible to them. The analysis of the UTCI index correlating with mortality in Europe shows that deaths occur in the southern Europe (Italy, Portugal, Greece, Spain) for the categories of moderate and strong stress. Heat load has an impact on increasing mortality as conditions become more thermally stressful (Di Napoli et al., 2018). The intensity of heat stress in Poland was assessed by means of the UTCI index on the basis of daily mortality and the weather data for the period from 1991 to 2000. There is an increase in mortality on the days with strong and very strong heat stress, in relation to no thermal stress days, of 12% and 47%, respectively (Błażejczyk et al., 2017). Biothermal conditions and mortality rates show that in southern Europe during the summer thresholds of Physiological Subjective Temperature (PST) increase when moving from the west to the east. The general finding is that the population of central Europe is more sensitive to thermal stimuli in the summer than the citizens in the south of the continent (Błażejczyk and McGregor, 2008). The statement about the PST index is in support of the comparison of UTCI index with other thermal indices, including PST index, which correlates very well with UTCI index (Błażejczyk et al., 2012). Human sensitivity to extreme heat weather can also be seen in the impact of heat waves on daily mortality in Belgrade during the summer, when the strong correlation between heat waves and daily mortality can be observed in July 2007 (Stanojević et al., 2014). Similarly, Bogdanović et al. (2013) reported a significant short-term excess mortality on 16<sup>th</sup> July in Belgrade, when the maximum daily temperature exceeded 35°C, leading to 167 excess deaths (38% increase compared to the number of expected deaths) for nine consecutive days of heat, with a progressive return to almost normal mortality as the maximum temperature dropped below 35°C on 25<sup>th</sup> July.

The application of the standardized bioclimatic heat budget index UTCI help improve the understanding of biothermal conditions relative to heat wave. The daily UTCI<sub>14h</sub> heat stress becomes more extreme in terms of severity



and heat wave duration up to very strong heat stress (VSHS). The biothermal indices investigated for the three weather stations follow the trend of general warming. In the light of climatic changes and other negative factors resulting from this global phenomenon, it is becoming a true challenge to minimize their effects and improve living conditions in urban and rural areas (Stevović et al., 2017). However, the human heat budget indices are increasingly serving as a determinant of extreme weather condition (Di Napoli et al., 2018; Matzarakis et al., 2018; Theoharatos et al., 2010; Urban and Kyselý, 2014). UTCI index improves the understanding of extreme thermal stress, serving as one of the criteria for initiating heat alerts about extreme thermal stress in Serbia.

## 5. Conclusion

The present research of the human bioclimatic evaluation of UTCI thermal stress in Serbia is considered to be of great importance due to the identification of biothermal heat hazard and the study of the evolution of such indices regarding climate change. This evaluation aims at providing a comprehensive assessment of the human heat stress associated with UTCI index and heat waves defined by UTCI very strong heat stress and maximum air temperature.

The assessment of human biothermal conditions for the investigated period from 1998 to 2017 was provided for three synoptic stations – two urban (Niš and Novi Sad) and one station representing mountain areas of an altitude up to 1500 m (Zlatibor). All of them are located in different geographical areas in Serbia. It was found that morning, midday and average UTCI show significant occurrence of thermal heat stress categories during the summer months and increase in biothermal discomfort especially for UTCI<sub>14h</sub>. The results obtained in the study indicate an increase in biothermal discomfort associated with heat waves defined by UTCI<sub>14h</sub> “very strong heat stress” (VSHS) above 38°C and heat waves defined by maximum air temperature above 35°C.

The VSHS describes an alarming biothermal state and has occurred frequently in the last ten years, particularly in Niš and Novi Sad. The outcome of the study is UTCI<sub>14h</sub> index “very strong heat stress event” (VSHSE) as an indicator of biothermal heat hazard corresponding to heat waves defined by maximum air temperature. The most extreme heat waves (HWE and VSHSE) occurred in 2007, 2012, 2015 and 2017. Heat wave events (HWE) lasting 10 days in July 2007, 9 days in July 2012 and heat wave events (VSHSE) lasting 10 days in July 2007 and July 2015 and those lasting 12 days in July 2012, which occurred in Nis, are the events with maximum duration. Another certainly important finding is the fact that defined HWs occurred twice in September in 2015. **Undoubtedly, heat wave events are one of the natural hazards with increasing impact in urban areas related to higher population density. Considering these facts, it can be deduced that heat waves are becoming more frequent, stronger and longer. Thus, frequent heat waves since 2007 certainly indicate biothermal heat hazard. The longer events of bioclimatic discomfort could indicate more stressful effect on bioclimatic conditions for human health and wellbeing. However, the results of the study highlight the importance of UTCI index as a bioclimatic indicator of biothermal hazard in Serbia particularly if it can serve as one of the criteria for initiating heat warnings in Serbia.**

**Acknowledgements:** The paper represents the results of research on the National projects supported by Ministry of Education, Science and Technological Development, Republic of Serbia (No. III 47007; 176017 and 176008). The authors highly appreciate comments and suggestions of reviewers. UB contributed to the interpretation of the results and the improvement of the figures All authors contributed to the discussion and interpretation of the results.

**Author’s contribution:** Milica Pecelj designed the study, collected, analysed and interpreted the data, wrote the manuscript and processed the figures. Milica Lukić collected the data, supported the writing and processed the figures. Dejan Filipović carried out the management activities. Branko Protić participated in the technical support.

**Conflict of interest:** We claim that all authors agree to the submission of the manuscript. This manuscript has not been published before and is not concurrently being considered for publication elsewhere. This manuscript does not violate any copyright or other personal proprietary right of any person or entity and it contains no statements that are unlawful in any way.

## References

- ANSI/ASHRAE Standard 55, Thermal Environmental Conditions for Human Occupancy, (ANSI/ASHRAE Approved), [www.techstreet.com/ashrae/standards](http://www.techstreet.com/ashrae/standards), 2017.
- Basarin, B., Lukić, T., and Matzarakis, A.: Quantification and assessment of heat and cold waves in Novi Sad, Northern Serbia, *Int. J. Biometeorol.*, 60(1), 139-150, <https://doi.org/10.1007/s00484-015-1012-z>, 2016.
- Basarin, B., Lukić, T., Mesaroš, M., Pavić, D., Djordjević, J., and Matzarakis, A.: Spatial and temporal analysis of extreme bioclimatic condition in Vojvodina, Northern Serbia, *Int. J. Climatol.*, 38, 142-157, <https://doi.org/10.1002/joc.5166>, 2018.
- Błażejczyk, K.: BioKlima2.6 (version 2.6) – Universal tool for bioclimatic and thermophysiological studies, <https://www.igipz.pan.pl/Bioklima-zgik.html>, 1994.
- Błażejczyk, K.: New climatological and physiological model of the human heat balance outdoor (MENEX) and its application in bioclimatological studies in different scales, *Zeszyty IGIPZ PAN*, 28, 27-58, 1994.
- Błażejczyk, K. and McGregor, G.: Mortality in European cities and its relations to biothermal conditions, In: *Klimat i bioklimat miast*, edited by: Klysik, K., Wibig, J., Fortuniak, K.: Wydawnictwo Uniwersytetu Łódzkiego, Łódź, Poland, pp 313–324, 2008.
- Błażejczyk, K., Epstein, Y., Jendritzky, G., Staiger, H., and Tinz, B.: Comparison of UTCI to selected thermal indices. *Int. J. Biometeorol.*, 56(3), 515-535, <https://doi.org/10.1007/s00484-011-0453-2>, 2012.
- Błażejczyk, K., Jendritzky, G., Bröde, P., Fiala, D., Havenith, G., Epstein, Y., Psikuta, A., and Kampmann, B.: An introduction to the Universal Thermal Climate Index, *Geogr. Pol.*, 86(1), 5-10, <http://dx.doi.org/10.7163/GPol.2013.1>, 2013.
- Błażejczyk, K., Kuchcik, M., Błażejczyk, A., Milewski, P., and Szmyd, J.: Assessment of urban thermal stress by UTCI – experimental and modelling studies: an example from Poland, *DIE ERDE*, 145 (1-2), 16-33, doi: 10.12854/erde-145-3, 2014.

- Błażejczyk, A., Błażejczyk, K., Baranowski, J., and Kuchcik, M.: Heat stress mortality and desired adaptation responses of healthcare system in Poland, *Int. J. Biometeorol.*, 62(3), 307-318, <https://doi.org/10.1007/s00484-017-1423-0>, 2017.
- Bleta, A., Nastos, P.T., and Matzarakis, A.: Assessment of bioclimatic condition in Crete Island, Greece, *Reg. Environ. Change.*, 14(5), 1967-1981, <https://doi.org/10.1007/s10113-013-0530-7>, 2014.
- Bogdanović, D., Milošević, Z., Lazarević, K., Dolićanin, Z., Ranđelović, D., and Bogdanović, S.: The impact of the July 2007 heat wave on daily mortality in Belgrade, Serbia, *Cent Eur J Public Health*, 21,(3), 140-145, <https://doi.org/10.21101/cejph.a3840>, 2013.
- Bröde, P., Fiala, D., Błażejczyk, K., Holmér, I., Jendritzki, G., Kampmann, B., Tinz, B., and Havenith, G.: Deriving the operational procedure for the Universal Thermal Climate Index (UTCI), *Int. J. Biometeorol.*, 56(3), 481-494, <https://doi.org/10.1007/s00484-011-0454-1>, 2012.
- Brown, S. J., Caesar, J. and Ferro, C. A. T.: Global changes in extreme daily temperature since 1950, *J. Geophys. Res.*, 113(D5), 1-11, <https://doi.org/10.1029/2006JD008091>, 2008.
- Collins, D.A., Della-Marta, P.M., Plummer, N., and Trewin, B.C.: Trends in annual frequencies of extreme temperature events in Australia, *Aust. Met. Mag.*, 49, 277–292, 2000.
- Di Napoli, C., Pappenberg, F., and Cloke, H.L.: Assessing heat-related health risk in Europe via the Universal Thermal Climate Index (UTCI), *Int. J. Biometeorol.*, 62(7), 1155-1165, <https://doi.org/10.1007/s00484-018-1518-2>, 2018.
- Di Napoli, C., Pappenberger, F., and Cloke, H.L.: Verification of Heat Stress Thresholds for a Health-Based Heat-Wave Definition, *J. Appl. Meteorol. Climatol.*, 58(6), 177- 194, <https://doi.org/10.1175/jamc-d-18-0246.1>, 2019.
- Drljača, V., Tošić, I., and Unkašević, M.: An analysis of heat waves in Belgrade and Niš using the climate index, *J. Geogr. Inst. Jovan Cvijić SASA*, 59(1), 49-62, <https://doi.org/10.2298/IJGI0959049D>, 2009.
- Easterling, D.R., Horton, B., Jones, P.D., Peterson, T.C., Thomas, R.K., David, E.P., Salinger, M.J., Razuvayev, V., Plummer, N., Jamason, P., and Folland, C.K.: Maximum and minimum temperature trends for the globe. *Sci.*, 277(5324), 364–367, <https://doi.org/10.1126/science.277.5324.364>, 1997.
- Epstein, Y. and Moran, D.S.: Thermal comfort and heat stress indices, *Ind. Health*, 44(3), 388-398, <https://doi.org/10.2486/indhealth.44.388>, 2006.
- European Commission's (Eurostat) datasets: <https://ec.europa.eu/eurostat/web/gisco/geodata/reference-data/>, last access: 3<sup>rd</sup> February 2020.
- Fanger, P.O.: *Thermal Comfort: analysis and applications in environmental engineering*, Danish Technical Press, Copenhagen, Denmark, 1970.
- Giannopoulou, K., Livada, I., Santamouris, M., Saliari, M., Assimakopoulos, M., and Caouris, Y.: The influence of air temperature and humidity on human thermal comfort over the greater Athens area, *Sustain. Cities. Soc.*, 10, 184-194, <https://doi.org/10.1016/j.scs.2013.09.004>, 2014.
- Gocić, M. and Trajković, S.: Analysis of precipitation and drought data in Serbia over the period 1980–2010, *J. Hydrol.*, 494, 32-42, <https://doi.org/10.1016/j.jhydrol.2013.04.044>, 2013.
- Gocić, M. and Trajković, S.: Spatiotemporal characteristics of drought in Serbia, *J. Hydrol.*, 510, 110-123, <https://doi.org/10.1016/j.jhydrol.2013.12.030>, 2014.
- Höppe, P.: The physiological equivalent temperature - a universal index for the biometeorological assessment of the thermal environment, *Int. J. Biometeorol.*, 43, 71–75, <https://doi.org/10.1007/s004840050118>, 1999.
- IPCC, Special Report of the Intergovernmental Panel on Climate Change, in: *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*, edited by Field, C. B., Barros, V., Stocker, T.F., Qin, D., Dokken, D.J., Ebi, K.L., Mastrandrea, M.D., Mach, K.J., Plattner, G.K., Allen, S.K., Tignor, M. and Midgley, P.M., Cambridge University Press, Cambridge, UK, 582, 2012.
- ISO 8996: Ergonomics of the thermal environment - Determination of metabolic rate, <https://www.iso.org/standard/34251.html>, 2004.
- Jendritzki, G., De Dear, R. and Havenith, G.: UTCI-Why another thermal index? *Int. J. Biometeorol.*, 56, 421-428, <https://doi.org/10.1007/s00484-011-0513-7>, 2012.
- Landsberg, H. E.: *The Assessment of Human Bioclimate: a limited review of physical parameters*, World Meteorological Organization WMO, 331, Geneva, 1972.
- Lazić, L., Savić, S., and Tomić, Ž.: Analysis of the Temperature Characteristics and Trends in Novi Sad Area (Vojvodina, Serbia), *Geogr. Pannon.*, 10, 14-21, <https://doi.org/10.5937/GeoPan0610014L>, 2006.

- Lee, V., Zermoglio, F., and Ebi, K.: Heat waves and human health - emerging evidence and experience to inform risk management in a warming world, Technical Report, United States Agency for International Development USAID, Washington 2019
- Lukić, M., Pecelj, M., Protić, B., and Filipović, D.: An evaluation of summer discomfort in Niš (Serbia) using Humidex, *J. Geogr. Inst. Jovan Cvijić SASA.*, 69(2), 109-122, <https://doi.org/10.2298/IJGI1902109L>, 2019.
- Leščičen, I., Dolijan, D., Pantelić, M, and Popov, S.: Drought assessment in Vojvodina (Serbia) using k-means cluster analysis, *J. Geogr. Inst. Jovan Cvijić SASA.*, 69(1), 17-27, <https://doi.org/10.2298/IJGI1901017L>, 2019.
- Matzarakis, A., Muthers, S., and Rutz, F.: Application and comparison of UTCI and PET in temperate climate conditions, *Finisterra*, 48(98), 21–31., <https://doi.org/10.18055/Finis6453>, 2014.
- Masterton, J. and Richardson, F.A.: Humidex, a method of quantifying human discomfort due to excessive heat and humidity, Atmospheric Environment Service Canada, Ontario, Downsview, 1979.
- McGregor, G. and Vanos, J.: Heat: A primer for public health researchers, *Public Health*, 161, 138–146, <https://doi.org/10.1016/j.puhe.2017.11.005>, 2018.
- Meteorological Yearbook for the period from 1998 to 2017, Republic Hydrometeorological Service of Serbia, [http://www.hidmet.gov.rs/latin/meteorologija/klimatologija\\_godisnjaci.php](http://www.hidmet.gov.rs/latin/meteorologija/klimatologija_godisnjaci.php), last access: 15 July 2018.
- Milewski, P.: 2013. Application of the UTCI to the local bioclimate of Poland's Ziemia Kłodzka region, *Geogr. Pol.*, 86(1), 47-54, <https://doi.org/10.7163/GPol.2013.6.2013>.
- Milovanović, B., Radovanović, M., Stanojević, G., Pecelj, M., and Nikolić, J.: Klima Srbije (Climate of Serbia), In: *Geografija Srbije (Geography of Serbia)*, edited by: Radovanović, M., Geogr. Inst. Jovan Cvijić SASA, Belgrade, Serbia, pp 94-156, 863p, 2017.
- Malinović-Milićević S. Summer hazards in Novi Sad, *J. Geogr. Inst. Jovan Cvijić SASA.*, 63 (3), 335-344, DOI: 10.2298/IJGI1303335M, 2013.
- Malinovic-Milicevic, S., Radovanovic, M.M., Stanojevic, G., and Milovanović B.: Recent changes in Serbian climate extreme indices from 1961 to 2010. *Theor Appl Climatol* 124, 1089–1098, <https://doi.org/10.1007/s00704-015-1491-1>, 2016.
- Milovanović, M ., Gocić, M., and Trajković, S: Analysis of meteorological and agricultural droughts in Serbia, *Architecture and Civil Engineering*, 12(3), 253-264, <https://doi.org/10.2298/FUACE1403253M>, 2014.
- Nassiri, P., Monazzam, M.R., Golbabaee, F., Dehghan, S.F., Rafieepour, A., Morteza pour, A., Asghari, M.P.: Application of Universal Thermal Climate Index (UTCI) for assessment of occupational heat stress in open-pit mines, *Industrial Health*, 55 (5), 437-443, <https://doi.org/10.2486/indhealth.2017-0018>, 2017.
- Nastos, P. and Matzarakis, A.: The effect of air temperature and human thermal indices on mortality in Athens, Greece, *Theor. Appl. Climatol.*, 108, 591–599, <https://doi.org/10.1007/s00704-011-0555-0>, 2012.
- Nemeth, A.: Changing thermal bioclimate in some Hungarian cities, *Acta Climatologica et Chorologica*, 44–45, 93–101, 2011.
- Osczevski, R.J. and Bluestein, M.: The new wind chill equivalent temperature chart, *Bull. Am. Meteorol. Soc.*, 86, 1453–1458, <https://doi.org/10.1175/BAMS-86-10-1453>, 2005.
- Parsons, K.C.: Human thermal environments: the effects of hot, moderate, and cold environments on human health, comfort and performance, Taylor & Francis Group, Abingdon, England, UK, 2003.
- Papanastasiou, D.K., Melas, D., and Kambezidis, H.D.: Heat Waves characteristics and their relation to air quality in Athens, *Global. NEST. J.*, 16(5), 919-928, <https://doi.org/10.30955/gnj.001530>, 2014.
- Pecelj, M., Đordđević, A., Pecelj, M.R., Pecelj-Purković, J., Filipović, D., and Šećerov, V.: Biothermal conditions on Mt. Zlatibor based on thermophysiological indices, *Arch. Biol. Sci.*, 69(3), 455-461, <https://doi.org/10.2298/ABS151223120P>, 2017.
- Pecelj, M., Lukić, M., Vučićević, A., De Una-Alvarez, E., Esteves da Silva, C.G.J., Freinkin, I., Ciganović, S., and Bogdanović, U.: Geoecological evaluation of local surroundings for the purposes of recreational tourism, *J. Geogr. Inst. Jovan Cvijić SASA.*, 68(2), 215-231, <https://doi.org/10.2298/IJGI1802215P>, 2018.
- Peña-Gallardo, M., Vicente-Serrano, M.S., Domínguez-Castro, F., and Beguería, S.: The impact of drought on the productivity of two rainfed crops in Spain, *Nat. Hazards Earth Syst. Sci.*, 19, 1215–1234, <https://doi.org/10.5194/nhess-19-1215-2019>, 2019.
- Prokić, M.: Climate trends of temperature and precipitation in Nišava river valley (Serbia) for 1960-2015 period. *J. Geogr. Inst. Jovan Cvijić SASA.*, 68(1), 35-50, <https://doi.org/10.2298/IJGI1801035P>, 2018.
- Rahmstorf, S. and Coumou, D.: Increase of extreme events in a warming world, *P. Natl. Acad. Sci.*, 108(44), 17905–17909, <https://doi.org/10.1073/pnas.1101766108>, 2011.

- Robinson, J.P.: On the definition of a Heat Wave, *J. Appl. Meteor.*, 40, 762-775, [https://doi.org/10.1175/1520-0450\(2001\)040<0762:OTDOAH>2.0.CO;2](https://doi.org/10.1175/1520-0450(2001)040<0762:OTDOAH>2.0.CO;2), 2001.
- Siple, P. and Passel, C.F.: Measurements of dry atmospheric cooling in subfreezing temperatures, *Proc Am Philos Soc*, 89 (1), 177–199, [www.jstor.org/stable/985324](http://www.jstor.org/stable/985324), 1945.
- Spinoni, J., Naumann, G., Vogt, J., and Barbosa, P.: The biggest drought events in Europe from 1950 to 2012, *J. Hydrol. Reg. Stud.*, 3, 509-524, <https://doi.org/10.1016/j.ejrh.2015.01.001>, 2015.
- Stanojević, G., Stojićević, J., Spalević, A., and Kokotović, V.: The Impact of heat waves on daily mortality in Belgrade (Serbia) during summer, *Environ. Hazard.*, 13(4), 329-342, <https://doi.org/10.1080/17477891.2014.932268>, 2014.
- Seneviratne, S.I. and Nicholls, N.: Changes in Climate extremes and their impacts on the natural physical environment, in: *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaption*, edited by Field, C. B., Barros, V., Stocker, T.F., Qin, D., Dokken, D.J., Ebi, K.L., Mastrandrea, M.D., Mach, K.J., Plattner, G.K., Allen, S.K., Tignor, M. and Midgley, P.M., Cambridge University Press, Cambridge, UK, 109–230, 2012.
- Steadman, R.G.: A Universal Scale of Apparent Temperature, *J. Appl. Meteor.*, 23, 1674-1687, [https://doi.org/10.1175/1520-0450\(1984\)023<1674:AUSOAT>2.0.CO;2](https://doi.org/10.1175/1520-0450(1984)023<1674:AUSOAT>2.0.CO;2), 1984.
- Stevović, S., Mirjanić, S., and Đurić, N.: Sustainable urban environment and conflict of resources management, *Archives for Technical Sciences*, 17(1), 79-87, <https://doi.org/10.7251/afts.2017.0917.079S>, 2017.
- Stojićević, G., Basarin, B., and Lukić, T.: Detailed bioclimate analysis of Banja Koviljača (Serbia), *Geogr. Pannon.*, 20(3), 127-135, <https://doi.org/10.18421/GP20.03-01>, 2016.
- Theoharatos, G., Pantavou, K., Mavrakis, A., Spanou, A., Katavoutas, G., Efstathiou, P., Mpekas, P., and Asimakopoulos, D.: Heat waves observed in 2007 in Athens, Greece: Synoptic conditions, bioclimatological assessment, air quality levels and health effects, *Environ. Res.*, 110(2), 152-161, <https://doi.org/10.1016/j.envres.2009.12.002>, 2010.
- Toiika, K., Maheras, P., Pytharoulis, I., and Anagnostopoulou, C.: The anomalous low and high temperatures of 2012 over Greece - an explanation from a meteorological and climatological perspective, *Nat. Hazards Earth Syst. Sci.*, 14(3), 501–507, <https://doi.org/10.5194/nhess-14-501-2014>, 2014.
- Tomczyk, M.A.: Impact of atmospheric circulation on the occurrence of heat waves in southeastern Europe, *Idojarás.*, 120(4), 395–414, 2016.
- Tomczyk, M.A. and Owczarek M.: Occurrence of strong and very strong heat stress in Poland and its circulation conditions, *Theor. Appl. Climatol.*, 139, 893-905, <https://doi.org/10.1007/s00704-019-02998-3>, 2020.
- Tošić, I. and Unkašević, M.: Analysis of wet and dry periods in Serbia, *Int. J. Climatol.*, 34, 1357-1368, <https://doi.org/10.1002/joc.3757>, 2014.
- Unkašević, M. and Tošić, I.: An analysis of heat waves in Serbia, *Glob. Plane. Ghange.*, 65(1-2), 17-26, <https://doi.org/10.1016/j.gloplacha.2008.10.009>, 2009a.
- Unkašević, M. and Tošić, I.: Heat waves in Belgrade and Niš, *Geogr. Pannon.*, 13(1), 4-10, <https://doi.org/10.5937/GeoPan0901004U>, 2009b.
- Unkašević, M. and Tošić, I.: The maximum temperatures and heat waves in Serbia during the summer of 2007, *Clim. Change.*, 108(1-2), 207-223. <https://doi.org/10.1007/s10584-010-0006-4>, 2011.
- Unkašević, M. and Tošić, I.: Trends in temperature indices over Serbia: relationships to large-scale circulation patterns, *Int. J. Climatol.*, 33, 3152–3161, <https://doi.org/10.1002/joc.3652>, 2013.
- Unkašević, M. and Tošić, I.: Seasonal analysis of cold and heat waves in Serbia during the period 1949-2012, *Theor. Appl. Climatol.*, 120(1-2), 29-40, <https://doi.org/10.1007/s00704-014-1154-7>, 2015.
- Urban, A. and Kyselý J.: Comparison of UTCI with other thermal indices in the assessment of heat and cold effects on cardiovascular mortality in the Czech Republic, *Int. J. Environ. Res. Public Health*, 11(1), 952-967, <https://doi.org/10.3390/ijerph110100952>, 2014.
- Yaglou, C.P. and Minard, D.: Control of heat casualties at military training centers, *Am Med Assoc Arch Ind Health*, 16(4), 302–316, 1957.

**Table 1.** UTCI thermal stress classification

UTCI (°C)	> 46	38 to 46	32 to 38	26 to 32	9 to 26	0 to 9	0 to -13	-13 to -27	-27 to -40	< -40
Stress classes	extreme heat stress	very strong heat stress	strong heat stress	moderate heat stress	no thermal stress	slight cold stress	moderate cold stress	strong cold stress	very strong cold stress	extreme cold stress
Abbr.	EHS	VSHS	SHS	MHS	NTS	SLCS	MCS	SCS	VSCS	ECS

Source: Błażejczyk et al. 2013

**Table 2.** Definition of indices used in the study

Abbreviations	UTCI Indices	Definition
UTCI <sub>avg</sub>	Universal Thermal Climate Index	UTCI <sub>avg</sub> = f (t, f, v, t <sub>mrt</sub> )
UTCI <sub>07h</sub>	Universal Thermal Climate Index <sub>07h</sub>	UTCI <sub>07h</sub> = f (t <sub>7h</sub> , f <sub>7h</sub> , v <sub>7h</sub> , t <sub>mrt</sub> )
UTCI <sub>14h</sub>	Universal Thermal Climate Index <sub>14h</sub>	UTCI <sub>14h</sub> = f (t <sub>14h</sub> , f <sub>14h</sub> , v <sub>14h</sub> , t <sub>mrt</sub> )
Abbreviations	UTCI heat stress indices	Definition
SHS	Strong Heat Stress	SHS= UTCI (32°C to 38°C)
VSHS	Very Strong Heat Stress	VSHS = UTCI (38°C to 46°C)
VSHSE	Very Strong Heat Stress Event	minimum 5 VSHS days in a row
EHS	Extreme Strong Stress	EHS= UTCI >46°C
Abbreviations	Thermal indices	Definition
HD	Hot days	t <sub>max</sub> ≥35°C
HN	Hot nights	t <sub>min</sub> ≥20°C
HWE	Heat waves event	minimum 3 HD days in a row


t-air temperature; f-relative humidity; v-wind speed; tmrt-mean radiant temperature; tmax-maximum air temperature, tmin-minimum air temperature






1998	4	/	5	/	/	1998	7	/	5	/	/
1999	/	/	3, 4	/	/	1999	/	/	5	/	/
2000	3	/	4, 6	6	/	2000	/	/	6	7	/
2001	3	/	3, 3	/	/	2001	/	/	/	/	/
2002	6	/	/	/	/	2002	/	/	/	/	/
2003	4	/	/	/	/	2003	/	/	/	/	/
2004	3, 4	3	/	/	/	2004	5, 5	/	/	/	/
2005	6	/	/	/	/	2005	6	/	/	/	/
2006	/	/	3	/	/	2006	/	/	/	/	/
2007	3, 10, 4	6	6	/	/	2007	10	7	/	/	/
2008	/	/	4	3	3	2008	/	/	5	/	/
2009	/	/	3	/	/	2009	/	/	/	/	/
2010	/	/	4	/	/	2010	/	/	5	/	/
2011	3	4	3	/	/	2011	/	/	/	/	/
2012	9	6	4, 6	3, 5	/	2012	12	6	5, 6	5, 5	/
2013	/	/	7	4	/	2013	/	/	8	8	/
2014	/	/	/	/	/	2014	/	/	/	/	/
2015	7	9	6, 4	4, 5	5, 3	2015	10	8	7	6	5
2016	3	/	/	/	/	2016	/	/	/	/	/
2017	4, 4	/	6, 4	6, 3, 3	/	2017	/	/	6, 5	6	/

Note: pink cells have more than one HWE and VSHSE duration defined by numbers of the consecutive days



**Figure 1.** Relief map of Serbia with the areas studied: 1-Novı Sad, 2-Niř, 3-Zlatibor and a map of geographical location of Serbia [Maps were created using QGIS 3.8 software on the basis of the official European Commission's (Eurostat) datasets, available at <https://ec.europa.eu/eurostat/web/gisco/geodata/reference-data/>, (Map ratio: 1:1900000; Map projection: WGS 84/UTM, Zone 34N, the official national coordinate system)]



**Figure 2.** Monthly frequency of the days with different UTCI stress classes for mean daily ( $UTCI_{avg}$ ), morning ( $UTCI_{7h}$ ) and midday ( $UTCI_{14h}$ ) index during the period from July to September, 1998-2017: a) Niš, b) Novi Sad, c) Zlatibor. X-axis: time (months), y-axis: frequency (number of days)





Figure 3. The mean monthly UTCI<sub>14h</sub> index during the period (1998-2017). x-axis: time (years), y-axis: UTCI (°C)



**Figure 4.** Number of hot days (HD), hot nights (HN), strong heat stress (SHS) and very strong heat stress (VSHS) in Niš, Novi Sad and Zlatibor (July, Aug, Sep) 1998-2017. x-axis: time (years), y-axis: number of days

