

QUANTIFYING SEASONAL CORNICE DYNAMICS USING A TERRESTRIAL LASER SCANNER IN SVALBARD, NORWAY

Holt Hancock^{1,2}, Markus Eckerstorfer³, Alexander Prokop^{4,5,1}, Jordy Hendrikx⁶

¹Department of Arctic Geology, University Centre in Svalbard, N-9171 Longyearbyen, Norway

²Department of Geosciences, University of Oslo, N-0371 Oslo, Norway

³Earth Observation Group, NORCE, N-9294 Tromsø, Norway

⁴Department of Geodynamics and Sedimentology, University of Vienna, 1090 Vienna, Austria

⁵Snow Scan GmbH, Research, Engineering, Education, Stadlauerstrasse 31, 1220 Vienna, Austria

⁶Snow and Avalanche Lab, Department of Earth Sciences, Montana State University, 59717 Bozeman, Montana, USA

Correspondence to: Holt Hancock (holt.hancock@unis.no)

Abstract. Snow cornices develop along mountain ridges, edges of plateaus, and marked inflections in topography throughout regions with seasonal and permanent snow cover. Despite the recognized hazard posed by cornices in mountainous locations, limited modern research on cornice dynamics exists and accurately forecasting cornice failure continues to be problematic. Cornice failures and associated cornice fall avalanches comprise a majority of observed avalanche activity and endanger human life and infrastructure annually near Longyearbyen in central Svalbard, Norway. In this work, we monitored the seasonal development of the cornices along the plateaus near Longyearbyen with a terrestrial laser scanner (TLS) during the 2016/2017 and 2017/2018 winter seasons. The spatial resolution at which we acquired snow surface data with TLS enabled us to observe and quantify changes to the cornice systems in detail not previously achieved. We focused primarily on the evolution and failure of the lower cornice surfaces where accessibility has precluded previous research. We measured cornice accretion rates in excess of 10 mm hr⁻¹ during several accretion events coinciding with winter storms. We observed five cornice fall avalanche events following periods of cornice accretion and one event following a warm period with mid-winter rain. The results of our investigation provide quantitative reinforcement to existing conceptual models of cornice dynamics and illustrate cornice response to specific meteorological events. Our results demonstrate the utility of TLS for monitoring cornice processes and as a viable method for quantitative cornice studies in this and other locations where cornices are of scientific or operational interest.

1 Introduction

30 | Snow cornices are overhanging projections of snow ~~forming that form~~ due to the deposition of wind transported snow in the lee of ridgelines or sharp slope inflections (Montagne et al., 1968; Seligman, 1936). Cornices have attracted interest for their hydrologic implications (e.g. Anderton et al., 2004) and as agents of geomorphic change in periglacial environments (Eckerstorfer et al., 2013; Humlum et al., 2007), but are perhaps best recognized as a snow and avalanche hazard in mountainous terrain (Montagne et al., 1968; Vogel et al., 2012). Cornices pose an avalanche hazard when they fail either as a
35 | full cornice failure with the entire cornice detaching from the ground or as a partial failure with a smaller cornice mass separating from the rest of the cornice. The detached cornice blocks travel downslope under the influence of gravity and become a cornice fall avalanche by entraining loose surface snow or triggering a snow slab on the slope below (e.g. Vogel et al., 2012). In ski areas or where cornices and cornice fall avalanches endanger infrastructure, both explosives (Farizy, 2013; McCarty et al., 1986) and structural defenses (e.g. Montagne et al., 1968) are employed operationally to mitigate cornice
40 | hazards. Most cornice-related fatalities, however, occur in recreational backcountry settings and result from the victim's weight triggering cornice failure.

Despite the well-recognized hazards and operational challenges associated with cornices and cornice fall avalanches, specific cornice research is relatively scarce. Early cornice studies summarized by Vogel et al. (2012) focused on qualitative descriptions of cornice formation processes and resulting cornice structures (e.g. Montagne et al., 1968; Seligman, 1936).
45 | Later studies investigated mechanisms by which individual snow crystals adhere during cornice accretion (Latham and Montagne, 1970), the physical snow characteristics at various structural locations on individual cornices (Naruse et al., 1985), and the specific interactions between wind-drifted snow and cornice morphology during cornice formation (Kobayashi et al., 1988).

Recent work has refined the conceptual model of seasonal cornice dynamics established by these earlier studies primarily by
50 | employing time-lapse photography to examine cornice responses to the meteorological factors controlling the development and failure of cornices (Munroe, 2018; van Herwijnen and Fierz, 2014; Vogel et al., 2012). Vogel et al. (2012) observed cornice processes over two winter seasons on a single mountain slope in central Svalbard and proposed a conceptual model of seasonal cornice dynamics including cornice accretion, deformation, and failure. Their results indicated cornice accretion occurs during or immediately following winter storms with wind speeds in excess of 10 m s^{-1} from a direction perpendicular
55 | to the ridgeline, while cornice scouring resulted from strong winds oriented towards the cornice's leading edge. Smaller cornice failures observed in ~~this study by~~ Vogel et al. (2012) were clustered in June near the end of the snow season and coincided with increasing air temperatures and decreased snow strength. Less frequent failures in the earlier part of the winter often involved the entire cornice mass and resulted in the some of the largest cornice fall avalanches observed in the study.

60 Later work in an alpine setting also linked cornice accretion to strong winds during or soon after a snowfall and found the SNOWPACK wind drift index correlated well with cornice width estimates (van Herwijnen and Fierz, 2014). No cornice failures or cornice fall avalanches were observed in this study, however. Munroe (2018) used time-lapse photography to observe the growth and repeated failure of a cornice in Utah, USA. He also found cornice accretion to primarily coincide with periods of snowdrift. He divided the nineteen cornice fall avalanches observed in his study into two distinct groups:
65 snow-caused cornice fall avalanches where failure primarily resulted from snow loading on the cornice and temperature-caused avalanches where failure is related to rapid temperature increases presumably leading to destabilization of the cornice through the loss of snow strength.

We build upon the observational understanding and conceptual model of seasonal cornice dynamics established in these previous works by monitoring cornice systems in Longyeardalen – including one site previously examined by Vogel et al.
70 (2012) – with a terrestrial laser scanner (TLS). TLS – or ground-based LiDAR (Light detection and ranging) – is an active remote sensing technology with documented applications for observing and monitoring various slope processes and hazards including landslides (Jaboyedoff et al., 2012; Prokop and Panholzer, 2009), coastal cliff erosion (e.g. Caputo et al., 2018), and rock slope instability (Abellán et al., 2014). TLS is being increasingly employed in snow and avalanche research to map snow depth and snow depth change (e.g. Deems et al., 2013; Fey et al., 2019; Prokop, 2008; Schirmer et al., 2011). Other
75 specific snow-related applications include quantifying snow drift processes to verify physical models (Mott et al., 2011; Schön et al., 2015; Vionnet et al., 2014), observing avalanche activity to calibrate dynamic avalanche models (Prokop et al., 2015), assisting avalanche control operations (Deems et al., 2013), and as a tool ~~to~~ to plan and design snow-drift control fences to limit hazardous snow accumulation in avalanche release areas (Prokop and Procter, 2016).

We monitored cornice accretion, deformation, failure, and associated cornice fall avalanche activity near Longyearbyen,
80 Svalbard with TLS technology over two winter seasons (2016/2017 and 2017/2018). To our knowledge TLS has not been employed to specifically monitor cornice dynamics, so our primary objectives are to use the high spatial resolution snow surface data acquired via TLS to:

1. Demonstrate the utility of TLS to observe cornice processes
2. Observe and quantify cornice accretion, deformation, failure, and associated cornice fall avalanches and link these
85 processes to their controlling meteorological factors.
3. Use our findings to provide suggestions for forecasting cornice fall avalanches in this and other locations threatened by cornices.

2 Study Area

Figure 1 here

90 | The present study focuses on the cornices forming above Longyeardalen (“the Longyear valley”)~~the Longyear valley~~
(~~hereafter: Longyeardalen~~) in central Svalbard (Figure 1). Longyeardalen is a glacially sculpted, ~~u~~U-shaped valley with a
northeast/southwest oriented ~~valley~~ axis running approximately 3 km from the termini of two small mountain glaciers to a
fjord. The Gruvefjellet and Platåberget plateaus border Longyeardalen to the west and east, respectively, with Svalbard’s
administrative center, Longyearbyen, situated in the valley bottom. The Gruvefjellet and Platåberget slopes lie within the
95 | horizontally-bedded, lower-Tertiary aged Van Mijenfjord Group of sandstones and shales (Major, 2001). Resistant strata
within this group form the area’s extensive plateau topography. The entire region is underlain by continuous permafrost
ranging in thickness from 100 m near the coasts to over 500 m in the higher mountains (Humlum et al., 2003).

Figure 2 here

We investigated seasonal cornice dynamics and cornice fall avalanches along and under the Gruvefjellet and Platåberget
100 | plateau margins, respectively (Figure 2). The steep valley walls descending from the broad plateau summits (approximately
450 m elevation) are characterized in their upper portions by protruding resistant bedrock buttresses and transport couloirs
incised by fluvial and gravitational slope processes. The Gruvefjellet slope described in detail by Eckerstorfer et al. (2013)
consists of a 50-70 m near-vertical bedrock cliff situated ~~between~~ under the plateau margin and over a 40-50° slope that
serves as a slab avalanche release area. This broad slope transitions into the transport couloirs which in turn feed extensive
105 | avalanche fan deposits downslope. Similar morphology exists on the Platåberget slope, but the plateau margin transitions
directly into discrete 45-55° release areas leading into the couloirs and lacks the near-vertical bedrock face present on
Gruvefjellet.

Central Svalbard’s climate is cold and arid, with a mean annual air temperature of -4.6°C and mean annual precipitation of
191 mm at the Svalbard Airport automated weather station (AWS) for the 1981-2010 period of record (Førland et al., 2011).
110 | Combined mean winter (DJF) and mean spring (MAM) precipitation for 1981-2010 is 86 mm (Førland et al., 2011). Mean
winter air temperature for the same period is -11.7°C and mean spring air temperature is -8.3°C (Førland et al., 2011).
Rapidly increasing air temperatures in the winter and spring in response to decreased sea ice extent (Isaksen et al., 2016)
create difficulties establishing representative baseline temperature conditions, with recent reports indicating warming on the
order of 3-5°C for Svalbard as a whole from 1971 to 2017 (Hanssen-Bauer et al., 2019). Less clear changes are apparent in
115 | the precipitation trends (e.g. Hanssen-Bauer et al., 2019), but mid-winter rain-on-snow events are dramatically increasing in
frequency (e.g. Vikhamar-Schuler et al., 2016).

~~Svalbard’s climatic situation~~The climate of Svalbard prohibits the growth of woody vegetation, and snow distribution across
the landscape is strongly controlled by the wind (e.g. Jaedicke and Sandvik, 2002). Southeasterly winds generally prevail
across the region’s plateau mountains, but often switch to westerly or southwesterly during winter storms and are frequently
120 | redirected along the major valley axes at lower elevations (Christiansen et al., 2013). Winter weather in central Svalbard
fluctuates between extended periods of cold, stable high pressure punctuated by warm, wet low pressure systems conveyed

northwards along the North Atlantic cyclone track (Hanssen-Bauer et al., 1990; Rogers et al., 2005). This is reflected in the region's snow and avalanche climate, where the snowpack typically consists of persistent weak layers formed during high pressure interspersed with wind slabs or ice layers formed during snow storms or rain-on-snow events (Eckerstorfer and Christiansen, 2011a). Avalanche activity here displays a strong topographical and meteorological control, with direct action slab avalanches clustered around winter storms and the region's plateaus serving as source areas for the extensive cornice systems that contribute to frequent cornice fall avalanches (Eckerstorfer and Christiansen, 2011c).

3 Methods

3.1 Automated snow and weather data

We obtained wind and air temperature data from the Gruvefjellet automated weather station (AWS), precipitation data from the Svalbard Airport AWS, and a limited time series of snow depth data from a pair of ultrasonic snow depth sensors placed in avalanche release areas on Gruvefjellet and Platåberget during the 2017/2018 winter season (Figures 1 and 2). We defined the winter season as 1 December to 30 June for the purposes of this study. The Gruvefjellet AWS is located less than 500 m east of the Gruvefjellet cornice system at an elevation 464 m and records hourly meteorological data. The Svalbard Airport AWS is situated approximately 5 km northwest of the study area at 28 m and is the only weather station in the region with long-term precipitation measurements.

As part of the installation of a network of automated snow monitoring stations in Longyeardalen (Prokop et al., 2018) we mounted two ultrasonic snow depth sensors in avalanche release areas under the cornice systems in autumn 2017. These sensors were located at 350 m and 450 m elevation on Gruvefjellet and Platåberget, respectively (Figure 2). We employed the Campbell Scientific SR50A ultrasonic distance sensor to measure snow depth at each location. The snow sensors began recording reliable snow depth data on 15 November 2017 and continued until the end of the 2017/2018 season at ten-minute intervals.

3.2 Terrestrial laser scanning (TLS) and post-processing

We used a Riegl® Laser Measurement Systems VZ-6000 ultra-long range terrestrial laser scanner to repeatedly scan the Gruvefjellet and Platåberget cornice systems throughout the 2016/2017 and 2017/2018 winter seasons. The VZ-6000's 1064 nm operating wavelength is particularly well-suited for measuring snow surfaces, while the high scanning speed and measurement range up to 6 km with a 30 kHz pulse repetition rate ensured adequate data acquisition capabilities across the study area in a variety of atmospheric conditions (Riegl, 2019; Prokop, 2008).

We use data from 25 scans of Gruvefjellet and 22 scans of Platåberget during the duration of the study (Appendix I). Of these, one scan from each Gruvefjellet and Platåberget is a snow-free surface taken 16 September 2016. For Gruvefjellet, we acquired usable snow surface data from 18 scans during the 2016/2017 season and 7 scans during 2017/2018. We acquired

14 snow surface scans of Platåberget during 2016/2017 and seven scans during 2017/2018. The TLS was unfortunately damaged in late April 2018 and we were unable to acquire any scans after our final scan on 13 April 2018.

155 We pre-processed the raw point clouds in RiSCAN Pro, Riegl's proprietary data processing software. We established a suite of ground control points on both Platåberget and Gruvefjellet using a differential global positioning system (DGPS) which we used to georeferenced individual point point clouds. We then aligned repeated snow-covered scans to the snow-free scans established in September 2016 using these ground control points and the "Multi Station Adjustment" plugin in RiSCAN Pro following the approach outlined by Prokop and Panholzer (2009). ~~The 3D accuracy of this registration process ranged from <1 mm to 70 mm (Appendix 1).~~ We then manually filtered non-ground points or points above the snow surface. Finally, we 160 ~~applied an octree filter with a 0.10 m increment resampled the resulting point clouds to a 0.10 m grid~~ and exported to an XYZ text file.

We imported individual point clouds into CloudCompare (CloudCompare, 2019) , for further analyses (Figure 3). To create 2D cornice profile cross-sections, we extracted point cloud profile sections along manually defined axes using the polyline extraction tool native to CloudCompare (Figure 3c,d). This tool requires user-defined inputs for profile type, section 165 thickness, and maximum edge length which we set to "Both", 0.6 m, and 0.2 m, respectively. We then manually edited and digitized the resulting shapefiles in the ArcScene 3D Editing environment (ArcGIS 10.4.1) to create the vertical cornice profile schematics as 3D shapefiles.

We calculated representative volumes for selected areas from both the Platåberget and Gruvefjellet cornice systems using the 170 ~~"Compute 2.5D Volume"~~ tool in CloudCompare. This tool computes the volume between two 2.5D point clouds by rasterizing the point clouds to a specified grid size and then computing volumes based on the differences in a specified projection direction between the rasterized values (Figure 3 e,f,g,h). In our case, we rasterized our point clouds to a 1 m grid and calculated horizontal distance differences along the "X" projection direction, which in our georeferenced point clouds corresponds to east-west (i.e. the slope fall-lines). For each cornice system, we computed the volume of snow in 40 m x 8 m area of the plateau margin for each usable snow surface scan by subtracting the bare-earth surface from the scanned snow 175 surface (Figure 3 e,f,g). We chose this areal extent to maximize coverage of an individual cornice throughout its development during the season (i.e. to completely capture the vertical extension of the leading edge) while minimizing volume changes related to other snow on the slope.

Figure 3 here

180 We used the Multiscale Model to Model Cloud Comparison (M3C2) algorithm developed by Lague et al. (2013) and implemented as a plugin in CloudCompare to quantify changes to the cornices and snow surfaces on the slopes below in 3D. The M3C2 algorithm allows for direct comparison of point clouds in 3D and is specifically developed to handle 3D differences and detect changes to complex surfaces where both vertical and horizontal changes exist (Lague et al., 2013). This functionality requires the user to input the following parameters: the normal scale, the projection scale, and the

185 maximum depth (e.g. Lague et al., 2013; Watson et al., 2017). We selected a normal scale of 2 m oriented positively to the scan position (i.e. the normals “face” the scan position), a projection scale of 1 m, and a maximum depth of 10 m for all M3C2 calculations.

190 TLS-based snow surface measurement accuracy generally decreases with increasing distance from the scanner to the measured snow surface and is affected by the manner in which the laser beam interacts with the snow surface, the local terrain characteristics, the stability of the scanner while scanning, and the quality of the scan data registration process (Fey et al., 2019; Hartzell et al., 2017; Prokop et al., 2008). The relative accuracy – the deviation between measurements of an unchanged surface taken under different measurement conditions – can be assessed to quantify uncertainties related to both registration errors and positional errors from the interaction of the laser beam with the surface (Fey et al., 2019; Prokop and Panholzer, 2009). We assessed relative accuracy for our data by measuring M3C2 distances between each snow covered scan and the snow-free scan on a 10x10 m area of stable, snow-free rock faces near the cornices on both Gruvefjellet and
195 | Platåberget (Figure 2). ~~We were unfortunately limited to this single area on which to conduct accuracy assessments because all other surfaces near the cornices became snow covered at some point during the study.~~ We report relative accuracy for each snow-covered scan as the mean of all M3C2 distances on the 10x10 m area (Appendix 1). ~~This L~~locations for the relative accuracy assessment ~~w~~ere selected based on ~~their~~ its ability to remain functionally snow-free throughout the study and because ~~they~~ it were ~~was~~ not otherwise used in the registration process. As both registration and positional errors can be
200 | spatially variable across the scanned area (Fey et al., 2019; Hartzell et al., 2017; Prokop, 2008), we used ~~these~~ this locations in close proximity to the cornices of interest to best represent the relative accuracy near the cornices. Mean M3C2 distance values are smaller than 80 mm for all scans, with standard deviations ranging from <10 mm to 72 mm (Appendix 1). Uncertainty associated with the relative volume metric, calculated by multiplying the relative accuracy of each scan by the surface area considered in the volume calculations (369 m²) thus ranged from less than 1 m³ to 28 m³ (Appendix 1).

205 3.3 Supplemental Observational Data

We relied on snow and avalanche observations from Platåberget and Gruvefjellet from the Norwegian Water Resources and Energy Directorate’s (NVE) online observation platform regObs (www.regobs.no) to supplement our TLS observations. Local observers conduct snow and avalanche assessments on the Gruvefjellet and Platåberget slopes on a sub-weekly basis, so we were able to much better constrain avalanche cycle timing than with the temporal resolution available from the TLS
210 data.

4 Results

4.1 Seasonal summaries of meteorological conditions and cornice dynamics

4.1.1 2016/2017

215 We compare seasonal meteorological conditions (Figure 4) with cross-sectional cornice profiles derived from eight scanned snow surfaces on Gruvefjellet and seven surfaces on Platåberget. We selected these profiles from a pool of 18 usable scans from Gruvefjellet and 14 from Platåberget (Appendx 1) to represent key points in the development of the cornice systems.

Small cornices had accumulated on Gruvefjellet by 2 December 2016. Maximum horizontal cornice growth prior to this scan occurred in the vicinity of profile GF2, where both vertical and horizontal cornice growth exceeded 1 m from the edge of the plateau (Figure 5). The representative cornice volume in the vicinity of profile GF1 already approached 200 m³.
220 Temperatures remained below freezing over the next month, and daily averaged wind speeds exceeded 10 m s⁻¹ only on 29 December 2016. By 12 January 2017, the representative cornice volume on Gruvefjellet had more than tripled relative to early December to over 600 m³. Horizontal cornice extension along the Gruvefjellet cornice system exceeded 4 m in most locations, with maximum horizontal extension near profile GF1 exceeding 5 m (Figure 5). The representative cornice volume of just over 300 m³ from the Platåberget cornices on the same date show considerably less cornice growth (Figure 4).

225 *Figure 4 here*

Heavy snowfall followed by strong westerly winds preceded several cornice fall avalanches on 21 January on Platåberget (Figure 4, Table 1). Representative cornice volume on Platåberget nearly doubled from roughly 300 m³ to over 600 m³ between the 12 January and 21 January scans. Horizontal accretion on profile PB2 exceeded 3.5 m, resulting in an accretion rate of 17 mm hr⁻¹ (Table 2). The cornice represented by profile PB1 failed, triggering a cornice fall avalanche (Size D2, R3
230 [after](#) American Avalanche Association (2016)) which reached the road at the foot of the slope. The failure plane represented by the 21 January profile on PB1 does not extend back to the 12 January surface, suggesting newly accreted snow comprised the bulk of the failure (Figure 5). Cornices on Gruvefjellet experienced comparably minor changes during this event, with the representative volumes decreasing by just 30 m³ and minimal changes evident in the profiles (Figure 5).

Table 1 here

235 A major accretion event in mid-February 2017 followed several weeks of unseasonably high temperatures at cornice elevation during early February (Figure 4). Locally heavy snowfall and strong easterly winds accompanying a vigorous winter storm impacted the region between 19 February and 21 February. Profile GF1's horizontal extension increased by nearly 3 m between the 17 February and 24 February scans resulting in horizontal accretion rates exceeding 15 mm hr⁻¹ (Table 2). The representative volume increased by approximately 100 m³ during the same timeframe. The strong easterly
240 winds stripped the Platåberget cornice system on the windward side of the valley reflected by the abrupt decrease of over 100 m³ to the representative volume there.

Figure 5 here

Representative volumes for both cornice systems gradually increased in the following month, and profiles from 21 March 2017 show considerable rounding and downslope creep of the cornices' leading edges in profiles GF1 and GF3 (Figure 5).

245 Cornices continued to grow on Platåberget, with horizontal growth exceeding 2 m on portions of the PB1 and PB2 profiles and PB3's vertical extent increasing by over 2 m. The Platåberget cornices did not deform downslope to the same degree as the Gruvefjellet cornices during this time period. A representative volume decrease of over 500 m³ (roughly 50% of the volume) on Gruvefjellet in April is related to a major cornice failure near profile GF1, while Platåberget's representative volume increased by 150 m³ in an accretion event near the end of the month (see Section 4.2.1). Considerable cornice
250 accretion is evident in all cornice profiles between 21 March and 1 May except for profiles GF1 and PB2 where we documented cornice failures. Representative volumes continued to increase in early May as light precipitation coincided with continued subzero temperatures. Representative cornice volume on Gruvefjellet gradually increased through 31 May and then dramatically decreased with the onset of sustained positive temperatures at the Gruvefjellet AWS. Cornices on Platåberget continued to accrete through the 18 May scan before beginning to melt away between 18 May and 9 June.

255 **4.1.2 2017/2018**

Figure 6 here

We gathered seven scanned snow surfaces from both Gruvefjellet and Platåberget for the 2017/2018 season with which to compare to meteorological conditions. Cornice development during the 2017/2018 winter season differed considerably from 2016/2017 despite relatively similar seasonal meteorological conditions (Table 3). Gruvefjellet profiles from 15 December
260 2017 show over of 5 m horizontal cornice growth in all profiles, and representative volume approached 1000 m³ (Figures 6 and 7). Contrastingly, the Platåberget plateau margin remained functionally free of snow. Cornices continued to grow over the following five weeks on Gruvefjellet up to the 24 January scan, with profiles GF1 and GF3 reaching their maximum horizontal extensions for the season of nearly 7 m and over 8 m, respectively (Figure 7). Cornice fall avalanches observed on 13 January 2018 are evident in the decreased cornice extension in GF2 in the 24 January scan relative to the 15
265 December 2017 surface and were associated with positive air temperatures and rain at cornice elevation (Table 1). Profiles on Platåberget on 24 January 2018 do not show cornice development, with snow accumulating relatively parallel to the underlying topography.

Table 3 here

Representative volume doubled on Platåberget between the 31 January and 22 February scans from 400 m³ to 900 m³. This
270 coincided with a 0.34 m increase in snow depth at the snow sensor during a snowstorm on 5 February and 6 February where 14 mm of precipitation was measured at the Svalbard Airport AWS (Figure 6). Cornice system changes were more minimal on Gruvefjellet, with a subtle increase of 100 m³ in representative volume. Measured snow depth on Gruvefjellet increased

from 1.45 m on 31 January to a maximum of 1.77 m on 13 February, before slowly decreasing back to 1.48 m by 22 February (Figure 6). A minor decrease in horizontal extension (<1 m) and slight downslope deformation exhibited in profile GF1 are the main observed changes to the cornice profiles between 31 January and 22 February (Figure 7).

Snow depths increased by 0.20 m and 0.28 m on Gruvefjellet and Platåfjellet, respectively, on 26 and 27 February 2018 as over 7 mm precipitation was recorded at the airport (Figure 7). A marked increase in representative volume of 230 m³ on Platåberget between the 22 February and 2 March scans coincides with an increase in snow depth of 0.28 cm over 26 and 27 February. Although a small cornice is evident in profile PB3 on 2 March, increased volume during this time illustrates slope-normal snow depth increase rather than cornice accretion in the representative volume area in the vicinity of PB2 (Figure 7). On Gruvefjellet, downslope creep of the cornice masses continued, with maximum vertical deformation close to 0.80 m for the leading edge of profile GF1. A winter storm on 18 March 2018 resulted in cornice failures seen in both GF1 and GF2 and decreased representative volume on Gruvefjellet, while scouring reduced volume during this time on Platåberget (see Section 4.2.2). Minimal further changes are evident in season's final scan (due to scanner failure) taken 13 April 2018.

Figure 7 here

4.2 Case Studies

4.2.1 Cornice accretion and failure in April 2017

Figure 8 here

290 We documented three periods of cornice fall avalanche activity with TLS data in April 2017. In the first, a small portion of the cornice between profiles GF1 and GF2 failed on 9 April following a period of precipitation falling as snow and easterly winds in excess of 10 m s^{-1} (Figure 8a, Figure 9a annotation 1). The cornice represented by profile GF1 then failed completely on 21 April 2017 coincident with trace precipitation falling as snow and two days of moderate to strong easterly winds (Figure 8b, Figure 10). Negative M3C2 distances displaying changes to the Gruvefjellet cornice system between the
295 21 March 2017 and 25 April 2017 scans show the largest portion of the failed cornice along the axis of profile GF1 (Figure 9a, annotation 2). This failure extended northwards almost 40 m along the cornice. Negative M3C2 distances on the vertical rock face immediately downslope of both the 9 April and 21 April cornice failures show how the falling cornice blocks remove snow from the rock face before impacting avalanche release areas below (Figure 9a, annotation 3). Here, cornice impact craters and small slab avalanche releases are apparent in negative M3C2 distances (Figure 9a, annotation 4).
300 Lower on the slope, the cornice fall avalanche deposition – complete with intact cornice chunks in the avalanche debris – is apparent in strongly positive M3C2 distances (Figure 9a, annotation 5). Other positive M3C2 distances along the cornices (Figure 9a, annotation 6) and horizontal and vertical extent increases on profiles GF2 and GF3 (Figure 10) show cornice accretion occurred elsewhere along Gruvefjellet during this time span. The easterly winds stripped the cornices on Platåberget, evidenced by representative volume decreases of nearly 200 m^3 and vertical extension decreases of up to 1.5 m
305 at profile PB3 (Figure 10).

Figure 9 here

A warm winter storm accompanied by 4.5 mm of precipitation, southwesterly winds, and air temperatures approaching 0°C at cornice level resulted in a period of major cornice accretion and associated cornice fall avalanche activity on the Platåberget cornice system between the 25 April 2017 and 1 May 2017 scans (Figure 8c). Widespread cornice failures are
310 shown by negative M3C2 distances along the Platåberget plateau margin (Figure 9b, annotation 1). These failures coincide with positive M3C2 distances in excess of 1.5 m indicative of cornice accretion elsewhere along the plateau margin (Figure 9b, annotation 2). Profile PB3, for example, experienced over a meter of increased vertical cornice extension (Figure 10) just south of a cornice failure shown in the M3C2 distances (Figure 9b, annotation 3). In profile PB1, 2 m maximum increases in horizontal extension resulted in accretion rates of 17 mm hr^{-1} (Table 2). The semi-vertical profile
315 surface shown in profile PB2 (Figure 10) combined with the M3C2 distance decreases in the profile's immediate surroundings (Figure 9b, annotation 4). indicate cornice failure here occurred after some vertical cornice accretion, as the failure plane extends above the cornice roof from the 25 April snow surface. Cornice blocks released from this cornice and the one immediately to the north poured over cliffs further downslope and gouged impact craters (Figure 9b, annotation 5)

before releasing slab avalanches lower on the slope (Figure 9b, annotation 6). Minimal changes to the Gruvefjellet cornices
320 occurred during this event.

Figure 10 here

4.2.2 Cornice accretion and failure in March 2018

Figure 11 here

A storm in mid-March 2018 punctuated a month of otherwise stable weather and resulted in cornice fall avalanches on
325 Gruvefjellet (Figure 11a). From 15 March to 19 March, 5.6 mm of precipitation accumulated at the Airport AWS, snow
depths at the Gruvefjellet sensor increased by a maximum of 18 cm while those at the Platåberget sensor decreased by
approximately 0.25 m, and strong winds blew from the ENE for 24 hours on 17 March – 18 March. Two large cornice
failures on Gruvefjellet visible as strongly negative M3C2 distances near profile GF1 and slightly to the north (Figure 12a,
annotation 1) triggered avalanches on the slope below (Figure 12a, annotation 2). Similar to the morphology observed in the
330 April 2017 cornice fall avalanches, the failed cornice blocks stripped snow off the vertical rock face and created impact
craters while entraining snow as they moved downslope. The cornice chunks from these cornice failures also remained intact
throughout the event and ran further than the rest of the avalanche debris (Figure 12a, annotation 3). A cornice block
approximately 5 m in horizontal extension detached from the cornice represented by profile GF1, while a smaller (<1 m
horizontal extension) piece detached near GF2 (Figure 13). The GF3 profile did not fail, but over 1 m of snow accreted
335 vertically on the cornice's leading edge. By contrast, Platåberget's plateau margin lost snow, with over snow depth decreases
in excess of 0.20 m measured at the snow station and strongly negative M3C2 distances across the upper portion of the
Platåberget release areas (Figure 12b, annotation 1).

Figure 12 here

Figure 13 here

340 **5 Discussion**

5.1 Seasonal cornice dynamics

TLS-derived cornice data from the 2016/2017 and 2017/2018 winter seasons provide quantitative reinforcement to the
conceptual models of cornice dynamics developed in previous studies (e.g. Montagne et al., 1968; Vogel et al., 2012). In
these models, cornices accrete through relatively discrete events and begin to deform under their own weight before either
345 failing or melting away towards the end of the snow season.

Our data show cornices can rapidly accrete at any point in the snow season given abundant snow available for wind
transport, wind speeds sufficient to mobilize surface snow, and wind directions oriented relatively perpendicular to the

ridgeline. We documented accretion rates in excess of 15 mm hr^{-1} at various times throughout the 2016/2017 season on each side of the valley, which can be considered a minimum given the relatively poor temporal constraints on the accretion events provided by the TLS data. Each of these periods of accretion coincided with measured precipitation at the airport, wind speeds in excess of 5 m s^{-1} , and wind directions roughly placing the plateau margin in the lee. The relatively small proportion of the winter seasons characterized by meteorological conditions conducive for accretion suggests just a few accretion events play a key role in cornice development each season (Table 3). Asynchronous cornice responses on Gruvefjellet and Platåberget to specific weather events further illustrate the importance of wind direction in controlling cornice dynamics. During the February 2017 event, for example, cornices on Gruvefjellet rapidly accreted and gained volume with plentiful snow available for transport and strong easterly winds. Cornices on Platåberget lost volume, however, as they were eroded by the same easterly winds. Similar out of phase behavior was exhibited in late April 2017, when precipitation and westerly winds resulted in considerable cornice growth on Platåberget accompanied by slight decreases to horizontal and vertical extension in profiles GF2 and GF3 and minimal representative volume changes near profile GF1. Following initial accretion, the cornices' leading edges begin to deform downslope. Deformation becomes more pronounced later in the season, presumably as increased air temperatures and solar radiation begin to warm the snow, decreasing the stiffness of the cornices and increasing creep (e.g. Schweizer et al., 2003). Further accretion events can then be superimposed on this deformation as the season progresses, with short accretion events interspersed by longer periods of downslope creep. This can be seen in the minor increases in horizontal extension and continued downslope deformation in GF1 and GF3 through the latter portion of the 2017/2018 season (Figure 7). Cornice accretion and downslope deformation can also occur almost simultaneously with air temperatures approaching or even exceeding freezing at cornice level, as evidenced by the rapid accretion and downslope creep shown in profile PB1 for the 25 April – 1 May scan interval (Figure 10).

While meteorological conditions control the specific timing of cornice accretion and downslope deformation, the underlying topography appears to act as a fundamental control on cornice structure and seasonal cornice dynamics. The presence of the steep bedrock face directly beneath the Gruvefjellet plateau margin limits the support provided by the underlying topography compared to the gentler sloping Platåberget margin. The result is a more overhung cornice structure on Gruvefjellet, while Platåberget's topography allows for more slope-normal snow accumulation. Profiles PB1 and PB2 failed to develop cornices at all during the 2017/2018 winter season (Figure 7). The presence of cornices with horizontal extension approaching 5 m in these locations during the 2016/2017 winter season (Figure 5), however, shows the topography can support cornice development given the right meteorological conditions. Differences in meteorological conditions between the 2016/2017 and 2017/2018 winter seasons may provide a partial explanation for differing seasonal snow cover responses on Platåberget (Table 3). Winds in excess of 5 m s^{-1} – a conservative estimate for threshold wind speeds required to mobilize loose snow (Li and Pomeroy, 1997) – from the western quadrant conducive to cornice accretion on Platåberget were slightly less prevalent during the 2017/2018 season, and these winds also coincided with precipitation roughly half as frequently as during the

2016/2017 season. Easterly winds exceeding 5 m s^{-1} were considerably more prevalent during the 2017/2018 season which may have increased cornice scouring or limited snow available for transport –and thus accretion – on Platåberget. Nevertheless, the meteorological differences between the two winter seasons are subtle enough when compared to the noteworthy differences in cornice dynamics to [that](#) suggest specific interactions between meteorology and topography not necessarily captured by our analyses meaningfully impact cornice development.

Topography also seems to control the relative size of cornice failures. Vogel et al. (2012) describe a “geomorphologically determined sedimentary step approximately 3 m below the plateau that most likely acts as the cornice pivot point” on Gruvefjellet. This pivot point is most evident in profile GF1, where in both winter seasons the downslope creep of the overhung cornice beyond this pivot point ultimately became overburdened during an accretion event and caused the cornice to fail completely. The cornice represented by GF1 has the least topographic support and developed the most overhanging cornice structure of the specific cornices we investigated, [andwhile](#) also failed completely both seasons. By contrast, the topographic support provided by Platåberget does not promote overhanging cornices to the same degree, instead promoting a thicker slope-normal snowpack which in itself supports the cornice structure. Here, observed cornice failures such as that shown in PB2 during the 25 April - 1 May 2017 scan interval (Figure 10) are limited to the recently accreted snow and did not involve the entire cornice mass. Similarly, profile GF2 failed in March 2018 within hours of profile GF1’s full failure, but involved a much smaller portion of the cornice predating the 2 March scan – potentially related to increased topographic support to this cornice relative to GF1 (Figure 13).

5.2 Cornice fall avalanches

Previous work has differentiated cornice fall avalanche types by the inferred mechanism of cornice failure – either via increased snow load from accretion or decreased snow strength in the cornice related to increased snow and air temperatures. Five of the six cornice fall avalanche events observed in this study coincided with winter storms leading to accretion just prior to cornice failure (Table 1). This is in contrast to previous findings from this location, where no cornice failures were observed in direct response to snow loading caused by a snow storm (Vogel et al., 2012). The lone cornice fall avalanche event we cannot link to cornice accretion occurred in January 2018. This event coincided with heavy precipitation, but positive temperatures at the Gruvefjellet AWS and decreasing snow depths at the Gruvefjellet snow sensor indicate this precipitation fell as rain (Figure 6). Our truncated TLS observation record in late-spring 2018 unfortunately omits the May-June period found by Vogel et al. (2012) to be critical for air temperature induced cornice failures in this location, but observational records throughout this time do not indicate further cornice fall avalanches. Accretion’s role in determining cornice failure is also reflected in the asynchronous timing of cornice failures on Gruvefjellet and Platåberget during our study. None of the observed avalanche events included activity on both Gruvefjellet and Platåberget simultaneously as would be expected with air temperature induced failures, with avalanches instead occurring only on the leeward aspect.

Observed cornice fall avalanche size appears to be controlled largely by the snow conditions in the underlying release area. Cornice fall avalanches on Gruvefjellet follow a pattern exemplified by the April 2017 case study in which the cornice fails and removes snow from the steep bedrock face below as it descends before impacting the release areas at the base of the cliff
415 (e.g. Figure 9a). The cornice block can then, depending on the snow conditions in the release area, entrain snow from its impact crater and the avalanche path below or trigger a larger slab avalanche. Cornice failures near profile GF3 in both April 2017 and March 2018 triggered small slab avalanches, but the majority of the avalanche debris resulted from entrainment as the cornice blocks bounced downslope.

Platåberget's topography promotes slightly different avalanche dynamics. The gentler slope at the plateau edge allows snow
420 to accumulate directly under the cornices such that failed cornice masses land directly on the snow to be released as an avalanche. Release areas on Platåberget collect snow during accretion events much more efficiently than those on Gruvefjellet, where blowing snow mass losses due to suspension are promoted by the separation created between the cornices and the release areas by the bedrock cliff. Accumulation in the upper release areas on Platåberget coinciding with accretion events primes these locations for slab avalanche release with even small cornice failures. Relatively small cornice
425 failures triggering larger slab avalanches on Platåberget in April 2017 resulted in comparable magnitude avalanches (D2, R2-R3) to those releasing from much larger cornice failures but less entrainable snow on Gruvefjellet in March 2018 (Figure 9b and Figure 13a).

5.3 Hazard management implications

Cornice fall avalanches are the most common avalanche type observed in the portion of central Svalbard surrounding our
430 study area where the broad plateau summits and steep valley walls of Longyeardalen's topography are recurrent across the region (Eckerstorfer and Christiansen, 2011c). Cornice fall avalanches observed in this study thus represent processes occurring elsewhere throughout central Svalbard – and to a lesser extent other locations throughout the world– and provide an opportunity to reinforce existing forecasting frameworks with detailed cornice data. The conceptual model of avalanche hazard in North America treats cornice failure both as an individual avalanche problem to be considered by forecasters and
435 as a potential trigger when assessing the likelihood of other avalanche types releasing in a given forecasting area and time period (Statham et al., 2018). Cornice fall avalanche hazard assessments should thus consider both the likelihood of cornice failure and the nature of the snow conditions in the release area to best judge cornice fall avalanche hazard. Our limited dataset, especially in the absence of multiple air temperature induced failures, is insufficient to make broad generalizations linking cornice failure type and resulting cornice fall avalanche activity. As a specific example, however, fairly widespread
440 wind slab avalanche activity throughout the region accompanied each of the accretion-induced avalanche events observed in this work. The conditions leading to cornice accretion – strong winds and available snow for wind transport – also promote the development of wind slab problems. Thus, when conditions are favorable for cornice accretion and accretion-induced
cornice failures, conditions are also favorable for the ~~possible~~ development of more widespread – and potentially more

sensitive – slab avalanche problems. In this scenario, the chance of a cornice failure triggering a secondary slab avalanche would rise, subsequently amplifying the cornice fall avalanche hazard by also increasing the expected size of the resulting cornice fall avalanche. Furthermore, in all cornice fall avalanches observed on Gruvefjellet the main cornice blocks travelled further downslope than the rest of the avalanche debris. This pattern is apparent on larger failures on Platåberget as well, but is in some cases less obvious, likely due to the smaller cornice blocks being functionally indistinguishable from the avalanche debris. While the dataset presented here is insufficient to draw more quantitative conclusions regarding the runout distance of these cornice blocks, hazard management strategies should consider the destructive potential and extended runout of these blocks relative to the other entrained snow.

5.4 Uncertainties and limitations

The TLS data acquisition and processing techniques employed in this work allowed us to illustrate and quantify changes to the observed cornice systems in detail not previously achieved, but our results and subsequent interpretations are nonetheless limited by several factors. Measurement uncertainties specifically related to measuring snow surfaces with TLS are well-discussed in previous research (Deems et al., 2015; Prokop, 2008), but we introduced additional uncertainty to our results and interpretations due to the scan timing. Our TLS data acquisition scheme involved time-intensive manual input, so we were unable to achieve the temporal resolution required to better constrain individual accretion and cornice failure events. Decreasing time between scans would allow for more continuous and robust accretion rate calculations and could better constrain failure and avalanche snow surfaces, especially pre-event. Sufficiently decreasing the between-scan interval to a sub-daily resolution for such applications would likely require some degree of automation, and future work should consider employing a permanently-installed TLS acquiring data automatically similar to systems employed for mining applications or slope stability assessments.

Uncertainties in cornice volume calculations are also affected by occasionally lengthy inter-scan intervals. Volume changes corresponding to specific meteorological conditions are in these cases aggregated across the entire scan interval, making disentangling the specific contributions to volume changes difficult. These conceptual uncertainties are magnified by the technical uncertainties related to TLS data acquisition. The TLS accuracy is of increased importance for volume quantification as measurement uncertainties are propagated throughout the volume calculation process. However, calculated volume uncertainties (Appendix 1) are sufficiently low to instill a degree of confidence in the volume calculation process presented here. Finally, volume calculations are perhaps least robust in this study for times when the lack of obvious cornice structure makes calculating volumes particularly challenging (e.g. Platåberget during the 2017/2018 season).

Our experimental design focused on investigating the evolution and failure of the lower cornice surfaces from scan positions underneath the cornices where accessibility has precluded previous research. These scan positions did not, however, allow for systematic monitoring of the cornice roof. The orientation of the cornices' leading edges frequently shielded the cornice roof from the scanner, and our profiles often do not include the complete cornice roof. This also has implications for

representative volume calculations, as uncertainty in the location of the cornice roof can result in inaccurate horizontal difference calculations in these specific locations. By failing to capture the cornice roof in our data, we also limit comparisons with earlier work on Gruvefjellet relating downslope cornice deformation and cornice failure to the appearance of tension cracks between the cornice roof and the plateau anchoring point (Vogel et al., 2012). Future work should pair TLS data with some form of tension crack observation, and approaches combining TLS and UAV photogrammetry present intriguing possibilities for future work in this and other locations.

TLS was shown to be a particularly suitable remote sensing tool for cornice monitoring in Svalbard where we were able to obtain useful data during the early winter seasons when the polar night precludes direct visual observation and cornice photography. Svalbard's unique environmental characteristics – such as the polar night – limit to a degree the applicability of our results to lower latitudes where more diurnal variations in radiation and temperature may influence cornice dynamics in ways not represented in Svalbard (e.g. Munroe, 2018). It is also unclear how representative the two winter seasons for which we present data are for the cornice systems in Longyeardalen, as previous research has also noted considerable differences in cornice dynamics between seasons (Vogel et al., 2012). Continued cornice monitoring in this and other lower latitude settings would help clarify such uncertainties.

490 **6 Conclusions**

We monitored seasonal cornice dynamics and associated cornice fall avalanche activity with a TLS over two winter seasons in high-Arctic Svalbard. The spatial and temporal resolution at which we acquired snow surface data with the TLS allowed us to quantify changes to the cornices with sub-decimeter accuracy. These data provide quantitative reinforcement to existing conceptual models of cornice dynamics and further strengthen the validity of these models. Notable quantitative contributions from this work include documentation of conservatively calculated horizontal accretion rates well in excess of 10 mm hr⁻¹ and a methodology for calculating cornice volumes from TLS data.

This study demonstrated the viability of TLS methods for monitoring cornice dynamics. TLS methods for obtaining snow surface data are suitable in Svalbard where the long polar night precludes data acquisition via other methods (e.g. photogrammetry), but techniques presented in this work are also suitable for cornices in other, lower latitude environments. Future work should investigate automated TLS data acquisition as an avenue to improve the temporal resolution of the measurements and better constrain cornice dynamics to specific meteorological conditions.

Our findings show complex interactions between topography, wind speed and direction, snow available for transport, existing snowpack, and cornice structure govern the growth, failure, and associated avalanche activity of the cornices in Longyeardalen. In particular, we show cornices rapidly accrete given winds strong enough to mobilize surface snow from a direction roughly perpendicular to the plateau edge, placing the cornices in the lee. Our findings also reinforce previous work indicating an increased likelihood of cornice failure and associated avalanche activity during these periods of cornice

accretion. This is encouraging for hazard managers seeking to forecast cornice fall avalanches, as anticipating the relatively infrequent conditions leading to cornice accretion can help predict periods of elevated cornice fall avalanche hazard. We observed the largest failures in our dataset in areas with minimal topographic support, demonstrating knowledge of the topography underlying the cornices can be beneficial when considering the specific location of cornice failure. Nevertheless, our limited dataset of cornice failures hinders conclusions drawn from this work, and continued work in a variety of environments is needed to better understand the specific mechanisms and dynamics of cornice fall avalanches.

515 *Data Availability.* Data for the Svalbard Airport AWS are available through Norwegian Meteorological Institute's online data accessibility platform (www.eklima.no). Data from the Gruvefjellet AWS are freely available through the University Centre in Svalbard via www.unis.no/resources/weather-stations/. All TLS, snow depth, and other auxiliary data can be obtained by contacting the corresponding author by email (holt.hancock@unis.no).

520 *Author contributions.* HH was responsible for the majority of the data acquisition, analyses, and interpretation of the results. ME helped develop the conceptual framework for the study and contextualized and interpreted the results within a broader snow and avalanche perspective. AP provided technical guidance with regards to TLS data acquisition and analysis techniques and assisted in the development of the study's technical framework in addition to assisting in data acquisition. JH provided advice and supervision relating to study design, data analysis, and interpretation of the results. HH and ME were responsible for manuscript preparation with input from AP and JH.

525 *Competing interests.* The authors declare they have no conflicts of interest.

Acknowledgements. We thank Dr. Christine Fey and Dr. Jeffrey Munroe for their thorough and constructive reviews which greatly improved this work. Dr. Andreas Günther is thanked for serving as editor for this work.

References

- Abellán, A., Oppikofer, T., Jaboyedoff, M., Rosser, N. J., Lim, M., and Lato, M. J.: Terrestrial laser scanning of rock slope instabilities, *Earth Surface Processes and Landforms*, 39, 80-97, doi:10.1002/esp.3493, 2014.
- 530 American Avalanche Association: *Snow, Weather and Avalanches: Observation Guidelines for Avalanche Programs in the United States* (3rd ed). Victor, ID, 104 pp., 2016.
- Anderton, S. P., White, S. M., and Alvera, B.: Evaluation of spatial variability in snow water equivalent for a high mountain catchment, *Hydrological Processes*, 18, 435-453, doi:10.1002/hyp.1319, 2004.
- 535 Caputo, T., Marino, E., Matano, F., Somma, R., Troise, C., and De Natale, G.: Terrestrial Laser Scanning (TLS) data for the analysis of coastal tuff cliff retreat: application to Coriglio cliff, Naples, Italy, 2018, 61, doi:10.4401/ag-7494, 2018.
- Christiansen, H. H., Humlum, O., and Eckerstorfer, M.: Central Svalbard 2000-2011 meteorological dynamics and periglacial landscape response, *Arctic, Antarctic, and Alpine Research*, 45, 6-18, doi:10.1657/1938-4246-45.16, 2013.
- 540 Deems, J. S., Painter, T. H., and Finnegan, D. C.: Lidar measurement of snow depth: a review, *Journal of Glaciology*, 59, 467-479, doi:10.3189/2013JoG12J154, 2013.
- Deems, J. S., Gadowski, P. J., Vellone, D., Evanczyk, R., LeWinter, A. L., Birkeland, K. W., and Finnegan, D. C.: Mapping starting zone snow depth with a ground-based lidar to assist avalanche control and forecasting, *Cold Regions Science and Technology*, 120, 197-204, doi:10.1016/j.coldregions.2015.09.002, 2015.
- Eckerstorfer, M., and Christiansen, H. H.: The "High Arctic Maritime Snow Climate" in Central Svalbard, *Arctic, Antarctic, and Alpine Research*, 43, 11-21, doi:10.1657/1938-4246-43.1.11, 2011a.
- 545 Eckerstorfer, M., and Christiansen, H. H.: Topographical and meteorological control on snow avalanching in the Longyearbyen area, central Svalbard 2006-2009, *Geomorphology*, 134, 186-196, doi:10.1016/j.geomorph.2011.07.001, 2011c.
- Eckerstorfer, M., Christiansen, H. H., Rubensdotter, L., and Vogel, S.: The geomorphological effect of cornice fall avalanches in the Longyeardalen valley, Svalbard, *The Cryosphere*, 7, 1361-1374, doi:10.5194/tc-7-1361-2013, 2013.
- 550 Farizy, B.: Cornice control by helicopter, *International Snow Science Workshop, Grenoble - Chamonix Mont-Blanc*, 2013, 845-847.
- Fey, C., Schattan, P., Helfricht, K., and Schöber, J.: A compilation of multitemporal TLS snow depth distribution maps at the Weisssee snow research site (Kaunertal, Austria), *Water Resources Research*, 55, 5154-5164, doi:10.1029/2019wr024788, 2019.
- Førland, E., Benestad, R., Hanssen-Bauer, I., Haugen, J. E., and Skaugen, T. E.: Temperature and precipitation development at Svalbard 1900-2100, *Advances in Meteorology*, 2011, doi:10.1155/2011/893790, 2011.
- 555 Hanssen-Bauer, I., Solås, M. K., and Steffensen, E.: *The Climate of Spitsbergen*, Det Norske Meteorologiske Institutt, 40 pp., 1990.
- Hanssen-Bauer, I., Førland, E., Hisdal, H., Mayer, S., Sando, A., and Sorteberg, A.: *Climate in Svalbard 2100 - a knowledge base for climate adaptation*, Norwegian Centre for Climate Services/2019, 207, 2019.
- Hartzell, P. J., Gadowski, P. J., Glennie, C. L., Finnegan, D. C., and Deems, J. S.: Rigorous error propagation for terrestrial laser scanning with application to snow volume uncertainty, *Journal of Glaciology*, 61, 1147-1158, doi:10.3189/2015JoG15J031, 2017.
- 560 Humlum, O., Instanes, A., and Sollid, J. L.: Permafrost in Svalbard: a review of research history, climatic background and engineering challenges, *Polar Research*, 22, 191-215, 2003.
- Humlum, O., Christiansen, H. H., and Juliussen, H.: Avalanche-derived rock glaciers in Svalbard, *Permafrost and Periglacial Processes*, 18, 75-88, 2007.
- 565 Isaksen, K., Nordli, Ø., Førland, E. J., Lúpikasza, E., Eastwood, S., and Niedźwiedz, T.: Recent warming on Spitsbergen—Influence of atmospheric circulation and sea ice cover, *Journal of Geophysical Research: Atmospheres*, 121, 11,913-911,931, doi:10.1002/2016JD025606, 2016.
- Jaboyedoff, M., Oppikofer, T., Abellán, A., Derron, M.-H., Loye, A., Metzger, R., and Pedrazzini, A.: Use of LIDAR in landslide investigations: a review, *Natural Hazards*, 61, 5-28, doi:10.1007/s11069-010-9634-2, 2012.
- Jaedicke, C., and Sandvik, A. D.: High resolution snow distribution data from complex Arctic terrain: a tool for model validation, *Natural Hazards and Earth System Science*, 2, 147-155, 2002.
- 570 Kobayashi, D., Ishikawa, N., and Nishio, F.: Formation process and direction distribution of snow cornices, *Cold Regions Science and Technology*, 15, 131-136, doi:10.1016/0165-232X(88)90059-6, 1988.
- Lague, D., Brodu, N., and Leroux, J.: Accurate 3D comparison of complex topography with terrestrial laser scanner: Application to the Rangitikei canyon (N-Z), *ISPRS Journal of Photogrammetry and Remote Sensing*, 82, 10-26, doi:10.1016/j.isprsjprs.2013.04.009, 2013.
- 575 Latham, J., and Montagne, J.: The Possible Importance of Electrical Forces in the Development of Snow Cornices, *Journal of Glaciology*, 9, 375-384, doi:10.3189/S0022143000022899, 1970.
- Li, L., and Pomeroy, J. W.: Estimates of Threshold Wind Speeds for Snow Transport Using Meteorological Data, *Journal of Applied Meteorology*, 36, 205-213, doi:10.1175/1520-0450(1997)036<0205:EOTWSF>2.0.CO;2, 1997.
- 580 Major, H., Haremo, P., Dallmann, W.K., Andresen, A., Salvigsen, O.: *Geological map of Svalbard 1: 100,000, sheet C9G Adventdalen*, Norsk Polarinstitutt, 2001.

- McCarty, D., Brown, R., and Montagne, J.: Cornices: their growth, properties, and control, International Snow Science Workshop, Lake Tahoe, CA, USA, 1986, 41-45,
- 585 Montagne, J., McPartland, J., Super, A., and Townes, H.: The Nature and control of snow cornices on the Bridger Range, Southwestern Montana, Alta Avalanche Study Center, Miscellaneous Report, 1968.
- Mott, R., Schirmer, M., and Lehning, M.: Scaling properties of wind and snow depth distribution in an Alpine catchment, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- Munroe, J. S.: Monitoring snowbank processes and cornice fall avalanches with time-lapse photography, *Cold Regions Science and Technology*, 154, 32-41, doi:10.1016/j.coldregions.2018.06.006, 2018.
- 590 Naruse, R., Nishimura, H., and Maeno, N.: Structural Characteristics of Snow Drifts and Cornices, *Annals of Glaciology*, 6, 287-288, 1985.
- Prokop, A.: Assessing the applicability of terrestrial laser scanning for spatial snow depth measurements, *Cold Regions Science and Technology*, 54, 155-163, doi:10.1016/j.coldregions.2008.07.002, 2008.
- 595 Prokop, A., Schirmer, M., Rub, M., Lehning, M., and Stocker, M.: A comparison of measurement methods: terrestrial laser scanning, tachymetry and snow probing for the determination of the spatial snow-depth distribution on slopes, *Annals of glaciology*, 49, 210-216, doi:10.3189/172756408787814726, 2008.
- Prokop, A., and Panholzer, H.: Assessing the capability of terrestrial laser scanning for monitoring slow moving landslides, *Natural Hazards and Earth Systems Science*, 9, 1921-1928, doi:10.5194/nhess-9-1921-2009, 2009.
- 600 Prokop, A., Schön, P., Singer, F., Pulfer, G., Naaim, M., Thibert, E., and Soruco, A.: Merging terrestrial laser scanning technology with photogrammetric and total station data for the determination of avalanche modeling parameters, *Cold Regions Science and Technology*, 110, 223-230, 2015.
- Prokop, A., and Procter, E. S.: A new methodology for planning snow drift fences in alpine terrain, *Cold Regions Science and Technology*, 132, 33-43, doi:10.1016/j.coldregions.2016.09.010, 2016.
- 605 Prokop, A., Hancock, H., Praz, M., and Jahn, E.: Slope scale avalanche forecasting in the arctic (Svalbard), International Snow Science Workshop, Innsbruck, Austria, 2018, 1035-1039,
- Riegl VZ-6000 Datasheet: http://www.riegl.com/uploads/tx_pxprriegl/downloads/RIEGL_VZ-6000_Datasheet_2019-05-28_01.pdf, 2019.
- Rogers, J. C., Yang, L., and Li, L.: The role of Fram Strait winter cyclones on sea ice flux and on Spitsbergen air temperatures, *Geophysical Research Letters*, 32, 2005.
- 610 Schirmer, M., Wirz, V., Clifton, A., and Lehning, M.: Persistence in intra-annual snow depth distribution: 1. Measurements and topographic control, *Water Resources Research*, 47, 2011.
- Schön, P., Prokop, A., Vionnet, V., Guyomarc'h, G., Naaim-Bouvet, F., and Heiser, M.: Improving a terrain-based parameter for the assessment of snow depths with TLS data in the Col du Lac Blanc area, *Cold Regions Science and Technology*, 114, 15-26, doi:10.1016/j.coldregions.2015.02.005, 2015.
- Schweizer, J., Bruce Jamieson, J., and Schneebeli, M.: Snow avalanche formation, *Reviews of Geophysics*, 41, 2003.
- 615 Seligman, G.: Snow structure and ski fields: being an account of snow and ice forms met with in nature and a study on avalanches & snowcraft., Macmillan, London, 1936.
- Statham, G., Haegeli, P., Greene, E., Birkeland, K., Israelson, C., Tremper, B., Stethem, C., McMahon, B., White, B., and Kelly, J.: A conceptual model of avalanche hazard, *Natural Hazards*, 90, 663-691, doi:10.1007/s11069-017-3070-5, 2018.
- 620 van Herwijnen, A., and Fierz, C.: Monitoring snow cornice development using time-lapse photography, *Proceedings of the International Snow Science Workshop*, 2014, Banff, Canada, 2014, 865-869,
- Vikhamar-Schuler, D., Isaksen, K., Haugen, J. E., Tommervik, H., Luks, B., Schuler, T. V., and Bjerke, J. W.: Changes in winter warming events in the Nordic Arctic region, *Journal of Climate*, 29, 6223-6244, doi:10.1175/jcli-d-15-0763.1, 2016.
- Vionnet, V., Martin, E., Masson, V., Guyomarc'h, G., Naaim Bouvet, F., Prokop, A., Durand, Y., and Lac, C.: Simulation of wind-induced snow transport and sublimation in alpine terrain using a fully coupled snowpack/atmosphere model, *Cryosphere*, 8, p. 395 - p. 415, doi:10.5194/tc-8-395-2014, 2014.
- 625 Vogel, S., Eckerstorfer, M., and Christiansen, H.: Cornice dynamics and meteorological control at Gruevfjellet, Central Svalbard, *The Cryosphere*, 6, 157-171, doi:10.5194/tc-6-157-2012, 2012.
- Watson, C. S., Quincey, D. J., Smith, M. W., Carrivick, J. L., Rowan, A. V., and James, M. R.: Quantifying ice cliff evolution with multi-temporal point clouds on the debris-covered Khumbu Glacier, Nepal, *Journal of Glaciology*, 63, 823-837, doi:10.1017/jog.2017.47,
- 630 2017.

Tables

Table 1. Summary of avalanche cycles.

Event date	Area	Trigger	Pre-event scan date	Post-event scan date	# observed cornice fall avalanches
21 January 2017	Platåberget	Accretion	12 January 2017	22 January 2017	>3
09 April 2017	Gruvefjellet	Accretion	21 March 2017	25 April 2017	1
21 April 2017	Gruvefjellet	Accretion	21 March 2017	25 April 2017	1
29 April 2017	Platåberget	Accretion	25 April 2017	01 May 2017	>3
14 January 2018	Gruvefjellet	Temperature (rain)	15 December 2017	24 January 2018	1
18 March 2018	Gruvefjellet	Accretion	2 March 2018	23 March 2018	2

635

Table 2. Summary of well-constrained accretion events


Area	Pre-event scan date and time (UTC)	Post-event scan date and time (UTC)	Between scan interval (hrs)	Profile with max horizontal accretion	Max horizontal accretion (m)	Accretion rate (mm hr ⁻¹)
Platåberget	12 January 2017 1930	21 January 2017 2100	217.5	PB2	3.6	17
Gruvefjellet	17 February 2017 0900	24 February 2017 1100	170	GF1	2.9	17
Platåberget	25 April 2017 1315	01 May 2017 0945	140.5	PB1	2.0	14

Table 3. Seasonal summaries. All parameters are measured at the Gruvefjellet AWS except for precipitation, which is measured at the Svalbard Airport AWS.


	2016/2017	2017/2018
Mean seasonal air temperature	-9.3	-7.5
Accumulated precipitation (mm)	125.6	124.5
% hours in season with accretion winds on Platåberget	5.7	5.1
% hours in season with accretion winds on Platåberget and daily precipitation > 0.2 mm	3.0	1.7
% hours in season with accretion winds on Gruvefjellet	13.5	21.7
% hours in season with accretion winds on Gruvefjellet and daily precipitation > 0.2 mm	4.0	4.1
Accretion winds on Platåberget: wind speed > 5 m s ⁻¹ , 225° < wind direction < 315°		
Accretion winds on Gruvefjellet: wind speed > 5 m s ⁻¹ , 45° < wind direction < 135		

640


Figures



645 | **Figure 1.** Overview of Longyeardalen and key locations, including automated weather stations (AWS), mentioned in the text. Contour lines in panel (a) are spaced at 100 m. The location and direction from which the photo in panel (b) was taken is indicated by **POV** in (a). The location and extent of the Gruvefjellet and Platåberget study sites are indicated in (a) and (b) with green and red shading, respectively. Locations of scan positions SP1, SP2, SP3, and SP4 as well as the orientation of the scanner at each position are also indicated.



650 **Figure 2.** Overview of the cornice systems and locations of the primary spatial data employed in this work. Panel (a) shows Gruvefjellet
 from SP1 taken on 21 March 2017, with the white rectangle approximating the 600 m horizontal extent of panel (b), and the red square
 showing the area used to assess relative accuracy. Panel (b) indicates the location of the 2D cross-sectional profiles GF1, GF2, and GF3, in
 addition to outlining the area used to calculate representative cornice volume in red. Panel (c) shows Platåberget from SP4 taken on 24
 655 May 2017, with the white rectangle approximating the 600 m horizontal extent of panel (d), and the red square showing the area used to
 assess relative accuracy. Panel (d) indicates the location of the 2D cross-sectional profiles PB1, PB2, and PB3, in addition to outlining the
 area used to calculate representative cornice volume in red. Snow depth sensor locations are indicated in all panels with the black dot.



660 **Figure 3.** Visualization of the point cloud processing methods in CloudCompare. Panel (a) shows a photo of the cornice represented by
 profile GF1 on 21 March 2017. Panel (b) shows the same surface as represented by the 0.10 m point cloud. The manually defined axis of
 GF1 is indicated by the white line. Panels (c) and (d) show the 21 March scanned surface and extracted profile from two vantage points.
 Panel (e) displays both the 21 March (colored points) and bare-earth (white points) surfaces oriented parallel to the projection direction,
 with the 21 March profile (green) and bare-earth profile (white) also indicated. Panels (f) and (h) display similar data but with the surfaces
 oriented roughly perpendicular to the projection direction (shown with red arrows), and panel (h) shows a cross-section of the surface
 665 shown in panels (e) and (f). The 1 m grid showing the horizontal differences between the 21 March and bare-earth scan is displayed in
 panel (g). All scale bars are in meters





Figure 4. Summary of the representative cornice volume progression and meteorological conditions for the 2016/2017 winter season. Wind speed and air temperature are daily averaged values from the Gruvefjellet AWS, and precipitation data are daily values from the Svalbard Airport AWS measured at 0600 UTC. Shaded blue vertical bars indicate well-constrained cornice accretion periods for which we were able to calculate horizontal cornice accretion rates (Table 2). Shaded grey vertical bars indicate 48 hour periods with observed noteworthy cornice fall avalanche activity (Table 1).

670



675 **Figure 5.** 2D cornice profiles showing cornice progression for selected scan dates throughout the 2016/2017 winter season. Each profile is labeled as it referred to in the text and corresponds to the location and POV depicted in Figure 2





Figure 6. Summary of the representative cornice volumes and meteorological conditions for the 2017/2018 winter season. Wind speed and air temperature are daily averaged values from the Gruvefjellet AWS, precipitation data are daily values from the Svalbard Airport AWS measured at 0600 UTC, and snow depths are daily averages from the snow sensors on Gruvefjellet and Platåfjellet. Shaded grey vertical bars indicate 48 hour periods with observed noteworthy cornice fall avalanche activity (Table 2).

680



685 **Figure 7.** 2D cornice profiles showing cornice progression for 2017/2018 winter season scan dates. Each profile is labeled as it referred to in the text and corresponds to the location and POV depicted in Figure 2.




Figure 8. Meteorological summary of the April 2017 case study. Wind speed, wind direction, and air temperature are hourly values from the Gruvefjellet AWS, and precipitation data are daily values from the Svalbard Airport AWS measured at 0600 UTC. Colored vertical lines in the time series indicate the scan timing corresponding to the profiles in Figure 9. Vertical grey bars marked (a), (b), and (c) correspond to 48 hour time periods with noteworthy avalanche activity discussed in the text.

690






Figure 9. M3C2 distances displaying changes to the snow cover on Gruvefjellet between the 21 March and 25 April 2018 scans (a) and on Platåberget between the 25 April and 1 May 2018 scans (b). Red rectangles in both panels indicate the locations of the cornice profiles. Specific snow surface features are annotated as they are referred to in the text.




695

Figure 10. Cornices profiles illustrating cornice dynamics during the April 2017 case study, with each profile labeled as it is referred to in the text. Dashed lines indicate interpolated data where overhanging cornice structure shadowed the snow surface from the TLS.



700 **Figure 11.** Meteorological summary of the March 2018 case study. Wind speed, wind direction, and temperature are hourly values from the Gruvefjellet AWS, and precipitation data are daily values from the Svalbard Airport AWS measured at 0600 UTC. Colored vertical lines in the time series indicate the scan timing corresponding to the profiles in Figure 12, and the grey vertical bar annotated with (a) corresponds to the 48 hour time period with noteworthy avalanche activity mentioned in the text.



705 **Figure 12.** M3C2 distances displaying changes to the snow cover on Gruvefjellet (a) and Platåberget (b) between the 2 March and 23 March 2018 scans. Red rectangles in both panels indicate the locations of the cornice profiles. Specific snow surface features are annotated as they are referred to in the text, and snow depth sensors are marked and labeled.




Figure 13. Cornices profiles illustrating cornice dynamics during the March 2018 case study, with each profile labeled as it is referred to in the text. Dashed lines indicate interpolated data where overhanging cornice structure shadowed the snow surface from the TLS.

Appendix 1

<u>Date and Time (UTC)</u>	<u>Area</u>	<u>Scan Position</u>	<u>Mean Relative Error (m)</u>	<u>Standard Deviation (m)</u>	<u>Representative Volume (m³)</u>	<u>Volume Uncertainty</u>
16 September 2016 12:00	Gruvefjellet	S1	NA	NA	NA	NA
16 September 2016 13:00	Platåberget	S3	NA	NA	NA	NA
02 December 2016 10:15	Gruvefjellet	S1	0.036	0.001	179	13.173
05 December 2016 12:50	Gruvefjellet	S1	0.019	0.016	159	6.827
12 January 2017 17:30	Gruvefjellet	S4	0.047	0.028	649	17.343
12 January 2017 19:30	Platåberget	S3	0.067	0.039	328	24.612
21 January 2017 21:00	Platåberget	S3	0.032	0.044	619	11.956
22 January 2017 14:00	Gruvefjellet	S2	0.030	0.025	633	11.144
27 January 2017 08:50	Gruvefjellet	S1	0.024	0.016	689	8.930
03 February 2017 08:00	Platåberget	S4	0.000	0.044	688	0.148
14 February 2017 14:00	Platåberget	S4	0.066	0.039	1145	24.170
14 February 2017 15:00	Gruvefjellet	S1	0.022	0.020	789	7.970
17 February 2017 09:00	Gruvefjellet	S1	0.032	0.030	823	11.771
17 February 2017 10:00	Platåberget	S4	0.055	0.039	1190	20.111
22 February 2017 10:45	Platåberget	S3	0.042	0.037	1072	15.350
24 February 2017 11:00	Gruvefjellet	S1	0.027	0.031	970	9.926
12 March 2017 16:00	Platåberget	S4	0.003	0.046	1224	1.033
12 March 2017 17:30	Gruvefjellet	S2	0.034	0.027	1043	12.435
21 March 2017 13:10	Gruvefjellet	S1	0.031	0.027	1117	11.402
21 March 2017 14:10	Platåberget	S4	0.064	0.042	1255	23.616
25 April 2017 10:00	Gruvefjellet	S1	0.068	0.027	581	25.018
25 April 2017 13:15	Platåberget	S4	0.007	0.042	1291	2.731
01 May 2017 09:45	Platåberget	S4	0.045	0.071	1440	16.753
01 May 2017 10:25	Gruvefjellet	S1	0.034	0.020	591	12.472
08 May 2017 10:15	Platåberget	S4	0.015	0.045	1563	5.535
09 May 2017 08:25	Gruvefjellet	S1	0.041	0.023	654	15.166
18 May 2017 12:05	Gruvefjellet	S1	0.076	0.016	677	28.044
18 May 2017 13:00	Platåberget	S4	0.011	0.039	1620	3.948
31 May 2017 10:40	Gruvefjellet	S1	0.061	0.020	693	22.546
01 June 2017 13:15	Platåberget	S4	0.033	0.062	1599	11.993
09 June 2017 12:20	Gruvefjellet	S1	0.018	0.015	485	6.642
09 June 2017 13:30	Platåberget	S4	0.014	0.039	1379	5.203
14 June 2017 14:35	Gruvefjellet	S1	0.005	0.010	401	1.845

Formatted Table

Appendix 1 continued...

<u>Date and Time (UTC)</u>	<u>Area</u>	<u>Scan Position</u>	<u>Reported Registration Error</u>	<u>Mean Relative Error (+- m)</u>	<u>Standard Deviation</u>	<u>Representative Volume (m³)</u>	<u>Volume Uncertainty</u>
15 December 2017 10:20	Gruvefjellet	S1	0.013	0.018	0.024	929	6.458
15 December 2017 11:00	Platåberget	S4	0.060	0.016	0.039	5	5.978
24 January 2018 11:20	Gruvefjellet	S1	0.000	0.049	0.031	938	18.081
24 January 2018 12:25	Platåberget	S4	0.032	0.069	0.062	419	25.277
31 January 2018 15:45	Gruvefjellet	S1	0.025	0.009	0.024	947	3.321
31 January 2018 17:00	Platåberget	S4	0.044	0.070	0.052	415	25.978
22 February 2018 10:30	Gruvefjellet	S1	0.017	0.020	0.025	1051	7.528
22 February 2018 11:30	Platåberget	S4	0.067	0.032	0.056	902	11.734
02 March 2018 11:55	Gruvefjellet	S1	0.004	0.003	0.037	1031	1.144
02 March 2018 12:30	Platåberget	S4	0.001	0.024	0.072	1130	8.819
23 March 2018 13:15	Gruvefjellet	S1	0.001	0.041	0.023	539	15.092
23 March 2018 14:00	Platåberget	S4	0.013	0.032	0.049	889	11.845
13 April 2018 11:00	Gruvefjellet	S1	0.012	0.024	0.021	593	8.745
13 April 2018 13:20	Platåberget	S4	0.010	0.000	0.044	1053	0.037

715

Response to RC1

We appreciate Dr. Christine Fey for reviewing our work and for her constructive comments to improve this manuscript. We first respond to issues and suggestions provided directly in Dr. Fey's interactive comment and subsequently respond to comments and corrections included in Dr. Fey's supplement to the interactive comment. Reviewer comments are displayed below in bold, author responses are in standard text.

Responses to the interactive comment:

It would be helpful to add the tiepoints used for registration of the scans to the figures.

We agree visual tiepoint – and tieobject – display in the figures would enhance this work. Unfortunately, we were unable to use the same tiepoints and tieobjects to register each of the scans. This is primarily due to changing snow cover rendering certain tiepoints and tieobjects unusable as snow free surfaces at different times throughout the two winter seasons. Thus, we drew on a suite of over 50 tieobjects to adequately register all scans included in this work, and we believe including all tiepoints and tieobjects we employed would overly clutter the figures. Additionally, some of the tiepoints were located on buildings not included in the figures' extent.

To assess the accuracy only one area is used. The registration error can vary significantly between different areas. Because of this, the mean relative error is not representative for the entire scan. Either the authors enhance the accuracy assessment to more stable areas, which might be difficult in the case that there are no other snow free areas. The other option is to explain in the text why only one area was used. However, the research question focuses on changes in the magnitude of meters and in this case the accuracy assessment is not so important for the process understanding of cornices.

We struggled with this issue ourselves. The concern again, to which Dr. Fey alludes in her review, is the dearth of adequately large snow free areas in the vicinity of the cornices with which to use for accuracy assessments. One option would have been to report accuracies from building walls near the foot of the slope. Accuracy near these buildings is generally very good – and almost always better than near the cornices, as the building roofs and walls served as good surfaces to use in the registration process. Since we agree with Dr. Fey that the accuracy assessment is not so important for the process understanding of cornices, we elected to display accuracy assessment from the only stable, snow free area near the cornices because:

- Accuracy assessments from this location are most representative of scan accuracies near the cornices, and
- Reported accuracies from this location are typically worse than in other locations in the scanning domain (i.e. the relative accuracies assessed from building walls would not be representative of the accuracy near the cornices).

As suggested by Dr. Fey, we have added language in lines 195 – 196 clarifying our decision to use just a single area for accuracy assessments.

More critical I evaluate the volume estimation of cornices. Here, i) the TLS data uncertainties play a more important role in quantification and ii) it's very challenging to calculate the distance and volume of cornices at scoured areas iii) data gaps are causing significant uncertainties in volume quantification. The problem of data gaps is mentioned and the other points could be enhanced in the discussion. Since the process interpretation in this paper is mainly based on the shape of cornices taken from 2D profiles and distance changes of the snowpack it might be better to study the volume calculation of cornices in detail and publish in another research work.

All three of the issues raised with our volume estimation of cornices are valid concerns. We have specifically addressed these concerns in lines 464 – 471 of the discussion rather than eliminating these analyses altogether from the paper. We agree another research work specifically studying cornice volume changes is needed, but based on the data we currently have gathered and Reviewer 2's positive responses we have chosen to leave these analyses in this work, acknowledging the uncertainty in our calculations.

Responses to the supplement to the comment:

Minor grammatical comments and suggestions have been corrected as suggested by the reviewer.

Revisions from the supplement affecting the content of the manuscript:

Line 156 – The values given from the RiScan MSA are not the 3D accuracy and does only reflect the distance between the point correspondences used at the ICP. I would neglect this value because it says nothing about the registration quality.

True, and we did not explain this adequately. Based on these recommendations, however, we will remove both this sentence and the MSA values from Appendix 1.

Line 158 – You mean the point cloud was thinned by blockthinning? The term “grid” is misleading with raster data.

We've updated the sentence with more specific and less misleading language. Thank you.

Line 207 – Headings in chapter 4 can be more meaningful which would help the reader to understand the structure of the paper faster.

We've updated the headings in chapter 4 to “Seasonal summaries of meteorological conditions and cornice dynamics”, which will hopefully help with readability and clarity.

Line 635 – A legend would be helpful for faster understanding of the figure.

We've added a legend for the symbology where appropriate and removed redundant language from the figure caption.

Lines 663 and 667 – Revise the color scheme of the profile to distinguish between the dates.
We've changed the color scheme of the profiles to hopefully enhance readability.

Lines 683 and 687 – Revise the color scheme of the distance change to allow a better interpretation of the magnitude of change.

We've adjusted the figure to make interpretation of the changes to the snow surface on the slope below the cornices easier. See the detailed response to RC2 regarding Figures 9 and 12.

Response to RC2

We are grateful for Dr. Jeffery Munroe's review and constructive comments to improve this manuscript. We respond to his comments below. Reviewer comments are displayed below in bold, author responses are in standard text.

Specific comments:

All of the figures are relevant and helpful to the reader, but I think a few of them should be changed. Specifically, Figures 5 and 7 are difficult to read because the colors used to represent the snow surface at the different times are too similar. Both of these figures are really important – they nicely present the data and allow clear visual distinctions to be made between scans, between slopes, and between the two winters. Improving their readability with more contrasting colors is a necessary step that will greatly help comprehension of the reader.

Thanks. We have struggled with the color schemes for this figure for a while. To address these concerns, we have attempted to render the diagrams with more contrasting colors than the greyscale we had originally selected. Although we acknowledge readability is still not perfect, we thought it was important to include all selected profiles in the seasonal profile progressions rather than eliminate some profiles in the interest of legibility.

I found Figures 9 and 12 difficult for a similar reason. While one might think that the bright red and dark blue colors representing the extremes of the change spectrum would be visible against the grayscale hilshade, the differences are actually really subtle. The figures are both important because they illustrate just how sensitive the TLS method is to even small changes in the snow surface. Unfortunately, the areas that changed are just really hard to see – even with the arrows drawing attention to specific regions in the images. I'm not sure what to recommend here. It is possible that a different color scheme would work better. Another possibility would be to keep a large figure representing the overview of the slope, and having a series of enlargements of small areas (keyed back to boxes in the overview figure) that showed the detected changes in a more obvious, zoomed-in way. Just as in my comment above, these figures are critical to presenting your data and supporting the following discussion – it would be great if they could be made even more compelling.

We agree the changes were too difficult to discern with the color scale as originally presented in the figure. The issue was attempting to display more subtle changes to the snow surface on the slopes below the cornices with a quantitative M3C2 distance scale suitable to represent the much greater changes to the cornices themselves (scale from -5 m to 5 m). To address this issue, we have changed the scale from -2 m to 2 m to better represent the changes to slope below the cornices. This hopefully helps differentiation between specific snow surface features, while also drawing attention to the dramatic changes on the cornices. We've added "greater than" or "less than" symbology to the scale labels to show that M3C2 distance changes on the cornices exceed the variability captured by this scale (also evident in the text and Figures 5 and 7).

Technical corrections:

Line 30 – “...projections of snow that form due to...”

Changed.

Line 40 – “...Cornice hazards.”

Fixed.

Line 76 – Would “designing” be a better word than “planning” here?

Yes – we altered the sentence to hopefully improve clarity.

Line 90 – I usually capitalize “U-shaped valley.”

We have changed the punctuation to reflect this.

Line 90 – “...oriented axis running...”

Changed, thank you.

Line 95 – Is there any information about the thickness of the continuous permafrost?

Yes, thank you for pointing out this omission. We’ve added the information from Humlum et al. (2003).

Line 101 – “consists of a 50-70 m, near-vertical bedrock cliff situated under the plateau margin and above...”

Thanks.

Line 115 – “The climate of Svalbard prohibits...”

Changed.

Line 139 – by “reliable” snow depth data do you mean the start of the seasonal snow accumulation? Or is this the date at which the snowpack exceeded a certain minimum thickness necessary for accurate measurement?

We meant the date at which the snowpack became deep enough to overcome to local ground surface roughness and become “smooth” enough to generate reliable measurements. Since this point is largely irrelevant for the larger purposes of this paper, we have removed the word “reliable” from the sentence to avoid confusion.

Line 154 – “...we used to georeferenced individual...”

Thanks.

Line 225 – I’m not sure what “(Size D2, R3)” means.

This refers to the destructive / relative size classifications used to characterize avalanche sizes.

We’ve left this description in for now, but it can be easily removed if the editor finds it confusing or detrimental to the manuscript’s clarity. We have also added a reference to the recording standards for avalanche sizes.

Line 370-372 – I would include reference to Figure 7 and Figure 5 here.
Good point, thanks. Figure references added.

Line 380 – “...to suggest that specific interactions...”
Changed.

Line 387 – “...cornices we investigated, and also failed completely both seasons.”
Changed, thanks.

Line 439 – “...also favorable for the development of more...”
Thanks.

Line 629 – “...was taken is indicated by POV in ...”
Fixed, thank you!