



1 **Precipitation stable isotopic signatures of tropical cyclones in Metropolitan**
2 **Manila, Philippines show significant negative isotopic excursions**

3
4 Dominik Jackisch¹, Bi Xuan Yeo², Adam D. Switzer^{1,2}, Shaoneng He¹, Danica Linda M.
5 Cantarero³, Fernando P. Siringan³ and Nathalie F. Goodkin^{1,2,4}

6
7 ¹ Earth Observatory of Singapore, Nanyang Technological University, Singapore
8 639798

9 ² Asian School of the Environment, Nanyang Technological University, Singapore
10 639798

11 ³ Marine Science Institute, University of the Philippines Diliman, Quezon City 1101,
12 Philippines

13 ⁴ American Museum of Natural History, New York 10024, USA

14
15 Correspondence to: Adam D. Switzer (aswitzer@ntu.edu.sg)

16
17 **Abstract**

18
19 Tropical cyclones have devastating impacts on the environment, economies, and societies,
20 and may intensify in the coming decades due to climate change. Stable water isotopes serve
21 as tracers of the hydrological cycle, as the fractionation process may leave distinct
22 precipitation isotopic signatures. Here we present a record of daily precipitation isotope
23 measurements from March 2014 to October 2015 for Metropolitan Manila, which is a first of
24 a kind dataset for the Philippines and Southeast Asia, and analyze if there is an isotopic
25 response to tropical cyclones. The most negative shift in $\delta^{18}\text{O}$ value (-13.84 ‰) leading to a
26 clear isotopic signal was caused by Typhoon Rammasun, which directly hit Metropolitan
27 Manila. The average $\delta^{18}\text{O}$ value of precipitation associated with tropical cyclones is -10.24 ‰,
28 whereas the mean isotopic value for rainfall associated with non-cyclone events is -5.29 ‰.
29 Further, the closer the storm track to the sampling site, the more negative the isotopic values,
30 indicating that in-situ isotope measurements can provide a direct linkage between isotopes
31 and typhoon activities in the Philippines.



32 **1. Introduction**

33

34 The Philippine archipelago, with its fast-growing population clustered along the coastline, is
35 one of the most vulnerable countries to climate change. It is especially prone to the
36 devastating effects of tropical cyclones (Cinco et al., 2014). Thus, it is considered a hotspot
37 region for hydrometeorological disasters (Cruz et al., 2013; Takagi and Esteban, 2016). There
38 is a clear need for developing a better understanding of tropical cyclone (TC) dynamics and
39 cyclone histories in the context of prediction that may allow government agencies to
40 implement proper mitigation and adaptation policies. Nine TCs per year made landfall on
41 average between 1951 to 2013 in the Philippines, while for that time period the average
42 number of TCs reaching Philippine waters is 19.4 per year (Cinco et al., 2016). Changing
43 climate and associated warming of the surface ocean, will likely increase the number and
44 intensity of tropical cyclones in the future (Emanuel, 2005; Webster and Holland, 2005;
45 Woodruff et al., 2013).

46

47 The Philippines were struck by several devastating TCs in recent years (Table 1). Typhoon
48 Haiyan (2013) which tracked over the Visayas has been the costliest TC to date (~ 2.06 billion
49 USD in 2013), with strong winds and intense storm surges inundating coastal areas resulting
50 in more than 6000 fatalities (Alojado and Padua, 2015; Lagmay et al., 2015; Soria et al., 2016).
51 Typhoon Rammasun, which made landfall in July 2014, is ranked number 3 with ~ 880 million
52 USD in 2014 (Alojado and Padua, 2015; NDRRMC, 2014). Eighty percent of the strongest
53 typhoons making landfall in the Philippines over the last three decades developed during
54 higher than average sea surface temperatures (SST), which supports growing evidence that
55 TC numbers are projected to rise in the future due to an increase in global temperatures
56 (Guan et al., 2018; Webster and Holland, 2005; Takagi and Esteban, 2016). For example, SST
57 was found to be anomalously high and reaching 29.6 °C during the formation of Typhoon
58 Haiyan (Takagi and Esteban, 2016). The average Philippines' ocean SST we have calculated for
59 the period 1945 to 2014 (basin between 6° – 18° N, 120° – 140° E) is ~ 28.5 °C based on
60 National Oceanic and Atmospheric Administration Extended Reconstructed Sea Surface
61 Temperature Dataset, Version 5 (NOAA ERSST v5) (Takagi and Esteban, 2016). By the end of
62 the 21st century, average typhoon intensity in the low-latitude northwestern Pacific is
63 predicted to increase by 14 % due to warming ocean temperatures (Mei et al., 2015).



64

65 A few studies have demonstrated the potential to investigate tropical cyclones using stable
66 water isotopes (e.g. Good et al., 2014; Lawrence et al., 2002; Munksgaard et al., 2015). Stable
67 water isotopes ($\delta^2\text{H}$ and $\delta^{18}\text{O}$) serve as dynamic tracers of hydrological processes and can
68 provide insights into the water and energy budgets of TCs (Good et al., 2014; Lawrence and
69 Gedzelman, 1996). For regions with general TC occurrence, significantly lower $\delta^2\text{H}$ and $\delta^{18}\text{O}$
70 are associated with TC rainfall due to strong fractionation processes, compared to other
71 tropical rain events (Lawrence, 1998; Lawrence and Gedzelman, 1996). Furthermore, $\delta^2\text{H}$ and
72 $\delta^{18}\text{O}$ have been used successfully to interpret TC history from paleoarchives, such as tree rings
73 and speleothems. For instance, tree-ring cellulose isotope proxies have recorded the recent
74 220 years of cyclones in the southeastern USA (Miller et al., 2006); similarly, high-resolution
75 isotopic analysis of tree-rings from the eastern US revealed the occurrence of hurricanes in
76 2004 (Li et al., 2011); a 23-year stalagmite record from Central America was used to
77 reconstruct past TC activity (Frappier et al., 2007), and isotope signals from a 800-year
78 stalagmite record were used to reconstruct past TC frequencies in northeastern Australia
79 (Nott et al., 2007). Interpretation of TC history from paleoarchives is based on the fact that
80 TCs leave distinct isotopic signatures on precipitation, possibly providing information on TC's
81 evolution and structure (Lawrence et al., 2002).

82

83 The depletion in isotopes is attributed to the high condensation levels, strong isotopic
84 exchanges between inflowing water vapour and falling raindrops in cyclonic rainfall bands,
85 resulting in a temporal decrease of isotopic values throughout a rain event (i.e. amount effect)
86 (Lawrence, 1998; Lawrence and Gedzelman, 1996). Isotopic depletion is further enhanced by
87 TC's thick, deep clouds, relatively large storm size and longevity (Lawrence, 1998).
88 Furthermore, while isotopic depletion increases inwards towards the eye wall of the storm
89 (Lawrence and Gedzelman, 1996), isotope ratios inside the inner eye wall region are relatively
90 enriched, likely due to an intensive isotopic moisture recharge with heavy isotopes from sea
91 spray (Fudeyasu et al., 2008; Gedzelman et al., 2003). These findings are based on work
92 conducted in the 1990s in Puerto Rico and on the southern and eastern coasts of the United
93 States. More recently, these previous findings have been confirmed by studying TCs which
94 occurred in a few other regions, such as in China or Australia (Chakraborty et al., 2016;
95 Fudeyasu et al., 2008; Good et al., 2014; Munksgaard et al., 2015; Xu et al., 2019).



96

97 The above-mentioned studies are geographically limited to a few locations globally, with no
 98 studies in Southeast Asia and the Philippines in particular. In this manuscript, we present the
 99 first such study for the Philippines, with daily isotope measurements of precipitation from
 100 Metropolitan Manila (the National Capital Region) spanning from March 2014 to October
 101 2015. During the study period, nine tropical cyclones passed by or made landfall within 500
 102 km of the sampling site (Fig. 1). The major objective of this research is to understand if there
 103 is an isotopic response of precipitation to TC activities in the Philippines, and if so – what
 104 signal do we measure and how is it represented spatially? Further, we aim to understand the
 105 isotopic variation with distance from the TC track. Our findings provide a baseline dataset for
 106 reconstruction of typhoon activities using stable isotopes and contribute to a better
 107 understanding of past and future TC activities in the Philippines.

108

109

110 **2. Materials and methods**

111

112 **2.1 Site description**

113

114 The Philippines is a Southeast Asian country comprising more than 7000 islands located in the
 115 Northwest Pacific between 4° 40' N and 21° 10' N, and 116° 40' E and 126° 34' E (Fig. 1). The
 116 country experiences an average annual rainfall of about 2000 mm, influenced by two
 117 monsoon seasons, the northeast monsoon from November to April and the southwest
 118 monsoon from May to October (Cinco et al., 2014). About 35 % of the annual rainfall is related
 119 to TC activity, while that number rises to about 50 % for Luzon and decreases to 4 % for the
 120 southern island of Mindanao (Cinco et al., 2016). Part of the rainfall amount in the Philippines
 121 is of orographic nature due to north-south oriented mountain ranges of more than 1000 m
 122 spanning the largest islands of Luzon and Mindanao (Villafuerte et al., 2014). The majority of
 123 the steadily growing population in the Philippines (101 million) live in densely populated, low-
 124 elevation areas close to the coastlines (Cinco et al., 2014, 2016; Philippine Statistics Authority,
 125 2017).

126

127



128 **2.2 Isotopic data**

129

130 In total, 186 daily precipitation samples were collected from 10 March 2014 to 26 October
 131 2015 using a PALMEX collector (Gröning et al., 2012) at the University of the Philippines
 132 Diliman (14.654° N, 121.068° E), located in Quezon City, which is part of Metropolitan Manila.
 133 Samples were collected daily at 10 am, and transferred without headspace to 30-ml HDPE
 134 bottles made by AZLON (www.azlonplastics.com) for storage prior to analysis. Samples were
 135 sent to the Earth Observatory of Singapore, Nanyang Technological University, Singapore and
 136 were analyzed for stable isotopes using a Picarro L1240-*i* laser spectroscopy instrument
 137 (www.picarro.com). We followed the procedures described by Van Geldern and Barth (2012)
 138 for post-run corrections and calibration. Three in-house water standards used for calibration
 139 include KONA (0.02 ‰ of $\delta^{18}\text{O}$; 0.25 ‰ of $\delta^2\text{H}$), TIBET (-19.11 ‰ of $\delta^{18}\text{O}$; -143.60 ‰ of $\delta^2\text{H}$),
 140 and ELGA (-4.25 ‰ of $\delta^{18}\text{O}$; -27.16 ‰ of $\delta^2\text{H}$). They are calibrated against the international
 141 reference water VSMOW2 and SLAP2. Long-term analysis of our QA/QC standards yields
 142 precision of 0.04 ‰ for $\delta^{18}\text{O}$ and 0.2 ‰ for $\delta^2\text{H}$.

143

144 **2.3 Cyclone track data**

145

146 The International Best Track Archive for Climate Stewardship (IBTrACS) dataset contains
 147 global TC best-track data, and is a joint effort of various regional meteorological institutions
 148 and centers that are part of the World Meteorological Organization (WMO). The data is
 149 publicly available (ftp://eclipse.ncdc.noaa.gov/pub/ibtracs/), and comprises information on
 150 storm eye/center with its coordinates, wind speed, and pressure, etc., with a temporal
 151 resolution of six hours (Knapp et al., 2010; Rios Gaona et al., 2018). Apart from visualization
 152 of cyclone paths, we used the dataset to calculate the spatial distance between the storm's
 153 eye coordinates and our sampling site.

154

155 **2.4 Satellite precipitation data**

156

157 We used the IMERG Version 5 Final daily product, a remotely-sensed precipitation dataset
 158 from satellites (https://disc.gsfc.nasa.gov/SSW/) to highlight cyclonic tracks and precipitation
 159 patterns of several TC's passing by Metropolitan Manila, and to identify which rainfall events



were not affected by cyclonic activity, and instead were associated with local or other regional convection activities. Such datasets are beneficial as they provide quasi-global grid-based rainfall estimates for land and the oceans (Pom  on et al., 2017). The Integrated Multi Satellite Retrievals for GPM (IMERG) from the Global Precipitation Measurement (GPM) programme with a fine 0.1-degree grid size (Huffman et al., 2017) has been available since March 2014, and provides precipitation data in different temporal resolutions, such as half-hourly or daily. Such satellite rainfall data has been previously utilized to show TC tracks and related rainfall intensities (Rios Gaona et al., 2018; Villarini et al., 2011).

2.5 Rainfall data

Daily rainfall data was obtained from the Philippine Atmospheric, Geophysical and Astronomical Services Administration (PAGASA), which maintains a rainfall monitoring station about 2.7 km away from our sampling site. The data is freely available for the period 2013 to 2017, and can be accessed on the Philippines Freedom of Information website (www.foi.gov.ph).

3. Results

3.1 Precipitation isotope evolution during TC events

The stable isotope composition during the 19 months study period spanning from 10 March 2014 to 26 October 2015 shows large seasonal isotopic variability in Metropolitan Manila. One hundred and eighty-six daily precipitation samples have been collected and analyzed in total (Fig. 2). $\delta^{18}\text{O}$ ranges from 4 ‰ to -13.84 ‰, and $\delta^2\text{H}$ from 16.84 ‰ to -99.1 ‰. The highest $\delta^{18}\text{O}$ value of 4 ‰ was observed on 9 April 2014 during the annual dry period, whereas the lowest $\delta^{18}\text{O}$ value of -13.84 ‰ was observed on 16 September 2014 in association with TC activity. The mean $\delta^{18}\text{O}$ of precipitation at the study site is -5.29 ‰ for non-TC rain systems, while TCs, as large regional convective systems, have the potential to cause a change in δ -values of up to almost 9 ‰ relative to the mean. The average $\delta^{18}\text{O}$ value of the nine TCs that tracked within <500 km from the sampling site is -10.24 ‰ (STDEV of 2.11), a factor of 2 larger than the mean from non-TC precipitation (average is -5.29 ‰, STDEV of 2.64).



192

193 Overall, precipitation isotopes associated with TCs mark the lower range of $\delta^{18}\text{O}$ values during
 194 the study period. Especially during the 2014 season, most of the very low precipitation
 195 isotope values occurred throughout passage of TCs. For instance, Rammasun led to the lowest
 196 δ -value (Fig. 2, point a) of the whole study period, while other TCs such as Fung-Wong (Fig.2,
 197 point c), Kalmaegi (Fig. 2, point b), or Hagupit (Fig. 2, point d) caused other negative
 198 excursions in isotopic values. The 2015 season is characterized by on average a slightly higher
 199 isotopic enrichment during the rainfall intensive summer months. Nonetheless, a similar
 200 noticeable isotope signal is visible with low $\delta^{18}\text{O}$ isotopes, clustered along the lower end of
 201 the sample range, for example, caused by Linfa (Fig. 2, point f) or Koppu (Fig. 2, point i).
 202 However, relatively negative isotope samples (Fig. 2) also originated from non-TC rainfall
 203 systems. Those events are discussed below.

204

205 Out of the nine TCs that occurred within a 500 km radius from the sampling site, Rammasun
 206 and Kalmaegi left clearly observable, distinct isotopic signatures during their approach and
 207 dissipation, which we will therefore present in more detail in the next paragraphs. Typhoon
 208 Hagupit (Fig. 2, point d) similarly lead to a clear isotopic evolution pattern during its time of
 209 occurrence in the Philippines and is shown in the supplementary (S1).

210

211 Typhoon Rammasun's rainfall intensities based on the IMERG precipitation data together
 212 with its track from IBTrACKS is shown in Fig. 3a. Typhoon Rammasun stands out in our study
 213 period as it moved straight towards the National Capital Region of the Philippines, resulting
 214 in a direct hit. Rammasun, locally named Glenda, made landfall in the Bicol region of southern
 215 Luzon on 15 July, with wind speeds of about 160 km/h. On 16 July, it passed south of
 216 Metropolitan Manila 50 km from our sampling site, with maximum winds of 130 km/h,
 217 gradually losing strength over land. As Rammasun approached on 15 July, the precipitation
 218 has shown relatively high $\delta^{18}\text{O}$ of -4 ‰ while rainfall was weak (Fig. 4a). On 16 July, $\delta^{18}\text{O}$
 219 shifted to -13.84 ‰, while the typhoon's track was the closest to our sampling site and rainfall
 220 amount was high. As Rammasun moved away, precipitation isotopes became more positive,
 221 and rainfall amount decreased. The characteristic isotopic evolution related to Rammasun's
 222 distance and rainfall intensities as a function of time can be seen in Fig. 5a, where the different
 223 radii indicate the distance to the sampling site, and the strong isotopic depletion observed on



224 16 July is also evident. As Rammasun with its storm center tracked towards the northwest
225 and away from Metropolitan Manila, our precipitation samples were relatively isotopically
226 enriched for the following two days.

227

228 Typhoon Kalmaegi, locally named Luis, was the first typhoon to make landfall in the
229 Philippines two months after Rammasun. Kalmaegi reached typhoon intensity on 13
230 September, making landfall the following day in northern Luzon, with maximum wind speeds
231 of about 120 km/h. Kalmaegi tracked relatively far away from the sampling site (about 350
232 km), but the accumulated rainfall it produced was centered south of the track, placing it
233 considerably closer to the National Capital Region (Fig. 3b). Despite the distance of the eye
234 from the sampling site, a characteristic isotopic pattern was visible, with the most negative
235 $\delta^{18}\text{O}$ value of -11.39 ‰ on 15 September, coincident with the highest rainfall (Fig. 4b). The
236 following day, $\delta^{18}\text{O}$ values returned to higher values with the increase in distance from the
237 eye. This is also seen in a spatial representation in Fig. 5b.

238

239

240 4. Discussion

241 4.1 Stable isotopes of precipitation – a possible proxy for TCs

242

243 As stable water isotopes fractionate during the physical process of evaporation and
244 condensation, they serve as effective tracers in the hydrological cycle (Dansgaard, 1964; He
245 et al., 2018; Risi et al., 2008; Tremoy et al., 2014). Here, we have demonstrated that stable
246 water isotopes can possibly be used to identify TC activity in the Southeast Asian region by
247 excursions in $\delta^{18}\text{O}$, providing evidence and supporting the hypothesis that TCs may leave a
248 clear isotopic signal in the Philippines. The strong isotopic depletion is due to high
249 condensation efficiencies in cyclonic convective rain bands, leading to extensive
250 fractionation. This is particularly pronounced in intense, large-scale TCs (Lawrence, 1998;
251 Lawrence and Gedzelman, 1996). In the previous section, we have presented our findings of
252 precipitation isotope ratios associated with typhoon activities affecting Metropolitan Manila
253 during the study period of March 2014 to October 2015. Based on our time series, we
254 therefore argue that for the Philippines, the lowest measured isotope value likely indicates



the occurrence of a TC, such as is the case for Typhoon Rammasun (Fig. 2). Similarly, other anomalously low $\delta^{18}\text{O}$ values at our site are caused by TC making landfall or passing by.

257

Individual TCs (Rammasun and Kalmaegi) were characterized by consistent isotopic excursions to very negative values in a range of up to -9‰ compared to the mean isotopic value of -5.29‰ (Fig. 4 and 5). A TC approaching the sampling site had relatively higher isotope values than at its later stages when it was closest to the site in Metropolitan Manila. When at its closest, strong rainfall together with increased fractionation depleted precipitation isotopes, leading to a distinct drop in isotope value. Such a strong negative isotopic shift in precipitation has been previously observed in other regions (Fudeyasu et al., 2008; Lawrence and Gedzelman, 1996; Munksgaard et al., 2015; Xu et al., 2019). As the TC moved away and rainfall intensities weakened, $\delta^{18}\text{O}$ in precipitation became again more positive, likely due to evaporative effects (Munksgaard et al., 2015; Xu et al., 2019).

268

As the strongest TC in terms of wind speeds, damage costs, and fatalities, Typhoon Rammasun reduced the isotope values most during our study period, to -13.84‰ . Similarly, Typhoon Kalmaegi lead to extensive damage and caused a significantly negative excursion in precipitation isotopes to -11.39‰ , suggesting that the lowest isotope values might indicate the occurrence of the strongest TC at that time at our site in the Philippines. We note that our isotopic measurements are similar to observations elsewhere. For example, the range of $\delta^{18}\text{O}$ values caused by Typhoon Shanshan affecting the subtropical Ishigaki island was -6 to -13‰ , (Fudeyasu et al., 2008); Tropical Cyclone Ita led to a range of -4.8 to -20.2‰ in northeastern Australia (Munksgaard et al., 2015); several TCs which made landfall in Texas resulted in isotope values from -3.9 to -14.3‰ (Lawrence and Gedzelman, 1996); or hurricanes that affected Puerto Rico and southern Texas were found to deplete $\delta^{18}\text{O}$ up to -18‰ (Lawrence, 1998). The lowest value resulting from Typhoon Phailin on the Andaman Islands was reported to be -5.5‰ , and Typhoon Lehar depleted the precipitation sample to -17.1‰ (Chakraborty et al., 2016). For TCs within a distance of up to 500 km from the sampling site at the University of the Philippines Diliman in Metropolitan Manila we measured an isotopic range of -7.7‰ (Typhoon Koppu) to -13.84‰ (Typhoon Rammasun). Despite the overall comparability to our measurements, differences exist. The lowest values observed in some studies are considerably more negative than at our site (Lawrence, 1998; Munksgaard



287 et al., 2015). However, we attribute these differences to a variety of features, such as the
288 specific climatic condition at each site, differences in temperature, humidity, and altitude or
289 latitude, which are likely contributing factors to the observed isotopic variation by altering
290 isotopic fractionation. Further, rainout history, location of typhoon tracks, topography,
291 respective strength of each TC, as well as its distance to the sampling site most likely have a
292 significant influence as well (Fudeyasu et al., 2008; Good et al., 2014; Munksgaard et al., 2015;
293 Xu et al., 2019).

294

295 In order to assess why other very low isotopic excursions occurred on various days (Fig. 2) we
296 used IMERG satellite precipitation data. IMERG data with its fine spatio-temporal resolution
297 allows the identification of convective rainfall areas and the passage of TCs and other rain
298 systems. Our analysis shows that precipitation events with anomalously low isotope signals
299 unassociated with TCs are largely related to local, strong convective rainfall events or large
300 scale and slow-moving rain areas passing over the National Capital Region. Therefore, the
301 degree of convection is responsible to produce the other observed low $\delta^{18}\text{O}$ value outliers
302 that are not related to cyclone rainfall, as strong convection and long stratiform rainfall leads
303 to intense fractionation (He et al., 2018; Risi et al., 2008; Tremoy et al., 2014). Contrarily, we
304 speculate that the more positive isotope values which cluster along the higher end of the
305 sample spectrum around 0 ‰, are associated with local, short convective rainfall events and
306 light intensity rain as confirmed with IMERG satellite precipitation data. Additionally, the
307 PAGASA rain gauge data indicates that rainfall amounts are very low during days with such
308 very enriched isotope samples, such as 0.3 mm/day for the highest recorded sample of 4 ‰
309 on 9 April 2014. Interestingly, TCs at our site were found to be related with low isotope values
310 together with high rainfall amounts (Fig. 2), while the majority of other low isotopic values
311 unassociated with TCs were characterized by on average lesser rainfall amounts. This possibly
312 indicates that TCs in the Philippines, besides using for instance modern-day satellite or radar
313 data, can be detected using these two parameters, i.e. strong isotopic depletion coupled with
314 high rainfall amounts.

315

316 The aforementioned local convective precipitation events have the potential to induce a
317 signal of very negative $\delta^{18}\text{O}$ values which are not related to TC activities. We therefore label
318 such a signal as a “false non-TC signal”, as it is induced by non-TC rainfall. This results in the



fact that TCs occurring during our study period do not entirely cluster along the lowest range of isotope values as seen in figure Fig. 2. Nevertheless, Typhoon Rammasun caused a clear drop in isotopes and stands out in the dataset. This might be the case because Rammasun's track and heavy rainfall comes in closest proximity (50 km) to the sampling site. Other TCs occurring within the 500 km radius did not lead to such a clear negative isotopic signature, likely because these typhoons did not pass the sampling site at all or heavy rainfall occurred elsewhere within the TC rainfall system (see S 2 for their tracks and accumulated rainfall areas). Some of these TCs have intense rainfall areas over other parts of the Philippines and are characterized by a variable track, likely influenced by land interactions. Land interaction reduces TC strength and can lead to rain out caused by orographic effects induced by the north-south oriented mountain ranges (Park et al., 2017; Xie and Zhang, 2012; Xu et al., 2019). Especially Typhoon Koppu rained out before making landfall and abruptly changed its track, instead of passing by the Metropolitan Manila. Similarly, Typhoon Mekkhala's intense rainfall occurred along the eastern coasts, before it started to dissipate. Evidently, due to these factors the isotope values associated with those TCs were not as negative as during Rammasun. Therefore, a TC which is relatively far away from the sampling site causes an isotope signal that is not as clear and as negative, thus averaging out between the other low values from rain systems unassociated with TC.

337

338 **4.2 Distance of TCs from Metropolitan Manila**

339

Our observations provide details on spatial distance from collection site towards TCs' centers, as our findings indicate that the distance from the storm center to the sampling site impacts isotopic value. The TCs' distance of up to 500 km to sampling site and the precipitation isotope value ($r=0.55$, $n=16$, $p\text{-value} < 0.0001$, 99% confidence level) are correlated. This relationship weakens with an increase in the distance from the sampling site: a distance of 500 to 1000 km yields an r of 0.2 ($n=19$, $p\text{-value}=0.019$), the distance of 1000 to 1500 km yields an r of 0.18 ($n=24$, $p\text{-value}=0.087$), while a 1500 to 2000 km distance results in an r of 0.1 ($n=21$, $p\text{-value}=0.65$). This suggests that a TC more than 500 km away from the sampling site has no influence on precipitation isotopes (Munksgaard et al., 2015). Thus, the closer the TC is to the sampling site, the more negative the isotope signal and the larger the δ -change. This relationship might provide information on storm structure and intensity, as the intensity



351 increases with proximity of the TC to the sampling location. We thus confirm that the isotope
 352 value at our location is a function of the closest approach of the storm's center to the
 353 sampling site (Lawrence and Gedzelman, 1996).

354

355 Figure 6 displays all the precipitation samples associated with TC presence and activities
 356 within a 2000 km radius from Metropolitan Manila, and further highlights the relationship
 357 between distance and isotopic depletion, additionally providing a spatial indication of TC's
 358 quadrants and their tracks relative to the location of the sampling site. Strongest depletion
 359 occurs within the 500 km radius. However, two relatively negative outliers are located within
 360 a 1000 to 1500 km radius in the northwest quadrant. These two samples were taken during
 361 the passage of tropical storm Kujira on 22nd and 23rd of June 2015 (Fig. 2), which was more
 362 than 1000 km away from Metropolitan Manila travelling east along the coast of Vietnam as
 363 seen with IBTrACKS data. We investigated these two samples with IMERG satellite
 364 precipitation data and identified them as a part of a mesoscale system, with strong convective
 365 cells delivering intense rainfall, leading to distinct isotopic depletion and inducing a "false
 366 non-TC signal" of very negative rainfall unassociated with TC activity.

367

368 **4.3 Cyclone track's rainfall intensity**

369

370 IMERG satellite precipitation data also identify that the highest rainfall intensities occur at the
 371 left side of the TC track for all the TC within the 500 km radius, except for Hagupit and
 372 Mekkhala, which are more complex cases (Fig. 3a, b, supplementary S 2). This is in contrast
 373 to the results from Villarini et al. (2011), stating that the largest rainfall accumulation appears
 374 on the right side of the hurricane tracks. They also noted that large rainfall amounts occur far
 375 away from the storm's track, which we can confirm and quantify with our observations. The
 376 largest rainfall totals vary in a range of 50 to 150 km away from the storm's center depending
 377 on the TC. For Kalmaegi the intense rainfall areas are up to 150 km away from the storm's
 378 center. These areas with the highest rainfall totals should most likely coincide with the most
 379 negative isotope value, indicating that the strongest depletion occurs in the outer cyclonic
 380 rain bands. This is consistent with previous findings (Gedzelman et al., 2003; Lawrence and
 381 Gedzelman, 1996; Munksgaard et al., 2015). However, Fudeyasu et al. (2008) observed the
 382 highest isotope values in the inner eye wall, i.e. in close proximity to the storm's center. We



could not investigate this further as no TC passed by our site in a distance of about 20 km, which is the size of a typical typhoon's eye (Weatherford and Gray, 1988).

4.4 Implications for paleoclimate studies

Isotope proxies from paleoarchives such as tree rings and speleothems have been utilized to reconstruct past cyclone activities (Frappier, 2013; Frappier et al., 2007; Miller et al., 2006; Nott et al., 2007). For instance, stalagmites yielded a record of weekly temporal resolution with negative isotopic excursions related to TC activity (Frappier et al., 2007). Such a high temporal resolution from stalagmites makes our in-situ measurements very comparable, highlighting the potential to use both in conjunction. Similarly, high-resolution tree ring isotope analysis identified the occurrence of Hurricane Ivan and Hurricane Frances in 2004, which both resulted in the lowest observed precipitation isotope values for that year (Li et al., 2011). However, the aforementioned paleoclimate studies suffer from uncertainty regarding parameters such as TC intensity and distance to the storm's center affecting the isotope signal. With our study, we provide further information on these parameters as we hypothesize that immediate proximity of a TC results in very low $\delta^{18}\text{O}$. Therefore, we might aid with a better interpretation of paleoarchives. Moreover, these paleoclimate studies are limited in number and only focus on a few regions affected by TCs, such as Central America and the Southeastern USA (Frappier et al., 2007; Miller et al., 2006). However, more studies investigating paleoarchives related to typhoon footprints covering different regions and countries would provide a better understanding of past TC activity, ultimately resulting in better and more accurate climate reconstructions. TC projections related to climate change could also be improved, which is especially relevant for decision makers dealing with TC related impacts and damages. Our in-situ isotope measurements provide baseline data input in an understudied tropical region, providing isotopic data of TC occurrence and quantifying the isotopic depletion associated with TC activity. Further, our 19-month dataset suggests that the lowest measured isotope value at the Philippines study site is associated with TC activity, resulting in the distinct negative isotopic shift in the time series (Fig. 2). As rain out history, topography, distance of track or rainfall unassociated with TCs can induce a weak or "false non-TC signal", it is important to choose stalagmites or trees as archives based on their location, ideally covering a spatial gradient thus capturing a TC in its full size.



5. Conclusions

As presented with our dataset, a strong, high-energy TC with a track directly approaching and hitting the sampling site leads to a clear isotopic signal in a time series in the Philippines. If the TC is further away, such as more than 500 km from the site, or heavy TC rainfall occurred elsewhere prior of making landfall, the signal is not as clear and might average out between other rainfall events. Other strong convective rainfall events unassociated with TCs may result in similarly low isotope values, and we label these as a weak or “false non-TC signal”. Therefore, our data suggests that distance of TC to the sampling site is a key factor in controlling the isotope signal and that such a spatial component needs to be considered when interpreting the isotope signal. However, a longer time series isotope record would help to better constrain controlling factors, such as the influence of topography on high-energy TCs. To what extent mountain ranges and low elevation coastal areas shape the TC induced isotope signal needs further investigation. Based on our findings we conclude that the location of sample collection needs to be chosen strategically. Ideally, several rainwater collection stations should be operated, covering a wide geographical range such as stretching from northern Luzon to its south. If such a spatial gradient was covered, a TC would likely be captured in its full size. Consequently, we aim to expand our time series spatially and temporally.

Previous studies conducted in other regions found that TCs can leave detectable isotopic signals of very negative $\delta^{18}\text{O}$ values in precipitation (Good et al., 2014; Munksgaard et al., 2015; Xu et al., 2019). Daily precipitation isotope samples confirm the hypothesis that TC activities using isotopes can also be identified in the tropical Philippines. A total of 186 daily precipitation samples spanning 10 March 2014 to 26 October 2015 from Metropolitan Manila were analyzed for their isotopic composition, resulting in seasonal isotopic variability and in TC related isotopic signatures. The mean isotopic value for the study period is -5.29 ‰ for rain events unassociated with TC, whereas the average TC induced isotope value is -10.24 ‰ for TCs occurring within 500 km. The lowest recorded value is -13.84 ‰, which is a δ -change of almost -9 ‰ compared to the mean, and it was sampled during the closest approach of Typhoon Rammasun to the National Capital Region of the Philippines. Similarly, individual TCs



such as the intense and costly Rammasun that struck the Philippines in July 2014 or Kalmaegi left characteristic isotopic signatures. During their approach, $\delta^{18}\text{O}$ values were relatively high but once they moved closer to the collection site the isotopes became more depleted alongside increasing rainfall amounts. Once they moved away their remnants lead again to higher values. The distance of TC center to sampling site plays a key role in determining the observed isotope signals. Correlation between isotopes and distance of up to 500 km was found, though this relationship significantly weakens with increasing distance. Information on storm structure and intensity can be derived from the interconnectedness of distance and isotopic depletion, due to the fact that strong rainfall leads to increased isotopic fractionation (He et al., 2018; Tremoy et al., 2014; Xu et al., 2019). The closer the TC is to the sampling location, the stronger the rainfall intensities and the more negative the $\delta^{18}\text{O}$ in precipitation. Additionally, we found that the degree of convection can induce a “false non-TC signal” of very low isotope values not associated with TC activity. Other factors which limit the strength and clarity of the isotope signal are distance of TC towards the sampling side, rain out history, TC track and topography. Our dataset is the first of such record in the Philippines and provides much needed data in scarcely sampled Southeast Asia. It can be used as a baseline in paleoclimate studies reconstructing past TC history, in conjunction with tree ring and speleothem datasets, as our data suggest that for Metropolitan Manila the lowest measured isotope value is caused by typhoon activity. A higher precipitation sampling frequency on sub-daily levels at several locations would yield more detailed constraints on TC parameters such as storm structure, which we aim to realize in the future.

Data availability

The underlying research data can be accessed via the supplementary document.

Author Contributions

Dominik Jackisch analyzed the data and wrote the manuscript. Bi Xuan Yeo contributed to data analysis and improved the manuscript. Adam D. Switzer conceived the idea, reviewed



479 and improved the manuscript. Shaoneng He provided advice, reviewed and improved the
480 manuscript. Danica Cantarero and Fernando P. Siringan collected the precipitation samples
481 and improved the manuscript. Nathalie F. Goodkin reviewed and improved the manuscript.

482

483

484 **Competing interests**

485

486 The authors declare that they have no conflict of interest.

487

488

489 **Acknowledgments**

490

491 This study is supported by the National Research Foundation Singapore and the Singapore
492 Ministry of Education under the Research Centres of Excellence Initiative. It is Earth
493 Observatory of Singapore contribution no. 188. This study is also the part of IAEA Coordinated
494 Research Project (CRP Code: F31004) on “Stable isotopes in precipitation and palaeoclimatic
495 archives in tropical areas to improve regional hydrological and climatic impact model” with
496 IAEA Research Agreement No. 17980.

497

498

499

500

501

502

503

504

505

506

507

508

509

510



511 **References**

512

- 513 Alojado, D. and Padua, D. M.: Costliest Typhoons of the Philippines (1947 - 2014), [online]
 514 Available from:
 515 [https://www.typhoon2000.ph/stormstats/WPF_CostliestTyphoonsPhilippines_2015E](https://www.typhoon2000.ph/stormstats/WPF_CostliestTyphoonsPhilippines_2015Ed.pdf)
 516 [d.pdf](https://www.typhoon2000.ph/stormstats/WPF_CostliestTyphoonsPhilippines_2015Ed.pdf) (Accessed 17 September 2019), 2015.
- 517 Chakraborty, S., Sinha, N., Chattopadhyay, R., Sengupta, S., Mohan, P. M. and Datye, A.:
 518 Atmospheric controls on the precipitation isotopes over the Andaman Islands, Bay of
 519 Bengal, *Sci. Rep.*, 6, 1–11, doi:10.1038/srep19555, 2016.
- 520 Cinco, T. A., de Guzman, R. G., Hilario, F. D. and Wilson, D. M.: Long-term trends and
 521 extremes in observed daily precipitation and near surface air temperature in the
 522 Philippines for the period 1951-2010, *Atmos. Res.*, 145–146, 12–26,
 523 doi:10.1016/j.atmosres.2014.03.025, 2014.
- 524 Cinco, T. A., de Guzman, R. G., Ortiz, A. M. D., Delfino, R. J. P., Lasco, R. D., Hilario, F. D.,
 525 Juanillo, E. L., Barba, R. and Ares, E. D.: Observed trends and impacts of tropical
 526 cyclones in the Philippines, *Int. J. Climatol.*, 36(14), 4638–4650, doi:10.1002/joc.4659,
 527 2016.
- 528 Cruz, F. T., Narisma, G. T., Villafuerte, M. Q., Cheng Chua, K. U. and Olaguera, L. M.: A
 529 climatological analysis of the southwest monsoon rainfall in the Philippines, *Atmos.*
 530 *Res.*, 122, 609–616, doi:10.1016/j.atmosres.2012.06.010, 2013.
- 531 Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16(4), 436–468,
 532 doi:10.3402/tellusa.v16i4.8993, 1964.
- 533 Emanuel, K.: Increasing destructiveness of tropical cyclones over the past 30 years, *Nature*,
 534 436(7051), 686–688, doi:10.1038/nature03906, 2005.
- 535 Frappier, A. B.: Masking of interannual climate proxy signals by residual tropical cyclone
 536 rainwater: Evidence and challenges for low-latitude speleothem paleoclimatology,
 537 *Geochemistry, Geophys. Geosystems*, 14(9), 3632–3647, doi:10.1002/ggge.20218,
 538 2013.
- 539 Frappier, A. B., Sahagian, D., Carpenter, S. J., González, L. A. and Frappier, B. R.: Stalagmite
 540 stable isotope record of recent tropic cyclone events, *Geology*, 35(2), 111–114,
 541 doi:10.1130/G23145A.1, 2007.
- 542 Fudeyasu, H., Ichiyangi, K., Sugimoto, A., Yoshimura, K., Ueta, A., Yamanaka, M. D. and



- 543 Ozawa, K.: Isotope ratios of precipitation and water vapor observed in typhoon
- 544 Shanshan, J. *Geophys. Res. Atmos.*, 113(12), 1–9, doi:10.1029/2007JD009313, 2008.
- 545 Gedzelman, S., Lawrence, J., Gamache, J., Black, M., Hindman, E., Black, R., Dunion, J.,
- 546 Willoughby, H. and Zhang, X.: Probing Hurricanes with Stable Isotopes of Rain and
- 547 Water Vapor, *Mon. Weather Rev.*, 131(6), 1112–1127, doi:10.1175/1520-
- 548 0493(2003)131<1112:PHWSIO>2.0.CO;2, 2003.
- 549 Van Geldern, R. and Barth, J. A. C.: Optimization of instrument setup and post-run
- 550 corrections for oxygen and hydrogen stable isotope measurements of water by
- 551 isotope ratio infrared spectroscopy (IRIS), *Limnol. Oceanogr. Methods*, 10, 1024–
- 552 1036, doi:10.4319/lom.2012.10.1024, 2012.
- 553 Good, S. P., Mallia, D. V., Lin, J. C. and Bowen, G. J.: Stable isotope analysis of precipitation
- 554 samples obtained via crowdsourcing reveals the spatiotemporal evolution of
- 555 superstorm sandy, *PLoS One*, 9(3), doi:10.1371/journal.pone.0091117, 2014.
- 556 Gröning, M., Lutz, H. O., Roller-Lutz, Z., Kralik, M., Gourcy, L. and Pölsenstein, L.: A simple
- 557 rain collector preventing water re-evaporation dedicated for $\delta^{18}\text{O}$ and $\delta^2\text{H}$ analysis
- 558 of cumulative precipitation samples, *J. Hydrol.*, 448–449, 195–200,
- 559 doi:10.1016/j.jhydrol.2012.04.041, 2012.
- 560 Guan, S., Li, S., Hou, Y., Hu, P., Liu, Z. and Feng, J.: Increasing threat of landfalling typhoons
- 561 in the western North Pacific between 1974 and 2013, *Int. J. Appl. Earth Obs. Geoinf.*,
- 562 68(7), 279–286, doi:10.1016/j.jag.2017.12.017, 2018.
- 563 He, S., Goodkin, N. F., Jackisch, D., Ong, M. R. and Samanta, D.: Continuous real-time
- 564 analysis of the isotopic composition of precipitation during tropical rain events:
- 565 Insights into tropical convection, *Hydrol. Process.*, 32(11), 1531–1545,
- 566 doi:10.1002/hyp.11520, 2018.
- 567 Huffman, G. J., Bolvin, D., Braithwaite, D., Hsu, K., Joyce, R., Kidd, C., Nelkin, E. J.,
- 568 Sorooshian, S., Tan, J. and Xie, P.: Algorithm Theoretical Basis Document (ATBD) of
- 569 Integrated Multi-satellite Retrievals for GPM (IMERG), version 4.6, Nasa [online]
- 570 Available from:
- 571 https://pmm.nasa.gov/sites/default/files/document_files/IMERG_ATBD_V4.6.pdf
- 572 (Accessed 11 September 2019), 2017.
- 573 Knapp, K. R., Levinson, D. H., Kruk, M. C., Howard, J. H. and Kossin, J. P.: The International
- 574 Best Track Archive for Climate Stewardship (IBTrACS), *Bull. Am. Meteorol. Soc.*, 91,



- 575 363–376, doi:10.1007/978-90-481-3109-9_26, 2010.
- 576 Lagmay, A. M. F., Agaton, R. P., Bahala, M. A. C., Briones, J. B. L. T., Cabacaba, K. M. C., Caro,
 577 C. V. C., Dasallas, L. L., Gonzalo, L. A. L., Ladiero, C. N., Lapidez, J. P., Mungcal, M. T.
 578 F., Puno, J. V. R., Ramos, M. M. A. C., Santiago, J., Suarez, J. K. and Tablazon, J. P.:
 579 Devastating storm surges of Typhoon Haiyan, *Int. J. Disaster Risk Reduct.*, 11, 1–12,
 580 doi:10.1016/j.ijdr.2014.10.006, 2015.
- 581 Lawrence, J. R.: Isotopic spikes from tropical cyclones in surface waters: Opportunities in
 582 hydrology and paleoclimatology, *Chem. Geol.*, 144(1–2), 153–160,
 583 doi:10.1016/S0009-2541(97)00090-9, 1998.
- 584 Lawrence, J. R. and Gedzelman, S. D.: Low stable isotope ratios of tropical cyclone rains,
 585 *Geophys. Res. Lett.*, 23(5), 527–530, 1996.
- 586 Lawrence, J. R., Gedzelman, S. D., Gamache, J. and Black, M.: Stable isotope ratios:
 587 Hurricane Olivia, *J. Atmos. Chem.*, 41(1), 67–82, doi:10.1023/A:1013808530364,
 588 2002.
- 589 Li, Z. H., Labbé, N., Driese, S. G. and Grissino-Mayer, H. D.: Micro-scale analysis of tree-ring
 590 $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ on α -cellulose spline reveals high-resolution intra-annual climate
 591 variability and tropical cyclone activity, *Chem. Geol.*, 284(1–2), 138–147,
 592 doi:10.1016/j.chemgeo.2011.02.015, 2011.
- 593 Mei, W., Xie, S. P., Primeau, F., McWilliams, J. C. and Pasquero, C.: Northwestern Pacific
 594 typhoon intensity controlled by changes in ocean temperatures, *Sci. Adv.*, 1(4), 1–8,
 595 doi:10.1126/sciadv.1500014, 2015.
- 596 Miller, D. L., Mora, C. I., Grissino-Mayer, H. D., Mock, C. J., Uhle, M. E. and Sharp, Z.: Tree-
 597 ring isotope records of tropical cyclone activity., *Proc. Natl. Acad. Sci. U. S. A.*,
 598 103(39), 14294–14297, doi:10.1073/pnas.0606549103, 2006.
- 599 Munksgaard, N. C., Zwart, C., Kurita, N., Bass, A., Nott, J. and Bird, M. I.: Stable Isotope
 600 Anatomy of Tropical Cyclone Ita, North-Eastern Australia, April 2014, *PLoS One*,
 601 10(3), 1–15, doi:10.1371/journal.pone.0119728, 2015.
- 602 NDRRMC: SitRep No. 38 re Effects of Typhoon “Pablo” (Bopha), [online] Available from:
 603 http://www.ndrrmc.gov.ph/attachments/article/2245/SitRep_No_38_Effects_of_Typ
 604 [hoon_PABLO_as_of_25DEC2012_0600H.pdf](http://www.ndrrmc.gov.ph/attachments/article/2245/SitRep_No_38_Effects_of_Typ) (Accessed 18 September 2019), 2012.
- 605 NDRRMC: Final Report re Effects of Typhoon “Glenda” (RAMMASUN), [online] Available
 606 from:



- 607 [http://ndrrmc.gov.ph/attachments/article/1293/Effects_of_Typhoon_Glenda_\(RAM](http://ndrrmc.gov.ph/attachments/article/1293/Effects_of_Typhoon_Glenda_(RAM)
 608 [MASUN\)_Final_Report_16SEP2014.pdf](http://ndrrmc.gov.ph/attachments/article/1293/Effects_of_Typhoon_Glenda_(RAM) (Accessed 17 September 2019), 2014.
- 609 NDRRMC: Final Report re Preparedness Measures and Effects of Typhoon “Lando” (I.N.
 610 Koppu), [online] Available from:
 611 [http://ndrrmc.gov.ph/attachments/article/2607/FINAL_REPORT_re_Preparedness_](http://ndrrmc.gov.ph/attachments/article/2607/FINAL_REPORT_re_Preparedness_Measures_and_Effects_of_Typhoon_LANDO_KOPPU_as_of_14_-_21OCT2015.pdf)
 612 [Measures_and_Effects_of_Typhoon_LANDO_KOPPU_as_of_14_-_21OCT2015.pdf](http://ndrrmc.gov.ph/attachments/article/2607/FINAL_REPORT_re_Preparedness_Measures_and_Effects_of_Typhoon_LANDO_KOPPU_as_of_14_-_21OCT2015.pdf)
 613 (Accessed 17 September 2019), 2015.
- 614 Nott, J., Haig, J., Neil, H. and Gillieson, D.: Greater frequency variability of landfalling tropical
 615 cyclones at centennial compared to seasonal and decadal scales, *Earth Planet. Sci.*
 616 *Lett.*, 255, 367–372, doi:10.1016/j.epsl.2006.12.023, 2007.
- 617 Park, M. S., Lee, M. I., Kim, D., Bell, M. M., Cha, D. H. and Elsberry, R. L.: Land-based
 618 convection effects on formation of tropical cyclone Mekkhala (2008), *Mon. Weather*
 619 *Rev.*, 145(4), 1315–1337, doi:10.1175/MWR-D-16-0167.1, 2017.
- 620 Philippine Statistics Authority: Philippine Population Surpassed the 100 Million Mark
 621 (Results from the 2015 Census of Population), [online] Available from:
 622 <http://www.psa.gov.ph/population-and-housing/node/120080> (Accessed 15
 623 September 2019), 2017.
- 624 Poméon, T., Jackisch, D. and Diekkrüger, B.: Evaluating the performance of remotely sensed
 625 and reanalysed precipitation data over West Africa using HBV light, *J. Hydrol.*, 547,
 626 doi:10.1016/j.jhydrol.2017.01.055, 2017.
- 627 Rios Gaona, M. F., Villarini, G., Zhang, W. and Vecchi, G. A.: The added value of IMERG in
 628 characterizing rainfall in tropical cyclones, *Atmos. Res.*, 209, 95–102,
 629 doi:10.1016/j.atmosres.2018.03.008, 2018.
- 630 Risi, C., Bony, S., Vimeux, F., Descroix, L., Ibrahim, B., Lebreton, E., Mamadou, I. and Sultan,
 631 B.: What controls the isotopic composition of the African monsoon precipitation?
 632 Insights from event-based precipitation collected during the 2006 AMMA field
 633 campaign, *Geophys. Res. Lett.*, 35(24), 1–6, doi:10.1029/2008GL035920, 2008.
- 634 Soria, J. L. A., Switzer, A. D., Villanoy, C. L., Fritz, H. M., Bilgera, P. H. T., Cabrera, O. C.,
 635 Siringan, F. P., Maria, Y. Y. S., Ramos, R. D. and Fernandez, I. Q.: Repeat storm surge
 636 disasters of typhoon haiyan and its 1897 predecessor in the Philippines, *Bull. Am.*
 637 *Meteorol. Soc.*, 97(1), 31–48, doi:10.1175/BAMS-D-14-00245.1, 2016.
- 638 Takagi, H. and Esteban, M.: Statistics of tropical cyclone landfalls in the Philippines: unusual



- 639 characteristics of 2013 Typhoon Haiyan, *Nat. Hazards*, 80(1), 211–222,
 640 doi:10.1007/s11069-015-1965-6, 2016.
- 641 Tremoy, G., Vimeux, F., Soumana, S., Souley, I. and Risi, C.: Clustering mesoscale convective
 642 systems with laser-based water vapor d18O monitoring in Niamey (Niger), *J.*
 643 *Geophys. Res. Atmos. Res.*, 5079–5103, doi:10.1002/2013JD020968.Received, 2014.
- 644 Villafuerte, M. Q., Matsumoto, J., Akasaka, I., Takahashi, H. G., Kubota, H. and Cinco, T. A.:
 645 Long-term trends and variability of rainfall extremes in the Philippines, *Atmos. Res.*,
 646 137, 1–13, doi:10.1016/j.atmosres.2013.09.021, 2014.
- 647 Villarini, G., Smith, J. A., Baeck, M. L., Marchok, T. and Vecchi, G. A.: Characterization of
 648 rainfall distribution and flooding associated with U.S. landfalling tropical cyclones:
 649 Analyses of Hurricanes Frances, Ivan, and Jeanne (2004), *J. Geophys. Res. Atmos.*,
 650 116(23), 1–19, doi:10.1029/2011JD016175, 2011.
- 651 Weatherford, C. L. and Gray, W. M.: Typhoon Structure as Revealed by Aircraft
 652 Reconnaissance. Part I: Data Analysis and Climatology, *Mon. Weather Rev.*, 116(5),
 653 1032–1043, doi:10.1175/1520-0493(1988)116<1032:TSARBA>2.0.CO;2, 1988.
- 654 Webster, P. J., Holland, G. J., Curry, J. A. and Chang, H.-R.: Changes in Tropical Cyclone
 655 Number, Duration, and Intensity in a Warming Environment, *Science* (80-.), 309(4),
 656 1844–1846, doi:10.1126/science.1116448, 2005.
- 657 Woodruff, J. D., Irish, J. L. and Camargo, S. J.: Coastal flooding by tropical cyclones and sea-
 658 level rise, *Nature*, 504(7478), 44–52, doi:10.1038/nature12855, 2013.
- 659 Xie, B. and Zhang, F.: Impacts of typhoon track and Island topography on the heavy rainfalls
 660 in Taiwan associated with Morakot (2009), *Mon. Weather Rev.*, 140(10), 3379–3394,
 661 doi:10.1175/MWR-D-11-00240.1, 2012.
- 662 Xu, T., Sun, X., Hong, H., Wang, X., Cui, M., Lei, G., Gao, L., Liu, J., Lone, M. A. and Jiang, X.:
 663 Stable isotope ratios of typhoon rains in Fuzhou , Southeast China , during 2013 –
 664 2017, *J. Hydrol.*, 570, 445–453, doi:10.1016/j.jhydrol.2019.01.017, 2019.
- 665
 666
 667
 668
 669
 670



Figures and Captions




Figure 1 Metropolitan Manila sampling site and TC tracks of 2014 and 2015 seasons. Three different sized circles indicate the distance to the sampling site with the outermost one being 500 km in radius. Cyclone tracks are color coded according to the typhoon classification from Regional Specialized Meteorological Center (RSMC) Tokyo. Cyclones in gray color refer to TC outside the 500 km radius.




Figure 2 Complete time series of 186 precipitation samples taken between 10 March 2014 to 26 October 2015. $\delta^{18}\text{O}$ data points associated with TC activity are colored in red. Other anomalously low $\delta^{18}\text{O}$ values were investigated using IMERG satellite precipitation data. a: Rammasun 16/07/14, -13.84 ‰, 83 mm. b: Kalmaegi 15/09/14, -11.39 ‰, 85 mm. c: Fung-Wong 20/09/14, -12.16 ‰, 175 mm. d: Hagupit 8-9/12/14, -9.88 ‰, -10.62 ‰, 40 mm. e: Mekkhala 19/1/15, -10.77 ‰, 22 mm. f: Linfa 07/07/15, -8.5 ‰, 63 mm. g: Twelve 23/07/15, -7.7 ‰, 68 mm. h: Mujigae 01/10/15, -7.5 ‰, 51 mm. i: Koppu 19-20/10/15, -8.7 ‰, -9.72 ‰, 38 mm, 26 mm. 1: storm passing by 19/06/14, -10.44 ‰, 6 mm. 2: large rain areas 27/08/14, -12.5 ‰, 21 mm 3: storm passing by 15/11/14 -10.58 ‰, 3 mm. 4: large rain areas, 22-23/06/15 -10.76 ‰, -11.52 ‰, 2 mm, 4 mm. 5: heavy rainfall 13/08/15, -9.96 ‰, 80 mm. 6: heavy rainfall 18/08/15, -9.26 ‰, 13 mm. 7: local convection 16/09/15, -8.28 ‰, 47 mm.




Figure 3 Accumulated precipitation from IMERG satellite data and TC tracks from IBTrACKS for a) Rammasun with precipitation accumulation for 14-17 July 2014, b) Kalmaegi with accumulated precipitation for 12-15 September 2014. Made with base layers from Natural Earth.




Figure 4 Isotopic signature from TCs during their passage to the Metropolitan Manila sampling site. $\delta^{18}\text{O}$ (blue color), distance from storm center to sampling location (green) and daily rainfall amount (red). a) Rammasun, b) Kalmaegi




Figure 5 Spatio-temporal evolution of $\delta^{18}\text{O}$ isotopes. Centered on Metropolitan Manila collection site, different radii provide information on distance between storm's center to Metropolitan Manila. $\delta^{18}\text{O}$ values are color coded. a) Rammasun, b) Kalmaegi




Figure 6 Spatio-temporal variation of isotopes related to TC activity within 2000 km, with different radii indicating the distance towards Metropolitan Manila. $\delta^{18}\text{O}$ values are color coded.



799 Tables

800

801 **Table 1 Costliest typhoons in the Philippines.** Two devastating typhoons, Rammasun and Koppu (ranking 3 and 7),
 802 occurred during our study period and made landfall. Damage in USD based on each time of TC occurrence (not adjusted to
 803 current inflation rates).

| Rank | Name (local name) | Category (Saffir Simpson scale) | Period of occurrence | Damage in USD | Fatalities | Part of our dataset |
|--|-------------------|---------------------------------|----------------------|--------------------|------------|---------------------|
| 1. | Haiyan (Yolanda) | Category 5 | 2-11 November 2013 | ~ 2.06 billion USD | ~ 6000 | No |
| 2. | Bopha (Pablo) | Category 5 | 2-10 December 2012 | ~ 977 million USD | 1067 | No |
| 3. | Rammasun (Glenda) | Category 5 | 12-17 July 2014 | ~ 880 million USD | 106 | Yes |
| 7. | Koppu (Lando) | Category 4 | 12-21 October 2015 | ~ 310 million USD | 62 | Yes |
| References: Alojado and Padua, 2015; Lagmay et al., 2015; NDRRMC, 2012, 2014, 2015; Soria et al., 2016 | | | | | | |

804