



Brief communication: The role of using precipitation or river discharge data when assessing global coastal compound flooding

Emanuele Bevacqua¹, Michalis I. Voudoukas², Theodore G. Shepherd¹, and Mathieu Vrac³

¹Department of Meteorology, University of Reading, Reading, United Kingdom

²European Commission, Joint Research Centre (JRC), Ispra, Italy.

³Laboratoire des Sciences du Climat et de l'Environnement, CNRS/IPSL, Gif-sur-Yvette, France.

Correspondence: Emanuele Bevacqua (e.bevacqua@reading.ac.uk)

Abstract. Interacting storm surges and high water-runoff can cause compound flooding (CF) in low-lying coasts and river estuaries. The large-scale CF hazard has been typically studied using proxies such as the concurrence of storm surge extremes either with precipitation or with river discharge extremes. Here the impact of the choice of such proxies is addressed employing state-of-the-art global datasets. Although being proxies of diverse physical mechanisms, we find that the two approaches show similar CF spatial patterns. However, deviations increase with the catchment size and our findings indicate that CF in long rivers (catchment $\gtrsim 5\text{-}10,000\text{ Km}^2$) is more accurately analysed using river discharge data. The precipitation-based assessment allows for considering local rainfall-driven CF, and CF in small rivers not resolved by large-scale datasets.

1 Introduction

Compound flooding (CF) happens in low-lying coastal areas due to the interaction of high precipitation runoff and high sea level. The combination of the two hazards can cause larger damages than those caused by the either of hazards in isolation, and recent events have occurred in, e.g., Mozambique (2019), Texas (US, 2017), the Shoalhaven Estuary (Australia, 2016), Ravenna (Italy, 2015), Cork (Ireland, 2009), and Lymington (United Kingdom, 1999) (Couasnon et al., 2019; Zscheischler et al., 2018; Kumbier et al., 2018; Bevacqua et al., 2017; Olbert et al., 2017; Hendry et al., 2019). Practitioners and the scientific community are becoming more aware of CF risk, and there were several recent studies addressing the phenomenon at local (Bevacqua et al., 2017; Kumbier et al., 2018; van den Hurk et al., 2015) or larger scales (Wahl et al., 2015; Bevacqua et al., 2019a; Couasnon et al., 2019; Ward et al., 2018; Ganguli and Merz, 2019a, b). Recent advances in large-scale sea level and river discharge modelling allowed the generation of sub-daily time-series of water levels along the global coastline (Voudoukas et al., 2018; Muis et al., 2016), thus enabling continental CF assessments (Ward et al., 2018; Bevacqua et al., 2019a; Couasnon et al., 2019).

CF can be the result of different mechanisms depending on the local topography and meteorology. According to Wahl et al. (2015), CF is possible to occur when: (1) the joint occurrence of high river discharge and storm surge in estuarine regions may elevate water levels to a point where flooding is initiated or its impacts exacerbated; (2) a destructive storm surge, which already caused widespread flooding, is followed by rainfall, as the latter can drive additional flooding, even if it is not an extreme event on its own; and (3) a moderate storm surge occurs which does not directly cause flooding, but is high enough to fully block



or slow down gravity-fed storm water drainage, and as a result precipitation causes flooding. In addition, CF may occur if (4)
25 precipitation falls on wet soil that is saturated by a preceding storm surge (Bevacqua et al., 2019a).

Quantifying the actual CF, i.e. the water level resulting from the combination of pluvial/fluviial flooding and high sea level,
is challenging. For example, quantifying and interpreting CF in the vicinity of rivers (mechanism 1) requires water level
measurements in the river mouth, which are rare, probably because most gauges are installed to monitor either riverine or
marine processes (Bevacqua et al., 2017; Paprotny et al., 2018). Model-based data are also limited because only recently have
30 modellers started considering CF. Statistical and hydrodynamic modelling approaches integrating fluvial and sea-level flooding
have been developed and applied recently at the local scale (Bevacqua et al., 2017; van den Hurk et al., 2015; Kumbier et al.,
2018); however, at large scale, these approaches are only now being developed. Similarly, explicit study of the actual CF water
level due to pluvial flooding and storm surges (mechanisms 2-4) has not received much attention so far, to our knowledge,
which may also be due to the scarcity of data. Thus, to gain information on the CF hazard at the regional, continental, or global
35 scales, scientists usually employ proxies of flood hazard, e.g. the probability of potential CF (Bevacqua et al., 2019a; Ward
et al., 2018; Wahl et al., 2015; Couasnon et al., 2019; Paprotny et al., 2018). Quantifying potential CF, under the present or
future climate, is useful as it allows identifying potential hotspots of CF hazard. Then, more detailed assessments of the local
CF risk can be carried out at such hotspots, using computationally intensive methodologies that integrate all the hydrological
and meteorological sources of flooding and their physical interaction (Wahl et al., 2015).

40 The large-scale assessment of potential CF includes the analysis of the probability (or return periods) of concurring extreme
values of CF drivers. Two main approaches exist, focussing - respectively - on the analysis of the probability of concurring
high values of sea level and river discharge, or of sea level and precipitation. Through considering river discharge, the first
approach takes into account CF in estuaries and deltas, i.e. serving as a proxy of CF mechanism 1. The second approach,
based on the analysis of accumulated precipitation around the coast when the high sea levels occur, can represent CF due to
45 local precipitation extremes, i.e. related to mechanisms 2, 3, and 4. Given that precipitation is among the main drivers of river
discharge, the two proxies are correlated to a certain extent and the use of precipitation can thereby allow quantifying CF
potential also in certain river estuaries. However, the correlation between the two proxies can be sometimes poor, especially
in locations where river discharge is strongly influenced by other factors such as snowmelt, evaporation, and accumulated
precipitation over previous weeks or months (Blöschl et al., 2019).

50 Given the scarcity and heterogeneous distribution of in situ data (Ward et al., 2018; Couasnon et al., 2019; Wu et al., 2018),
scientists have started to employ model data - of river, storm surge, and precipitation - to assess the potential CF hazard (Ward
et al., 2018; Bevacqua et al., 2019a; Wu et al., 2018; Couasnon et al., 2019; Wu et al., 2018; Paprotny et al., 2018; Bevacqua
et al., 2019b). Against the foregoing background, the present study aims to assess whether a precipitation based CF assessment
can be used as a surrogate for potential CF in estuaries. To that end we use coherent global model datasets of storms surges
55 (including wave effects) (Vousdoukas et al., 2018), precipitation (Beck et al., 2017b), and river discharge (Couasnon et al.,
2019; Eilander, 2019) and conduct a first global comparison of the results obtained through the two approaches, keeping all
the other methodological aspects identical.



2 Data

We analyse the period 1979-2015. We consider river discharge daily maxima from a publicly available global dataset (Eilander, 2019; Couasnon et al., 2019). The dataset was the result of hydrological model simulations forced with temperature and potential evaporation derived from ERA-Interim, and with precipitation from the MSWEPv1.2 dataset (Couasnon et al., 2019). The latter is obtained by merging gauge, satellite, and reanalysis data (including ERA-Interim); more information can be found in Beck et al. (Beck et al., 2017b).

Precipitation is taken from the same MSWEPv1.2 dataset used to simulated river discharges and consists of daily data over a 0.25° grid. On each day we consider accumulated precipitation amounts within a 3-day centered window. This enables us to account for precipitation occurring just before and after midnight of the storm surge day (Bevacqua et al., 2019a; Martius et al., 2016), and to consider different mechanisms causing CF (Wahl et al., 2015; Bevacqua et al., 2019a).

Storm surges and waves were simulated with the hydrodynamic model DFLOW FM (Vousdoukas et al., 2017, 2018) and the wave model Wavewatch III (Mentaschi et al., 2017; Vousdoukas et al., 2017, 2018). The wave model was forced by 6-hourly wind, while DFLOW-FM was also forced by sea level pressure fields, both available from the ERA-Interim reanalysis (Dee et al., 2011). The effects of tropical cyclones (TCs) were considered in the reanalysis through storm surge simulations forced by downscaled atmospheric fields from all recorded TCs and by considering satellite-observed TC wave extremes (Vousdoukas et al., 2018). Astronomical tides are not considered in this analysis in order to focus on the meteorological component of the sea level, and excluding the stochastic coupling with tide induced water level variations. We consider daily water level maxima from the combined result of storm surges and wave setup (hereinafter mentioned as storm surges) according to Vousdoukas et al. (2018).

We analyse CF only around river mouth locations whose nearest precipitation and storm surge grid points lie within a distance of 75 km (Couasnon et al., 2019).

2.1 Methods

We assess the potential CF hazard via bivariate return periods of concurring extreme events (Vandenberghe et al., 2011) of the variables X and Y , i.e. storm surge and precipitation (CF_{prec}) or storm surge and river discharge (CF_{river}). Extremes of the individual variables (x_{ext} and y_{ext}) are defined as the associated α -year return levels. We use different α in the following, though we present the main results for $\alpha=5$; images for $\alpha=2$ are shown in the supplementary material. Return levels are obtained through fitting a generalized extreme value distribution to the annual maxima of the individual variables (Coles et al., 2001). Annual maxima are defined based on adjacent windows centred on the month where the climatological river discharge average is the highest, rather than from January-December windows (such a window definition reduces the chance of selecting two consecutive annual maxima belonging to the same river discharge extreme event, and therefore leads to a more robust definition of the return levels).

CF bivariate return periods are computed following the methodology presented by Bevacqua et al. (2019a). The CF return period computation is based on the bivariate distribution of the variables of interest (X, Y), which is estimated semi-empirically



to allow for robust estimation. For a given location, we select pairs whose individual values are simultaneously larger than the individual 95th percentiles (x_{sel} and y_{sel}), and we model these pairs via a copula-based distribution. If the defined thresholds result in a small group of selected pairs, we lower the 95th percentile selection threshold to guarantee having a sufficient amount of selected pairs, i.e. at least 20. Clusters of pairs separated by less than 3 days were considered as part of the same event represented by the maximum X and Y values observed in the cluster. The return period is defined as:

$$T(x_{ext}, y_{ext}) = \frac{\mu}{P((x > x_{ext} \text{ and } y > y_{ext}) | (x > x_{sel} \text{ and } y > y_{sel}))} = \frac{\mu}{1 - u_{X_{ext}} - u_{Y_{ext}} + C_{XY}(u_{X_{ext}}, u_{Y_{ext}})} \quad (1)$$

where μ is the average time elapsing between the selected pairs, $u_{X_{ext}} = F_X(x_{ext})$, F_X is the marginal cumulative distribution of the excesses over the selection threshold (accordingly for Y), and C_{XY} is the copula modelling the dependence between the selected pairs. We model the marginal distributions of X and Y beyond the selection thresholds by a Generalised Pareto Distribution (GPD). We fit copulas from the families Gaussian, t, Clayton, Gumbel, Frank, Joe, BB1, BB6, BB7, BB8 to (u_X, u_Y) (obtained via empirical marginal cumulative distribution function (Vandenberghe et al., 2011)); then we select the best ranked family according to the Akaike information criterion. We fit copulas and marginal distributions via a maximum likelihood estimator (using the *VineCopula* (Schepsmeier et al., 2016) and *ismev* (Heffernan et al., 2016) R-packages). We test the goodness of fit of copulas and marginals via the Cramer-von-Mises criterion (via the *eva* (Bader and Yan, 2016) and *VineCopula* (Schepsmeier et al., 2016) R-packages respectively).

When referring to the assessment of whether the CF return periods based on river discharge (T_{river}) are statistically different from those based on precipitation (T_{prec}) or not, we use the concept of statistical compatibility, recently introduced by Amrhein et al. (Amrhein et al., 2019). We compute the centred 95% (2.5-97.5%) confidence interval of T_{prec} on the basis of 600 resampled bivariate time series of precipitation and storm surge (each of them built randomly combining observed 1 calendar year bivariate time series (Bevacqua et al., 2019b)). T_{river} is regarded as being statistically compatible with T_{prec} if T_{river} lies within the 95% confidence interval of T_{prec} , and incompatible otherwise.

We qualitatively investigate how the two assessments compare for different classes of catchment size. To do so, we rank the rivers based on their catchment size and divide them into groups having the same sample size; for each group we compute different statistics to compare the two assessments: Spearman correlation of T_{river} and T_{prec} , ratio T_{river}/T_{prec} , and percentage of locations with T_{river} compatible to T_{prec} . This binning procedure provides equally robust statistics for each bin and shows similar results for small variations in the bin size.

2.1.1 Results and discussion

The spatial patterns of the potential CF return periods based on either precipitation (T_{prec}), or river discharge (T_{river}) are very similar (Fig. 1; Fig. A1 is identical but shows results based on extremes defined considering 2-year return levels). The results for clusters of locations with the 5% lowest CF return periods are also similar in the two assessments (Fig. A4 and A5). These hotspot regions are mainly found along the US and central American coasts, Central Chile, Madagascar, the southern North Atlantic coasts, and southern Japan.




Figure 1. Present-day (1980-2014) potential compound flooding probability based on precipitation and on river discharge. Return periods of CF defined as co-occurring extremes (5-year return levels) of the CF drivers. Return period of co-occurring (a) storm surge (including waves) and precipitation (accumulated 3-day centred) extremes; (b) storm surge (including waves) and river discharge extremes. Major rivers are shown in light blue.

While the spatial patterns of the CF return periods obtained from the two approaches are very similar, their relative differences can be substantial, especially at certain locations (Fig. 2 and A2). Given that the return period computation procedure involves several uncertainty factors (e.g. bivariate model fitting and definition of the return levels), we test the hypothesis that the return period based on the river discharge is statistically compatible (at 95% confidence level) with that based on



precipitation. When defining extremes based on the 5-year return levels, the river-based return period is compatible with the precipitation-based value in about 82% of the locations (Fig. 2c; 76% for 2-year return levels: Fig. A2c). The spatial distribution of locations where T_{river} is not statistically compatible with T_{prec} does not seem to follow a clear spatial pattern, though it appears that T_{river} is lower than T_{prec} in northern Europe and in the tropics. The latter are areas where CF is unlikely. T_{river} is higher than T_{prec} along the Gulf of Mexico (Fig. 2c and Fig. A2c). Compatible T_{river} and T_{prec} are found but with large discrepancies in the tropics and above 60deg North (Fig. 2b), consistently with the high uncertainty of these large CF return periods that do not allow for detecting potential differences between T_{river} and T_{prec} .

The spatial association of the return periods map obtained from the two assessments is shown in Fig. 3a and A3a. The Spearman correlation between T_{river} and T_{prec} is ~ 0.7 , and increases as the return level threshold employed to define the extremes decreases; e.g., 0.66 and 0.73 for the 5- and 2-year return level, respectively (Fig. 3a and A3a). This trend in the correlation is consistent with the higher uncertainties characterizing larger return periods.

Although some of the differences between the two assessments may be driven by uncertainties in the return period estimation, several physical processes, e.g., topography-dependent, may be involved. The Spearman correlation between the two assessments decreases (p-value < 0.0024) as the catchment size increases (Fig. 3c). The latter suggests that - for larger catchments - different processes may cause either positive or negative deviations between T_{river} and T_{prec} .

Water levels in the mouth of small rivers are expected to be largely influenced by the precipitation around the coast (Hendry et al., 2019; van den Hurk et al., 2015; Bevacqua et al., 2017), while around large rivers inland hydrological processes are usually dominant. Therefore, we qualitatively investigate whether the actual difference between the two assessments (quantified as T_{river}/T_{prec}) depends on the size of the catchment (Hendry et al., 2019). For example, the T_{river}/T_{prec} ratio, defined using the 5-year return level, tends to increase with the catchment size (Fig. 3b; see the deviation of the scatter plot of T_{river} vs T_{prec} from the identity line emerging with the increasing dimension of the catchment). For CF return periods defined on 2-year return level extremes the median T_{river}/T_{prec} is near unity for most of the catchments, even though T_{river}/T_{prec} increases slightly for larger catchments (blue line in Fig. 3d). For higher return levels, the median T_{river}/T_{prec} increases with the size of the catchment (Fig. 3d), indicating that the CF assessments based on precipitation and river discharge differ largely for large rivers. For all return levels, the median T_{river}/T_{prec} is near unity for rivers whose catchment size is up to about 5-10,000 Km² (Fig. 3d), i.e. about 75% of the rivers presently analysed (Fig. 3f). Binning the rivers per catchment size we find that the variance of the T_{river}/T_{prec} ratio within each bin tends to be smaller for small than for large rivers (Fig. 3d, shown by the dashed line for the 2-year return level). However, it is important to highlight that there is substantial variance (Fig. 3d for the 2-year return level extremes), and that for a river of any catchment size, the associated T_{river}/T_{prec} can be either small or large. Overall, the similarity of T_{river} and T_{prec} for small rivers is highlighted by the fact that for small rivers it is more likely that T_{river} is statistically compatible with T_{prec} (Fig. 3e). For example, Fig. 3e (4-year return level curve) shows that T_{river} is statistically compatible with T_{prec} for $\sim 82\%$ of the small rivers (catchment size $< 5,000$ Km²) and for $\sim 75\%$ of large rivers (catchment size $> 50,000$ Km²). The decrease in the compatibility of T_{river} and T_{prec} with the catchment size (Fig. 3e) is statistically significant for return levels smaller than the 6-year return level (p-value < 0.022). However, for high return levels this decrease is not prominent, most likely due to the large uncertainty associated with longer return periods. Fig. 3e indicates




Figure 2. Difference between the present-day (1980-2014) potential compound flooding probability based on precipitation and on river discharge. Ratio (T_{river}/T_{prec}) of CF return period of concurring river discharge and storm surge extremes (T_{river}), to CF return period of concurring precipitation (accumulated 3-day centred) and storm surge extremes (T_{prec}). Panel (a) shows the ratio where T_{river} is statistically incompatible (at 95% confidence level, see methods) with T_{prec} . Panel (b) shows the ratio where T_{river} is statistically compatible with T_{prec} , while panel (c) shows the coastline fraction where this happens (binned every 5° of latitude). In (c), the dashed grey line shows the 95% level.




Figure 3. Comparison of the potential compound flooding probability based on precipitation and on river discharge. (a) Scatterplot (based on Fig. A1) of CF return periods based on river discharge (T_{river}) and on precipitation (T_{prec}), for extremes defined based on 5-year return levels. Black contours represent the isolines of the kernel density containing from 10% to 90% of the (T_{river}, T_{prec}) pairs. The identity line is shown in red. (b) Empirical probability density function of the variables (T_{river}, T_{prec}) conditioned on the catchment size; the thickness of the isolines increases with the catchment size, such that each bin considers 1/5 of the total number of analysed rivers (3178). Each contour line (isolines of the kernel density) contains about 50% of the (T_{river}, T_{prec}) pairs. (c-e) Statistics comparing T_{river} and T_{prec} as a function of the catchment size and of the return levels used to define the CF univariate extremes (see legend in panel (a)). (c) Spearman correlation between the maps of T_{river} and T_{prec} . (d) T_{river}/T_{prec} ratio. The dashed line is the centered 68% (16th-84th percentiles) confidence interval of the ratio based on the 2-year return levels. A non linear Y-axis for values below 1 is plotted such that the specular cases, e.g., ratio $r=2$ ($T_{river} = 2 \cdot T_{prec}$) and $r=1/2$ ($T_{prec} = 2 \cdot T_{river}$) (see Fig. 3b) appear symmetric with respect to $r=1$ ($T_{river} = T_{prec}$). (To obtain the plot, the ratio is defined as $r = T_{river}/T_{prec}$; then if $r < 1$, r is transformed to $r = -T_{prec}/T_{river} + 2$. Now, e.g., $r=2$ and $r=0$ represent specular cases, therefore, where the y-axis is $r=0,1,2,\dots$, we can write $r=1/2,1/3,1/4,\dots$). See Fig. A3b for the plot with standard axis. (e) Percentage of locations where T_{river} is statistically compatible with T_{prec} at the 95% confidence level. (f) Probability density function of the catchment size of the analysed rivers; 50th, 75th, and 95th percentiles of the distribution are shown in red. In (c-e), tick lines are obtained through regressing the investigated statistical values to the natural logarithm of the mean size of the catchment bins. All lines are regressed using linear regression, except from the median of the ratio in (d) where a spline function is used. The slopes of the linear regressions are all significant in (c) (p -value < 0.0024), and significant up to the 6-year return level in (e) (p -value < 0.022).



the discrepancy for large catchments in T_{river}/T_{prec} being greater for low return levels, whereas Fig. 3d indicates the opposite; this is also likely caused by the larger uncertainty of return periods of higher return levels, that does not allow for detecting potential differences between large values of T_{river} and T_{prec} .

165 The differences between T_{river} and T_{prec} are not only controlled by the catchment size, but can be the result of several other factors. During the hydrological modelling, input data artefacts, or model inaccuracies, among others, introduce uncertainty which may contribute to the observed differences. Another important contribution should arise from the uncertainties in the bivariate return period estimation (Bevacqua et al., 2019a; Wahl et al., 2015), which can contribute to both positive and negative deviations between T_{river} and T_{prec} . Moreover, the catchment response time to precipitation depends also on rock and soils
170 catchment permeability (Hendry et al., 2019). Finally, river discharge is influenced not only by local coastal precipitation, but also by the weather over the previous weeks or months over the catchment including evaporation and potentially snowmelt (Couasnon et al., 2019; Bevacqua et al., 2017).

Clearly, the diversity of the physical processes leading to CF_{prec} and CF_{river} is very relevant and can cause both positive and negative differences between T_{river} and T_{prec} (Blöschl et al., 2019). For example, for any given catchment, a slow
175 catchment response time may either increase or decrease the T_{river}/T_{prec} ratio. In locations where cyclones cause frequent concurring storm surge and widespread coastal precipitation, it is not guaranteed that CF_{river} probability will be also high. For example, if the rainfall in the catchment upstream needs time to reach the coast, long enough for the storm surge to recede, then CF_{river} will be unlikely and in this case the T_{river}/T_{prec} ratio will be high (Klerk et al., 2015; Ward et al., 2018). In contrast, where the CF_{prec} is unlikely because different weather systems cause precipitation and storm surge extremes, a relatively slow
180 catchment response time may sometimes allow for high river discharge and storm surge to concur if, e.g., a first cyclone causes precipitation driving high river discharge reaching the coast when a second cyclone drives a storm surge (contributing to low T_{river}/T_{prec}) (Bevacqua et al., 2019a).

In addition the presence of a pronounced annual cycle in river discharge can modulate the river discharge driven CF hazard, and thus the T_{river}/T_{prec} ratio. In regions where CF_{prec} is likely, CF_{river} may be unlikely if the storm surge season does not
185 coincide with the season of the high river discharge (Ward et al., 2018) (high T_{river}/T_{prec}). In contrast, where the CF_{prec} is unlikely, CF_{river} may be more likely if the storm surge season coincides with the high river discharge season (low T_{river}/T_{prec}). In addition, precipitation extremes are typically short in duration, while river discharge extremes are less dynamic. Although events exceeding the α -year return level threshold occur both for precipitation and river discharge on average every α years, the number of days with high river discharge can be larger than the number of days with high precipitation amount. From a
190 statistical point of view, the above mechanism alone would make CF_{river} more likely than CF_{prec} (low T_{river}/T_{prec}). (This effect is even more pronounced in catchments with a slow response time and especially in areas where different weather systems cause precipitation and storm surge extremes (Bevacqua et al., 2019b).) The above effect is weakened as the return level threshold used to define extremes increases; due to the shorter duration of river discharge extremes, the above considerations justify T_{river}/T_{prec} increasing with the return level choice (Fig. 3d).

195 While the relevance of these mechanisms may depend on the local climate, they are expected to be less relevant in very small rivers where the catchment response time is small, and thus autocorrelation in the river time series is smaller. Consistently, we



find that T_{river} and T_{prec} tend to match more in smaller catchments. However, a higher agreement for small rivers might also arise from the relatively coarse spatial resolution of the data which would attenuate differences between precipitation and river discharges in small rivers. Overall, we find that independently of the catchment size, T_{river} tends to be higher than T_{prec} on average (Fig. 3d), suggesting that the mechanisms causing $T_{river} > T_{prec}$ may be more likely or relevant. Apparently this is a general remark and not a universal law since there are also several locations where T_{prec} exceeds T_{river} .

The presented results are based on state-of-the-art model data which have been validated and discussed in previous papers (Vousdoukas et al., 2018; Beck et al., 2017a; Couasnon et al., 2019; Bevacqua et al., 2019a; Muis et al., 2016). However, the above mentioned validation efforts did not include areas where field measurements are scarce, like parts of South America, Africa, and Asia (Ward et al., 2018). At high latitudes, ice and snow dynamics apply a certain control to both river hydrology (Yamazaki et al., 2014) and wave and storm surge dynamics (Vousdoukas et al., 2017). However, such processes are not resolved by the global models used to generate the forcing datasets used in the present study. For these reasons, the present findings should be interpreted with care, especially in northern regions (Couasnon et al., 2019).

The two approaches investigated here provide information only on the *potential* for CF. The actual CF occurrence depends also on the local topography which can favour or not the interaction between the CF drivers; also, concurrent but not hydrologically-interacting storm surge and pluvial or fluvial flooding are relevant as they can, e.g., limit the ability to respond to emergency, and amplify the impacts that the two hazards would have caused if they occurred in isolation (Martius et al., 2016; Barton et al., 2016; Zscheischler et al., 2019). Moreover, while the two approaches investigated here are supposed to represent different CF mechanisms, separating the CF mechanisms in this way could be misleading, as CF may happen due to a combination of river discharge, local rainfall and associated surface runoff, together with high sea level. For example, in July 2019, New Orleans (Louisiana, US) (Vahedifard et al., 2016) was simultaneously threatened by the tropical Storm Barry causing local rainfall and storm surge around the coast, and by extremely high water discharge from the Mississippi River which lasted from March to July. Local CF risk assessment at the local scale should be carried out via complex hydrological modelling that can take into account the complex mechanisms causing CF, including storm surges, waves, astronomical tides, and when necessary not only fluvial or pluvial flooding but also their combination.

3 Conclusions

We conduct two global-scale potential CF hazard assessments, using either storm surge and precipitation, or storm surge and river discharge model data, and compare how the choice of precipitation versus river discharge as covariate with storm surge affects the results. We find that the two approaches result in similar spatial CF hazard patterns, which tend to deviate as the river catchment size increases.

Due to data scarcity, current large-scale CF assessments rely on approaches and model-based datasets similar to those used here. This study indicates that for these large-scale assessments, a precipitation-based CF analysis can provide satisfactory information on the CF potential in estuaries of small and medium size rivers (catchment smaller than about 5-10,000 Km²). Moreover, a precipitation-based CF analysis allows for assessing the CF hazard arising from the interaction of local coastal



230 rainfall and storm surges where no rivers exist, or along the mouths of small rivers often not represented in global river datasets.
Naturally, employing river discharge data should always be preferred to using precipitation when studying both the large and
local scale CF in estuaries, when data are available. The importance of using river discharge data is even greater in estuaries of
long rivers.

Data availability. Precipitation data is available on request online (<https://ec.europa.eu/jrc/en/publication/mswep-3-hourly-025-global-gri>
235 dded-precipitation-1979-2014-merging-gauge-satellite-and-reanalysis). Sea level data are available at <https://data.jrc.ec.europa.eu/collecti>
on/liscoast (further inquiries should be addressed to MIV). River discharge data were obtained from the dataset “Paired time series of daily
discharge and storm surge” are available at: <https://doi.org/10.5281/zenodo.3258007>.




Figure A1. The same as Fig. 1, but the extremes are defined as 2-year return levels.

Appendix A: Supporting figures




Figure A2. The same as Fig. 1, but the results are based on extremes defined considering 2-year return levels.




Figure A3. (a) The same as Fig. 3a, but the results are based on extremes defined considering 2-year return levels (the figure is based on Fig. A1). (b) As Fig. 3d, but with linear y-axis.




Figure A4. Locations with the lowest potential compound flooding probability based on precipitation and on river discharge. Locations with return periods below the 10th and 5th percentile are shown in black and red, respectively. The extremes are defined considering 5-year return levels. CF return periods are based on precipitation in (a) and on river discharge in (b).




Figure A5. The same as Fig. A4, but the results are based on extremes defined considering 2-year return levels.



240 *Author contributions.* EB initiated the study, carried out the data analysis, and drafted the manuscript. EB designed the study development with contributions from MIV. EB and MIV worked on the final manuscript version. MIV performed the storm surge runs. All the authors gave conceptual inputs during the data analysis, discussed the results, and commented on the manuscript.

Competing interests. The authors declare no competing interests.

245 *Acknowledgements.* EB acknowledges financial support from the European Research Council grant ACRCC (project 339390). EB acknowledge the European COST Action DAMOCLES (CA17109). MV acknowledges financial support from the EUPHEME project and CoCliServ project, which are part of ERA4CS, an ERA-NET initiated by JPI Climate and co-funded by the European Union (grant no. 690462). The authors would like to thank Lorenzo Mentaschi and Evangelos Voukouvalas who contributed in the generation of the storm surge time series, and Dirk Eilander for creating and publicly sharing the dataset “Paired time series of daily discharge and storm surge” which was fundamental for developing this study.



References

- 250 Amrhein, V., Greenland, S., and McShane, B.: Scientists rise up against statistical significance, 2019.
- Bader, B. and Yan, J.: eva: Extreme Value Analysis with Goodness-of-Fit Testing, R package version, 0.2.4, 2016.
- Barton, Y., Giannakaki, P., Von Waldow, H., Chevalier, C., Pfahl, S., and Martius, O.: Clustering of regional-scale extreme precipitation events in southern Switzerland, *Monthly Weather Review*, 144, 347–369, 2016.
- Beck, H. E., van Dijk, A. I., de Roo, A., Dutra, E., Fink, G., Orth, R., and Schellekens, J.: Global evaluation of runoff from ten state-of-the-art hydrological models, *Hydrology and Earth System Sciences*, 21, 2881–2903, 2017a.
- 255 Beck, H. E., Van Dijk, A. I., Levizzani, V., Schellekens, J., Gonzalez Miralles, D., Martens, B., and De Roo, A.: MSWEP: 3-hourly 0.25 global gridded precipitation (1979-2015) by merging gauge, satellite, and reanalysis data, *Hydrology and Earth System Sciences*, 21, 589–615, 2017b.
- Bevacqua, E., Maraun, D., Haff, I. H., Widmann, M., and Vrac, M.: Multivariate statistical modelling of compound events via pair-copula constructions: analysis of floods in Ravenna (Italy), *Hydrology and Earth System Sciences*, 21, 2701, <https://doi.org/10.5194/hess-21-2701-2017>, 2017.
- 260 Bevacqua, E., Maraun, D., Voudoukas, M. I., Voukouvalas, E., Vrac, M., Mentaschi, L., and Widmann, M.: Higher probability of compound flooding from precipitation and storm surge in Europe under anthropogenic climate change, *Science advances*, 5, eaaw5531, <https://doi.org/10.1126/sciadv.aaw5531>, 2019a.
- 265 Bevacqua, E., Voudoukas, M. I., Zappa, G., Hodges, K., Mentaschi, L., Maraun, D., and Shepherd, T. G.: Global projections of compound flooding from storm surges and precipitation under anthropogenic climate change, in preparation, 2019b.
- Blöschl, G., Hall, J., Viglione, A., Perdigão, R. A., Parajka, J., Merz, B., Lun, D., Arheimer, B., Aronica, G. T., Bilibashi, A., et al.: Changing climate both increases and decreases European river floods, *Nature*, 573, 108–111, 2019.
- Coles, S., Bawa, J., Trenner, L., and Dorazio, P.: An introduction to statistical modeling of extreme values, vol. 208, Springer, 2001.
- 270 Couasnon, A., Eilander, D., Muis, S., Veldkamp, T. I. E., Haigh, I. D., Wahl, T., Winsemius, H., , and Ward, P. J.: Measuring compound flood potential from river discharge and storm surge extremes at the global scale and its implications for flood hazard, *Nat. Hazards Earth Syst. Sci. Discuss.*, in review, <https://doi.org/https://doi.org/10.5194/nhess-2019-205>, <https://doi.org/10.5194/nhess-2019-205>, 2019.
- Dee, D. P., Uppala, S., Simmons, A., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M., Balsamo, G., Bauer, P., et al.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Quarterly Journal of the royal meteorological society*, 137, 553–597, 2011.
- 275 Eilander, D.: Paired time series of daily discharge and storm surge, Zenodo, <https://doi.org/https://doi.org/10.5281/zenodo.3258007>, <https://doi.org/10.5281/zenodo.3258007>, 2019.
- Ganguli, P. and Merz, B.: extreme coastal Water Levels exacerbate fluvial flood Hazards in northwestern europe, *Scientific reports*, 9, 1–14, 2019a.
- 280 Ganguli, P. and Merz, B.: Trends in Compound Flooding in Northwestern Europe During 1901–2014, *Geophysical Research Letters*, 46, 10 810–10 820, 2019b.
- Heffernan, J., Stephenson, A., and Gilleland, E.: Ismev: an introduction to statistical modeling of extreme values, R package version, 1.41, <https://CRAN.R-project.org/package=ismev>, 2016.



- Hendry, A., Haigh, I. D., Nicholls, R. J., Winter, H., Neal, R., Wahl, T., Joly-Laugel, A., and Darby, S. E.: Assessing the characteristics and drivers of compound flooding events around the UK coast, *Hydrology and Earth System Sciences*, 23, 3117–3139, <https://doi.org/10.5194/hess-23-3117-2019>, 2019.
- Klerk, W.-J., Winsemius, H., van Verseveld, W., Bakker, A., and Diermanse, F.: The co-occurrence of storm surges and extreme discharges within the Rhine–Meuse Delta, *Environmental Research Letters*, 10, 035 005, 2015.
- Kumbier, K., Carvalho, R. C., Vafeidis, A. T., and Woodroffe, C. D.: Investigating compound flooding in an estuary using hydrodynamic modelling: a case study from the Shoalhaven River, Australia, *Natural Hazards and Earth System Sciences*, 18, 463, 2018.
- Martius, O., Pfahl, S., and Chevalier, C.: A global quantification of compound precipitation and wind extremes, *Geophysical Research Letters*, 43, 7709–7717, 2016.
- Mentaschi, L., Voudoukas, M. I., Voukouvalas, E., Dosio, A., and Feyen, L.: Global changes of extreme coastal wave energy fluxes triggered by intensified teleconnection patterns, *Geophysical Research Letters*, 44, 2416–2426, 2017.
- Muis, S., Verlaan, M., Winsemius, H. C., Aerts, J. C., and Ward, P. J.: A global reanalysis of storm surges and extreme sea levels, *Nature communications*, 7, 2016.
- Olbert, A. I., Comer, J., Nash, S., and Hartnett, M.: High-resolution multi-scale modelling of coastal flooding due to tides, storm surges and rivers inflows. A Cork City example, *Coastal Engineering*, 121, 278–296, 2017.
- Paprotny, D., Voudoukas, M. I., Morales-Nápoles, O., Jonkman, S. N., and Feyen, L.: Compound flood potential in Europe, *Hydrol. Earth Syst. Sci. Discuss*, 2018, 1–34, 2018.
- Schepsmeier, U., Stoeber, J., Brechmann, E. C., Graeler, B., Nagler, T., and Erhardt, T.: VineCopula: Statistical inference of vine copulas, R package version, 2.0.5, 2016.
- Vahedifard, F., AghaKouchak, A., and Jafari, N. H.: Compound hazards yield Louisiana flood, *Science*, 353, 1374, 2016.
- van den Hurk, B., van Meijgaard, E., de Valk, P., van Heeringen, K.-J., and Gooijer, J.: Analysis of a compounding surge and precipitation event in the Netherlands, *Environmental Research Letters*, 10, 035 001, <https://doi.org/10.1088/1748-9326/10/3/035001>, 2015.
- Vandenberghe, S., Verhoest, N., Onof, C., and De Baets, B.: A comparative copula-based bivariate frequency analysis of observed and simulated storm events: A case study on Bartlett-Lewis modeled rainfall, *Water Resources Research*, 47, 2011.
- Voudoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., and Feyen, L.: Extreme sea levels on the rise along Europe’s coasts, *Earth’s Future*, 5, 304–323, 2017.
- Voudoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., Jevrejeva, S., Jackson, L. P., and Feyen, L.: Global probabilistic projections of extreme sea levels show intensification of coastal flood hazard, *Nature communications*, 9, 2360, 2018.
- Wahl, T., Jain, S., Bender, J., Meyers, S. D., and Luther, M. E.: Increasing risk of compound flooding from storm surge and rainfall for major US cities, *Nature Climate Change*, 5, 1093–1097, 2015.
- Ward, P. J., Couasnon, A., Eilander, D., Haigh, I. D., Hendry, A., Muis, S., Veldkamp, T. I., Winsemius, H. C., and Wahl, T.: Dependence between high sea-level and high river discharge increases flood hazard in global deltas and estuaries, *Environmental Research Letters*, 13, 084 012, 2018.
- Wu, W., McInnes, K., O’grady, J., Hoeke, R., Leonard, M., and Westra, S.: Mapping Dependence Between Extreme Rainfall and Storm Surge, *Journal of Geophysical Research: Oceans*, 123, 2461–2474, 2018.
- Yamazaki, D., O’Loughlin, F., Trigg, M. A., Miller, Z. F., Pavelsky, T. M., and Bates, P. D.: Development of the global width database for large rivers, *Water Resources Research*, 50, 3467–3480, 2014.



Zscheischler, J., Westra, S., Hurk, B. J., Seneviratne, S. I., Ward, P. J., Pitman, A., AghaKouchak, A., Bresch, D. N., Leonard, M., Wahl, T., et al.: Future climate risk from compound events, *Nature Climate Change*, p. 1, 2018.

Zscheischler, J., Martius, O., Westra, S., Bevacqua, E., Raymond, C., Horton, R., van den Hurk, B., AghaKouchak, A., Jézéquel, A., Mahecha, M. D., Maraun, D., Ramos, A. M., Ridder, N., Thiery, W., and Vignotto, E.: A typology of compound weather and climate events, Submitted, 2019.

325