

# Supplementary Material – Exploring the added value of a long-term multidisciplinary dataset in drought research - a drought catalogue for southwestern Germany dating back to 1801

Mathilde Erfurt<sup>1</sup>, Georgios Skiadaresis<sup>2</sup>, Erik Tijdeman<sup>3</sup>, Veit Blauhut<sup>4</sup>, Jürgen Bauhus<sup>2</sup>, Rüdiger Glaser<sup>1</sup>, Julia Schwarz<sup>2</sup>, Willy Tegel<sup>5</sup>, Kerstin Stahl<sup>4</sup>

<sup>1</sup>Environmental Social Science and Geography, Albert-Ludwigs-University, Freiburg i. Br., 79085, Germany


<sup>2</sup>Chair of Silviculture, Albert-Ludwigs-University, Freiburg i. Br., 79085, Germany

<sup>3</sup>Institute of Geography, Ruprecht-Karls-University, Heidelberg, 69117, Germany


<sup>4</sup>Environmental Hydrological Systems, Albert-Ludwigs-University, Freiburg i. Br., 79085, Germany

<sup>5</sup>Chair of Forest Growth and Dendroecology, Albert-Ludwigs-University, Freiburg i. Br., 79085, Germany


Correspondence to: mathilde.erfurt@geographie.uni-freiburg.de




**Figure S1: The combined chronology.**



**Figure S2: The oak chronology.**



**Figure S3: The fir chronology.**



**Figure S4: The spruce chronology.**

The two meteorological drought indices (SPI and SPEI), calculated for different accumulation periods and for different months of the year, correlated strongly with each other (Figures S.5 and S.6). As expected, series of SPEI and SPI for the same accumulation periods (e.g. 3 months) and of the same months (e.g. June) showed nearly perfect correlations ( $r > 0.9$ ). The relationship between the same meteorological drought indices for the same months but for different accumulation periods was weaker (e.g. between SPEI-3 of June and SPEI-6 of June). Similarly, strong correlations ( $r > 0.7$ ) were found between streamflow percentiles of the two considered rivers in BW and for both accumulation periods examined (Mar-Nov and Jun-Nov). Less pronounced relationships were observed among the tree-ring chronologies, except for the strong correlation between the combined chronology and the oak chronology in both 40 year periods. The two conifer chronologies (spruce and fir) were strongly correlated in both periods ( $r=0.68$  and  $0.67$  in the early and later period respectively).

Apart from the expected relationships between indicators belonging to the same groups, strong correlations were also observed between indices belonging to different groups. Streamflow percentiles correlated most strongly with long-term accumulation periods (12 and 24 months) of meteorological indices in both the early and later period ( $r > 0.6$ ). Correlations between streamflow and SPI/ SPEI-6 were slightly weaker and even weaker for indices calculated for an accumulation period of three months. Tree-ring chronologies showed overall weak correlations with streamflow percentiles with the exception of the oak and the combined tree-ring chronologies which were significantly correlated with the two streamflow series from the Rhine River. However, the combined tree-ring chronology as well as the oak chronology showed the strongest correlations with short-term meteorological drought indicators in both periods. In the early period, high correlations were observed between these two chronologies and the series of SPEI/SPI-3 of August and SPEI/SPI-6 of September. In the later period, these correlations were weaker and even absent in the case of SPEI/SPI-3, while at the same time the strongest correlations were observed with SPI/SPEI-6 of June. A similar change, although weaker in strength, is observed for the spruce chronology for the two periods. Apart from a weak correlation with SPEI-24 of December, no relationship between the fir chronology and meteorological drought indicators was found. However, fir growth showed weak but significant correlation in the later period with short-term (3 and 6 months) SPI and SPEIs of June.

Correlation matrix 1901–2011




Figure S5: Correlations between the different indices over their common period (1901-2011).





Figure S6: Correlations between the different indices and for two different time periods (1901 to 1940 and 1972 to 2011).

## Similarity Between Datasets 1901–2011



**Figure S7: Quantification of similarities in extreme drought event occurrence between the different pairs of indices (Eq. 1) over their common period (1901-2011).**

**Table S1: The tree-ring datasets used in this study. Information on the location, species, elevation, sources and basic descriptive statistics.**

| Site Code       | Lat   | Lon  | Species                       | Elevation (m) | Online source   | N series | eps  | snr   | first | last | N years |
|-----------------|-------|------|-------------------------------|---------------|---|----------|------|-------|-------|------|---------|
| BAME            | 49.49 | 9.78 | Quercus spp.                  | 220.6         | no  | 210      | 0.98 | 46.96 | 1731  | 2014 | 284     |
| B-W_ABI         | 48.7  | 9.03 | Abies alba Mill.              | -             | no  | 115      | 0.77 | 3.27  | 1029  | 1952 | 924     |
| B-W_QUE         | 48.7  | 9.03 | Quercus spp.                  | -             | no  | 404      | 0.94 | 15    | 1242  | 2014 | 773     |
| ECO             | 48.11 | 7.89 | Quercus spp.                  | 250           | no  | 40       | 0.88 | 7.03  | 1921  | 2016 | 96      |
| EEM             | 48.09 | 7.85 | Quercus spp.                  | 212           | no  | 42       | 0.95 | 19.1  | 1941  | 2016 | 76      |
| EEO             | 48.09 | 7.85 | Quercus spp.                  | 212           | no  | 40       | 0.92 | 11.37 | 1896  | 2016 | 121     |
| EPPI            | 49.13 | 8.92 | Quercus spp.                  | 200           | no  | 88       | 0.94 | 15.01 | 1770  | 2014 | 245     |
| EWM             | 48.09 | 7.84 | Quercus spp.                  | 197           | no  | 40       | 0.95 | 19.73 | 1956  | 2016 | 61      |
| EWO             | 48.11 | 7.8  | Quercus spp.                  | 197           | no  | 39       | 0.95 | 18.58 | 1915  | 2017 | 103     |
| FEM             | 48.04 | 7.85 | Quercus spp.                  | 237           | no  | 41       | 0.98 | 57.7  | 1923  | 2016 | 94      |
| FEO             | 48.04 | 7.84 | Quercus spp.                  | 237           | no  | 40       | 0.94 | 16.63 | 1813  | 2016 | 204     |
| Fir_Schwa<br>tz | 48.02 | 7.95 | Abies alba Mill.              | -             | no  | 60       | 0.89 | 7.76  | 1900  | 2016 | 117     |
| Fir_<br>Schwatz | 47.97 | 8.88 | Abies alba Mill.              | -             | no  | 60       | 0.97 | 30.37 | 1848  | 2012 | 165     |
| FNM             | 48.04 | 7.89 | Quercus spp.                  | 291           | no  | 40       | 0.94 | 15.66 | 1926  | 2016 | 91      |
| FNO             | 48.05 | 7.88 | Quercus spp.                  | 291           | no  | 40       | 0.93 | 13.25 | 1783  | 2016 | 234     |
| FORS            | 49.16 | 8.58 | Quercus spp.                  | 112           | no  | 44       | 0.92 | 10.79 | 1811  | 2014 | 204     |
| FWO             | 48.05 | 7.83 | Quercus spp.                  | 226           | no  | 40       | 0.95 | 19.6  | 1836  | 2016 | 181     |
| germ041w        | 47.8  | 8.08 | Picea abies (L.)<br>H. Karst. | 1200          | <a href="https://www.ncdc.noaa.gov/paleo-search/study/4631">https://www.ncdc.noaa.gov/paleo-search/study/4631</a> | 13       | 0.97 | 27.89 | 1932  | 1992 | 61      |
| germ042w        | 48.08 | 7.68 | Picea abies (L.)<br>H. Karst. | 440           | <a href="https://www.ncdc.noaa.gov/paleo-search/study/4452">https://www.ncdc.noaa.gov/paleo-search/study/4452</a> | 16       | 0.89 | 8.33  | 1903  | 1995 | 93      |
| germ043w        | 48.09 | 7.68 | Abies alba Mill.              | 440           | <a href="https://www.ncdc.noaa.gov/paleo">https://www.ncdc.noaa.gov/paleo</a>                                     | 16       | 0.9  | 9.37  | 1926  | 1995 | 70      |



|          |       |      |                               |      |   |    |      |       |      |      |     |  |
|----------|-------|------|-------------------------------|------|---|----|------|-------|------|------|-----|--|
|          |       |      |                               |      | -<br>search/study/445<br>1  |    |      |       |      |      |     |  |
| germ044w | 47.83 | 7.7  | Picea abies (L.)<br>H. Karst. | 390  | <a href="https://www.ncdc.noaa.gov/paleo-search/">https://www.ncdc.noaa.gov/paleo-search/</a>                               | 20 | 0.93 | 13.52 | 1890 | 1995 | 106 |  |
| germ045w | 47.86 | 7.7  | Abies alba Mill.              | 390  | <a href="https://www.ncdc.noaa.gov/paleo-search/">https://www.ncdc.noaa.gov/paleo-search/</a>                               | 20 | 0.94 | 16.37 | 1894 | 1995 | 102 |  |
| germ046w | 47.8  | 7.75 | Picea abies (L.)<br>H. Karst. | 930  | <a href="https://www.ncdc.noaa.gov/paleo-search/?dataType=Id=18">https://www.ncdc.noaa.gov/paleo-search/?dataType=Id=18</a> | 20 | 0.95 | 17.06 | 1844 | 1995 | 152 |  |
| germ047w | 47.85 | 7.75 | Abies alba Mill.              | 930  | <a href="https://www.ncdc.noaa.gov/paleo-search/?dataType=Id=18">https://www.ncdc.noaa.gov/paleo-search/?dataType=Id=18</a> | 20 | 0.91 | 9.61  | 1844 | 1995 | 152 |  |
| germ048w | 47.8  | 7.98 | Picea abies (L.)<br>H. Karst. | 1320 | <a href="https://www.ncdc.noaa.gov/paleo-search/study/4645">https://www.ncdc.noaa.gov/paleo-search/study/4645</a>           | 20 | 0.97 | 34.78 | 1871 | 1995 | 125 |  |
| germ050w | 48.03 | 8.35 | Picea abies (L.)<br>H. Karst. | 880  | <a href="https://www.ncdc.noaa.gov/paleo-search/study/4721">https://www.ncdc.noaa.gov/paleo-search/study/4721</a>           | 20 | 0.95 | 20.48 | 1897 | 1994 | 98  |  |
| germ051w | 48    | 8.35 | Abies alba Mill.              | 880  | <a href="https://www.ncdc.noaa.gov/paleo-search/study/4720">https://www.ncdc.noaa.gov/paleo-search/study/4720</a>           | 20 | 0.98 | 54.45 | 1898 | 1994 | 97  |  |
| germ052w | 47.8  | 8.03 | Picea abies (L.)<br>H. Karst. | 1250 | <a href="https://www.ncdc.noaa.gov/paleo-search/">https://www.ncdc.noaa.gov/paleo-search/</a>                               | 20 | 0.92 | 11.39 | 1756 | 1995 | 240 |  |

|          |       |       |                               |     |   |    |      |       |      |      |     |  |
|----------|-------|-------|-------------------------------|-----|---|----|------|-------|------|------|-----|--|
|          |       |       |                               |     | search/study/463<br>2   |    |      |       |      |      |     |  |
| germ053  | 47.85 | 7.78  | Picea abies (L.)<br>H. Karst. | 490 | https://www.ncd<br>c.noaa.gov/paleo<br>-<br>search/study/454<br>5 | 20 | 0.96 | 23.08 | 1881 | 1995 | 115 |  |
| germ054  | 47.86 | 7.78  | Abies alba Mill.              | 490 | https://www.ncd<br>c.noaa.gov/paleo<br>-<br>search/study/454<br>4 | 20 | 0.9  | 8.97  | 1864 | 1995 | 132 |  |
| germ055w | 47.69 | 7.75  | Picea abies (L.)<br>H. Karst. | 940 | https://www.ncd<br>c.noaa.gov/paleo<br>-<br>search/study/465<br>2 | 20 | 0.95 | 18.63 | 1867 | 1995 | 129 |  |
| germ056w | 47.78 | 7.75  | Abies alba Mill.              | 940 | https://www.ncd<br>c.noaa.gov/paleo<br>-<br>search/study/465<br>1 | 20 | 0.86 | 6.17  | 1841 | 1995 | 155 |  |
| germ16   | 50.25 | 10.25 | Picea abies (L.)<br>H. Karst. | 550 | https://www.ncd<br>c.noaa.gov/paleo<br>-search/                   | 16 | 0.97 | 27.88 | 1830 | 1955 | 126 |  |
| germ17   | 49.48 | 10.58 | Picea abies (L.)<br>H. Karst. | 410 | https://www.ncd<br>c.noaa.gov/paleo<br>-search/                   | 10 | 0.79 | 3.66  | 1841 | 1972 | 132 |  |
| germ18   | 48.02 | 8.5   | Picea abies (L.)<br>H. Karst. | 770 | https://www.ncd<br>c.noaa.gov/paleo<br>-search/                   | 10 | 0.77 | 3.35  | 1906 | 1973 | 68  |  |
| germ3    | 47.82 | 7.77  | Abies alba Mill.              | 910 | https://www.ncd<br>c.noaa.gov/paleo/<br>study/2703                | 31 | 0.97 | 37.85 | 1868 | 1976 | 109 |  |
| LCM      | 49.64 | 8.65  | Quercus spp.                  | 293 | no  | 40 | 0.9  | 8.5   | 1818 | 2016 | 199 |  |
| LCO      | 49.65 | 8.68  | Quercus spp.                  | 293 | no  | 40 | 0.96 | 23.28 | 1873 | 2016 | 144 |  |
| LCY      | 49.65 | 8.67  | Quercus spp.                  | 293 | no  | 40 | 0.96 | 22.71 | 1976 | 2016 | 41  |  |

|                    |       |      |                               |       |    |     |      |       |      |      |     |
|--------------------|-------|------|-------------------------------|-------|----|-----|------|-------|------|------|-----|
| LNM                | 49.72 | 8.52 | Quercus spp.                  | 96    | no | 40  | 0.97 | 31.39 | 1948 | 2016 | 69  |
| LNO                | 49.72 | 8.52 | Quercus spp.                  | 96    | no | 40  | 0.97 | 32.44 | 1861 | 2016 | 156 |
| LSM                | 49.59 | 8.56 | Quercus spp.                  | 100   | no | 40  | 0.95 | 20.22 | 1909 | 2016 | 108 |
| LSO                | 49.58 | 8.58 | Quercus spp.                  | 100   | no | 40  | 0.96 | 23.63 | 1823 | 2016 | 194 |
| LWM                | 49.71 | 8.53 | Quercus spp.                  | 93    | no | 40  | 0.95 | 20.76 | 1904 | 2016 | 113 |
| LWO                | 49.72 | 8.54 | Quercus spp.                  | 93    | no | 42  | 0.92 | 12.02 | 1845 | 2016 | 172 |
| MUND               | 49    | 9.21 | Quercus spp.                  | 222.8 | no | 90  | 0.94 | 15.66 | 1797 | 2014 | 218 |
| OFFE               | 48.48 | 7.95 | Quercus spp.                  | 156.9 | no | 52  | 0.94 | 15.36 | 1826 | 2014 | 189 |
| rapp               | 49.24 | 9.1  | Quercus spp.                  | 240.8 | no | 17  | 0.58 | 1.38  | 1789 | 2018 | 230 |
| REUT               | 48.51 | 9.2  | Quercus spp.                  | 404.4 | no | 130 | 0.93 | 12.26 | 1608 | 2014 | 407 |
| sins               | 49.25 | 8.89 | Quercus spp.                  | 156.5 | no | 13  | 0.64 | 1.78  | 1815 | 2014 | 200 |
| Spruce_<br>Schwatz | 48.43 | 8.23 | Picea abies (L.)<br>H. Karst. | -     | no | 44  | 0.93 | 12.98 | 1848 | 2012 | 165 |
| UNTE               | 48.77 | 9.57 | Quercus spp.                  | 473   | no | 23  | 0.76 | 3.23  | 1820 | 2014 | 195 |
| URBA               | 48.81 | 9.58 | Quercus spp.                  | 268.3 | no | 138 | 0.94 | 16.56 | 1703 | 2014 | 312 |
| WIDD               | 49.33 | 9.43 | Quercus spp.                  | 311.8 | no | 61  | 0.93 | 12.74 | 1784 | 2014 | 231 |
| WITT               | 49.61 | 9.84 | Quercus spp.                  | 264.9 | no | 233 | 0.98 | 54.58 | 1769 | 2014 | 246 |