

**General comments** The quality of the paper, as well as the clarity in the presentation of results have substantially improved. All of the comments and questions raised have been well answered, although some of them the corresponding modifications were not included in the text. There are still minor corrections to be included before publication.

We thank Reviewer for his/her carefully reading and suggestion for the manuscript and we are sorry about the missing modifications in the previous version of the text.

#### Specific comments

**Figure 2 – In the answers to the reviewers it is mentioned that Figure 2 is suppressed from the manuscript since the authors agreed that it adds no relevant physical information. However, the figure and references to it are still present in the manuscript.**

The figure was removed and the referring sentence « An example of the operational measurements which started on January 2012 are provided in Figure 2 with figures ranging from 2.2~m to 2.6~m. ».

**L226-L227 – The loss of skill in I WV representation after 12 h needs further explanation. Rephrase to “... correspond to the time of the early stage of convection, which is less easy to predict and introduces inhomogeneities in the water vapour distribution due to precipitation processes (condensation, evaporation, cold pool formation, etc.).”**

The explanation was added in the text.

**L243-247 – The introduction to the SEVIRI results is missing and the paragraph seems to be disconnected from the context of the section. Please introduce the paragraph properly indicating that additional experiments were performed even though the results are not shown.**

The paragraph was written as follows: “In addition, AN and FG were compared with radiances from SEVIRI imager, on board the geostationary satellite Meteosat Second Generation, MSG), which is sensitive to moisture (channels WV 6.2  $\mu$ m for upper-troposphere and 7.3  $\mu$ m for mid-troposphere). This imager indeed represents another important source of humidity information, especially over oceans where no information from GNSS nor radiosondes is available. These humidity channel data are assimilated for each AROME-WMED analyses. The study of the FG and AN statistics for the REANA, OPERGNSS and NO GNSS shows no impact between the various experiments for these observations (not, shown, Table in supplement file).”

**The nomenclature is still confusing in some parts of the paper since, even if the different experiments and validation procedure are well defined in Sections 2.4 and 2.5, not always the nomenclature is respected. For example, in Section 3.1, often “analysis” is written, instead of AN, which can be confusing. Additionally, every time that a confusion between the AN and FG simulations is possible, a suffix to the experiment should be included as EXPERIMENT-FG and EXPERIMENT-AN. For example NOGNSS-FG or NOGNSS-AN. This occurs continuously throughout the text, please correct.**

The modifications were made in the text according the suggestion of the reviewer.

**In Figures 10, 11, 13, 15 and 16 add a label indicating the forecast range period (6 to 30h and 30 to 54h) to aid the comprehension of the figures.**

Label added.

**In Figure 12 labels indicating that the left panel corresponds to FG and the right to AN are needed to aid the differentiating of the graphs.**

FG and AN added.

**In Figures 14, 15 and 16 a label indicating the forecast range (3h) is needed.**

Labels added.

**The paragraph between Lines 348 and 354 needs more references to existing literature, otherwise the statements are not supported. There have been several other publications for this case study.**

References to Duffourg et al (2016) and to Martinet et al (2017) have been added in the text.

Several indications from the last review have not been corrected. Even if in the answers to the reviewers the authors agreed on changing them, none of the following corrections were considered:

- L241 – Change lower panel for right panel.

Corrected

- Section 3.3 – The explanation about the biases for high precipitation values in Figure 7 is missing. The answers given by the authors were “The loss of skill for OPERGNSS simulation for larger daily precipitation from the 8 3-h forecast is surely due to the lower number of occurrences for heavier precipitations. “. Still this information was not included in the new version of the manuscript.

Corrected

- IWC was not replaced for I WV

Corrected

- Iberic Peninsula was not corrected for Iberian Peninsula

Corrected

- Figure 2 still is in the manuscript.

The figure was removed.

Conclusions – The relevant messages are somewhat hidden due to some confusing sentences that should be corrected:

L366-L371 – Change for “Two reanalyses were produced after the HyMeX autumn campaign for the first SOP. The first one was carried out after the field campaign to provide reanalyses simulations of the period enjoying an updated AROMEWMED version. The second one, performed a few years after, was done to improve the quality of the first reanalyses, assimilating as many data as possible from the experimental field campaign (i.e. lidar and dropsonde humidity profiles) or from reprocessed data (such as GNSS-ZTD, wind profiles, high-vertical resolution radiosondes and Spanish doppler radars) using the latest AROME version.”

Done

L380-L382 – “... which includes all data sets. In total, the amount of denied data ranged between 0.15 % and 1.86 % of the total number of observations, depending on the OSE and the impact was thus expected to be small. Our study finds a small positive impact...”

Done

L385-L386 – The explanation of the reprocessed GNSS data is not clear. This information has to do with the OSEs NOGNSS and OPERGNSS but here in the conclusion it is not clear. Please review.

The explanation was modified as followin : Our study finds a small positive impact on humidity forecast at short term ranges of the reprocessed GNSS ground based zenithal total delay assimilation compared with NOGNSS. This data set is evenly distributed over the AROME-WMED domain and provided at each analysis time information on integrated water vapour. The impact of the data reprocessing for GNSS zenithal total delays compared to the processing made in real-time was also studied by comparing REANA with OPERGNSS. Even though a positive impact is observed with REANA compared to OPERGNSS, this improvement is not statistically significant.

L388-L289 – The information about the validation against RASTA is not clear. Please rephrase. Two are the problems in this sentence. The first is that it is stated that “no impact from Lidar data was found”, but then it is said that “except when comparing to RASTA”. The second problem is that the use of acronyms at this point should be avoided to aid a quick

**comprehension of the conclusions. It would be better writing “independent airborne radar observations” instead of RASTA.**

We replaced the sentence with « The sole impact of the assimilation of lidar data is found when comparing with independent airborne radar observations located over the Mediterranean Sea. »

### **Technical corrections**

**Abstract (L06-07) – Needs rephrasing “...Which are important to better describe the humidity entering mesoscale convective systems, were assimilated”.**

We rephrase as following : Among them, observations related to the low level humidity flow were assimilated. Such observations are important for the description of the feeding of the convective mesoscale systems with humidity (Duffourg and Ducrocq, 2011, Bresson et al., 2012 and Ricard et al.,2012).

**L56-L58 – It is important to mind the scope of the paper explaining that that the four observation systems evaluated convey information either on the moisture distribution or on wind circulation. Hence, effectively it is an assimilation aiming at correcting the moisture flux for the SOP1 period. To express this condition more clearly rephrase for “These observation data sets, assimilated in the AROME-WMED reanalysis of SOP1 provided information on the atmospheric humidity distribution and the wind circulation, hence describing the moisture flux of the low-level flow.”**

Done

**L70-L71 – For clarity rephrase to “These experiments consist of removing one observation data set to compare its forecast quality with the reanalysis simulations, where all observations are assimilated ”.**

Done

**Figure 1 – For readers, focusing only on figures, change “LEANDRE” in legend for “LEANDRE (lidar)”. Like this at a glimpse, the reader knows what observation type is LEANDRE. For readers unfamiliar with the observations from HyMeX.**

Done

**L112-L113 – Correct typo. Parenthesis within parenthesis.**

Global Navigation Satellite System was removed as GNSS was defined earlier in the text.

**L125 – Rephrase “were assimilated in AROME-WMED (see Fig. 1)”.**

Done

**L135 – Correct typo. Brackets within brackets.**

Done

**L219 – “Comparison to the Integrated Water Vapour”.**

Done

**L222 – “the RMSE of the differences (Fig.3, right panel).**

Done

**\*L227 – “... in data assimilation (Figure 4).”**

Done

**L227 – Analysis should be AN, see specific comments.**

Done

**L228-L229 – “... during the assimilation process, with differences between the OSEs.”.**

Done

**\*L231-232”... 3-h later. The lowest differences are obtained with REANA-FG and the largest ones with NOGNSS-FG. OPERGNSS-FG differences are close to REANA...”.**

Done

**L236-L238 – Rephrase: “... data versus OPER-GNSS appears between 700 and 850 hPa.”**

“The largest improvement of the assimilation of GNSS data is found between 600 and 850 hPa, NOGNSS-FG being larger than REANA-FG and OPER-GNSS-FG in this part of the atmosphere. In addition, OPER-GNSS-FG is larger than REANA-FG between 700 and 850 hPa showing a slight benefit of assimilating REPROC-GNSS instead of OPER-GNSS.”

“

**L230-L238 – Add suffixes AN and FG, as in Figure 4 (see specific comments). Adding suffixes helps the reading comprehension.**

Done

**L239-L240 – Analysis should read AN (see specific comments).**

Done

**L256-L257 – Starting a new paragraph is not advised since the explanation of this bias corresponds to the problem described before. It should read “... and OPERGNSS (4.8 mm). This bias could be explained...”.**

Corrected.

**Table 3 – Change order of columns, it should be: Mean ZTD (m) | AN ZTD Correlation | FG ZTD Correlation.**

Done

**L264 – “It is worth reminding...”.**

Done

**L289 “... when compared to ...”.**

Corrected.

**L289 – Correct to “to values averaged”.**

Done

**L290 – Correct to “itself”.**

Done

**L294 – “... wind (not shown).”.**

Done

**L310 – “... differences at 300 hPa (left panel).”.**

Done

**L324-L325 – Sentence reads odd, reformulate.**

The sentence was changed into: « No significant impact has been noticed over the HyMeX domain. However, when focusing over the Iberian Peninsula, the ETS for the 24-h accumulated precipitation from the sum of the 8 3-h precipitation background forecast (Figure 13, former Figure 14) is improved with the assimilation of Spanish radar data. »

**L341 – “... the Mediterranean Sea...”.**

Done

# Data assimilation impact studies with the AROME WMED reanalysis of the HyMeX SOP1

Nadia Fourrié<sup>1</sup>, Mathieu Nuret<sup>1</sup>, Pierre Brousseau<sup>1</sup>, and Olivier Caumont<sup>1</sup>

<sup>1</sup>CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

**Correspondence:** Nadia Fourrié (nadia.fourrie@meteo.fr)

## Abstract.

This study was performed in the framework of HyMeX (Hydrological cycle in the Mediterranean Experiment) which aimed to study the heavy precipitation that regularly affects the Mediterranean area. A reanalysis with a convective-scale model AROME-WMED was performed which assimilated most of all available data for a 2 month period corresponding to the first

5 Special Observation Period of the field campaign (Fourrié et al., 2019). ~~Among them, observations related to the low level humidity flow which are important for the description of the feeding of the convective mesoscale systems with humidity (Duffourg and Duerœq, 2011, Bresson et al., 2012 and Ricard et al., 2012), were assimilated. Among them, observations related to the low level humidity flow were assimilated. Such observations are important for the description of the feeding of the convective mesoscale systems with humidity (Duffourg and Ducrocq, 2011, Bresson et al., 2012 and Ricard et al., 2012).~~

10 Among them there were a dense reprocessed network of high quality Global Navigation Satellite System (GNSS) Zenithal Total Delay (ZTD) observations, reprocessed data from wind profilers, lidar-derived vertical profiles of humidity (ground and airborne) and Spanish radar data. The aim of the paper is to assess the impact of the assimilation of these four observation types on the analyses and the forecasts from the 3h forecast range (first guess) up to the 48-h forecast range. In order to assess this impact, several OSEs or also-called denial experiments, were carried out by removing one single data set from the observation

15 data set assimilated in the reanalysis.

Among the evaluated observations, it is found that the ground-based GNSS ZTD data set provides the largest impact on the analyses and the forecasts as it represents an evenly spread and frequent data set providing information at each analysis time over the AROME-WMED domain. The impact of the reprocessing of GNSS ZTD data also improves the forecast quality but this impact is not statistically significant. The assimilation of the Spanish radar data improves the 3-h precipitation forecast

20 quality as well as the short term (30-h) precipitation forecasts but this impact remains located over Spain. Moreover, marginal impact from wind profilers was observed on wind background quality. No impacts have been found regarding lidar data as they represent a very small data set, mainly located over the sea.

## 1 Introduction

Heavy precipitation regularly affects the Mediterranean area with huge damages and sometimes casualties. One of the aims

25 of the Hydrological cycle in the Mediterranean Experiment (HyMeX ; Drobinski et al. (2014)) was to study the high impact

weather events, especially during the first Special Observation Period one (SOP1, Ducrocq et al. (2014)), which took place in the autumn 2012 (5 September - 6 November 2012) in northwestern Mediterranean. The importance of an accurate description of the low-level humidity flow, which feeds the mesoscale systems, was shown in previous studies (Duffourg and Ducrocq, 2011; Bresson et al., 2012; Ricard et al., 2012). This is why during this period research observations were deployed over the north-western Mediterranean area. These observations aimed at a better description of the humidity and wind fields. As an example, water vapour lidars were deployed in Candillargues and Menorca island (pink dots in Figure 1). Particular attention was also paid to the control of data quality.

Another important element to better understand the key processes related to the high precipitation and their forecasting is the convective scale modeling. Since many years, such numerical weather prediction models have been implemented in operations to enhance the forecast quality. In addition, the forecast quality depends on their initial atmospheric conditions, which are determined with data assimilation system.

For the HyMeX SOP1 campaign, an AROME (Application of Research to Operations at MEsoscale, Seity et al., 2011) version was developed and ran in real-time to forecast and study heavy precipitation in this region: the AROME-WMED (western Mediterranean) model (Fourrié et al., 2015). This model is centered over the western Mediterranean basin and includes a data assimilation system, which provides every 3 hours an analysis of the meteorological situation. In the framework of the Innovative Observing and Data Assimilation Systems for severe weather events in the Mediterranean (IODA-MED) project, two reanalyses were performed after the campaign (Fourrié et al., 2019) with the aim of providing new references for process studies. The first one intended to provide a homogeneous data set of atmospheric fields (which was not the case in real-time version due to a system upgrade in the middle of the SOP1). The second one included in addition a maximum of observations deployed during SOP1 field campaign with a more recent version of the model. The latter will be considered in this study.

Among the research observations assimilated in AROME-WMED reanalysis were the humidity profiles from ground based and airborne lidars. Reprocessing after the campaign was also performed for the wind profiler data (Saïd et al., 2016) and the ground based GNSS (Global Navigation Satellite System) zenithal total delays (ZTD) (Bock et al., 2016) to improve data quality and filter out bad data.

Previous impact studies were already performed for this type of observations in other contexts. For example, Bielli et al. (2012); Grzeschik et al. (2008) tested the impact of the assimilation of water vapour lidars in meso-scale models and found a positive impact of such an assimilation up to the 24-h forecast range. Benjamin et al. (2004) studied the impact of a wind profiler network and obtained a positive impact on short-range (3–12 h) forecasts. Concerning the GNSS data, Mahfouf et al. (2015) showed systematic improvements of the atmospheric humidity short-range forecasts and of the structure and the location of precipitation in the AROME models as found previously in a heavy precipitation context (Boniface et al., 2009). These results agree well with previous studies performed in other NWP models (Macpherson et al., 2008; Gutman et al., 2004)

As previously mentioned, an accurate description of the low-level humidity flow is required to well simulate the evolution of the mesoscale system. The aim of the study presented here is to quantify the impact of four observation systems on the quality of precipitation simulation. [These observation data sets, assimilated in the AROME-WMED reanalysis of SOP1 provided information on the atmospheric humidity distribution and the wind circulation, hence describing the moisture flux of the low-level](#)

flow. ~~These observation data sets were assimilated in the AROME-WMED reanalysis of SOP1 and provided information on this low-level flow.~~ The observations are the reprocessed ZTD from the ground based GNSS (Bock et al., 2016), the humidity profiles from ground based and airborne lidars (Chazette et al., 2016 and Di Girolamo et al., 2016), reprocessed wind profiler data (Saïd et al., 2016) and the Doppler winds and reflectivities from the Spanish radars. To achieve this, a number of denial 65 data assimilation experiments, consisting in removing one observation type, were carried out during the 2-month period of SOP1.

The paper is arranged as follows. Section 2 describes the AROME-WMED configurations, the observations data sets and the denial experiments. Section 3 assesses the impact of the ground-based GNSS data assimilation on the analyses, the background and on the forecast quality during SOP1. Section 4 provides information on the impact of other observation types (i. e. wind 70 profilers, lidars and Spanish radars). Section 5 focusses on the impact of all these data on the IOP 16a case study. Finally, conclusions are given in Section 6.

## 2 Sensitivity study description and validation methodology

### 2.1 "Denial" Experiment Methodology

To study the contribution of the observations on the analysis and forecast quality of the heavy precipitating events of the SOP1, 75 denial experiments have been devised. ~~These experiments consist of removing one observation data set and to compare the forecast quality with the one originating from assimilating all the observations.~~ These experiments consist of removing one observation data set to compare its forecast quality with the reanalysis simulations, where all observations are assimilated. Here, denial experiments were conducted on the following four observation types: the ground-based GNSS ZTD, the wind profilers, the water vapour lidars and Spanish radars. They were performed with the AROME-WMED model.

### 80 2.2 AROME-WMED configuration

The different AROME-WMED model configurations are described in Fourrié et al. (2015, 2019) and rely on the operational limited area model AROME (Seity et al., 2011; Brousseau et al., 2016) version running at Météo-France since 2008. At the time of the SOP1 campaign, analyses were performed at 2.5 km horizontal resolution every 3 hours with a three dimensional variational data assimilation (3D-Var, Brousseau et al., 2011). The AROME-WMED version used in this study as the reference 85 is the second reanalysis one, named hereafter REANA. An extensive description of this reanalysis can be found in Fourrié et al. (2019). The main components are recalled here. The REANA dataset has a 2.5 km horizontal resolution and the model has 60 vertical levels from 10 m above the surface to 1 hPa. Deep convection is explicitly resolved and one-moment microphysical scheme with five classes of hydrometeors is used (Pinty and Jabouille, 1998; Caniaux et al., 1994).

Initial atmospheric states of AROME-WMED come every 3 hours from 3D-Var analyses assimilating observations within a 90 +/- 1h30 assimilation window. This systems analyzes the two components of horizontal wind, temperature, specific water-vapor humidity and surface-pressure fields on the model grid at full resolution. The other prognostic model fields (turbulent kinetic

Observations type	amount	percentage
Satellites	8,663,312	53.00%
Surface stations	2,485,620	15.21%
Radar	1,942,539	11.88%
Spanish radars	97,847	0.6%
Aircraft	1,413,313	8.65%
Radiosondes	1,319,523	8.07%
GNSS ZTD	302,191	1.85%
Wind profiler	191,012	1.17%
Lidars	19,470	0.12%
Total	16,346,191	100%


**Table 1.** Sorted amounts of assimilated data in REANA over the SOP1 period (5 September-5 November 2012).

energy, pressure departure from hydrostatic, vertical divergence and specific content of five condensed water species) are not updated by the analyses but copied from the background. The background error statistics are climatological. Based on the Berre (2000) multivariate formulation, cross-covariances between errors for different physical quantities are represented using 95 scale-dependent statistical regressions, including an extra balance relationship for specific humidity. The background error statistics have been calculated using forecast differences from a AROME-WMED Ensemble data assimilation (Brousseau et al. (2011) approach) over a 15-day period of the HyMeX SOP1 (17 to 31 October 2012) to be representative of the encountered meteorological conditions of the SOP1 in average. More details on these background error covariances are available in Fourrié et al (2019). Lateral boundary conditions are hourly provided by the global NWP ARPEGE (Courtier et al., 1991) forecasts 100 which also benefited from a maximum of assimilated observation with longer cutoff analyses. Each day at 00UTC, a 54 hour forecast is run. Conventional observations (from radiosondes, aircraft, surface stations, wind profiler, GNSS ZTDs), radar data and satellite observations (infrared and microwave radiances, atmospheric motion vectors and ocean surface winds from scatterometers) were assimilated.

Table 1 presents the distribution of assimilated data in REANA as a function of observation types. Satellite data represent 105 the majority of observations. This can be explained by the fact that the IASI sensor provides 44 channels per observation point. Surface observations provide 15.21% of assimilated data. Aircraft and radiosondes give similar amount of data (around 8%). GNSS ZTD represent 1.85% of the total and wind profilers 1.17%. Special efforts were made to assimilated non operational data types such as Lidar water vapour profiles and Spanish radar data. Humidity data from Lidar contribute very few with 0.12% of assimilated data. Radar data represent 11.88% of the total amount of assimilated data and Spanish ones only 0.6%.

### 110 2.3 Description of the studied observing systems

As mentioned above, four observing systems were studied. The location of these observations is shown in Figure 1, excepting the ground based GNSS ZTD location which is available in Figure 4 of Fourrié et al. (2019).



**Figure 1.** Location of observations considered in this study, with the exception of GNSS Zenithal Total Delays. Wind profilers are depicted with blue squares, ground based lidars with pink dots, assimilated Leandre II airborne profiles with purple dots, Spanish radars with orange symbols. Red symbols correspond to the French radar locations.

### 2.3.1 GNSS Zenithal Total Delays

GNSS ZTD provides useful information on precipitable water and pressure at a high temporal frequency and in all weather conditions. In REANA2, we considered here reprocessed data (REPRO-GNSS in the following) with a homogeneous reprocessing using a single software and more precise satellite orbits position and clocks (Bock et al., 2016), which were available for the whole SOP1. Additional data were also considered compared to the operational and data set available in near real-time. This data set, called hereafter OPER-GNSS, is provided by E-GVAP (EUMETNET EIG GNSS ([Global Navigation Satellite System](#)) water VApour Programme) and ZTD data for one reception station may be available for more than 10 processing centres. These ZTD data are assimilated according the methodology described in Mahfouf et al. (2015). The model equivalent is computed with the following equation (Mahfouf et al., 2015):

$$ZTD = 10^{-6} \int_0^{z_{top}} (k_1 \frac{p}{T} + k_3 \frac{e}{T^2}) dx \quad (1)$$

where  $p$  is the pressure,  $T$  the temperature,  $e$  the water vapour pressure,  $k_1 = 0.776 \text{ Pa}^{-1} \text{ K}$ , and  $k_3 = 3730 \text{ Pa}^{-1} \text{ K}^2$ ,  $x$  is the height above the ground and  $z_{top}$  is the height of the model top. After a monitoring of the difference between observations and model equivalent, observations with good statistics are selected in a 'white list'. ZTD data are also bias corrected and

an updated bias correction for each GNSS station was also computed in the REANA2 version. They are finally assimilated if they pass the first guess quality control which rejects data too far from the model background. Only one observation per 3-h assimilation and per surface station is assimilated for each analysis. Please refer to Mahfouf et al. (2015) for more information on the data assimilation of GNSS ZTD in AROME.

130 **2.3.2 Wind profilers**

Data from eight wind profiler radars (sounding in VHF or UHF bands) were assimilated in AROME-WMED (see Figure 1). These profilers provided vertical profiles of wind vector, turbulence, precipitation and the height of the atmospheric boundary layer and tropopause (Saïd et al., 2016). The measure principle is described in Annex 1 of Saïd et al. (2016)'s paper. Profilers measure the Doppler radial spectra of the returned signal backscattered by various types of targets. In order to derive the three 135 components of the wind, most of the HyMeX profilers use five beams. These data were available for the whole SOP1 in real-time and have been reprocessed after the SOP1 by Saïd et al. (2016) with an improved quality control to remove spurious data. Here, observations from 8 wind radars (UHF and VHF) mainly located in the South of France, in Corsica and Menorca (Figure 1) were considered. These observations are assimilated as vertical profiles of horizontal wind.

**2.3.3 Lidars**

140 During SOP1, ground based and airborne lidars were operated. The mobile Water vapour and Aerosol Raman Lidar (WALI, Chazette et al, 2016) operates with an emitted wavelength of 354.7 nm. This instrument was operated at a site close to Ciutadella (western part of Menorca located by 39 59 07 N and 3 50 13 E). Mixing ratio profiles were delivered with a resolution of 15 m for the 0 m - 6000 m altitude range. A detailed description of this instrument can be found in Chazette et al. (2016). The raw 145 vertical resolution of the data is 75 m but for assimilation above 2000 m, resolution was thinned starting from 75 m to 450 m above 5000 m.

The second ground based lidar, the BASIL instrument (Di Girolamo et al., 2016) was located in Candillargues in the South of France. The original data resolution is 30 m but data were thinned at 60 m below 1000 m, increasing up to 420 m above 4000 m in the assimilation. For WALI, 292 mixing ratio profiles were assimilated in REANA, covering the period 17 September 2012 - 03 UTC to 27 October 2012 - 21 UTC, whereas for BASIL, 172 profiles were assimilated, covering the period 10 September 150 2012 - 09 UTC to 5 November 2012 15 UTC.

Concerning Leandre II lidar (Chazette et al., 2016) on board ATR aircraft, data were available for 22 analysis slots (512 assimilated profiles), covering the period 11 September 2012 09 UTC to 25 October 2012 21 UTC. Profiles with a 150 m vertical resolution were thinned at a 15 km horizontal resolution and are mainly located over the Mediterranean Sea (Figure 1).

**2.3.4 Spanish radars**

155 Doppler radial winds and reflectivities from five Spanish radars, located in Barcelona, Valencia, Almeria, Murcia, Palma de Mallorca and provided by AEMET were assimilated in REANA. After a strict quality control check to exclude data with gross

errors, only the three lowest elevations have been considered for the assimilation. Doppler wind are assimilated in the 3D-Var of AROME according the method described by Montmerle and Faccani (2009) and reflectivity data are assimilated as pseudo-observations of relative humidity profiles as proposed in Caumont et al. (2010) and implemented in Wattrelot et al. (2014).

160

Several procedures are applied to raw data in order to avoid as much as possible erroneous measurements entering the minimization. An observation operator allows to simulate radial Doppler winds measurements from the model horizontal wind based on Caumont and Ducrocq (2008). Only measurements performed within 150 km of the radar are considered due to the broadening of the beam with increasing distance and the lack of reliability. An observation error variance proportional to their 165 distance from the radar is applied in the minimization. Reflectivities are not directly assimilated but they are used to retrieve pseudo-observations of relative humidity from surrounding simulated reflectivity profiles through a unidimensional Bayesian inversion. A horizontal thinning on the data (Doppler winds and retrieved profiles of relative humidity) is performed to avoid horizontal correlation of observation errors: only one profile, having the most important number of elevations that passed the quality control, is selected in each  $15 \times 15 \text{ km}^2$  box.

170 **2.4 Description of the experiments**

Experiment name	Description	Difference (%) in the number of assimilated data
REANA	AROME-WMED reanalysis (2nd), see Fourrié et al. (2019)	
NOGNSS	REANA - reprocessed GNSS ZTD	-1.86%
OPERGNSS	NOGNSS + operational GNSS ZTD	-1.04%
NOLIDAR	REANA - LIDAR	-0.15%
NOWPROF	REANA - wind profilers	-1.12%
NORADSPAIN	REANA - Spanish radars	-0.6%

**Table 2.** Description of the data denial experiments discussed in this study and difference (in %) in the number of assimilated data compared to the reanalysis REANA.

Table 2 summarizes the names of the denial experiments and the observations considered. Five experiments were conducted over the 2-month period of SOP1 (from 5 September 2012 to 5 November 2012). They all used the same configuration of AROME-WMED, the differences lying in the observations assimilated. For each experiment, it differs only one observation type from the reanalysis (REANA) used as the reference. This allows to evaluate the impact of this observation type on the 175 analysis and the forecast. Among the five experiments, two experiments deal with the impact of GNSS ZTD. The first one, NOGNSS is obtained by removing the REPROC-GNSS ZTD from the assimilation. The second, called OPERGNSS, aimed to evaluate the impact of the REPROC-GNSS data set provided by Bock et al. (2016) compared to the OPER-GNSS data set provided by E-GVAP. The E-GVAP data set was thus assimilated in replacement of the Bock et al. (2016)'s one in OPERGNSS.

180 The NOLIDAR experiment is the run with no airborne nor ground-based Lidar data in the data assimilation. The NOWPROF experiment is obtained by removing the wind profiler data and the NORADSPAIN experiment was run without any data from the five Spanish radars. 97847 observations, representing 0.6% of the total number of assimilated observations, were removed in the NORADSPAIN experiments

As shown in Table 2, the largest differences in terms of number of assimilated observations are obtained with NOGNSS which leads to a 1.85% difference in the number of assimilated data.

185 **2.5 Validation protocol**

As a first step, the performance of the data assimilation system is validated by comparing the various Analysis (AN) and First-Guess (FG) values against available observations which can be independent from REANA (i.e. not assimilated) or on the contrary assimilated in REANA. One of the key tool used to evaluate the performance of the assimilation system is to examine the FG departure (O-FG) and the AN departure (O-A) in terms of mean and root-mean square (RMS) values, O standing for 190 Observation with the other assimilated observations. Statistics of departures are computed at the observation location.

Those statistics were also computed using few available independent data. The first source comes from the vessel Marfret-Niolon, which was an instrumented commercial ship of opportunity, cruising regularly between the southern France harbour of Marseille and two Algerian harbours (Algiers and Mostagadem). Please refer to Figure 14 of Fourrié et al. (2019) for the 195 trajectories of the vessel during SOP1. Two autonomous systems were installed in order to provide atmospheric and oceanic measurements, in the context of the HyMeX Long Observation Period (LOP). A GNSS antenna was installed at the front on the vessel Marfret-Niolon for the duration of the HyMeX campaign. [An example of the operational measurements which started on January 2012 are provided in Figure ?? with figures ranging from 2.2 m to 2.6 m.](#) The data were post-processed in kinematic Precise Point Positioning with the software provided by Natural Resources Canada (Kouba and Héroux, 2001) and using high-resolution products provided by the International GNSS Service.

200 The second source of independent data comes from wind data obtained from an airborne Doppler cloud-profiler radar named RASTA (Radar Airborne System Tool for Atmosphere (Bouniol et al., 2008; Protat et al., 2009; Delanoë et al., 2013)) that flew 45 days during SOP1. This airborne radar was on board the Falcon 20 research aircraft. It allows the documentation of the microphysical properties and the horizontal components of the wind field in terms of vertical profiles.

205 The operational data assimilation monitoring procedure also provides FG and AN departure statistics for assimilated observations in the experiments, which are described in a companion paper (Fourrié et al., 2019).

210 In a second step, the forecast (range between +3 to +54 hours) quality is assessed in terms of surface parameters and precipitation scores. The surface parameters (temperature and relative humidity at 2 m and wind at 10 m) come from the HyMeX database which provides surface synoptic observations available over the AROME-WMED domain, together with additional hourly observations from Météo-France, AEMET and MeteoCat mesoscale networks. Some of these observations were assimilated to produce surface analyses. For the evaluation of the precipitation quality, the dense surface data set rain gauge network available in the HyMeX data base (V4 version, DOI:10.6096/MISTRALS-HyMeX.904) has been used. Scores

of 3 hourly accumulated precipitation from all analysis times on a given day are compared to the corresponding observed 24-h accumulated precipitation.

The evaluation of the various denial experiments is compared with the reference REANA run. This allows to get the impact 215 of each considered observation type on the analysis and the forecast.


### 3 Impact of GNSS data on the analysis and first-guess quality

This section investigates the impact of assimilating the ground-based GNSS ZTD data on the numerical weather prediction model analysis and subsequent forecast quality. This data set represents the largest one in terms of the number of studied 220 observation types, even though in the end it represents only a small fraction of assimilated data (1.85%) in the analyses (Table 1). As seen in Table 1, satellite data are the most numerous, followed by surface stations data, radar data from the French network, aircraft data and radiosondes ones. Even if surface data provide information only for one level, the network is very dense over France and was reinforced in other countries like Spain or Italy. The other observation types provide information at different levels all along the vertical.

#### 3.1 Impact on moisture field

225 Comparison to the Integrated Water Vapour (IWV) from the reprocessed GNSS observations (not independent from REANA as the information from this data set is assimilated in this experiment) indicates that the best correlation, as expected, is obtained for REANA (around 0.99), the second one being OPERGNSS (around 0.975) and the last one NOGNSS (around 0.96), as shown in Figure 2. This result is confirmed when computing the RMSE of the differences (Figure 2, right panel). A weak diurnal cycle of the scores is noticed with a maximum correlation around 09 UTC and a minimum around 15 UTC. Concerning 230 the standard deviation of the differences, they are lower during the 3-9 UTC period and larger in the afternoon. These minimum of correlation and maximum of standard deviation correspond to the time of the early stage of the convection, which is less easy to predict and introduces inhomogeneities in the water vapour distribution due to precipitation processes (condensation, evaporation, cold pool formation, etc.).

We then discuss the result of the statistics for the analysis and first-guess against radiosonde observations, which represents 235 a reference data set in data assimilation (Figure 3). First of all, as expected, the AN RMS difference (solid lines) are smaller than the FG difference (dashed lines) for the three simulations showing the expected behaviour of the minimisation during the assimilation process, with differences between the OSEs (Figure 3). No impact can be seen on the AN RMS differences. The absence of impact can be explained by the fact that radiosondes are reference observations for assimilation and all the analyses are very constrained by these observations. However, a small positive impact is present on the FG RMS difference 3-h later. 240 The lowest differences are obtained with REANA-FG and the largest ones with NOGNSS-FG. OPER-GNSS-FG differences are close to REANA-FG but slightly larger, showing on the one hand that the assimilation of GNSS data is beneficial (OPER-GNSS data set or REPROC-GNSS one) and on the other hand that the reprocessing of the data brings a small improvement in the comparison of FG with humidity of radiosondes. This shows that the modifications in the analysis brought by the GNSS at



**Figure 2.** Correlation (left panel) and standard deviations (right panel, in  $\text{kg}/\text{m}^2$ ) of integrated water vapour (IWV) content from reprocessed GNSS observations (Bock et al., 2016) and AROME-WMED analyses (REANA in black NOGNSS in red and OPERGNSS in blue) as a function of analysis slot (hours).


other places than radiosondes ones are beneficial and kept during the 3-h forecast. The largest improvement of the assimilation of GNSS data is found between 600 and 850 hPa, NOGNSS-FG being larger than REANA-FG and OPER-GNSS-FG in this part of the atmosphere. In addition, OPER-GNSS-FG is larger than REANA-FG between 700 and 850 hPa showing a slight benefit of assimilating REPROC-GNSS instead of OPER-GNSS.

The various AN mean departures are very close to each other, with slight negative values in the lower and mid troposphere (analysis too moist), as displayed in Figure 3 right panel. Mean FG departures are larger and homothetic, with stronger values for the REANA-FG, being the signature of a weak moist bias in the corresponding analysis for the lower troposphere. The less biased first guess is the one from the NOGNSS experiment.


In addition, AN and FG were compared with radiances from SEVIRI imager, on board the geostationary satellite Meteosat Second Generation, MSG, which is sensitive to moisture (channels WV  $6.2 \mu\text{m}$  for upper-troposphere and  $7.3 \mu\text{m}$  for mid-troposphere). This imager indeed represents another important source of humidity information, especially over oceans where no information from GNSS nor radiosondes is available. These humidity channel data are assimilated for each AROME-WMED analyses. The study of the FG and AN statistics for the REANA, OPERGNSS and NO GNSS shows no impact between the various experiments for these observations (not, shown, Table in supplement file).

The correlation between the various AROME-WMED ZTD AN and corresponding independent (not assimilated) Marfret-Niolon observations is higher for REANA than for NOGNSS and even for OPERGNSS (Figure 4). There is a correlation maximum around 09 UTC, and a minimum around 15 UTC. The mean ZTD is quite similar in all experiments, with a maximum at 09 UTC and a minimum around 00 UTC. A moist bias is found in all simulations when compared to the mean observation in grey shown in Figure 4. The magnitude of this relative positive (moist) bias is around 0.5 percent. Table 3 shows the mean correlation of REANA, NOGNSS and OPERGNSS AN and FG with Marfret-Niolon observations. The higher correlation is obtained with REANA for both AN and FG. When comparing the mean value of ZTD at the Marfret-Niolon places, the closest

265 value to the observed one is obtained with REANA, even if a small moist bias is observed (0.9 mm). This bias is larger for NOGNSS (3.6 mm) and OPERGNSS (4.8 mm). **This could be explained by the fact that there few assimilated observations over the sea which results in a more biased model.** Although the sample size of Marfret-Niolon data set is rather small (around 1000 collocations), this is an original result and makes clear that the REANA experiment produces the best reanalysis, and the best 3-hour forecasts.




**Figure 3.** Root mean square differences (left hand side panel) and mean (right hand side panel) for Analysis (AN solid lines) and First-Guess (FG, dashed lines) departures against assimilated radio-sounding observations for mixing ratio (g/kg); REANA in black, NOGNSS in red and OPERGNSS in blue.)



**Figure 4.** Correlation of the differences between zenithal total delays (ZTD) between REANA (black), NOGNSS (red), OPERGNSS (blue) analyses and corresponding Marfret-Niolon independent observations as a function of analysis time in the left panel; mean value of ZTD (m) in the right panel, the grey line corresponding to observations.

Experiments	Mean ZTD (m)	AN ZTD Correlation	FG ZTD Correlation
REANA	2.4617	0.967	0.961
NOGNSS	2.4642	0.961	0.958
OPERGNSS	2.4654	0.962	0.958
OBS	2.4606		

**Table 3.** Mean zenithal total delays (ZTD) for REANA, NOGNSS and OPERGNSS analyses and Marfret-Niolon observations (first column), correlation of ZTD between REANA, NOGNSS and OPERGNSS analyses and corresponding Marfret-Niolon observations computed over the 8 analysis slots and correlation between ZTD forecasted by AROME-WMED at the 3-hour forecast range and observations from Marfret-Niolon.



**Figure 5.** Location of RASTA observations during the HyMeX Special Observation Period 1. Coloured dots represent the number of wind data available per profile.

### 270 3.2 Impact on wind field

Analysis and First-guess quality has been evaluated against RASTA (Radar Airborne System Tool for Atmosphere) Doppler winds (Borderies et al., 2019a). This airborne radar was on board Falcon 20 aircraft and provided 33083 wind observations over the Mediterranean area as illustrated in Figure 5, where only few wind data from conventional observations are available. **It is worth reminding** that the data from this instrument were not assimilated in REANA. This data set thus represents an additional 275 independent information for the evaluation of our denial experiments.

Table 4 provides the RMS errors (RMSE) for wind calculated with these data. The RMSE for background and analysis are lower in REANA than in the other two experiments. The analysis RMSE for OPERGNSS is lower than the one for NOGNSS.


As GNSS observations do not provide any wind information, the improvement observed in wind field can be explained by the effect of mass field information assimilation on the wind field, essentially created during model integration. There is 280 indeed a little coupling between these fields during the analysis (Borderies et al., 2019b). This indirect effect was already

demonstrated by Wattrelot et al. (2014), for example, who noted a positive impact on the wind field when assimilating pseudo-observations of relative humidity. Lindskog et al. (2017) also reported—but did not show—a positive impact on wind forecasts when assimilating ZTD data.

Experiment	AN RMSE	FG RMSE
REANA	5.59	5.87
NOGNSS	5.63	5.97
OPERGNSS	5.60	5.92

**Table 4.** Analysis (AN) and First Guess (FG) Root Mean Square Errors (RMSE) for the wind (in m/s) computed with respect to RASTA observations (sample size 33083 observations) for REANA, NOGNSS and OPER GNSS experiments.

### 3.3 Impact on short-range precipitation



**Figure 6.** Equitable Threat Score (ETS) for the 24-h accumulated precipitation from the sum of the eight 3-h forecasts used as background of the data assimilation cycle each day of the period from 5 September to 5 November 2012. Results for REANA are displayed in black, for NOGNSS in red and OPERGNSS in blue. Dots indicate that the difference between the curves and the REANA curve as a reference is statistically significant at a 0.95 confidence threshold using a Bootstrap test.

285 Figure 6 shows that the Equitable Threat Score (ETS) of the 24-h accumulated precipitation computed with the sum of the 3-h precipitation from the 8 analysis times is improved with the assimilation of GNSS ZTD data compared to the NOGNSS experiment. It represents an evaluation of the background quality. The difference is statistically significant for each threshold. When comparing the assimilation of REANA to OPERGNSS, the ETS for precipitation is slightly better with the reprocessed data set but the differences are not significant except for the 40 mm/day threshold. [The loss of skill for OPERGNSS simulation](#)


Parameter	REANA	NOGNSS	OPERGNSS
Correlation (1-24h)	0,962	0,957	0,957
Correlation (25-48h)	0,922	0,917	0,919
Correlation (49-54h)	0,906	0,902	0,905
Standard deviation (forecast -observation, 1-24h)	0,0152	0,0164	0,0160
Standard deviation (forecast -observation, 25-48h)	0,0221	0,0226	0,0223
Standard deviation (forecast -observation, 49-54h)	0,0244	0,0249	0,0244

**Table 5.** Correlation and standard deviation of ZTD (in m) between AROME-WMED forecasts and reprocessed GNSS observations averaged over forecast ranges.

290 for larger daily precipitation from the 8 3-h forecast is surely due to the lower number of occurrences for heavier precipitations. Overall, the background quality is improved with the assimilation of GNSS observations and the data reprocessing brings improvement in terms of precipitation from 3-hour forecast even though this benefit is not significant.

#### 4 Impact of GNSS data on medium term forecast


295 The impact of the GNSS data has also been assessed for longer forecast ranges (3 to 54-h). The effect of the assimilation of the GNSS data on the correlation with IWV from the GNSS data set is maximal for the analysis and decreases up to the 30-h forecast range (Figure 7) as the general impact of the initial conditions on the forecast performances reduces. A similar behaviour is found with the standard deviations of the differences between observed **IWV** and simulated one from the three experiments.



**Figure 7.** Correlation (left hand side panel) and standard deviations (right hand side panel,  $\text{kg}/\text{m}^2$ ) of integrated water vapour content between AROME-WMED forecasts and reprocessed GNSS observations (Bock et al., 2016) as a function of forecast range (hours).

Compared to the observed ZTD from the Marfret-Niolon ship, the signal is more noisy because of a smaller dataset but 300 when compared to values averaged over the forecast ranges (Table 5), the correlation for the NOGNSS is lower than REANA and OPERGNSS, which provides itself lower correlation than REANA. The standard deviations are higher for the NOGNSS forecasts. In addition, a decrease of the correlation (respectively an increase of the standard deviation) is seen for forecast range over 24-h.

The forecast quality has also been evaluated against surface data. No impact was found on temperature at 2 m or on 10 m 305 wind (not shown). A small impact was found on relative humidity at 2 m (Figure 8). A reduction of the bias is noticed with REANA during the first 9-h of the forecast compared to OPER GNSS and NOGNSS. From 12-h onwards the results for REANA and OPERGNSS are similar. Regarding the standard deviation, it is smaller for REANA between 0 and 9-h than for NOGNSS and OPER GNSS and between 21 and 27-h forecast range than for NOGNSS. This difference represents more than 2 % of improvement. For the other forecast ranges the differences are lower than 1%.




**Figure 8.** Bias (forecast minus observations, dashed lines) and standard deviations (solid lines) computed with relative humidity at 2 m (left hand side panel) and relative root mean square differences (% right hand side panel) with respect to REANA.


310 The impact of the assimilation of GNSS data on the 24-h accumulated precipitation from the forecast initialized at 06 UTC  
315 is less clear. The improvement of the GNSS data reprocessing compared to the real time data set is beneficial for all thresholds  
except for the 2mm/day (where the ETS is better for OPERGNSS) and is statistically significant for moderate thresholds (10  
and 20 mm/day, Figure 9). The difference between REANA ETS and NOGNSS ETS values is not significant. When examining  
scores for precipitation forecasts between 30-h and 54-h, there is a small significant degradation of the ETS for the 2 mm/day  
with the NOGNSS experiment and a small improvement with the OPERGNSS for the 40 mm/day (Figure 10).

## 5 Other impact studies

As previously mentioned we performed other impact studies with wind profilers, lidar data and Spanish radar data.




**Figure 9.** Equitable Threat Score of the 24-h accumulated precipitation from the 6-30 hour forecast range of the long forecast initialized at 00 UTC each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). Dots indicate that the difference between the curves is statistically significant.



**Figure 10.** Equitable Threat Score of 24 h accumulated precipitation from the 30 to 54 hour forecast range of the long forecast initialized at 00 UTC each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). Dots indicate that the difference between the curves is statistically significant.

## 5.1 Wind profilers

No impact of the assimilation of wind profiler data is found except on wind field. Small impact is noticed in terms of wind  
320 RMS differences of background and analysis departures for radiosondes, aircraft and satellite winds (Figure 11). The largest impact is a decrease of -0.08 m/s for the radiosonde FG RMS differences at 300 hPa ([left panel](#)). Concerning the AN RMS differences, the improvement (SATOBI) or degradation (AIREP and TEMP) are very small. The largest value obtained at 200 hPa are due to the small number of data available for the computation.




**Figure 11.** First-Guess (left plot) and analysis (right plot) RMS differences (REANA-NOWPROF experiments) computed against TEMP (black), AIREP (red) and SATOB (green) observations for the zonal wind component (m/s); negative value correspond to a positive impact of wind profiler.

A small improvement of REANA compared to NOWPROF, but not significant (Figure 12), appears on the ETS of the 24 h  
325 accumulated precipitation accumulated from the 6 to 30 hour forecast ranges.

## 5.2 Ground-based and airborne lidar data

As discussed in Section 2.2, humidity profiles retrieved from ground-based and air-borne lidars have been assimilated in the REANA experiment. In Figure 1, the trajectories of all ATR-42 flights are plotted, together with the localization of the two ground-based lidars. The denial NOLIDAR experiment results are very close to the reanalysis ones (Table 6) as these data  
330 represent very few additional data located over ocean where few observations are available for the comparison. No impact of the Lidar data is found when comparing the various analyzed ZTD to the Marfret-Nilon corresponding observations. These results agree with the Bielli et al. (2012) study where no significant impact where found on the 24-h accumualted precipitation.




**Figure 12.** Equitable Threat Score of 24 h accumulated precipitation from the 6 to 30 hour forecast ranges of the long forecast starting at 00 UTC each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). The lack of dots indicates that the difference between the curves is not significant. REANA is plotted in black and NOWPROF in red.

	REANA	NOLIDAR
Correlation for the forecast 0h	0.968	0.960
Correlation for the forecast 1-24 h	0.962	0.961
Correlation for the forecast 25-48 h	0.923	0.924
Correlation for the forecast 49-54 h	0.906	0.907
Standard deviation for the forecast 0h	0.0144	0.0167
Standard deviation for the forecast 1-24 h	0.0152	0.0154
Standard deviation for the forecast 25-48 h	0.0221	0.0220
Standard deviation for the forecast 49-54 h	0.0243	0.0243

**Table 6.** Correlation and standard deviation of Zenithal Total Delays (in m) between AROME-WMED forecasts from 00UTC and reprocessed GNSS observations averaged over forecast ranges (0 h, 1-24 h, 25-48 h and 49-54 h).

### 5.3 Spanish radars


No significant impact has been noticed over the HyMeX domain. However, when focusing over the Iberian Peninsula, the ETS 335 for the 24-h accumulated precipitation from the sum of the 8 3-h precipitation background forecast (Figure 13) is improved with the assimilation of Spanish radar data. This impact also remains in longer forecast ranges as the ETS for the 24-h precipitation accumulation between 6-h and 30-h forecast ranges is improved with the assimilation of Spanish radars for thresholds between 0.5 and 20 mm/24h (Figure 14). This impact does not remain at longer forecast ranges (Figure 15). These results are in good 340 agreement with Wattrelot et al. (2014) study which found an improvement of the short term precipitation forecast scores. However contrary to the aforementioned study, we obtained a significant improved of the 24-h precipitation accumulation 345 between 6-h and 30-h forecast ranges over the Iberian Peninsula. Even if we do not obtain significant impact at the HyMeX domain scale but a significant one over the Iberian Peninsula, it is interesting to remind that the assimilation of Spanish radar data in AROME-WMED was made on a research mode as only French radars were assimilated at the time of the HyMeX campaign and the reanalyses. These data represent only 0.6% of the assimilated data. This is three times less than REPROC-GNSS data.




**Figure 13.** Equitable Threat Score (ETS) for the 24-h accumulated precipitation obtained from the sum of eight 3-h forecasts used as background of the data assimilation cycle each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). Results for REANA are displayed in black, for NORADSPAIN in red. Dots indicate that the difference between the curves is statistically significant.

## 6 IOP16 case study

During HyMeX SOP1, IOP16a was dedicated to HPE that occurred over Cévennes-Vivarais (CV) in France and later on, in Italy (IOP16b) on 25-26 October; this event was associated with locally flash-flooding and several casualties. This off-shore



**Figure 14.** Equitable Threat Score (ETS) for the 24-h accumulated precipitation from the 6 to 30 hour forecast ranges initialized at 00 UTC each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). Results for REANA are displayed in black, for NORADSPAIN in red. Dots indicate that the difference between the curves is statistically significant.



**Figure 15.** Equitable Threat Score (ETS) for the 24-h accumulated precipitation from the 30 to 54 hour forecast ranges initialized at 00 UTC each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). Results for REANA are displayed in black, for NORADSPAIN in red. Dots indicate that the difference between the curves is statistically significant.


convection case is well documented in Duffourg et al. (2016). On the 26th - 00 UTC active convection was occurring over  
350 Catalonia; this area of intense convective activity crossed the Gulf of Lion reaching the French Mediterranean coast around 06 UTC and later on, and then hitting the Italian Ligurian coast in the evening. It is well known that the associated convective systems are usually fed with moisture, during their early stage over the warm Mediterranean Sea. A moist conditionally unstable south-western flux is therefore found in the lower troposphere (Figure 16) with a low-level jet by the Candillargues radar around 09-12 UTC, associated to a slow evolving weak pressure low (around 995 hPa) localized over southern France  
355 on the 26th mid-day. Moreover, low level convergence is reinforced by the complex orography (Cévennes ridge of the Massif-Central and Alps in France) triggering convection. An upper south-westerly wind jet is observed above 500 hPa (Figure 16); in the evening of the 25th the wind rotates to the west on the CV area as shown by the Candillargues UHF radar.

During 25th and 26th October period, many deep convective systems developed over the Northwestern Mediterranean (Duffourg et al., 2016). Although observed accumulated surface precipitation from Friday 26th at 18 UTC to Saturday Oct. 27th  
360 at 06 UTC over southern France only reached around 150 mm in 24h (Martinet et al., 2017), very strong hourly rates (near 50 mm/1h) were recorded, with intense river discharges (Ardèche, Gardons and Gapeau rivers for example). Such intense rainfall amounts led to local flash-floods and 2 casualties in the Var region. In fact as shown in Figure 17, three local precipitation maxima appear on the observed 24-hour accumulated rainfall amount (25th October - 06 UTC to 26th October - 06 UTC) on the Mediterranean costal area of France and Italy (Liguria Tuscany region); a first elongated one in the Cévennes area (more  
365 than 150 mm, M1) and a small second one close to the coast (around 100 mm, M2).

Figure 18 shows the 24-h accumulated precipitation between the 6-h and 30-h forecasts for the different experiments considered in this study. The REANA 24-hour accumulated rainfall (from 06 h to 30 h forecast range) simulation agrees to the observations for both M1 and M2 systems. The NOLIDAR experiment is very close to REANA, this is consistent with the fact that the amount of additional lidar data is fairly small in REANA when compared to NOLIDAR. The strongest impact is found  
370 when no GNSS data are assimilated (NOGNSS run): M1 and M2 are strongly underestimated; surprisingly the OPERGNSS experiment leads to an accurate forecast of M2, but underestimates the southwestward extension of M1. Finally a strong negative impact is found with the NOWPROF simulation which misses M2 and does not reproduce correctly M1. Over Italy, the gain brought by the observations is not so evident but it is quite well known in data impact studies that the assimilation of observation does not always improve the forecast at each analysis time but in overall.

## 375 7 Conclusions


The AROME-WMED model was originally developed to study and forecast heavy-precipitating Mediterranean events during the Special Observation Periods (SOPs) of the HyMeX programme. Two reanalyses were produced after the HyMeX autumn campaign for the first SOP. The first one was carried out after the field campaign to provide reanalyses simulations of the period enjoying an updated AROMEWMED version. The second one, performed a few years after, was done to improve  
380 the quality of the first reanalyses, assimilating as many data as possible from the experimental field campaign (i.e. lidar and




**Figure 16.** Time-height cross-section of the wind measured by the Candillargues UHF radar for IOP16. Horizontal wind components are represented by the black arrows (meteorological convention), and wind speed in colour.

dropsonde humidity profiles) or from reprocessed data (such as GNSS-ZTD, wind profiles, high-vertical resolution radiosondes and Spanish doppler radars) using the latest AROME version.

Previous studies such as Duffourg and Ducrocq (2011) Ricard et al. (2012) or Bresson et al. (2012), have shown the importance of an accurate description of the low-level moist flow feeding mesoscale convective systems. In this study the impact of 385 various data set related to humidity and wind on the forecast quality from this comprehensive reanalysis is investigated over the 2-month period. Many data sets of the Special Observation Period 1 of the HyMeX campaign have been considered here. The reprocessed GNSS data set (Bock et al. (2016)) were removed and replaced with the operational data set used in the real-time AROME-WMED version. We examined the humidity data provided by ground based and airborne lidars. The impact of the 390 reprocessed wind profilers and the Spanish radar data was also evaluated. The impact of these data sets was assessed through Observing system Experiments which consist of removing the data sets and to compare forecast quality from these denial experiments to a reference which includes all data sets. The selected data sets were research observations (water vapour lidars)



**Figure 17.** 24 h accumulated precipitation (mm) between 26 October 06 UTC and 27 October 2012 at 06 UTC over the AROME-WMED domain (upper plot) and zoom over the Cevennes region (left lower plot) and over North of Italy (right lower plot).



**Figure 18.** Same as Figure 20 but for 24h accumulated precipitation forecasted by (a) REANA, (b) NOGNSS, (c) NOLIDAR (d) OPERGNSS and (e) NOWPROF experiments, from the 06 h to 30 h forecast ranges.

or reprocessed data (from ground based GNSS receivers or wind profilers). In total, the amount of denied data ranged between 0.15 % and 1.86 % of the total number of observations, depending on the OSE and the impact was thus expected to be small. Our study finds a small positive impact. The selected data sets were research observations (water vapour lidars) or reprocessed data (from ground based GNSS receivers or wind profilers). They represented a modest part of the assimilated data amounts and their impact was thus expected to be small.

Our study finds a small positive impact on humidity forecast at short term ranges of the reprocessed GNSS ground based zenithal total delay assimilation compared with NOGNSS. This data set is evenly distributed over the AROME-WMED domain and provided at each analysis time information on integrated water vapour. The impact of the data reprocessing for GNSS zenithal total delays compared to the processing made in real-time was also studied by comparing REANA with OPERGNSS. Even though a positive impact is observed with REANA compared to OPERGNSS, this improvement is not statistically significant. Given the impact of ground-based GNSS, there is also an interest in continuing work to assimilate GNSS data over ocean surfaces.

Small impacts on wind fields were also observed for wind profilers. The sole impact of the assimilation of lidar data is found when comparing with independent airborne radar observations located over the Mediterranean Sea. Since this data

set represents a very small fraction of assimilated data, this may explain the absence of impact. In addition they were not assimilated at their full available temporal frequency but just once every 3 hours.

Spanish radar data assimilation improves the short term quality of the background as noticed on the 24-h accumulated precipitation of the eight 3-h background forecasts for each day but only over the [Iberian](#) Peninsula with no clear impact over  
410 the HyMeX domain. It is interesting to stress that this impact remains during the first 30-h of the forecast but without any remote impact over the rest of the AROME-WMED domain. More impact could possibly be obtained if the data were provided with additional scan elevations.

With the examination of the impact of the assimilation of 4 different data sets over a two-month period in the meso-scale AROME-WMED, our study shows that it is required to have well spatially distributed and frequent data sets such as the GNSS  
415 ZTD data set to get, with its assimilation, an overall impact in terms of analysis and forecast skills. This result agrees with the findings of Mahfouf et al. (2015) who show that the assimilation of GNSS systematically improves the atmospheric humidity short-range forecasts despite the small fraction of GNSS observations assimilated in AROME. A high temporal availability and a regular horizontal distribution are both needed to get a significant impact on the forecast scores. Moreover, it is interesting to process as precisely as possible a maximum of GNSS data in real time and to have bias-corrected observations valuable for  
420 data assimilation. In addition, GNSS data available on ship seems to be promising to increase the coverage over ocean (Fan et al., 2016). When the data set is available frequently but not well spread over the model domain such as the Doppler winds and reflectivities from the Spanish radars or winds from profiler radars, its assimilation may lead to a positive impact on the precipitation forecast but it remains local. Finally, marginal impact from local and sporadic data sets such as humidity profiles from water-vapour Lidars can be obtained but it is not visible on "global scores". To get a material impact on the forecast in  
425 a mesoscale model from a set of observation through its data assimilation, our study suggests to select data sets which are frequently available at each analysis time and also well spread over the domain.

The impact of the above mentioned data could be further improved. For example, the impact of GNSS in AROME-France has been recently improved with the use of variational bias correction in replacement of the static bias correction used in this  
430 study (P Moll, personal Communication). In addition radar data from foreign countries are now assimilated in AROME since July 2020. The distribution of these data by OPERA (the EUMETNET Radar programme) allows to get data of high quality in the data assimilation and thus to increase their impact in the AROME model Martet et al. (2019). With a more frequent data assimilation cycle, making use of observations at higher temporal frequency as it is the case with the current AROME-France model (Brousseau et al., 2016), it is likely that surface observations or the remote sensing data such as radars, GNSS or SEVIRI available for each hourly analysis in this study would have a greater impact on analyses and forecasts. Moreover new  
435 data assimilation systems such as the 4D-Var or the 4D-Envar are under development for convective scale models (Gustafsson et al., 2018) and will allow to account for very frequent data. Therefore, they are expected to enhance the impact of observations available several times an hour. In the future, the impact of the Infra-Red Sounder onboard Meteosat Third Generation will benefit from these new data assimilation systems as this sounder will provide observations every 30 minutes over the AROME domain and especially over the oceans.

440 *Code availability.* The source code of AROME-WMED, derived from the operational AROME code cannot be obtained.

*Data availability.* The analyses and the forecast fields are available in the HyMeX database (<http://mistral.sedoo.fr/HyMeX/>, last access 19 August 2019). The final (second) reanalysis labelled REANA in this paper is available at <https://doi.org/10.14768/MISTRALS-HYMEX.1492> (Fourrié and Nuret, 2017).

445 *Author contributions.* NF and MN prepared and carried out all the numerical experiments of the reanalysis and the OSEs. They investigated the results, and wrote the paper with the help of all the coauthors. PBr and OC helped to investigate the results by performing diagnostics and verification computations.

*Competing interests.* The authors declare that they have no conflict of interest.

450 *Acknowledgements.* The authors would like to acknowledge the MISTRALS/HyMeX program and the funding from the ANR under contracts IODA-MED ANR-11-BS56-0005 and MUSIC ANR-14-CE01-0014. Jean-François Mahfouf and Véronique Ducrocq are acknowledged for their helpful comment on a previous version of the paper. The authors also thank the anonymous reviewers for their fruitful comments. The authors acknowledge principal investigators of data sets : Dr Patrick Chazette (WALI), Dr Cyrille Flamant (LEANDRE), Dr Paolo Di Girolamo (BASIL), Dr Frédérique Saïd (wind profilers), Dr Olivier Bock, Pierre Bosser (reprocessed GNSS data) and Dr Isabelle Taupier-Letage (data from opportunity ships).

## References

455 Benjamin, S. G., Schwartz, B. E., Szoke, E. J., and Koch, S. E.: The Value of Wind Profiler Data in U.S. Weather Forecasting, Bulletin of the American Meteorological Society, 85, 1871–1886, <https://doi.org/10.1175/BAMS-85-12-1871>, <https://doi.org/10.1175/BAMS-85-12-1871>, 2004.

Berre, L.: Estimation of Synoptic and Mesoscale Forecast Error Covariances in a Limited-Area Model, Monthly Weather Review, 128, 644–667, [https://doi.org/10.1175/1520-0493\(2000\)128<0644:EOSAMF>2.0.CO;2](https://doi.org/10.1175/1520-0493(2000)128<0644:EOSAMF>2.0.CO;2), 2000.

460 Bielli, S., Grzeschik, M., Richard, E., Flamant, C., Champollion, C., Kiemle, C., Dorninger, M., and Brousseau, P.: Assimilation of water-vapour airborne lidar observations: impact study on the COPS precipitation forecasts, Quarterly Journal of the Royal Meteorological Society, 138, 1652–1667, <https://doi.org/10.1002/qj.1864>, 2012.

Bock, O., Bosser, P., Pacione, R., Nuret, M., Fourrié, N., and Parracho, A.: A high-quality reprocessed ground-based GPS dataset for atmospheric process studies, radiosonde and model evaluation, and reanalysis of HyMeX Special Observing Period, Quart. J. Roy. Meteorol. Soc., 142, 56–71, <https://doi.org/10.1002/qj.2701>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.2701>, 2016.

465 Boniface, K., Ducrocq, V., Jaubert, G., Yan, X., Brousseau, P., Masson, F., Champollion, C., Chéry, J., and Doerflinger, E.: Impact of high-resolution data assimilation of GPS zenith delay on Mediterranean heavy rainfall forecasting, Annales Geophysicae, 27, 2739–2753, <https://doi.org/10.5194/angeo-27-2739-2009>, <https://www.ann-geophys.net/27/2739/2009/>, 2009.

Borderies, M., Caumont, O., Delanoë, J., Ducrocq, V., and Fourrié, N.: Assimilation of wind data from airborne Doppler cloud-profiling radar 470 in a kilometre-scale NWP system, Natural Hazards and Earth System Sciences, 19, 821–835, <https://doi.org/10.5194/nhess-19-821-2019>, <https://www.nat-hazards-earth-syst-sci.net/19/821/2019/>, 2019a.

Borderies, M., Caumont, O., Delanoë, J., Ducrocq, V., Fourrié, N., and Marquet, P.: Impact of airborne cloud radar reflectivity data assimilation on kilometre-scale numerical weather prediction analyses and forecasts of heavy precipitation events, Natural Hazards and Earth System Sciences, 19, 907–926, <https://doi.org/10.5194/nhess-19-907-2019>, <https://nhess.copernicus.org/articles/19/907/2019/>, 2019b.

475 Bouniol, D., Protat, A., Plana-Fattori, A., Giraud, M., Vinson, J.-P., and Grand, N.: Comparison of Airborne and Spaceborne 95-GHz Radar Reflectivities and Evaluation of Multiple Scattering Effects in Spaceborne Measurements, Journal of Atmospheric and Oceanic Technology, 25, 1983–1995, <https://doi.org/10.1175/2008JTECHA1011.1>, <https://doi.org/10.1175/2008JTECHA1011.1>, 2008.

Bresson, E., Ducrocq, V., Nuissier, O., Ricard, D., and de Saint-Aubin, C.: Idealized numerical simulations of quasi-stationary convective systems over the Northwestern Mediterranean complex terrain, Q.J.R. Meteorol. Soc., 138, 1751–1763, 2012.

480 Brousseau, P., Berre, L., Bouttier, F., and Desroziers, G.: Background error covariances for a convective-scale data-assimilation system : AROME-France 3D-Var, Quarterly Journal of the Royal Meteorological Society, 137, 409–422, <https://doi.org/10.1002/qj.750>, 2011.

Brousseau, P., Seity, Y., Ricard, D., and Léger, J.: Improvement of the forecast of convective activity from the AROME-France system, Quarterly Journal of the Royal Meteorological Society, 142, 2231–2243, <https://doi.org/10.1002/qj.2822>, 2016.

Caniaux, G., Redelsperger, J.-L., and Lafore, J.-P.: A numerical study of the stratiform region of a fast-moving squall line. Part I: General 485 description and water and heat budgets, Journal of the Atmospheric Sciences, 51(14), 2046–2074, 1994.

Caumont, O. and Ducrocq, V.: What should be considered when simulating Doppler velocities measured by ground-based weather radars?, J. Applied Meteor. Climatol., 47, 2256–2262, <https://doi.org/10.1175/2008JAMC1894.1>, 2008.

Caumont, O., Ducrocq, V., Wattrelot, E., Jaubert, G., and Pradier-Vabre, S.: 1D+3DVar assimilation of radar reflectivity data: a proof of concept, Tellus A: Dynamic Meteorology and Oceanography, 62, 173–187, <https://doi.org/10.1111/j.1600-0870.2009.00430.x>, <https://doi.org/10.1111/j.1600-0870.2009.00430.x>, 2010.

Chazette, P., Flamant, C., Shang, X., Totems, J., Raut, J.-C., Doerenbecher, A., Ducrocq, V., Fourrié, N., Bock, O., and Cloché, S.: A multi-instrument and multi-model assessment of atmospheric moisture variability over the western Mediterranean during HyMeX, Quarterly Journal of the Royal Meteorological Society, 142, 7–22, <https://doi.org/10.1002/qj.2671>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.2671>, 2016.

495 Courtier, P., Freydier, C., Rabier, F., and Rochas, M.: The ARPEGE Project at Météo-France, ECMWF Seminar Proceedings, 7, 193–231, 1991.

Delanoë, J., Protat, A., Jourdan, O., Pelon, J., Papazzoni, M., Dupuy, R., Gayet, J.-F., and Jouan, C.: Comparison of Airborne In Situ, Airborne Radar–Lidar, and Spaceborne Radar–Lidar Retrievals of Polar Ice Cloud Properties Sampled during the POLARCAT Campaign, Journal of Atmospheric and Oceanic Technology, 30, 57–73, <https://doi.org/10.1175/JTECH-D-11-00200.1>, <https://doi.org/10.1175/JTECH-D-11-00200.1>, 2013.

500 Di Girolamo, P., Flamant, C., Cacciani, M., Richard, E., Ducrocq, V., Summa, D., Stelitano, D., Fourrié, N., and Saïd, F.: Observation of low-level wind reversals in the Gulf of Lion area and their impact on the water vapour variability, Quarterly Journal of the Royal Meteorological Society, 142, 153–172, <https://doi.org/10.1002/qj.2767>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.2767>, 2016.

Drobinski, P., Ducrocq, V., Alpert, P., Anagnostou, E., Béranger, K., Borga, M., Braud, I., Chanzy, A., Davolio, S., Delrieu, G., Estournel, C., 505 Boubrahmi, N. F., Font, J., Grubišić, V., Gualdi, S., Homar, V., Ivančan-Picek, B., Kottmeier, C., Kotroni, V., Lagouvardos, K., Lionello, P., Llasat, M. C., Ludwig, W., Lutoff, C., Mariotti, A., Richard, E., Romero, R., Rotunno, R., Roussot, O., Ruin, I., Somot, S., Taupier-Letage, I., Tintore, J., Uijlenhoet, R., and Wernli, H.: HyMeX: A 10-Year Multidisciplinary Program on the Mediterranean Water Cycle, Bulletin of the American Meteorological Society, 95, 1063–1082, <https://doi.org/10.1175/BAMS-D-12-00242.1>, <https://doi.org/10.1175/BAMS-D-12-00242.1>, 2014.

510 Ducrocq, V., Braud, I., Davolio, S., Ferretti, R., Flamant, C., Jansa, A., Kalthoff, N., Richard, E., Taupier-Letage, I., Ayral, P.-A., Belamari, S., Berne, A., Borga, M., Boudevillain, B., Bock, O., Boichard, J.-L., Bouin, M.-N., Bousquet, O., Bouvier, C., Chiggiato, J., Cimini, D., Corsmeier, U., Coppola, L., Cocquerez, P., Defer, E., Delanoë, J., Girolamo, P. D., Doerenbecher, A., Drobinski, P., Dufournet, Y., Fourrié, N., Gourley, J. J., Labatut, L., Lambert, D., Coz, J. L., Marzano, F. S., Molinié, G., Montani, A., Nord, G., Nuret, M., Ramage, K., Rison, W., Roussot, O., Said, F., Schwarzenboeck, A., Testor, P., Baelen, J. V., Vincendon, B., Aran, M., and Tamayo, J.: HyMeX-SOP1: The 515 Field Campaign Dedicated to Heavy Precipitation and Flash Flooding in the Northwestern Mediterranean, Bulletin of the American Meteorological Society, 95, 1083–1100, <https://doi.org/10.1175/BAMS-D-12-00244.1>, <https://doi.org/10.1175/BAMS-D-12-00244.1>, 2014.

Duffourg, F. and Ducrocq, V.: Origin of the moisture feeding the Heavy Precipitating Systems over Southeastern France, Nat. Hazards Earth Syst. Sci., 11, 1163–1178, 2011.

Duffourg, F., Nuissier, O., Ducrocq, V., Flamant, C., Chazette, P., Delanoë, J., Doerenbecher, A., Fourrié, N., Di Girolamo, P., Lac, C., Legain, 520 Martinet, M., Saïd, F., and Bock, O.: Offshore deep convection initiation and maintenance during HyMeX IOP16a heavy precipitation event, Quarterly Journal of the Royal Meteorological Society, 142, 259–274, 2016.

Fan, S., Zang, J.-F., Peng, X.-Y., Wu, S.-Q., Liu, Y.-X., and Zhang, K.-F.: Validation of Atmospheric Water Vapor Derived from Ship-Borne GPS Measurements in the Chinese Bohai Sea, Terrestrial, Atmospheric and Oceanic Sciences, 27, 213, [https://doi.org/10.3319/TAO.2015.11.04.01\(A\)](https://doi.org/10.3319/TAO.2015.11.04.01(A)), 2016.

525 Fourrié, N., Bresson, E., Nuret, M. and Jany, C., Brousseau, P., Doerenbecher, A., Kreitz, M., Nuissier, O., Sevault, E., Bénichou, H., Amodei, M., and Pouponneau, F.: AROME-WMED, a real-time mesoscale model designed for the HyMeX special observation periods, Geoscientific Model Development, 8, 1919–1941, <https://doi.org/10.5194/gmd-8-1919-2015>, <https://www.geosci-model-dev.net/8/1919/2015/>, 2015.

Fourrié, N., Nuret, M., Brousseau, P., Caumont, O., Doerenbecher, A., Wattrelot, E., Moll, P., Bénichou, H., Puech, D., Bock, O., Bosser, P., Chazette, P., Flamant, C., Di Girolamo, P., Richard, E., and Saïd, F.: The AROME-WMED reanalyses of the first special observation period of the Hydrological cycle in the Mediterranean experiment (HyMeX), *Geoscientific Model Development*, 12, 2657–2678, <https://doi.org/10.5194/gmd-12-2657-2019>, <https://www.geosci-model-dev.net/12/2657/2019/>, 2019.

530 Grzeschik, M., Bauer, H.-S., Wulfmeyer, V., Engelbart, D., Wandinger, U., Mattis, I., Althausen, D., Engelmann, R., Tesche, M., and Riede, A.: Four-Dimensional Variational Data Analysis of Water Vapor Raman Lidar Data and Their Impact on Mesoscale Forecasts, *Journal of Atmospheric and Oceanic Technology*, 25, 1437–1453, <https://doi.org/10.1175/2007JTECHA974.1>, 2008.

535 Gustafsson, N., Janjić, T., Schraff, C., Leuenberger, D., Weissmann, M., Reich, H., Brousseau, P., Montmerle, T., Wattrelot, E., Bučánek, A., Mile, M., Hamdi, R., Lindskog, M., Barkmeijer, J., Dahlbom, M., Macpherson, B., Ballard, S., Inverarity, G., Carley, J., Alexander, C., Dowell, D., Liu, S., Ikuta, Y., and Fujita, T.: Survey of data assimilation methods for convective-scale numerical weather prediction at operational centres, *Quarterly Journal of the Royal Meteorological Society*, 144, 1218–1256, <https://doi.org/10.1002/qj.3179>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.3179>, 2018.

540 Gutman, S. I., Sahm, S. R., Benjamin, S. G., Schwartz, B. E., Holub, K. L., Stewart, J. Q., and Smith, T. L.: Rapid Retrieval and Assimilation of Ground Based GPS Precipitable Water Observations at the NOAA Forecast Systems Laboratory: Impact on Weather Forecasts, *Journal of the Meteorological Society of Japan. Ser. II*, 82, 351–360, <https://doi.org/10.2151/jmsj.2004.351>, 2004.

545 Kouba, J. and Héroux, P.: Precise Point Positioning Using IGS Orbit and Clock Products, *GPS Solutions*, <https://doi.org/https://doi.org/10.1007/PL00012883>, 2001.

Lindskog, M., Ridal, M., Thorsteinsson, S., and Ning, T.: Data assimilation of GNSS zenith total delays from a Nordic processing centre, *Atmospheric Chemistry and Physics*, 17, 13 983–13 998, <https://doi.org/10.5194/acp-17-13983-2017>, <https://acp.copernicus.org/articles/17/13983/2017/>, 2017.

550 Macpherson, S. R., Deblonde, G., Aparicio, J. M., and Casati, B.: Impact of NOAA Ground-Based GPS Observations on the Canadian Regional Analysis and Forecast System, *Monthly Weather Review*, 136, 2727–2746, <https://doi.org/10.1175/2007MWR2263.1>, <https://doi.org/10.1175/2007MWR2263.1>, 2008.

Mahfouf, J.-F., Ahmed, F., Moll, P., and Teferle, F. N.: Assimilation of zenith total delays in the AROME France convective scale model: a recent assessment, *Tellus A: Dynamic Meteorology and Oceanography*, 67, 26 106, <https://doi.org/10.3402/tellusa.v67.26106>, <https://doi.org/10.3402/tellusa.v67.26106>, 2015.

555 Martet, M., Guillaume, F., and Wattrelot, E.: Assimilation of OPERA radar data in AROME-FRANCE NWP MODEL: a challenge, 39th international conference on radar meterology, 16-20 September 2019,Nara Kasugano International Forum IRAKA, Nara, Japan., [https://cscenter.co.jp/icrm2019/program/data/abstracts/Session3B-02\\_2.pdf](https://cscenter.co.jp/icrm2019/program/data/abstracts/Session3B-02_2.pdf), lastaccesson19August2020., 2019.

Martinet, M., Nuissier, O., Duffourg, F., Ducrocq, V., and Ricard, D.: Fine-scale numerical analysis of the sensitivity of the HyMeX IOP16a heavy precipitating event to the turbulent mixing length parameterization, *Quarterly Journal of the Royal Meteorological Society*, 143, 3122–3135, 2017.

560 Montmerle, T. and Faccani, C.: Mesoscale Assimilation of Radial Velocities from Doppler Radars in a Preoperational Framework, *Monthly Weather Review*, 137, 1939–1953, <https://doi.org/10.1175/2008MWR2725.1>, <https://doi.org/10.1175/2008MWR2725.1>, 2009.

Pinty, J.-P. and Jabouille, P.: A mixed-phased cloud parameterization for use in a mesoscale non-hydrostatic model: simulations of a squall line and of orographic precipitation. *Proceeding, Conference on Cloud Physics*, Everett, WA., Amer. Meteor. Soc., pp. 217–220, 1998.

565 Protat, A., Bouniol, D., Delanoë, J., O'Connor, E., May, P. T., Plana-Fattori, A., Hasson, A., Görsdorf, U., and Heymsfield, A. J.: Assessment of Cloudsat Reflectivity Measurements and Ice Cloud Properties Using Ground-Based and Airborne Cloud Radar Observations,

Journal of Atmospheric and Oceanic Technology, 26, 1717–1741, <https://doi.org/10.1175/2009JTECHA1246.1>, <https://doi.org/10.1175/2009JTECHA1246.1>, 2009.

570 Ricard, D., Ducrocq, V., and Auger, L.: A Climatology of the Mesoscale Environment Associated with Heavily Precipitating Events over a Northwestern Mediterranean Area, *J. Appl. Meteor. Climatol.*, 51, 468–488, 2012.

Saïd, F., Campistron, B., Delbarre, H., Canut, G., Doerenbecher, A., Durand, P., Fourrié, N., Lambert, D., and Legain, D.: Offshore winds obtained from a network of wind-profiler radars during HyMeX, *Quarterly Journal of the Royal Meteorological Society*, 142, 23–42, <https://doi.org/10.1002/qj.2749>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.2749>, 2016.

575 Seity, Y., Brousseau, P., Malardel, S., Hello, G., Bénard, P., Bouttier, F., Lac, C., and Masson, V.: The AROME-France convective scale operational model., *Mon.Wea.Rev.*, 139, 976–991, 2011.

Wattrelot, E., Caumont, O., and Mahfouf, J.-F.: Operational Implementation of the 1D+3D-Var Assimilation Method of Radar Reflectivity Data in the AROME Model, *Monthly Weather Review*, 142, 1852–1873, <https://doi.org/10.1175/MWR-D-13-00230.1>, <https://doi.org/10.1175/MWR-D-13-00230.1>, 2014.