

Data assimilation impact studies with the AROME WMED reanalysis of the HyMeX SOP1

Nadia Fourrié¹, Mathieu Nuret¹, Pierre Brousseau¹, and Olivier Caumont¹

¹CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

Correspondence: Nadia Fourrié (nadia.fourrie@meteo.fr)

Abstract.

This study was performed in the framework of HyMeX (Hydrological cycle in the Mediterranean Experiment) which aimed to study the heavy precipitation that regularly affects the Mediterranean area. A reanalysis with a convective-scale model AROME-WMED was performed which assimilated most of all available data for a 2 month period corresponding to the first
5 Special Observation Period of the field campaign (Fourrié et al., 2019). Among them, observations related to the low level humidity flow which are important for the description of the feeding of the convective mesoscale systems with humidity (Duffourg and Ducrocq, 2011, Bresson et al., 2012 and Ricard et al., 2012), were assimilated. Among them there were a dense reprocessed network of high quality Global Navigation Satellite System (GNSS) Zenithal Total Delay (ZTD) observations, reprocessed data from wind profilers, lidar-derived vertical profiles of humidity (ground and airborne) and Spanish radar data.
10 The aim of the paper is to assess the impact of the assimilation of these four observation types on the analyses and the forecasts from the 3h forecast range (first guess) up to the 48-h forecast range. In order to assess this impact, several OSEs or also-called denial experiments, were carried out by removing one single data set from the observation data set assimilated in the reanalysis.

Among the evaluated observations, it is found that the ground-based GNSS ZTD data set provides the largest impact on the analyses and the forecasts as it represents an evenly spread and frequent data set providing information at each analysis time
15 over the AROME-WMED domain. The impact of the reprocessing of GNSS ZTD data also improves the forecast quality but this impact is not statistically significant. The assimilation of the Spanish radar data improves the 3-h precipitation forecast quality as well as the short term (30-h) precipitation forecasts but this impact remains located over Spain. Moreover, marginal impact from wind profilers was observed on wind background quality. No impacts have been found regarding lidar data as they represent a very small data set, mainly located over the sea.

1 Introduction

Heavy precipitation regularly affects the Mediterranean area with huge damages and sometimes casualties. One of the aims of the Hydrological cycle in the Mediterranean Experiment (HyMeX ; Drobinski et al. (2014)) was to study the high impact weather events, especially during the first Special Observation Period one (SOP1, Ducrocq et al. (2014)), which took place in the autumn 2012 (5 September - 6 November 2012) in northwestern Mediterranean. The importance of an accurate description
25 of the low-level humidity flow, which feeds the mesoscale systems, was shown in previous studies (Duffourg and Ducrocq,

2011; Bresson et al., 2012; Ricard et al., 2012). This is why during this period research observations were deployed over the north-western Mediterranean area. These observations aimed at a better description of the humidity and wind fields. As an example, water vapour lidars were deployed in Candillargues and Menorca island (pink dots in Figure 1). Particular attention was also paid to the control of data quality.

30 Another important element to better understand the key processes related to the high precipitation and their forecasting is the convective scale modeling. Since many years, such numerical weather prediction models have been implemented in operations to enhance the forecast quality. In addition, the forecast quality depends on their initial atmospheric conditions, which are determined with data assimilation system.

For the HyMeX SOP1 campaign, an AROME (Application of Research to Operations at MEscale, Seity et al., 2011)
35) version was developed and ran in real-time to forecast and study heavy precipitation in this region: the AROME-WMED (western Mediterranean) model (Fourrié et al., 2015). This model is centered over the western Mediterranean basin and includes a data assimilation system, which provides every 3 hours an analysis of the meteorological situation. In the framework of the Innovative Observing and Data Assimilation Systems for severe weather events in the Mediterranean (IODA-MED) project, two reanalyses were performed after the campaign (Fourrié et al., 2019) with the aim of providing new references for process
40 studies. The first one intended to provide a homogeneous data set of atmospheric fields (which was not the case in real-time version due to a system upgrade in the middle of the SOP1). The second one included in addition a maximum of observations deployed during SOP1 field campaign with a more recent version of the model. The latter will be considered in this study.

Among the research observations assimilated in AROME-WMED reanalysis were the humidity profiles from ground based and airborne lidars. Reprocessing after the campaign was also performed for the wind profiler data (Saïd et al., 2016) and
45 the ground based Global Navigation Satellite System (GNSS) zenithal total delays (ZTD) (Bock et al., 2016) to improve data quality and filter out bad data.

Previous impact studies were already performed for this type of observations in other contexts. For example, Bielli et al. (2012); Grzeschik et al. (2008) tested the impact of the assimilation of water vapour lidars in meso-scale models and found a positive impact of such an assimilation up to the 24-h forecast range. Benjamin et al. (2004) studied the impact of a wind profiler
50 network and obtained a positive impact on short-range (3–12 h) forecasts. Concerning the GNSS data, Mahfouf et al. (2015) showed systematic improvements of the atmospheric humidity short-range forecasts and of the structure and the location of precipitation in the AROME models as found previously in a heavy precipitation context (Boniface et al., 2009). These results agree well with previous studies performed in other NWP models (Macpherson et al., 2008; Gutman et al., 2004)

As previously mentioned, an accurate description of the low-level humidity flow is required to well simulate the evolution
55 of the mesoscale system. The aim of the study presented here is to quantify the impact of four observation systems on the quality of precipitation simulation. These observation data sets were assimilated in the AROME-WMED reanalysis of SOP1 and provided information on this low-level flow. The observations are the reprocessed ZTD from the ground based GNSS (Bock et al., 2016), the humidity profiles from ground based and airborne lidars (Chazette et al., 2016 and Di Girolamo et al., 2016), reprocessed wind profiler data (Saïd et al., 2016) and the Doppler winds and reflectivities from the Spanish radars. To

60 achieve this, a number of denial data assimilation experiments, consisting in removing one observation type, were carried out during the 2-month period of SOP1.

The paper is arranged as follows. Section 2 describes the AROME-WMED configurations, the observations data sets and the denial experiments. Section 3 assesses the impact of the ground-based GNSS data assimilation on the analyses, the background and on the forecast quality during SOP1. Section 4 provides information on the impact of other observation types (i. e. wind 65 profilers, lidars and Spanish radars). Section 5 focusses on the impact of all these data on the IOP 16a case study. Finally, conclusions are given in Section 6.

2 Sensitivity study description and validation methodology

2.1 "Denial" Experiment Methodology

To study the contribution of the observations on the analysis and forecast quality of the heavy precipitating events of the SOP1, 70 denial experiments have been devised. These experiments consist of removing one observation data set and to compare the forecast quality with the one originating from assimilating all the observations. Here, denial experiments were conducted on the following four observation types: the ground-based GNSS ZTD, the wind profilers, the water vapour lidars and Spanish radars. They were performed with the AROME-WMED model.

2.2 AROME-WMED configuration

75 The different AROME-WMED model configurations are described in Fourrié et al. (2015, 2019) and rely on the operational limited area model AROME (Seity et al., 2011; Brousseau et al., 2016) version running at Météo-France since 2008. At the time of the SOP1 campaign, analyses were performed at 2.5 km horizontal resolution every 3 hours with a three dimensional variational data assimilation (3D-Var, Brousseau et al., 2011). The AROME-WMED version used in this study as the reference is the second reanalysis one, named hereafter REANA. An extensive description of this reanalysis can be found in Fourrié et al. 80 (2019). The main components are recalled here. The REANA dataset has a 2.5 km horizontal resolution and the model has 60 vertical levels from 10 m above the surface to 1 hPa. Deep convection is explicitly resolved and one-moment microphysical scheme with five classes of hydrometeors is used (Pinty and Jabouille, 1998; Caniaux et al., 1994).

Initial atmospheric states of AROME-WMED come every 3 hours from 3D-Var analyses assimilating observations within a +/- 1h30 assimilation window. This sytems analyzes the two components of horizontal wind, temperature, specific water-vapor 85 humidity and surface-pressure fields on the model grid at full resolution. The other prognostic model fields (turbulent kinetic energy, pressure departure from hydrostatism, vertical divergence and specific content of five condensed water species) are not updated by the analyses but copied from the background. The background error statistics are climatological. Based on the Berre (2000) multivariate formulation, cross-covariances between errors for different physical quantities are represented using scale-dependent statistical regressions, including an extra balance relationship for specific humidity. The background error 90 statistics have been calculated using forecast differences from a AROME-WMED Ensemble data assimilation (Brousseau et

Observations type	amount	percentage
Satellites	8,663,312	53.00%
Surface stations	2,485,620	15.21%
Radars	1,942,539	11.88%
Spanish radars	97,847	0.6%
Aircraft	1,413,313	8.65%
Radiosondes	1,319,523	8.07%
GNSS ZTD	302,191	1.85%
Wind profiler	191,012	1.17%
Lidars	19,470	0.12%
Total	16,346,191	100%

Table 1. Sorted amounts of assimilated data in REANA over the SOP1 period (5 September-5 November 2012).

al. (2011) approach) over a 15-day period of the HyMeX SOP1 (17 to 31 October 2012) to be representative of the encountered meteorological conditions of the SOP1 in average. More details on these background error covariances are available in Fourri  et al (2019). Lateral boundary conditions are hourly provided by the global NWP ARPEGE (Courtier et al., 1991) forecasts which also benefited from a maximum of assimilated observation with longer cutoff analyses. Each day at 00UTC, a 54 hour
95 forecast is run. Conventional observations (from radiosondes, aircraft, surface stations, wind profiler, GNSS ZTDs), radar data and satellite observations (infrared and microwave radiances, atmospheric motion vectors and ocean surface winds from scatterometers) were assimilated.

Table 1 presents the distribution of assimilated data in REANA as a function of observation types. Satellite data represent the majority of observations. This can be explained by the fact that the IASI sensor provides 44 channels per observation point.
100 Surface observations provide 15.21% of assimilated data. Aircraft and radiosondes give similar amount of data (around 8%). GNSS ZTD represent 1.85% of the total and wind profilers 1.17%. Special efforts were made to assimilated non operational data types such as Lidar water vapour profiles and Spanish radar data. Humidity data from Lidar contribute very few with 0.12% of assimilated data. Radar data represent 11.88% of the total amount of assimilated data and Spanish ones only 0.6%.

2.3 Description of the studied observing systems

105 As mentioned above, four observing systems were studied. The location of these observations is shown in Figure 1, excepting the ground based GNSS ZTD location which is available in Figure 4 of Fourri  et al. (2019).

2.3.1 GNSS Zenithal Total Delays

GNSS ZTD provides useful information on precipitable water and pressure at a high temporal frequency and in all weather conditions. In REANA2, we considered here reprocessed data (REPRO-GNSS in the following) with a homogeneous repro-
110 cessing using a single software and more precise satellite orbits position and clocks (Bock et al., 2016), which were available




Figure 1. Location of observations considered in this study, with the exception of GNSS Zenithal Total Delays. Wind profilers are depicted with blue squares, ground based lidars with pink dots, assimilated Leandre II airborne profiles with purple dots, Spanish radars with orange symbols. Red symbols correspond to the French radar locations.

for the whole SOP1. Additional data were also considered compared to the operational and data set available in near real-time. This data set, called hereafter OPER-GNSS, is provided by E-GVAP (EUMETNET EIG GNSS (Global Navigation Satellite System) water VApour Programme) and ZTD data for one reception station may be available for more than 10 processing centres. These ZTD data are assimilated according the methodology described in Mahfouf et al. (2015). The model equivalent
115 is computed with the following equation (Mahfouf et al., 2015):

$$ZTD = 10^{-6} \int_0^{z_{top}} \left(k_1 \frac{p}{T} + k_3 \frac{e}{T^2} \right) dx \quad (1)$$

where p is the pressure, T the temperature, e the water vapour pressure, $k_1 = 0.776 \text{ Pa}^{-1} \text{ K}$, and $k_3 = 3730 \text{ Pa}^{-1} \text{ K}^2$, x is the height above the ground and z_{top} is the height of the model top. After a monitoring of the difference between observations and model equivalent, observations with good statistics are selected in a 'white list'. ZTD data are also bias corrected and
120 an updated bias correction for each GNSS station was also computed in the REANA2 version. They are finally assimilated if they pass the first guess quality control which rejects data too far from the model background. Only one observation per 3-h assimilation and per surface station is assimilated for each analysis. Please refer to Mahfouf et al. (2015) for more information on the data assimilation of GNSS ZTD in AROME.

2.3.2 Wind profilers

125 Data from eight wind profiler radars (sounding in VHF or UHF bands) were assimilated in AROME-WMED. These profilers provided vertical profiles of wind vector, turbulence, precipitation and the height of the atmospheric boundary layer and tropopause (Saïd et al., 2016). The measure principle is described in Annex 1 of Saïd et al. (2016)'s paper. Profilers measure the Doppler radial spectra of the returned signal backscattered by various types of targets. In order to derive the three components of the wind, most of the HyMeX profilers use five beams. These data were available for the whole SOP1 in real-time and
130 have been reprocessed after the SOP1 by Saïd et al. (2016) with an improved quality control to remove spurious data. Here, observations from 8 wind radars (UHF and VHF) mainly located in the South of France, in Corsica and Menorca (Figure 1) were considered. These observations are assimilated as vertical profiles of horizontal wind.

2.3.3 Lidars

During SOP1, ground based and airborne lidars were operated. The mobile Water vapour and Aerosol Raman Lidar (WALI, Chazette et al. (2016)) operates with an emitted wavelength of 354.7 nm. This instrument was operated at a site close to Ciutadella (western part of Menorca located by 39 59 07 N and 3 50 13 E). Mixing ratio profiles were delivered with a resolution of 15 m for the 0 m - 6000 m altitude range. A detailed description of this instrument can be found in Chazette et al. (2016). The raw vertical resolution of the data is 75 m but for assimilation above 2000 m, resolution was thinned starting from 75 m to 450 m above 5000 m.

140 The second ground based lidar, the BASIL instrument (Di Girolamo et al., 2016) was located in Candillargues in the South of France. The original data resolution is 30 m but data were thinned at 60 m below 1000 m, increasing up to 420 m above 4000 m in the assimilation. For WALI, 292 mixing ratio profiles were assimilated in REANA, covering the period 17 September 2012 - 03 UTC to 27 October 2012 - 21 UTC, whereas for BASIL, 172 profiles were assimilated, covering the period 10 September 2012 - 09 UTC to 5 November 2012 15 UTC.

145 Concerning Leandre II lidar (Chazette et al., 2016) on board ATR aircraft, data were available for 22 analysis slots (512 assimilated profiles), covering the period 11 September 2012 09 UTC to 25 October 2012 21 UTC. Profiles with a 150 m vertical resolution were thinned at a 15 km horizontal resolution and are mainly located over the Mediterranean Sea (Figure 1).

2.3.4 Spanish radars

Doppler radial winds and reflectivities from five Spanish radars, located in Barcelona, Valencia, Almeria, Murcia, Palma de Mallorca and provided by AEMET were assimilated in REANA. After a strict quality control check to exclude data with gross errors, only the three lowest elevations have been considered for the assimilation. Doppler wind are assimilated in the 3D-Var of AROME according the method described by Montmerle and Faccani (2009) and reflectivity data are assimilated as pseudo-observations of relative humidity profiles as proposed in Caumont et al. (2010) and implemented in Wattrelot et al. (2014).

155 Several procedures are applied to raw data in order to avoid as much as possible erroneous measurements entering the minimization. An observation operator allows to simulate radial Doppler winds measurements from the model horizontal wind based on Caumont and Ducrocq (2008). Only measurements performed within 150 km of the radar are considered due to the broadening of the beam with increasing distance and the lack of reliability. An observation error variance proportional to their distance from the radar is applied in the minimization. Reflectivities are not directly assimilated but they are used to retrieve pseudo-observations of relative humidity from surrounding simulated reflectivity profiles through a unidimensional Bayesian inversion. A horizontal thinning on the data (Doppler winds and retrieved profiles of relative humidity) is performed to avoid horizontal correlation of observation errors: only one profile, having the most important number of elevations that passed the quality control, is selected in each $15 \times 15 \text{ km}^2$ box.

2.4 Description of the experiments

Experiment name	Description	Difference (%) in the number of assimilated data
REANA	AROME-WMED reanalysis (2nd), see Fourrié et al. (2019)	
NOGNSS	REANA - reprocessed GNSS ZTD	-1.86%
OPERNSS	NOGNSS + operational GNSS ZTD	-1.04%
NOLIDAR	REANA - LIDAR	-0.15%
NOWPROF	REANA - wind profilers	-1.12%
NORADSPAIN	REANA - Spanish radars	-0.6%

Table 2. Description of the data denial experiments discussed in this study and difference (in %) in the number of assimilated data compared to the reanalysis REANA.

165 Table 2 summarizes the names of the denial experiments and the observations considered. Five experiments were conducted over the 2-month period of SOP1 (from 5 September 2012 to 5 November 2012). They all used the same configuration of AROME-WMED, the differences lying in the observations assimilated. For each experiment, it differs only one observation type from the reanalysis (REANA) used as the reference. This allows to evaluate the impact of this observation type on the analysis and the forecast. Among the five experiments, two experiments deal with the impact of GNSS ZTD. The first one, NOGNSS is obtained by removing the REPROC-GNSS ZTD from the assimilation. The second, called OPERGNSS, aimed to evaluate the impact of the REPROC-GNSS data set provided by Bock et al. (2016) compared to the OPER-GNSS data set provided by E-GVAP. The E-GVAP data set was thus assimilated in replacement of the Bock et al. (2016)’s one in OPERGNSS. The NOLIDAR experiment is the run with no airborne nor ground-based Lidar data in the data assimilation. The NOWPROF experiment is obtained by removing the wind profiler data and the NORADSPAIN experiment was run without any data from the five Spanish radars. 97847 observations, representing 0.6% of the total number of assimilated observations, were removed in the NORADSPAIN experiments

As shown in Table 2, the largest differences in terms of number of assimilated observations are obtained with NOGNSS which leads to a 1.85% difference in the number of assimilated data.

2.5 Validation protocol

180 As a first step, the performance of the data assimilation system is validated by comparing the various Analysis (AN) and First-Guess (FG) values against available observations which can be independent from REANA (i.e. not assimilated) or on the contrary assimilated in REANA. One of the key tool used to evaluate the performance of the assimilation system is to examine the FG departure (O-FG) and the AN departure (O-A) in terms of mean and root-mean square (RMS) values, O standing for Observation with the other assimilated observations. Statistics of departures are computed at the observation location.

185 Those statistics were also computed using few available independent data. The first source comes from the vessel Marfret-Niolon, which was an instrumented commercial ship of opportunity, cruising regularly between the southern France harbour of Marseille and two Algerian harbours (Algiers and Mostagadem). Please refer to Figure 14 of Fourrié et al. (2019) for the trajectories of the vessel during SOP1. Two autonomous systems were installed in order to provide atmospheric and oceanic measurements, in the context of the HyMeX Long Observation Period (LOP). A GNSS antenna was installed at the front
190 on the vessel Marfret-Niolon for the duration of the HyMeX campaign. An example of the operational measurements which started on January 2012 are provided in Figure 2 with figures ranging from 2.2 m to 2.6 m. The data were post-processed in kinematic Precise Point Positioning with the software provided by Natural Resources Canada (Kouba and Héroux, 2001) and using high-resolution products provided by the International GNSS Service.

The second source of independent data comes from wind data obtained from an airborne Doppler cloud-profiler radar named
195 RASTA (Radar Airborne System Tool for Atmosphere (Bouniol et al., 2008; Protat et al., 2009; Delanoë et al., 2013)) that flew 45 days during SOP1. This airborne radar was on board the Falcon 20 research aircraft. It allows the documentation of the microphysical properties and the horizontal components of the wind field in terms of vertical profiles.

The operational data assimilation monitoring procedure also provides FG and AN departure statistics for assimilated observations in the experiments, which are described in a companion paper (Fourrié et al., 2019).

200 In a second step, the forecast (range between +3 to +54 hours) quality is assessed in terms of surface parameters and precipitation scores. The surface parameters (temperature and relative humidity at 2 m and wind at 10 m) come from the HyMeX database which provides surface synoptic observations available over the AROME-WMED domain, together with additional hourly observations from Météo-France, AEMET and MeteoCat mesoscale networks. Some of these observations were assimilated to produce surface analyses. For the evaluation of the precipitation quality, the dense surface data set rain
205 gauge network available in the HyMeX data base (V4 version, DOI:10.6096/MISTRALS-HyMeX.904) has been used. Scores of 3 hourly accumulated precipitation from all analysis times on a given day are compared to the corresponding observed 24-h accumulated precipitation.

The evaluation of the various denial experiments is compared with the reference REANA run. This allows to get the impact of each considered observation type on the analysis and the forecast.




Figure 2. Evolution of the Zenithal total delay (ZTD, m) observed onboard the Marfret-Niolon ship during 25 October 2012.

210 3 Impact of GNSS data on the analysis and first-guess quality

This section investigates the impact of assimilating the ground-based GNSS ZTD data on the numerical weather prediction model analysis and subsequent forecast quality. This data set represents the largest one in terms of the number of studied observation types, event though in the end it represents only a small fraction of assimilated data (1.85%) in the analyses (Table 1). As seen in Table 1, satellite data are the most numerous, followed by surface stations data, radar data from the French
 215 network, aircraft data and radiosondes ones. Even if surface data provide information only for one level, the network is very dense over France and was reinforced in other countries like Spain or Italy. The other observation types provide information at different levels all along the vertical.

3.1 Impact on moisture field

Comparison to the Integrated Water Vapour (IWV) from the reprocessed GNSS observations (not independent from REANA as
 220 the information from this data set is assimilated in this experiment) indicates that the best correlation, as expected, is obtained for REANA (around 0.99), the second one being OPERGNSS (around 0.975) and the last one NOGNSS (around 0.96), as shown in Figure 3. This result is confirmed when computing the RMS of the differences. A weak diurnal cycle of the scores is noticed with a maximum correlation around 09 UTC and a minimum around 15 UTC. Concerning the standard deviation of the differences, they are lower during the 3-9 UTC period and larger in the afternoon. These minimum of correlation and
 225 maximum of standard deviation correspond to the time of the early stage of the convection.

We then discuss the result of the statistics for the analysis and first-guess against radiosonde observations, which represents a reference data set in data assimilation. First of all, as expected, the analysis RMS difference (solid lines) are smaller than the FG




Figure 3. Correlation (left panel) and standard deviations (right panel, in kg/m^2) of integrated water vapour (IWV) content from reprocessed GNSS observations (Bock et al., 2016) and AROME-WMED analyses (REANA in black NOGNSS in red and OPERGNSS in blue) as a function of analysis slot (hours).

difference (dashed lines) for the three simulations showing the expected behaviour of the minimisation during the assimilation process (Figure 4). No impact can be seen on the analysis RMS differences. The absence of impact can be explained by the fact that radiosondes are reference observations for assimilation and all the analyses are very constrained by these observations. However, a small positive impact is present on the FG RMS difference 3-h later. Lowest differences are obtained with REANA simulation, the largest ones with NOGNSS. The OPERGNSS differences are close to REANA one but slightly larger, showing on the one hand that the assimilation of GNSS data is beneficial (OPER-GNSS data set or REPROC-GNSS one) and on the other hand that the reprocessing of the data brings a small improvement in the comparison of FG with humidity of radiosondes. This shows that the modifications in the analysis brought by the GNSS at other places than radiosondes ones are beneficial and kept during the 3-h forecast. The largest improvement of the assimilation of GNSS data (OPER-GNSS data set and REPROC-GNSS one) is found between 600 and 850 hPa. In addition, a slight benefit of assimilating REPROC-GNSS data versus OPER-GNSS ones appears between 700 and 850 hPa.

The various analysis mean departures are very close to each other, with slight negative values in the lower and mid troposphere (analysis too moist), as displayed in Figure 4 lower panel. Mean first-guess departures are larger and homothetic, with stronger values for the REANA simulation, being the signature of a weak moist bias in the corresponding analysis for the lower troposphere. The less biased first guess is the one from the NOGNSS experiment.

Radiances from SEVIRI (on board the geostationary satellite Meteosat Second Generation, MSG), sensitive to moisture (channels WV $6.2 \mu\text{m}$ for upper-troposphere and $7.3 \mu\text{m}$ for mid-troposphere) are assimilated in AROME. They are an important source of humidity information, especially over oceans where no information from GNSS nor radiosondes is available. Basically no impact between the various experiments is found on the FG and AN statistics for these observations (not, shown, Table in supplement file).

The correlation between the various AROME-WMED ZTD AN and corresponding independent (not assimilated) Marfret-Niolon observations is higher for REANA than for NOGNSS and even for OPERGNSS (Figure 5). There is a correlation maximum around 09 UTC, and a minimum around 15 UTC. The mean ZTD is quite similar in all experiments, with a maximum at 09 UTC and a minimum around 00 UTC. A moist bias is found in all simulations when compared to the mean observation in grey shown in Figure 5. The magnitude of this relative positive (moist) bias is around 0.5 percent. Table 3 shows the mean correlation of REANA, NOGNSS and OPERGNSS AN and FG with Marfret-Niolon observations. The higher correlation is obtained with REANA for both AN and FG. When comparing the mean value of ZTD at the Marfret-Niolon places, the closest value to the observed one is obtained with REANA, even if a small moist bias is observed (0.9 mm). This bias is larger for NOGNSS (3.6 mm) and OPERGNSS (4.8 mm).

This could be explained by the fact that there few assimilated observations over the sea which results in a more biased model. Although the sample size of Marfret-Niolon data set is rather small (around 1000 collocations), this is an original result and makes clear that the REANA experiment produces the best reanalysis, and the best 3-hour forecasts.




Figure 4. Root mean square differences (left hand side panel) and mean (right hand side panel) for Analysis (AN solid lines) and First-Guess (FG, dashed lines) departures against assimilated radio-sounding observations for mixing ratio (g/kg); REANA in black, NOGNSS in red and OPERGNSS in blue.)

260 3.2 Impact on wind field

Analysis and First-guess quality has been evaluated against RASTA (Radar Airborne System Tool for Atmosphere) Doppler winds (Borderies et al., 2019a). This airborne radar was on board Falcon 20 aircraft and provided 33083 wind observations over the Mediterranean area as illustrated in Figure 6, where only few wind data from conventional observations are available.




Figure 5. Correlation of the differences between zenithal total delays (ZTD) between REANA (black), NOGNSS (red), OPERGNSS (blue) analyses and corresponding Marfret-Niolon independent observations as a function of analysis time in the left panel; mean value of ZTD (m) in the right panel, the grey line corresponding to observations.

Experiments	AN ZTD Correlation	Mean ZTD (m)	FG ZTD Correlation
REANA	0.967	2.4617	0.961
NOGNSS	0.961	2.4642	0.958
OPERNSS	0.962	2.4654	0.958
OBS		2.4606	

Table 3. Correlation of zenithal total delays (ZTD) between REANA, NOGNSS and OPERGNSS analyses and corresponding Marfret-Niolon observations computed over the 8 analysis slots (first column), mean ZTD for REANA, NOGNSS and OPERGNSS analyses and Marfret-Niolon observations and correlation between ZTD forecasted by AROME-WMED at the 3-hour forecast range and observations from Marfret-Niolon.

Worth to remind that the data from this instrument were not assimilated in REANA. This data set thus represents an additional independent information for the evaluation of our denial experiments.

Table 4 provides the RMS errors (RMSE) for wind calculated with these data. The RMSE for background and analysis are lower in REANA than in the other two experiments. The analysis RMSE for OPERGNSS is lower than the one for NOGNSS.

As GNSS observations do not provide any wind information, the improvement observed in wind field can be explained by the effect of mass field information assimilation on the wind field, essentially created during model integration. There is indeed a little coupling between these fields during the analysis (Borderies et al., 2019b). This indirect effect was already demonstrated by Wattrelot et al. (2014), for example, who noted a positive impact on the wind field when assimilating pseudo-observations of relative humidity. Lindskog et al. (2017) also reported—but did not show—a positive impact on wind forecasts when assimilating ZTD data.




Figure 6. Location of RASTA observations during the HyMeX Special Observation Period 1. Coloured dots represent the number of wind data available per profile.

Experiment	AN RMSE	FG RMSE
REANA	5.59	5.87
NOGNSS	5.63	5.97
OPERGNS	5.60	5.92

Table 4. Analysis (AN) and First Guess (FG) Root Mean Square Errors (RMSE) for the wind (in m/s) computed with respect to RASTA observations (sample size 33083 observations) for REANA, NOGNSS and OPER GNSS experiments.

3.3 Impact on short-range precipitation

Figure 7 shows that the Equitable Threat Score (ETS) of the 24-h accumulated precipitation computed with the sum of the 3-h precipitation from the 8 analysis times is improved with the assimilation of GNSS ZTD data compared to the NOGNSS experiment. It represents an evaluation of the background quality. The difference is statistically significant for each threshold. When comparing the assimilation of REANA to OPERGNSS, the ETS for precipitation is slightly better with the reprocessed data set but the differences are not significant except for the 40 mm/day threshold. Overall, the background quality is improved with the assimilation of GNSS observations and the data reprocessing brings improvement in terms of precipitation from 3-hour forecast even though this benefit is not significant.

4 Impact of GNSS data on medium term forecast

The impact of the GNSS data has also been assessed for longer forecast ranges (3 to 54-h). The effect of the assimilation of the GNSS data on the correlation with IWV from the GNSS data set is maximal for the analysis and decreases up to the




Figure 7. Equitable Threat Score (ETS) for the 24-h accumulated precipitation from the sum of the eight 3-h forecasts used as background of the data assimilation cycle each day of the period from 5 September to 5 November 2012. Results for REANA are displayed in black, for NOGNSS in red and OPERGNSS in blue. Dots indicate that the difference between the curves and the REANA curve as a reference is statistically significant at a 0.95 confidence threshold using a Bootstrap test.

285 30-h forecast range (Figure 8) as the general impact of the initial conditions on the forecast performances reduces. A similar behaviour is found with the standard deviations of the differences between observed IWC and simulated one from the three experiments.




Figure 8. Correlation (left hand side panel) and standard deviations (right hand side panel, kg/m^2) of integrated water vapour content between AROME-WMED forecasts and reprocessed GNSS observations (Bock et al., 2016) as a function of forecast range (hours).

Compared to the observed ZTD from the Marfret-Niolon ship, the signal is more noisy because of a smaller dataset but when comparing to values average over the forecast ranges (Table 5), the correlation for the NOGNSS is lower than REANA and OPERGNSS, which provides it-self lower correlation than REANA. The standard deviations are higher for the NOGNSS

290

Parameter	REANA	NOGNSS	OPERNSS
Correlation (1-24h)	0,962	0,957	0,957
Correlation (25-48h)	0,922	0,917	0,919
Correlation (49-54h)	0,906	0,902	0,905
Standard deviation (forecast -observation, 1-24h)	0,0152	0,0164	0,0160
Standard deviation (forecast -observation, 25-48h)	0,0221	0,0226	0,0223
Standard deviation (forecast -observation, 49-54h)	0,0244	0,0249	0,0244

Table 5. Correlation and standard deviation of ZTD (in m) between AROME-WMED forecasts and reprocessed GNSS observations averaged over forecast ranges.

forecasts. In addition, a decrease of the correlation (respectively an increase of the standard deviation) is seen for forecast range over 24-h.

The forecast quality has also been evaluated against surface data. No impact was found on temperature at 2 m or on 10 m wind. A small impact was found on relative humidity at 2 m (Figure 9). A reduction of the bias is noticed with REANA during the first 9-h of the forecast compared to OPER GNSS and NOGNSS. From 12-h onwards the results for REANA and OPERGNSS are similar. Regarding the standard deviation, it is smaller for REANA between 0 and 9-h than for NOGNSS and OPER GNSS and between 21 and 27-h forecast range than for NOGNSS. This difference represents more than 2 % of improvement. For the other forecast ranges the differences are lower than 1%.




Figure 9. Bias (forecast minus observations, dashed lines) and standard deviations (solid lines) computed with relative humidity at 2 m (left hand side panel) and relative root mean square differences (%) right hand side panel) with respect to REANA.

The impact of the assimilation of GNSS data on the 24-h accumulated precipitation from the forecast initialized at 06 UTC is less clear. The improvement of the GNSS data reprocessing compared to the real time data set is beneficial for all thresholds except for the 2mm/day (where the ETS is better for OPERGNSS) and is statistically significant for moderate thresholds

(10 and 20 mm/day, Figure 10). The difference between REANA ETS and NOGNSS ETS values is not significant. When examining scores for precipitation forecasts between 30-h and 54-h, there is a small significant degradation of the ETS for the 2 mm/day with the NOGNSS experiment and a small improvement with the OPERGNSS for the 40 mm/day (Figure 11).




Figure 10. Equitable Threat Score of the 24-h accumulated precipitation from the 6-30 hour forecast range of the long forecast initialized at 00 UTC each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). Dots indicate that the difference between the curves is statistically significant.




Figure 11. Equitable Threat Score of 24 h accumulated precipitation from the 30 to 54 hour forecast range of the long forecast initialized at 00 UTC each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). Dots indicate that the difference between the curves is statistically significant.

As previously mentioned we performed other impact studies with wind profilers, lidar data and Spanish radar data.

5.1 Wind profilers

No impact of the assimilation of wind profiler data is found except on wind field. Small impact is noticed in terms of wind RMS differences of background and analysis departures for radiosondes, aircraft and satellite winds (Figure 12). The largest
 310 impact is a decrease of -0.08 m/s for the radiosonde FG RMS differences at 300 hPa. Concerning the AN RMS differences, the improvement (SATOB) or degradation (AIREP and TEMP) are very small. The largest value obtained at 200 hPa are due to the small number of data available for the computation.




Figure 12. First-Guess (left plot) and analysis (right plot) RMS differences (REANA-NOWPROF experiments) computed against TEMP (black), AIREP (red) and SATOB (green) observations for the zonal wind component (m/s); negative value correspond to a positive impact of wind profiler.

A small improvement of REANA compared to NOWPROF, but not significant (Figure 13), appears on the ETS of the 24 h accumulated precipitation accumulated from the 6 to 30 hour forecast ranges.

315 5.2 Ground-based and airborne lidar data

As discussed in Section 2.2, humidity profiles retrieved from ground-based and air-borne lidars have been assimilated in the REANA experiment. In Figure 1, the trajectories of all ATR-42 flights are plotted, together with the localization of the two ground-based lidars. The denial NOLIDAR experiment results are very close to the reanalysis ones (Table 6) as these data represent very few additional data located over ocean where few observations are available for the comparison. No impact of
 320 the Lidar data is found when comparing the various analyzed ZTD to the Marfret-Niolon corresponding observations. These results agree with the Bielli et al. (2012) study where no significant impact where found on the 24-h accumulated precipitation.




Figure 13. Equitable Threat Score of 24 h accumulated precipitation from the 6 to 30 hour forecast ranges of the long forecast starting at 00 UTC each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). The lack of dots indicates that the difference between the curves is not significant. REANA is plotted in black and NOWPROF in red.

	REANA	NOLIDAR
Correlation for the forecast 0h	0.968	0.960
Correlation for the forecast 1-24 h	0.962	0.961
Correlation for the forecast 25-48 h	0.923	0.924
Correlation for the forecast 49-54 h	0.906	0.907
Standard deviation for the forecast 0h	0.0144	0.0167
Standard deviation for the forecast 1-24 h	0.0152	0.0154
Standard deviation for the forecast 25-48 h	0.0221	0.0220
Standard deviation for the forecast 49-54 h	0.0243	0.0243

Table 6. Correlation and standard deviation of Zenithal Total Delays (in m) between AROME-WMED forecasts from 00UTC and reprocessed GNSS observations averaged over forecast ranges (0 h, 1-24 h, 25-48 h and 49-54 h).

5.3 Spanish radars

No significant impact has been noticed over the HyMeX domain however, when focusing on the scores over the Iberian Peninsula, we obtained a positive and significant impact of the assimilation of Spanish radar data on the ETS for the 24-h accumulated precipitation from the sum of the 8 3-h precipitation background forecast (Figure 14). This impact also remains in longer forecast ranges as the ETS for the 24-h precipitation accumulation between 6-h and 30-h forecast ranges is improved with the assimilation of Spanish radars for thresholds between 0.5 and 20 mm/24h (Figure 15). This impact does not remain at longer forecast ranges (Figure 16). These results are in good agreement with Wattrelot et al. (2014) study which found an improvement of the short term precipitation forecast scores. However contrary to the aforementioned study, we obtained a significant improved of the 24-h precipitation accumulation between 6-h and 30-h forecast ranges over the Iberic Peninsula. Even if we do not obtain significant impact at the HyMex domain scale but a significant one over the Iberian Peninsula, it is interesting to remind that the assimilation of Spanish radar data in AROME-WMED was made on a research mode as only French radars were assimilated at the time of the HyMeX campaign and the reanalyses. These data represent only 0.6% of the assimilated data. This is three times less than REPROC-GNSS data.




Figure 14. Equitable Threat Score (ETS) for the 24-h accumulated precipitation obtained from the sum of eight 3-h forecasts used as background of the data assimilation cycle each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). Results for REANA are displayed in black, for NORADSPAIN in red. Dots indicate that the difference between the curves is statistically significant.

6 IOP16 case study

During HyMeX SOP1, IOP16a was dedicated to HPE that occurred over Cévennes-Vivarais (CV) in France and later on, in Italy (IOP16b) on 25-26 October; this event was associated with locally flash-flooding and several casualties. This off-shore




Figure 15. Equitable Threat Score (ETS) for the 24-h accumulated precipitation from the 6 to 30 hour forecast ranges initialized at 00 UTC each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). Results for REANA are displayed in black, for NORADSPAIN in red. Dots indicate that the difference between the curves is statistically significant.




Figure 16. Equitable Threat Score (ETS) for the 24-h accumulated precipitation from the 30 to 54 hour forecast ranges initialized at 00 UTC each day of the period from 5 September to 5 November 2012 computed over the AROME-WMED domain with rain gauges of the HyMeX database (version 4). Results for REANA are displayed in black, for NORADSPAIN in red. Dots indicate that the difference between the curves is statistically significant.

convection case is well documented in Duffourg et al. (2016). On the 26th - 00 UTC active convection was occurring over Catalonia; this area of intense convective activity crossed the Gulf of Lion reaching the French Mediterranean coast around 06
340 UTC and later on, and then hitting the Italian Ligurian coast in the evening. It is well known that the associated convective systems are usually fed with moisture, during their early stage over the warm Mediterranean sea. A moist conditionally unstable south-western flux is therefore found in the lower troposphere (Figure 17) with a low-level jet by the Candillargues radar around 09-12 UTC, associated to a slow evolving weak pressure low (around 995 hPa) localized over southern France on the 26th mid-day. Moreover, low level convergence is reinforced by the complex orography (Cévennes ridge of the Massif-Central and Alps
345 in France) triggering convection. An upper south-westerly wind jet is observed above 500 hPa (Figure 17); in the evening of the 25th the wind rotates to the west on the CV area as shown by the Candillargues UHF radar.

During 25th and 26th October period, many deep convective systems developed over the Northwestern Mediterranean. Although observed accumulated surface precipitation from Friday 26th at 18 UTC to Saturday Oct. 27th at 06 UTC over southern France only reached around 150 mm in 24h, very strong hourly rates (near 50 mm/1h) were recorded, with intense
350 river discharges (Ardèche, Gardons and Gapeau rivers for example). Such intense rainfall amounts led to local flash-floods and 2 casualties in the Var region. In fact as shown in Figure 18, three local precipitation maxima appear on the observed 24-hour accumulated rainfall amount (25th October - 06 UTC to 26th October - 06 UTC) on the Mediterranean coastal area of France and Italy (Liguria Tuscany region); a first elongated one in the Cévennes area (more than 150 mm, M1) and a small second one close to the coast (around 100 mm, M2).

355 Figure 19 shows the 24-h accumulated precipitation between the 6-h and 30-h forecasts for the different experiments considered in this study. The REANA 24-hour accumulated rainfall (from 06 h to 30 h forecast range) simulation agrees to the observations for both M1 and M2 systems. The NOLIDAR experiment is very close to REANA, this is consistent with the fact that the amount of additional lidar data is fairly small in REANA when compared to NOLIDAR. The strongest impact is found when no GNSS data are assimilated (NOGNSS run): M1 and M2 are strongly underestimated; surprisingly the OPERGNSS
360 experiment leads to an accurate forecast of M2, but underestimates the southwestward extension of M1. Finally a strong negative impact is found with the NOWPROF simulation which misses M2 and does not reproduce correctly M1. Over Italy, the gain brought by the observations is not so evident but it is quite well known in data impact studies that the assimilation of observation does not always improve the forecast at each analysis time but in overall.

7 Conclusions

365 The AROME-WMED model was originally developed to study and forecast heavy-precipitating Mediterranean events during the Special Observation Periods (SOPs) of the HyMeX programme. Two reanalyses were undertaken after the HyMeX autumn campaign for the first SOP. A first one was carried out just after the campaign to provide the same model configuration over the whole SOP1 period because a version upgrade of AROME-WMED occurred during the period. A second reanalysis, performed a few years after, accounted for as many data as possible from the experimental campaign (i.e., lidar and dropsonde humidity




Figure 17. Time-height cross-section of the wind measured by the Candillargues UHF radar for IOP16. Horizontal wind components are represented by the black arrows (meteorological convention), and wind speed in colour.

370 profiles) or from reprocessed data sets (such as GNSS ground station ZTD, wind profilers, high-vertical resolution radiosondes, and Spanish Doppler radars). It also benefited from a updated version of the AROME code.

Previous studies such as Duffourg and Ducrocq (2011) Ricard et al. (2012) or Bresson et al. (2012), have shown the importance of an accurate description of the low-level moist flow feeding mesoscale convective systems. In this study the impact of various data set related to humidity and wind on the forecast quality from this comprehensive reanalysis is investigated over the 2-month period. Many data sets of the Special Observation Period 1 of the HyMeX campaign have been considered here. The reprocessed GNSS data set (Bock et al. (2016)) were removed and replaced with the operational data set used in the real-time AROME-WMED version. We examined the humidity data provided by ground based and airborne lidars. The impact of the reprocessed wind profilers and the Spanish radar data was also evaluated. The impact of these data sets was assessed through Observing system Experiments which consist of removing the data sets and to compare forecast quality from these denial 380 experiments to a reference which includes all data sets. The selected data sets were research observations (water vapour lidars)




Figure 18. 24 h accumulated precipitation (mm) between 26 October 06 UTC and 27 October 2012 at 06 UTC over the AROME-WMED domain (upper plot) and zoom over the Cevennes region (left lower plot) and over North of Italy (right lower plot).




Figure 19. Same as Figure 20 but for 24h accumulated precipitation forecasted by (a) REANA, (b) NOGNSS, (c) NOLIDAR (d) OPERGNSS and (e) NOWPROF experiments, from the 06 h to 30 h forecast ranges.

or reprocessed data (from ground based GNSS receivers or wind profilers). They represented a modest part of the assimilated data amounts and their impact was thus expected to be small.

Our study finds a small positive impact on humidity forecast at short term ranges of the reprocessed GNSS ground based zenithal total delay assimilation. This data set is evenly distributed over the AROME-WMED domain and provided at each analysis time information on integrated water vapour. The impact of the data reprocessing was also studied and even if a positive impact is observed, this improvement is not statistically significant compared to the impact of the real-time data. Given the impact of ground-based GNSS, there is also an interest in continuing work to assimilate GNSS data over ocean surfaces.

Small impacts on wind fields were also observed for wind profilers. No impact from Lidar data was found except when comparing with RASTA data located over the Mediterranean Sea. Since this data set represents a very small fraction of assimilated data, this may explain the absence of impact. In addition they were not assimilated at their full available temporal frequency but just once every 3 hours.

Spanish radar data assimilation improves the short term quality of the background as noticed on the 24-h accumulated precipitation of the eight 3-h background forecasts for each day but only over the Iberic Peninsula with no clear impact over the HyMeX domain. It is interesting to stress that this impact remains during the first 30-h of the forecast but without any

395 remote impact over the rest of the AROME-WMED domain. More impact could possibly be obtained if the data were provided with additional scan elevations.

With the examination of the impact of the assimilation of 4 different data sets over a two-month period in the meso-scale AROME-WMED, our study shows that it is required to have well spatially distributed and frequent data sets such as the GNSS ZTD data set to get, with its assimilation, an overall impact in terms of analysis and forecast skills. This result agrees with the findings of Mahfouf et al. (2015) who show that the assimilation of GNSS systematically improves the atmospheric humidity short-range forecasts despite the small fraction of GNSS observations assimilated in AROME. A high temporal availability and a regular horizontal distribution are both needed to get a significant impact on the forecast scores. Moreover, it is interesting to process as precisely as possible a maximum of GNSS data in real time and to have bias-corrected observations valuable for data assimilation. In addition, GNSS data available on ship seems to be promising to increase the coverage over ocean (Fan et al., 2016). When the data set is available frequently but not well spread over the model domain such as the Doppler winds and reflectivities from the Spanish radars or winds from profiler radars, its assimilation may lead to a positive impact on the precipitation forecast but it remains local. Finally, marginal impact from local and sporadic data sets such as humidity profiles from water-vapour Lidars can be obtained but it is not visible on "global scores". To get a material impact on the forecast in a mesoscale model from a set of observation through its data assimilation, our study suggests to select data sets which are frequently available at each analysis time and also well spread over the domain.

The impact of the above mentioned data could be further improved. For example, the impact of GNSS in AROME-France has been recently improved with the use of variational bias correction in replacement of the static bias correction used in this study (P Moll, personal Communication). In addition radar data from foreign countries are now assimilated in AROME since July 2020. The distribution of these data by OPERA (the EUMETNET Radar programme) allows to get data of high quality in the data assimilation and thus to increase their impact in the AROME model Martet et al. (2019). With a more frequent data assimilation cycle, making use of observations at higher temporal frequency as it is the case with the current AROME-France model (Brousseau et al., 2016), it is likely that surface observations or the remote sensing data such as radars, GNSS or SEVIRI available for each hourly analysis in this study would have a greater impact on analyses and forecasts. Moreover new data assimilation systems such as the 4D-Var or the 4D-Envar are under development for convective scale models (Gustafsson et al., 2018) and will allow to account for very frequent data. Therefore, they are expected to enhance the impact of observations available several times an hour. In the future, the impact of the Infra-Red Sounder onboard Meteosat Third Generation will benefit from these new data assimilation systems as this sounder will provide observations every 30 minutes over the AROME domain and especially over the oceans.

Code availability. The source code of AROME-WMED, derived from the operational AROME code cannot be obtained.

425 *Data availability.* The analyses and the forecast fields are available in the HyMeX database (<http://mistrals.sedoo.fr/HyMeX/>, last access 19 August 2019). The final (second) reanalysis labelled REANA in this paper is available at <https://doi.org/10.14768/MISTRALS-HYMEX.1492> (Fourrié and Nuret, 2017).

Author contributions. NF and MN prepared and carried out all the numerical experiments of the reanalysis and the OSEs. They investigated the results, and wrote the paper with the help of all the coauthors. PBr and OC helped to investigate the results by performing diagnostics
430 and verification computations.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The authors would like to acknowledge the MISTRALS/HyMeX program and the funding from the ANR under contracts IODA-MED ANR-11-BS56-0005 and MUSIC ANR-14-CE01-0014. Jean-Francois Mahfouf and Véronique Ducrocq are acknowledged for their helpful comment on a previous version of the paper. The authors acknowledge principal investigators of data sets : Dr Patrick
435 Chazette (WALI), Dr Cyrille Flamant (LEANDRE), Dr Paolo Di Girolamo (BASIL), Dr Frédérique Saïd (wind profilers), Dr Olivier Bock, Pierre Bosser (reprocessed GNSS data) and Dr Isabelle Taupier-Letage (data from opportunity ships).

References

- Benjamin, S. G., Schwartz, B. E., Szoke, E. J., and Koch, S. E.: The Value of Wind Profiler Data in U.S. Weather Forecasting, *Bulletin of the American Meteorological Society*, 85, 1871–1886, <https://doi.org/10.1175/BAMS-85-12-1871>, <https://doi.org/10.1175/BAMS-85-12-1871>, 2004.
- Berre, L.: Estimation of Synoptic and Mesoscale Forecast Error Covariances in a Limited-Area Model, *Monthly Weather Review*, 128, 644–667, [https://doi.org/10.1175/1520-0493\(2000\)128<0644:EOSAMF>2.0.CO;2](https://doi.org/10.1175/1520-0493(2000)128<0644:EOSAMF>2.0.CO;2), 2000.
- Bielli, S., Grzeschik, M., Richard, E., Flamant, C., Champollion, C., Kiemle, C., Dorninger, M., and Brousseau, P.: Assimilation of water-vapour airborne lidar observations: impact study on the COPS precipitation forecasts, *Quarterly Journal of the Royal Meteorological Society*, 138, 1652–1667, <https://doi.org/10.1002/qj.1864>, 2012.
- Bock, O., Bosser, P., Pacione, R., Nuret, M., Fourrié, N., and Parracho, A.: A high-quality reprocessed ground-based GPS dataset for atmospheric process studies, radiosonde and model evaluation, and reanalysis of HyMeX Special Observing Period, *Quart. J. Roy. Meteorol. Soc.*, 142, 56–71, <https://doi.org/10.1002/qj.2701>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.2701>, 2016.
- Boniface, K., Ducrocq, V., Jaubert, G., Yan, X., Brousseau, P., Masson, F., Champollion, C., Chéry, J., and Doerflinger, E.: Impact of high-resolution data assimilation of GPS zenith delay on Mediterranean heavy rainfall forecasting, *Annales Geophysicae*, 27, 2739–2753, <https://doi.org/10.5194/angeo-27-2739-2009>, <https://www.ann-geophys.net/27/2739/2009/>, 2009.
- Borderies, M., Caumont, O., Delanoë, J., Ducrocq, V., and Fourrié, N.: Assimilation of wind data from airborne Doppler cloud-profiling radar in a kilometre-scale NWP system, *Natural Hazards and Earth System Sciences*, 19, 821–835, <https://doi.org/10.5194/nhess-19-821-2019>, <https://www.nat-hazards-earth-syst-sci.net/19/821/2019/>, 2019a.
- Borderies, M., Caumont, O., Delanoë, J., Ducrocq, V., Fourrié, N., and Marquet, P.: Impact of airborne cloud radar reflectivity data assimilation on kilometre-scale numerical weather prediction analyses and forecasts of heavy precipitation events, *Natural Hazards and Earth System Sciences*, 19, 907–926, <https://doi.org/10.5194/nhess-19-907-2019>, <https://nhess.copernicus.org/articles/19/907/2019/>, 2019b.
- Bouniol, D., Protat, A., Plana-Fattori, A., Giraud, M., Vinson, J.-P., and Grand, N.: Comparison of Airborne and Spaceborne 95-GHz Radar Reflectivities and Evaluation of Multiple Scattering Effects in Spaceborne Measurements, *Journal of Atmospheric and Oceanic Technology*, 25, 1983–1995, <https://doi.org/10.1175/2008JTECHA1011.1>, <https://doi.org/10.1175/2008JTECHA1011.1>, 2008.
- Bresson, E., Ducrocq, V., Nuissier, O., Ricard, D., and de Saint-Aubin, C.: Idealized numerical simulations of quasi-stationary convective systems over the Northwestern Mediterranean complex terrain, *Q.J.R. Meteorol. Soc.*, 138, 1751–1763, 2012.
- Brousseau, P., Berre, L., Bouttier, F., and Desroziers, G.: Background error covariances for a convective-scale data-assimilation system : AROME-France 3D-Var, *Quarterly Journal of the Royal Meteorological Society*, 137, 409–422, <https://doi.org/10.1002/qj.750>, 2011.
- Brousseau, P., Seity, Y., Ricard, D., and Léger, J.: Improvement of the forecast of convective activity from the AROME-France system, *Quarterly Journal of the Royal Meteorological Society*, 142, 2231–2243, <https://doi.org/10.1002/qj.2822>, 2016.
- Caniaux, G., Redelsperger, J.-L., and Lafore, J.-P.: A numerical study of the stratiform region of a fast-moving squall line. Part I: General description and water and heat budgets, *Journal of the Atmospheric Sciences*, 51(14), 2046–2074, 1994.
- Caumont, O. and Ducrocq, V.: What should be considered when simulating Doppler velocities measured by ground-based weather radars?, *J. Applied Meteor. Climatol.*, 47, 2256–2262, <https://doi.org/10.1175/2008JAMC1894.1>, 2008.
- Caumont, O., Ducrocq, V., Wattrelot, E., Jaubert, G., and Pradier-Vabre, S.: 1D+3DVar assimilation of radar reflectivity data: a proof of concept, *Tellus A: Dynamic Meteorology and Oceanography*, 62, 173–187, <https://doi.org/10.1111/j.1600-0870.2009.00430.x>, <https://doi.org/10.1111/j.1600-0870.2009.00430.x>, 2010.

- Chazette, P., Flamant, C., Shang, X., Totems, J., Raut, J.-C., Doerenbecher, A., Ducrocq, V., Fourrié, N., Bock, O., and Cloché, S.: A multi-
 475 instrument and multi-model assessment of atmospheric moisture variability over the western Mediterranean during HyMeX, *Quarterly Journal of the Royal Meteorological Society*, 142, 7–22, <https://doi.org/10.1002/qj.2671>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.2671>, 2016.
- Courtier, P., Freydier, C., Rabier, F., and Rochas, M.: The ARPEGE Project at Météo-France, *ECMWF Seminar Proceedings*, 7, 193–231, 1991.
- 480 Delanoë, J., Protat, A., Jourdan, O., Pelon, J., Papazzoni, M., Dupuy, R., Gayet, J.-F., and Jouan, C.: Comparison of Airborne In Situ, Airborne Radar–Lidar, and Spaceborne Radar–Lidar Retrievals of Polar Ice Cloud Properties Sampled during the POLARCAT Campaign, *Journal of Atmospheric and Oceanic Technology*, 30, 57–73, <https://doi.org/10.1175/JTECH-D-11-00200.1>, <https://doi.org/10.1175/JTECH-D-11-00200.1>, 2013.
- Di Girolamo, P., Flamant, C., Cacciani, M., Richard, E., Ducrocq, V., Summa, D., Stelitano, D., Fourrié, N., and Saïd, F.: Observation of low-
 485 level wind reversals in the Gulf of Lion area and their impact on the water vapour variability, *Quarterly Journal of the Royal Meteorological Society*, 142, 153–172, <https://doi.org/10.1002/qj.2767>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.2767>, 2016.
- Drobinski, P., Ducrocq, V., Alpert, P., Anagnostou, E., Béranger, K., Borga, M., Braud, I., Chanzy, A., Davolio, S., Delrieu, G., Estournel, C., Boubrahmi, N. F., Font, J., Grubišić, V., Gualdi, S., Homar, V., Ivančan-Picek, B., Kottmeier, C., Kotroni, V., Lagouvardos, K., Lionello, P., Llasat, M. C., Ludwig, W., Lutoff, C., Mariotti, A., Richard, E., Romero, R., Rotunno, R., Roussot, O., Ruin, I., Somot, S., Taupier-
 490 Letage, I., Tintore, J., Uijlenhoet, R., and Wernli, H.: HyMeX: A 10-Year Multidisciplinary Program on the Mediterranean Water Cycle, *Bulletin of the American Meteorological Society*, 95, 1063–1082, <https://doi.org/10.1175/BAMS-D-12-00242.1>, <https://doi.org/10.1175/BAMS-D-12-00242.1>, 2014.
- Ducrocq, V., Braud, I., Davolio, S., Ferretti, R., Flamant, C., Jansa, A., Kalthoff, N., Richard, E., Taupier-Letage, I., Ayrat, P.-A., Belamari, S., Berne, A., Borga, M., Boudevillain, B., Bock, O., Boichard, J.-L., Bouin, M.-N., Bousquet, O., Bouvier, C., Chiggiato, J., Cimini, D.,
 495 Corsmeier, U., Coppola, L., Cocquerez, P., Defer, E., Delanoë, J., Girolamo, P. D., Doerenbecher, A., Drobinski, P., Dufournet, Y., Fourrié, N., Gourley, J. J., Labatut, L., Lambert, D., Coz, J. L., Marzano, F. S., Molinié, G., Montani, A., Nord, G., Nuret, M., Ramage, K., Rison, W., Roussot, O., Saïd, F., Schwarzenboeck, A., Testor, P., Baelen, J. V., Vincendon, B., Aran, M., and Tamayo, J.: HyMeX-SOP1: The Field Campaign Dedicated to Heavy Precipitation and Flash Flooding in the Northwestern Mediterranean, *Bulletin of the American Meteorological Society*, 95, 1083–1100, <https://doi.org/10.1175/BAMS-D-12-00244.1>, <https://doi.org/10.1175/BAMS-D-12-00244.1>, 2014.
- 500 Duffourg, F. and Ducrocq, V.: Origin of the moisture feeding the Heavy Precipitating Systems over Southeastern France, *Nat. Hazards Earth Syst. Sci.*, 11, 1163–1178, 2011.
- Duffourg, F., Nuissier, O., Ducrocq, V., Flamant, C., Chazette, P., Delanoë, J., Doerenbecher, A., Fourrié, N., Di Girolamo, P., Lac, C., Legain, D., Martinet, M., Saïd, F., and Bock, O.: Offshore deep convection initiation and maintenance during HyMeX IOP16a heavy precipitation event, *Quarterly Journal of the Royal Meteorological Society*, 142, 259–274, 2016.
- 505 Fan, S., Zang, J.-F., Peng, X.-Y., Wu, S.-Q., Liu, Y.-X., and Zhang, K.-F.: Validation of Atmospheric Water Vapor Derived from Ship-Borne GPS Measurements in the Chinese Bohai Sea, *Terrestrial, Atmospheric and Oceanic Sciences*, 27, 213, [https://doi.org/10.3319/TAO.2015.11.04.01\(A\)](https://doi.org/10.3319/TAO.2015.11.04.01(A)), 2016.
- Fourrié, N., Bresson, E., Nuret, M. and Jany, C., Brousseau, P., Doerenbecher, A., Kreitz, M., Nuissier, O., Sevault, E., Bénichou, H., Amodéi, M., and Poupponneau, F.: AROME-WMED, a real-time mesoscale model designed for the HyMeX special observation periods, *Geoscientific Model Development*, 8, 1919–1941, <https://doi.org/10.5194/gmd-8-1919-2015>, <https://www.geosci-model-dev.net/8/1919/2015/>, 2015.

- Fourrié, N., Nuret, M., Brousseau, P., Caumont, O., Doerenbecher, A., Wattrelot, E., Moll, P., Bénichou, H., Puech, D., Bock, O., Bossier, P., Chazette, P., Flamant, C., Di Girolamo, P., Richard, E., and Saïd, F.: The AROME-WMED reanalyses of the first special observation period of the Hydrological cycle in the Mediterranean experiment (HyMeX), *Geoscientific Model Development*, 12, 2657–2678, <https://doi.org/10.5194/gmd-12-2657-2019>, <https://www.geosci-model-dev.net/12/2657/2019/>, 2019.
- Grzeschik, M., Bauer, H.-S., Wulfmeyer, V., Engelbart, D., Wandinger, U., Mattis, I., Althausen, D., Engelmann, R., Tesche, M., and Riede, A.: Four-Dimensional Variational Data Analysis of Water Vapor Raman Lidar Data and Their Impact on Mesoscale Forecasts, *Journal of Atmospheric and Oceanic Technology*, 25, 1437–1453, <https://doi.org/10.1175/2007JTECHA974.1>, 2008.
- Gustafsson, N., Janjić, T., Schraff, C., Leuenberger, D., Weissmann, M., Reich, H., Brousseau, P., Montmerle, T., Wattrelot, E., Bučánek, A., Mile, M., Hamdi, R., Lindskog, M., Barkmeijer, J., Dahlbom, M., Macpherson, B., Ballard, S., Inverarity, G., Carley, J., Alexander, C., Dowell, D., Liu, S., Ikuta, Y., and Fujita, T.: Survey of data assimilation methods for convective-scale numerical weather prediction at operational centres, *Quarterly Journal of the Royal Meteorological Society*, 144, 1218–1256, <https://doi.org/10.1002/qj.3179>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.3179>, 2018.
- Gutman, S. I., Sahm, S. R., Benjamin, S. G., Schwartz, B. E., Holub, K. L., Stewart, J. Q., and Smith, T. L.: Rapid Retrieval and Assimilation of Ground Based GPS Precipitable Water Observations at the NOAA Forecast Systems Laboratory: Impact on Weather Forecasts, *Journal of the Meteorological Society of Japan. Ser. II*, 82, 351–360, <https://doi.org/10.2151/jmsj.2004.351>, 2004.
- Kouba, J. and Héroux, P.: Precise Point Positioning Using IGS Orbit and Clock Products, *GPS Solutions*, <https://doi.org/https://doi.org/10.1007/PL00012883>, 2001.
- Lindskog, M., Ridal, M., Thorsteinsson, S., and Ning, T.: Data assimilation of GNSS zenith total delays from a Nordic processing centre, *Atmospheric Chemistry and Physics*, 17, 13 983–13 998, <https://doi.org/10.5194/acp-17-13983-2017>, <https://acp.copernicus.org/articles/17/13983/2017/>, 2017.
- Macpherson, S. R., Deblonde, G., Aparicio, J. M., and Casati, B.: Impact of NOAA Ground-Based GPS Observations on the Canadian Regional Analysis and Forecast System, *Monthly Weather Review*, 136, 2727–2746, <https://doi.org/10.1175/2007MWR2263.1>, <https://doi.org/10.1175/2007MWR2263.1>, 2008.
- Mahfouf, J.-F., Ahmed, F., Moll, P., and Teferle, F. N.: Assimilation of zenith total delays in the AROME France convective scale model: a recent assessment, *Tellus A: Dynamic Meteorology and Oceanography*, 67, 26 106, <https://doi.org/10.3402/tellusa.v67.26106>, <https://doi.org/10.3402/tellusa.v67.26106>, 2015.
- Martet, M., Guillaume, F., and Wattrelot, E.: Assimilation of OPERA radar data in AROME-FRANCE NWP MODEL: a challenge, 39th international conference on radar meteorology, 16-20 September 2019, Nara Kasugano International Forum IRAKA, Nara, Japan., https://cscenter.co.jp/icrm2019/program/data/abstracts/Session3B-02_2.pdf, lastaccesson19August2020., 2019.
- Montmerle, T. and Faccani, C.: Mesoscale Assimilation of Radial Velocities from Doppler Radars in a Preoperational Framework, *Monthly Weather Review*, 137, 1939–1953, <https://doi.org/10.1175/2008MWR2725.1>, <https://doi.org/10.1175/2008MWR2725.1>, 2009.
- Pinty, J.-P. and Jabouille, P.: A mixed-phased cloud parameterization for use in a mesoscale non-hydrostatic model: simulations of a squall line and of orographic precipitation. *Proceeding, Conference on Cloud Physics*, Everett, WA., Amer. Meteor. Soc., pp. 217–220, 1998.
- Protat, A., Bouniol, D., Delanoë, J., O’Connor, E., May, P. T., Plana-Fattori, A., Hasson, A., Görsdorf, U., and Heymsfield, A. J.: Assessment of Cloudsat Reflectivity Measurements and Ice Cloud Properties Using Ground-Based and Airborne Cloud Radar Observations, *Journal of Atmospheric and Oceanic Technology*, 26, 1717–1741, <https://doi.org/10.1175/2009JTECHA1246.1>, <https://doi.org/10.1175/2009JTECHA1246.1>, 2009.

- 550 Ricard, D., Ducrocq, V., and Auger, L.: A Climatology of the Mesoscale Environment Associated with Heavily Precipitating Events over a
Northwestern Mediterranean Area, *J. Appl. Meteor. Climatol.*, 51, 468–488, 2012.
- Saïd, F., Campistron, B., Delbarre, H., Canut, G., Doerenbecher, A., Durand, P., Fourrié, N., Lambert, D., and Legain, D.: Offshore winds
obtained from a network of wind-profiler radars during HyMeX, *Quarterly Journal of the Royal Meteorological Society*, 142, 23–42,
<https://doi.org/10.1002/qj.2749>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.2749>, 2016.
- 555 Seity, Y., Brousseau, P., Malardel, S., Hello, G., Bénard, P., Bouttier, F., Lac, C., and Masson, V.: The AROME-France convective scale
operational model., *Mon. Wea. Rev.*, 139, 976–991, 2011.
- Wattrelot, E., Caumont, O., and Mahfouf, J.-F.: Operational Implementation of the 1D+3D-Var Assimilation Method of Radar Reflectivity
Data in the AROME Model, *Monthly Weather Review*, 142, 1852–1873, <https://doi.org/10.1175/MWR-D-13-00230.1>, <https://doi.org/10.1175/MWR-D-13-00230.1>, 2014.