

The Impact of Drought on Soil Moisture Trends across Brazilian Biomes

¹Flavio Lopes Ribeiro, ²Mario Guevara, ²Alma Vázquez-Lule, ³Ana Paula Cunha, ³Marcelo Zeri, ²Rodrigo Vargas

¹University of Delaware, School of Public Policy and Administration, Disaster Research Center, Newark, DE, USA

²University of Delaware, Department of Plant and Soil Sciences, Newark, DE, USA

³National Center for Monitoring and Early Warning of Natural Disasters (CEMADEN), São José dos Campos, SP, Brazil

Correspondence to: Flavio Lopes Ribeiro

Abstract: Over the past decade, Brazil has experienced severe droughts across its territory, with important implications for soil moisture dynamics. Soil moisture variability has a direct impact on agriculture, water security, and ecosystem services. Nevertheless, there is currently little information on how soil moisture across different biomes respond to drought. In this study, we used satellite soil moisture data from the European Space Agency, from 2009 to 2015, to analyze differences in soil moisture responses to drought for each biome of Brazil: The Amazon, Atlantic Forest, Caatinga, Cerrado, Pampas and Pantanal. We found an overall soil moisture decline of -0.5%/year ($p < 0.01$) at the national level. At the biome-level, Caatinga presented the most severe soil moisture decline (-4.4% per year); whereas Atlantic Forest and Cerrado biomes showed no significant trend. The Amazon biome showed no trend but a sharp reduction of soil moisture from 2013 to 2015. In contrast, Pampas and Pantanal presented a positive trend (1.6 and 4.3 %/year, respectively). These trends are consistent with vegetation productivity trends across each biome. This information provides insights for drought risk reduction and soil conservation activities to minimize the impact of drought in the most vulnerable biomes. Furthermore, improving our understanding of soil moisture trends during periods of drought is crucial to enhance the national drought early warning system and develop customized strategies for adaptation to climate change in each biome.

1. Introduction

Drought is a natural and human-induced hazard common to all climate zones in the world (Sheffield and Wood, 2008), generally referred to as a sustained occurrence of below average water availability due to precipitation deficit and soil moisture decline (Magalhães, 2016). Precipitation deficit is the most studied driver of drought (Mishra and Singh 2010; Smith 2013, Villarreal et al., 2016) and has been furthering several drought indicators and models. However, precipitation-based indicators are limited in the assessment of social and environmental responses to the lack of rain and therefore not suitable for evaluating the impacts of drought when used alone. On the other hand, drought indicators based on soil moisture are not only key to understand the physical mechanisms of drought, but also useful for assessing how soil moisture decline can alter vegetation water availability and, consequently, agricultural production and ecosystem services (Smith 2013; NWS 2008).

Definição de estilo: Cabeçalho

Definição de estilo: Rodapé

Formatado: Espaçamento entre linhas: Múltiplos 1,83 lin.

Formatado: Português (Brasil)

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

Soil When soil moisture declines below critical water stress thresholds it reduces biomass production, soil respiration and the overall soil carbon balance (Bot and Benites 2005; Vargas et al., 2018). Low carbon in soils (due to lower biological activity) reduces its structural integrity and increases the risk of soil erosion, contributing to river silting, ineffective runoff control, and loss of soil nutrients (Al-Kaisi and Rattan 2017). Soil moisture is also crucial for addressing the negative impacts of climate change in water and land resources (Bossio 2017). Indeed, temporal variability of soil moisture in a given biome is needed an important variable for the characterization of the local climate (Legates et al. 2011) and a key indicator of changes in the biome's water cycle (Sheffield and Wood 2008; Rossato et al. 2017).

In this study, we use satellite data from the European Space Agency (ESA) to analyze the impact of drought on soil moisture across all Brazilian biomes: The Amazon, Atlantic Forest, Caatinga, Cerrado, Pampas and Pantanal. Considering that each biome has distinct climate, soil and vegetation characteristics, we hypothesize that they would respond differently to drought conditions (e.g., positive, negative or non-significant) and show up relevant information for drought management at national and regional levels.

In Brazil, most of the work on drought management has been focused in the semiarid region, well-known for its recurrent problems with droughts and water scarcity (Fig. 1) and where predominates the Caatinga biome. However, droughts have been reported all over Brazil, affecting all other biomes as well. In the period selected for this study (i.e., 2009 to 2015), there was a high number of municipalities declaring emergency and even public calamity due to drought across the country (Cunha et al. 2019), but the impacts on soil moisture at national scale and how each biome responds to drought are still unknown.

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita




Figure R1 A perspective of the Caatinga forest during the dry season at the ground level (A). A perspective of land use in the Caatinga biome during the wet season at the landscape level (B). An example of human intervention to river course that has an impact on water availability across the region (C). **Figure 1: The Caatinga biome (Pontes, 2012).**

Due to climate change, extreme events such as drought are expected to become more intense and recurrent in some regions of Brazil. Therefore, integrating satellite soil moisture data into early warning systems could contribute to more efficient drought risk management and promote data-driven climate change adaptation.

Nevertheless, studies on soil moisture variation have been conducted at a stand-scale due to challenges for measurements across spatial and temporal scales (Legates et al. 2011; Novick et al 2016). As a consequence, the lack of soil moisture information could lead to inaccurate assessment of drought conditions, underestimation of drought impacts, and incomplete resilience and adaptation plans. As droughts become more frequent and intense, it is important to enable monitoring of soil moisture trends and communicate the results at different levels (e.g., municipal, state, national, regional) and across different perspectives (e.g., environmental, social, and economic). At present, the most reliable source of soil moisture information at large-scales (i.e., global-to-continental scales) is satellite remote sensing (i.e., <https://smap.jpl.nasa.gov/>, <http://www.esa-soilmoisture-cci.org/>), which provides soil moisture estimates for the first 0-5 cm of soil depth (Liu et al. 2011). Even though the first layer of soil is

Formatado: Fonte: (Padrão) Times New Roman, 9 pt

Formatado: Fonte: 9 pt

Formatado: Fonte: Não Negrito

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

expected to be very dynamic because of its interaction with the atmosphere and deeper layers still represent an important water storage, especially in the Amazon and Cerrado biomes, soil moisture at the first 5cm is still a good predictor of land and atmosphere interactions. Analyzing a shallow soil layer can provide key information for the detection of soil aridity conditions that are directly related with the loss of soil biodiversity and, therefore, with soil productivity. Thus, soil moisture at the surface is directly affected by drought conditions and could be also used as an indicator (i.e., proxy) of the water contained at deeper layers. Since we cannot measure *in situ* soil moisture at high spatial resolution due to logistical constraints (i.e., because is expensive or time consuming), we propose the use of multiple satellite remote sensing sensors (e.g., from ESA or NASA) as an alternative to obtain drought-relevant information on soil moisture at the national scale. The use of satellite-derived soil moisture has been demonstrated for evaluating soil moisture dynamics from regional to national scales (Guevara and Vargas 2019). The study period (2009 – 2015) was marked by successive droughts across Brazil, registered and confirmed by different monitoring instruments such as the Integrated Drought Index (IDI), which combines the Standardized Precipitation Index (SPI) and the Vegetation Health Index (VHI) (Cunha et al., 2019) and Municipal Emergency Declarations all over the country.

Most of the work has been focused on the semiarid region of Brazil, in the Caatinga biome, for its well-known recurrent problems with droughts and water scarcity (Fig. 1). The purpose of this study is showing the advantages and disadvantages of integrating satellite soil moisture observations into drought monitoring across Brazil on a biome basis. We show the differential impact of drought on the soil moisture of different biomes at a national scale (using Brazil as a case study).

Main limitations are that satellite measurements of soil moisture provide indirect estimates of soil moisture across large areas of around >25km grids and that these estimates are representative only in the topsoil (eg., 0-5cm), and unfortunately do not provide a direct metric of soil water storage. While soil moisture at the surface is a key indicator of soil and atmosphere interactions, topsoil moisture does not account entirely for the water used by plants to grow. The capacity of plants to grow can be measured also with satellite information in the form of primary productivity estimates (Li et al., 2019). Therefore, we also explore the correspondence between satellite soil moisture and primary productivity trends for each biome in Brazil. Both soil moisture and vegetation productivity are ecosystem variables directly affected by drought conditions. Understanding how soil moisture and vegetation productivity on each biome is affected by drought conditions from different perspectives (in our case superficial soil moisture) is crucial to assess their resilience. It is also important to provide evidence-based orientations to drought mitigation and soil conservation plans.

However, droughts have been reported all over Brazil, affecting all other biomes as well. In the period selected for this study (i.e., 2009 to 2015), there was a high number of municipalities declaring emergency and even public calamity due to drought all over the country (Cunha et al., 2019), but the impacts on soil moisture at national scale and how each biome responds to drought are still unknown.

Formatado: Fonte: Itálico

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita



Figure 1: The Caatinga biome (Pontes, 2012).

In this study, we use satellite soil moisture data from the European Space Agency (ESA) to analyze the impact of drought across all Brazilian biomes: The Amazon, Atlantic Forest, Caatinga, Cerrado, Pampas and Pantanal. Brazil is the water richest country in the world, with estimated total freshwater resources of $8233 \text{ km}^3 \text{ yr}^{-1}$ (Holden 2014). Just in terms of comparison, Russia, second country in the list, has around $4507 \text{ km}^3 \text{ yr}^{-1}$ (Holden 2014). This great amount of freshwater resources, however, does not spare Brazil from suffering with recurrent droughts influenced by both natural and human processes (Loon et al. 2016), putting drought as the number one disaster in terms of economic losses and number of people affected in the country (CEPED 2012; CENAD 2014).

Considering that each biome has distinct climate, soil and vegetation characteristics, we hypothesize that they would respond differently to drought conditions (e.g., positive, negative or non-significant) and show up relevant information for drought management at national and regional levels. Considering that due to climate change, extreme events such as drought can become more intense and recurrent also in some regions in Brazil, understanding these differences and integrating satellite soil moisture data into early warning systems could also contribute to more efficient drought risk mitigation actions and promote data-driven climate change adaptation.

2. Methodology

2.1. Study area

Brazil is the largest country in Latin America with a total area of $8,456,510 \text{ km}^2$, located between $05^{\circ}10' \text{ N}$ to $33^{\circ}44' \text{ S}$ (IBGE, 2017). The continental dimension of the country implies a complex spatial heterogeneity of environmental conditions resulting in six main biomes: Amazon, Atlantic Forest, Caatinga, Cerrado, Pampas and Pantanal (Fig. 3a).

The Amazon biome is mainly characterized by rainforest areas (Overbeck et al. 2015). It represents 49.5% of Brazil's total area, or $4,196,943 \text{ km}^2$ (IBGE, 2019). It has an equatorial climate, with temperatures between 22°C and 28°C and torrential rains distributed throughout the year. The geomorphology of the Amazon biome is quite diverse, presenting plateaus, plains, and depressions. Soils are generally clayey, iron-rich and with high soil

Formatado: Fonte: Não Negrito

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

organic carbon content. The Amazon biome is well known for its biodiversity and its large number of rivers and water bodies, which account for the world's greatest surface green water reserves (IBGE 2004).

The Atlantic Forest biome covers 13% of the total area of Brazil (1,110,182 km²). It comprises an environmental heterogeneity that incorporates high elevations, valleys, and plains. The Atlantic rainforest occupies the whole continental Atlantic coast of Brazil. This biome has a subtropical climate in the south and a tropical climate in central and northeast portions. The Atlantic rainforest is characterized by heavy rainfall influenced by the proximity of the ocean and winds that blow inward over the continent (IBGE, 2004). Although it is just a small fraction of the size of the Amazon rainforest, the Atlantic Forest still harbors a range of biological diversity comparable to that of the Amazon biome (The Nature Conservancy, 2015), with high soil carbon reserves (Guevara et al., 2018). The Atlantic Forest is recognized as the most degraded biome of Brazil with only 12% of the original biome preserved (SECOM, 2012).

Caatinga is the driest biome of Brazil and comprises an area of 844,453 km² stretching over nine federal states and covering nearly 10% of the total area of Brazil (IBGE, 2019). Semiarid climate is predominant across this biome (BSh type) with an average annual rainfall below 800 mm (Alvares et al., 2013), but high temperatures influence high potential evapotranspiration rates that exceed 2,500mm/year (Campos, 2006). Overall, the Caatinga is characterized by reduced water availability and a very limited storage capacity of rivers, which are mainly intermittent, with just a few exceptions that are perennial through streamflow regulating reservoirs during the dry season (CENAD 2014). Caatinga soils are generally shallow (0-50 cm), with a bedrock that is commonly exposed to the surface, limiting water infiltration processes and the recharge of local aquifers (Cirilo, 2008).

The Cerrado is the second largest biome of Brazil, characterized by large savannas (Overbeck, et al 2015) covering 2,036,448 km², and representing 23.3% of the country (IBGE, 2019). It extends from the central south of Brazil until the north coastal strip, interposing between the Amazon, Pantanal, Atlantic Forest, and the Caatinga biomes (IBGE, 2004). The dominant climate in the Cerrado is warm tropical sub-humid, with only two distinct seasons, dry winters and wet summers with torrential rains (Overbeck et al. 2015). The annual precipitation in this region varies between 600-2200 mm, where the bordering areas with the Caatinga are the driest and the bordering areas with the Amazon rainforest the wettest. Soils are diverse and include a variety of dystrophic (low inherent fertility and/or strongly weathered profile), acidic, and aluminum-rich conditions. Currently, the Cerrado hosts the largest rural expansion in Brazil, resulting in environmental degradation, biodiversity loss, and soil erosion and limited water availability. It is classified as the most endangered savannah on the planet and one of the 34 global hotspots (Ioris, Irigaray and Girard 2014).

The Pampas biome is located at the extreme south of Brazil and covers 2.1% of Brazil's total area (176,496 km²). It is mainly characterized by grasslands and shrublands (Overbeck et al. 2015). The region has a wet subtropical climate, characterized by a rainy climate throughout the whole year, with hot summers and cold winters, where temperatures fall below freezing (IBGE 2004). The Pampas comprises an environmental set of different lithology types and productive soils (e.g., carbon-rich), mainly under flat and smooth undulating terrain surfaces.

Pantanal is the biome with the smallest territorial extension of Brazil, covering 1.8% (150.355 km²) of the country's total area (IBGE, 2004). It is located at the left margin of the Paraguay River and shared by Brazil, Bolivia and Paraguay.

The Pantanal is by a vast extent of poorly drained lowlands that experiences annual flooding from summer to fall months (January–May) (Assine and Soares, 2004). The climate of the Pantanal is hot and humid during the

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

summer and cold and dry in winter (Ioris, Irigaray and Girard 2014). Precipitation varies from 1000-1400 mm per year, and rains are predominant from November to April. Average annual temperature is 32°C, but the dry season (May to October) has an average temperature of 21°C and it is not uncommon to have >100 days without rain (Ioris, Irigaray and Girard 2014). In the last two decades, temperature in the Pantanal has consistently risen and more humid than normal events as well as dryer than normal events have both increased (Marengo et al 2010).

2.3. Environmental variability of Brazilian Biomes

We used 1x1 km environmental gridded data to characterize the environment variability of the biomes. Data was provided by worldgrids.org, an initiative of ISRIC – World Soil Information Institute. This dataset compiled information from: 1) digital terrain analysis to represent topographic gradients, 2) gridded climatology products (e.g., precipitation and temperature), 3) remote sensing imagery, to represent land cover and vegetation spatial variability, and 4) legacy soil or rock type maps. We used 110 layers derived from this dataset. A list of all available information contained in the worldgrids.org project is available at Reuter & Hengl (2012). We used multivariate statistics in the form of principal component analysis (PCA) to linearly decompose the worldgrids.org dataset and identify relationships among the major environmental characteristics of Brazilian biomes. PCA is an analysis where a group of potentially correlated variables are decomposed in orthogonal space and therefore uncorrelated principal components. PCA analysis is useful to reduce data dimensionality to avoid the potential effects of statistical redundancy (multicollinearity) in further interpretations. Here, we use the PCA as an exploratory technique to visualize/characterize/interpret the environmental variability of Brazil's biome and assume that environmental differences in the biomes could support the hypothesis of different soil moisture response to drought.

2.4. Municipal emergency declarations due to drought across Brazil

Municipal Emergency Declarations (**EDsMEDs**) are administrative tools to inform the federal government that the magnitude of the disaster has surpassed local public capacities to respond and manage the installed crisis. The recognition of **EDsMEDs** by the federal government is based on field visits (when possible) and technical analysis of social, economic and climatological data that can support the petition. In the case of drought, data analysis is generally based on, but not limited to, private agricultural losses, level of local reservoirs, and precipitation data combined. Once the federal government recognizes that there is indeed a disaster, it establishes a legal situation where federal funds can be used to assist the affected population and recover essential services disrupted by the disaster (National Secretary of Civil Defense and Protection of Brazil 2017).

To determine drought distribution across the six Brazilian biomes, we retrieved official **EDsMEDs** due to drought in Brazil from 2009 to 2015. This information is public and can be accessed [in on](#) the website of the Ministry of National Integration of Brazil-. First, we downloaded the historical series of **EDsMEDs** in Brazil from 2009 to 2015. Then, we isolated the municipalities who declared emergency or public calamity due to drought from all other disasters. The last step was to cross this data with the boundaries of the six Brazilian biomes and discover the intensity and distribution of drought in each biome during the study period.

2.5. Soil Moisture **and Primary Productivity** Trends across Brazil

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

To analyze soil moisture trends during a period of successive droughts (2009-2015) across Brazilian biomes, we acquired remotely sensed soil moisture information from the European Space Agency (Liu et al. 2011). This soil moisture product has a daily temporal coverage from 1978 to 2016 and a spatial resolution of 0.25 degrees (~27x27 km grids). To represent vegetation primary productivity we use estimates from the OCO-2-based SIF product (GOSIF) and linear relationships between SIF (Solar-induced chlorophyll fluorescence) and GPP (gross primary production) used by Li and Xiao (2019) to map GPP globally at a 0.05° spatial resolution and 8-day time step. We calculated monthly averages from ~~original~~ soil moisture ~~data~~ and primary vegetation datasets for further statistical analysis using only information between 2009 and 2015. All available information was harmonized into a geographical information system using the same projection system and spatial integrity.

2.6. Data Analysis

We based our statistical analysis in a regression matrix containing 10,000 representative random spatial locations (e.g., latitude and longitude) across the biomes of Brazil (Fig. 3b) which were selected using standard re-sampling techniques (i.e., bootstrapping). Over 30% of the area for every biome is represented in the random selection. We randomize our statistical sampling with the ultimate goal of maximizing the accuracy of the results. We used a representative sample for improving the visualization of points cloud and a better understanding of differences on the five biomes in the statistical multivariate space. Finally, we extracted to these random points the environmental data and the values of the available satellite soil moisture and primary productivity time series.

To detect trends on ~~monthly~~ satellite soil moisture ~~data~~ and primary productivity time series during the study period, we used median based linear models calculated for each point with available satellite data. These non-parametric analyzes are known as Theil – Sen regressions (Sen 1968; Theil 1992) with repeated medians (Siegel 1982). This method uses a robust estimator for each point in time, where the slopes between it and the other points are calculated (resulting n-1 slopes), and then the median and the significance of the trend are reported.

The satellite soil moisture source has intrinsic quality limitations across areas where vegetation has more water than soil (McColl et al. 2017), including areas across the lower Amazon watershed, the Pantanal or the Pampas biomes. For these areas we used the sparse points with available satellite soil moisture information and generated predictions of soil moisture trends based on geostatistical analyses, such variogram fitting and Ordinary-Kriging modeling. Ordinary-Kriging assumes that the target variable (soil moisture trends) is controlled by a random field (main reason why we base our analysis in a random sampling strategy) and that shows a quantifiable level of spatial structure and autocorrelation (Hiemstra et al. 2009). We performed an automatic variogram analysis to assess the spatial structure and autocorrelation of satellite soil moisture records. For the variogram analysis we computed the relationships between the distance of randomly distributed soil moisture observations and the accumulated variance of their respective values. We used the aforementioned relationships to predict the satellite soil moisture trend in areas where no data is available and also provided a spatial explicit measure of error following a geostatistical framework (Hiemstra et al. 2009, Llamas et al., 2020). In contrast, the primary productivity dataset used here has complete coverage across Brazil. We show both the interpolated maps of soil moisture trends and the trend map of the primary productivity of vegetation.

3. Results and Discussion

3.1. Drought in Brazil from 2009 to 2015


8
8

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

This analysis of Municipal emergency declarations (EDs) due to drought in Brazil (Emergency Declarations (MEDs)) confirmed that the period from 2009 to 2015 was, indeed, marked by successive droughts countrywide (Fig. 2). During this period, Brazil had a total of 12,508 declarations of emergency or public calamity due to drought all over its territory (Ministry of National Integration of Brazil 2018), which affected directly 33 million people and caused economic losses around US\$ 6,5 billion (EM-DAT 2018).

Proportionally, Caatinga is the biome with more EDs/MEDs per municipality, followed by the Atlantic Forest, Cerrado, Pampas and the Amazon respectively (Fig. 2). The only biome with no EDs/MEDs due to drought during this period is the Pantanal, which is a natural wetland that covers only 1.8% of the national territory (Overbeck et al. 2015).



Formatado: Cor da fonte: Preto
Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita




Figure 2: Percentage of municipalities declaring emergency or public calamity due to drought in Brazil from 2009 to 2015

When considering climatological data from the Integrated Drought Index (IDI), which combines the Standardized Precipitation Index (SPI) and the Vegetation Health Index (VHI), Cunha et al. (2019) discovered that since 1962, when drought events started to be recorded in Brazil, only between 2012 and 2014 droughts occurred concurrently in the six biomes of the country. The IDI also showed that the hydrological year of 2011/2012 (October 2011 to September 2012) was the driest of the historical series, except in the South region, where the Pampas biome is located. During the period of study (2009-2015), the most severe drought events occurred in the northeast region (where the Caatinga predominates), in the central west region (where the Cerrado predominates), and in the southeast region (where there is a mix of Cerrado and Atlantic Forest). Even though the climatological data from the IDI show some inconsistencies with the ~~EDsMEDs~~ per biome, in general terms, it reinforces that the study period was marked by simultaneous droughts across all biomes of Brazil.


3.2. Environmental gridded information of Brazilian Biomes

The environmental characterization of Brazilian biomes showed a clear differentiation of three major groups (Fig. 3a and b). These results support the expectation that drought would have a differential impact on soil moisture dynamics in each of the six biomes (see section 3.3). This expectation is supported because each biome shows

Formatado: Cor da fonte: Preto
Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

differences on the spatial configuration of environmental soil moisture drivers, as revealed by the PCA analysis (Fig. 3b) as described below.

From the 110 environmental layers of information we used to represent the major environmental conditions across Brazil (see list of available layers in <http://worldgrids.org/doku.php>), at least 50 principal components were needed to capture >80% of total variance. The first and second component explained >25% of variability (Fig. 3b) and the variables that represented most of the variance in the first and second components were the digital elevation model ($r=0.5$) and the topographic wetness index ($r=0.31$) respectively. These two variables are directly related to the spatial variability of soil moisture dynamics as seen in other regional studies (Guevara and Vargas 2019). Across these principal components (i.e., PC1 and PC2), we found a clear separation of three major groups of data in the statistical space (Fig. 3c). The Amazon biome forms the larger group of values, followed by another group composed mainly by the Atlantic forest and the Pampas. The Caatinga and Cerrado biomes form a third larger group and the remaining Pantanal show a close but independent variability (Fig. 3c). These groups are located on different quadrants of the plane between the first two PCs (Fig. 3c). Thus, these differences could influence soil moisture response in these major groups at the biome level.



Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita




Figure 3: (a) The six biomes of Brazil. (b) Plane of the first and second PCAs showing the orthogonal and environmental variability of Brazil's biomes and (c) Clustering results showing the main values of each biome dataset and their proximity across the planet between PCAs one and two.

Formatado: Fonte: Não Negrito

Formatado: Espaço Depois de: 0 pt, Espaçamento entre linhas: 1,5 linhas

3.3. Drought assessment: Soil Moisture Trends Across Brazilian Biomes

Our analysis of satellite soil moisture at national level showed a soil moisture decline of -0.5% per year ($p < 0.1$) in Brazil from 2009 to 2015 (Fig. 4). ~~The years with the most accentuated soil moisture decline were 2012 and 2015. Our data also shows two clear moments of soil moisture variation at the national level, which can be divided in two blocks before and after 2012 (Fig. 4). Before 2012, there was a greater variance of soil moisture (2009, 2010, 2011) with rapid increases followed by also rapid declines. After 2012, there is less variance in absolute values of soil moisture, with a short recovery phase of soil moisture values from 2012 to 2013 followed by an abrupt decline by the end of 2014 and 2015.~~

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita






Figure 4: Brazil soil moisture trend from 2009 to 2015

Formatado: Fonte: Não Negrito

When considering variations of soil moisture per biome, our data suggests that the largest soil moisture decline in Brazil was found in the Caatinga biome with a persistent negative trend (-4.4% in soil moisture per year ($p < 0.001$)) from 2009 to 2015 (Fig. 5a). In contrast, Amazon, Cerrado and Atlantic Forest biomes showed no significant trend on soil moisture. Pampas and Pantanal biomes showed a significant increase in soil moisture of 1.6% and 4.3% respectively per year ($p < 0.001$) during the same period (Fig. 5e and f). Thus, the combination of environmental variables and satellite soil moisture records was able to identify drought dominated areas such as Caatinga and Cerrado from water-surplus dominated areas, such as Pantanal and Pampas. These results are also useful to prevent agricultural risk from water failure (decline or surplus) and monitor important ecosystem services of large and more inaccessible areas such as the Amazon forest and the Cerrado (Fig. 3).

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita



Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita




Figure 5: Soil moisture trends across Brazil. (a) Caatinga (n=921), (b) Cerrado (n=2410), (c) Atlantic Forest (n=1394), (d) Amazon (n=4819), (e) Pampas (n=231), and (f) Pantanal (n=179). The values in every graph show the slope percentages of changes. Red solid line showed the mean trend and red dashed lines show the standard deviation trend. * (p<0.01)**

A closer analysis of satellite soil moisture trend in the Caatinga biome shows that this biome did not fully ~~recovered~~ recover from an ~~accentuate~~ accentuated soil moisture decrease in 2012 (Fig. 5a). After 2012, there was a slight recovery of soil moisture in 2013, yet a negative trend remains in the following years, most likely because

Formatado: Português (Brasil)

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

the below average annual precipitation from 2013 to 2015 (Cunha et al., 2019) coupled with human activities commonly found within the boundaries of this biome such as deforestation, unsustainable irrigation and water abstraction (Medeiros 2012; Travassos and De Souza, 2014). As highlighted by Cunha et al. (2015) intense drought events can reduce the vegetation resiliency, rendering plants to be more vulnerable to a recurring disturbance. Furthermore, the vegetation can be durably affected by a drought, if the drought is preceded by another dry year that could substantially reduce gross primary productivity and other ecosystem processes (Vargas, 2012).

Consistent with previous studies (Zeri et al. 2018) precipitation data indicates that the years 2011, 2012, 2014 and 2015 have been drier as compared to the previous decades. Marengo et al. (2017) also confirmed that, from 2012 to 2015, drought affected hundreds of cities and rural areas with devastating impacts on the agricultural production and water supply. On the human activities side, data from the National Institute of Spatial Research (INPE, 2018) reveals that 45% of the Caatinga biome is degraded and 7.2% of its soil is already exposed. In addition, the Caatinga has been exposed to continuous land cover changes and less than 1% of the region is a strictly protected area (Leal et al., 2005; Morim et al., 2013). Thus, our results: (a) provide insights to identify geographical areas that could be preserved due to its capacity for providing blue and green water; and (b) could be part of a monitoring system for optimizing the limited water inputs and supply in this semiarid ecosystem (i.e., for agricultural planning).

Persistent and prolonged soil moisture decline could also negatively affect Caatinga's biodiversity, one of the world's plant biodiversity centers (Leal et al. 2005). The vegetation and soils of the Caatinga are exposed to 8-10 dry months per year (Santos et al. 2014), and more than 90% of the Caatinga biome is non-forest vegetation. Just ~20% of the biome has native vegetation, which is better adapted to support drought events and store higher amounts of water (Santos et al. 2014; Overbeck et al. 2015). Tomasella et al. (2018) using NDVI values for high density vegetation and bare soil showed that recurrent droughts are accelerating the degradation and desertification processes in the Caatinga.

The combination of these regional factors together with the effect of teleconnections such as the ENSO (El Niño Southern Oscillation) and other land atmosphere interactions (Kouadio et al. 2012) make the Caatinga biome in Brazil the most vulnerable biome to the recurrent droughts and consequently, prolonged soil moisture deficit condition. (Marengo et al. 2017).

Therefore, we highlight the need to include urgent actions such as reforestation and efficient use of underground water into drought mitigation plans for this biome to reduce future soil moisture decline. It is noteworthy that this biome is already presenting agricultural deficits and desertification areas due to natural and anthropogenic phenomena (Nascimento and Alves 2008; Sheffield and Wood 2008; Medeiros, 2012; Travassos and De Souza 2014). As an example, while studying the desertification process in part of the Caatinga biome, D' Souza, Fernandes and Barbosa (2008) found high levels of social, economic, and technological vulnerabilities which could be directly associated with removal of the natural vegetation covering and forest fires for subsistence agriculture. These human induced changes on soil moisture in the Caatinga are also related with the occurrence of soil erosion and local desertification processes that influence low agricultural productivity due to diminish soil moisture and quality of the soil (Nascimento and Alves 2008).


The Atlantic Forest biome didn't show significant positive or negative trends in soil moisture variation during the studied period. It registered, however, the greatest ups and downs in soil moisture from 2009 to 2015, with high

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

peaks (2009, 2011 and 2013) followed by abrupt declines in a relatively short time period. After the most intense period of soil moisture decline in the Atlantic Forest (2009-2012), this biome quickly bounced back to previous levels of soil moisture, showing capacity to recover from intense soil moisture losses in less than 12 months.

The Amazon biome showed no significant trend of satellite soil moisture data during the analyzed period (Fig. 4d), probably due to data limitations (i.e., data gaps) associated with lack of satellite-derived information (see Methods section). Field-based evidence collected by Anderson et al. (2018) showed a wide range of impacts of drought on the Amazon forest structure and functioning (e.g.: widespread tree mortality and increased susceptibility to wildfires) in 2016 after the 2015 drought, which affected approximately 46% of the Brazilian Amazon biome. However, considering the size and differences in topography in the Amazon biome, the eastern and western areas of the Amazon ~~rain-forest~~rainforest may respond differently to drought due to differences in climate conditions and therefore, different sensibility to soil moisture decline. The western portion of the Amazon biome shows higher soil moisture values (and potentially positive soil moisture trends) than the eastern region (Fig. 6a and b). This result is consistent with previous findings describing differences in drought response from east and west portions of this biome (Duffy et al. 2015), suggesting that soil moisture conservation plans and drought mitigation strategies in the Amazon biome should consider the heterogeneity of the region and the different soil moisture feedback from the east and west portions of this biome.



Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita




Figure 6: Geostatistical analysis (Ordinary-Kriging with automatic variogram fitting) of satellite soil moisture across Brazil from 2009 to 2015. (a) The trend prediction of soil moisture 2009-2015. (b) The kriging variance (error map), (c) Variogram fitting parameters and spatial autocorrelation model (blue line) supporting the soil moisture prediction. The numbers around the blue line are the pairs of points available for the interpolation at a specific distance (x-axis)

Formatado: Espaço Depois de: 0 pt, Espaçamento entre linhas: 1,5 linhas

The Pampas biome showed a positive trend of ~1.6% per year ($p < 0.001$) during the analyzed period (Fig. 5e), but with three distinct periods. The year 2009 registered a recovery period of positive soil moisture trend followed by a steady soil moisture decline until its lowest point in the beginning of 2012. Then, this biome started a consistent recovery process surpassing previous values of soil moisture trend registered before 2013, showing great capacity to recover soil moisture after periods of drought. Cunha et al. (2019) showed that in 2012 most of the south region of Brazil presented drought conditions over an extensive area, with the highest intensity recorded in August 2012. This intense drought affected the water supply in the rural properties and the agricultural and livestock production.

Even though the Pampas has more than 60% of its biome degraded, especially for cattle raising (Santos and Silva 2012), our data shows that it is gradually increasing soil moisture even during a period of successive droughts across Brazil. Literature on soil moisture of the Pampas biome characterizes this biome as highly vulnerable to water and wind erosion (Roesch et al. 2009), making it susceptible to soil moisture decline (Duffy et al. 2015). On the other hand, extended flat landscapes, like the Pampas, show low lateral water transport as a result of low

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre: (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

surface runoff and slow groundwater fluxes, making this biome more suitable to accumulate surface water for long periods of time (Kuppel et al. 2015).

The Pantanal biome also showed a positive soil moisture trend of 4.3% per year ($p < 0.001$) from 2009 to 2015, the highest positive trend among all biomes. From 2009 to 2011, there were two extreme events characterized by sudden soil moisture increase immediately followed by abrupt soil moisture declines. After these two extreme events, a more stable and consistent positive soil moisture trend was registered from 2011 to 2014. Even though there was a subtle decline in the soil moisture by the end of 2014, this biome kept an overall positive trend during 2015.

The Pantanal and the Pampas biomes are both sub-humid aeolian plains, which make them more susceptible to experience flood events covering a significant fraction of the landscape for months or even years (Kuppel et al. 2015). Even though our data seems congruent with inundations registered in Pantanal in the beginning of 2011, when soil moisture trend reached its highest point for the Pantanal biome during the studied period, it did not capture a reduction of 81% of the total flooded area for the Pantanal biome in 2012, when there was a reduction of 18% in annual precipitation (Moraes, Pereira and Cardozo 2013). In contrast, our data showed a consistent positive trend throughout 2012, even though all months of the wet season in 2012 had a decrease in precipitation ranging from -28.6% in the beginning towards -12.1% in the end of the wet season (Moraes et al. 2013). These results suggest that, although the analyzed period is characterized by a sequence of dry spells across Brazil (Marengo et al. 2017), some areas such as the Pantanal region, were able to accumulate soil moisture during that time.

Detecting an increase in soil moisture does not mean that these biomes should receive less attention to drought and soil conservation plans. From 2009 to 2015, the Pampas had always a representative municipality declaring emergency due to drought and has constantly ~~report~~reported economic losses in the agricultural sector. The Pantanal, during the same period, was not directly impacted by drought at the municipal level, but the highly positive soil moisture trend deserves further understanding on how it impacts the local ecosystem, as well as agricultural practices and cattle raising with the ultimate goal to improve food security across Brazil.

Our results support our main hypothesis as we have found evidence that each of the six Brazilian biomes registered different soil moisture feedbacks to drought during the analyzed period (2009-2015). In practical terms, it means that drought response and mitigation plans, as well as soil conservation strategies should consider both differences among and within each biome of Brazil and concentrate efforts and resources to preserve or recover the regions with greater susceptibility to lose soil moisture during periods of drought. Confirming the value of satellite soil moisture signals monitoring drought related patterns, we observe the similar trends of soil moisture and the primary productivity of vegetation across Brazil.

3.4. Primary productivity trends across Brazil

We confirm the consistency of our results comparing trends of satellite soil moisture with trends calculated using the primary productivity (or GPP) datasets. Our results show that all biomes experienced positive and negative trends of vegetation productivity between the analyzed period of time (Fig. 7). We observe that the major surface of negative trends of primary productivity of vegetation is across the Caatinga biome and its intersection with both the Cerrado and Atlantic Forest biomes. Pampas and Pantanal are the biomes with higher surface of positive primary productivity trends (Fig. 7).

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita




Fig. 7. Trends of primary productivity of vegetation based on the GOSIF dataset between 2009-2015. Areas in black showed non-significant.

These results are consistent with the soil moisture trends described on each biome (Fig. 5). Caatinga is the biome with highest soil moisture decline and highest primary productivity decline. Cerrado and the Atlantic forest are biomes also experiencing decline in soil moisture and primary productivity. In contrast, the Pampa and Pantanal experienced an increase in soil moisture levels and increase in primary productivity rates (Fig. 8). Changes in primary productivity across the Amazon forest were less evident or not significant. Our results support the use of satellite soil moisture and primary productivity trends as accurate indicators of drought conditions across Brazilian biomes.

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita




Fig. 8 Primary productivity trends across Brazilian biomes based in the GOSIF-GPP product across the analyzed period of time (2009-2015).

4. Conclusion

The results of this research reveal an important environmental vulnerability to drought across Brazil. From 2009 to 2015, there was a national decline of soil moisture with a rate of 0.5% year⁻¹. Among all six biomes, Caatinga presented the most severe soil moisture decline (-4.4% year⁻¹), suggesting a need for immediate local soil and water conservation activities. The Atlantic Forest and Cerrado biomes showed no significant soil moisture trends but should be closely monitored for its importance to national food and water security and environmental balance. The Amazon biome also showed no soil moisture trend but a sharp reduction of soil moisture from 2013 to 2015. It is noteworthy that soil moisture from eastern and western portions of the Amazon biome may respond differently to drought. The western portion of the Amazon biome shows potentially more positive soil moisture trends than the eastern region. In contrast, the Pampas and the Pantanal biomes presented a positive soil moisture trend (1.6 and 4.3 % year⁻¹, respectively), which should also be constantly monitored considering the susceptibility of these biomes to floods. **Finally, this**

These results are consistent with primary productivity trends (Fig. 8), supporting the effectiveness of satellite soil moisture data to monitor drought impacts at a biome level. This study provides insights about the potential benefits of integrating satellite soil moisture data into drought monitoring and early warning systems and soil conservation plans at national and local levels.

Formatado: Fonte: Negrito

Formatado: Espaço Depois de: 8 pt, Espaçamento entre linhas: Múltiplos 1,08 lin.

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

Acknowledgments

FR acknowledges individual scholarship support from CNPq, Science without Borders program, Brazilian Federal Government. MG and AVL acknowledge individual fellowship support from CONACyT. RV acknowledges support from the National Science Foundation CIF21 DIBBs (Grant #1724843).

References

Al-Kaisi, M., Rattan, L.: Conservation Agriculture Systems to Mitigate Climate Variability Effects on Soil Health, in: Al-Kaisi, M. Lowery, B. (Eds). *Soil Health and Intensification of Agroecosystems*. Academic Press, 79-107, doi.org/10.1016/B978-0-12-805317-1.00004-X, 2017.

Anderson, L. O., Ribeiro Neto, G., Cunha, A. P., Fonseca, M. G., Mendes de Moura, Y., Dalagnol, R., et al. Vulnerability of Amazonian forests to repeated droughts. *Philos. Trans. R. Soc. B*. 373:20170411. doi: 10.1098/rstb.2017.0411, 2018

Assine, M.L., Soares, P.C.: Quaternary of the Pantanal, west-central Brazil. *Quatern Int* 114: 23–34, 2004.

Behling, H., Jeske-Pieruschka V., Schüller, L., Pillar, V.: Dinâmica dos campos no sul do Brasil durante o Quaternário Tardio. In Pillar VD, Müller SC, Castilhos ZMS Jacques AVA (eds). *Campos Sulinos: Conservação e Uso Sustentável da Biodiversidade*. Brasília: Ministério do Meio Ambiente, p. 13-25, 2009.

Bossio, D.: Soil Management – A Foundational Strategy for Conservation. The Nature Conservancy. Retrieved from <https://global.nature.org/content/soil-management-a-foundational-strategy-for-conservation?src=social.nature.facebook.main>, 2017.

Bot, A., Benites, J.: The importance of soil organic matter: key to drought-resistant soil and sustained food production. Rome: Food and Agriculture Organization of the United Nations, 2005.

Campos, J.N.: A gestão das águas e o desenvolvimento do Estado do Ceará: uma perspectiva histórica. T&C Amazônia, Ano IV, Num. 9, 2006.

CENAD - Centro Nacional de Gerenciamento de Desastres: Anuário Brasileiro de Desastres Naturais. Brasília, DF: Ministério da Integração Nacional and Secretaria Nacional de Proteção e Defesa Civil, 2014.

CEPED - Centro Universitário de Estudos e Pesquisas sobre Desastres: Atlas Brasileiro de Desastres Naturais 1991 a 2010: volume Brasil. Universidade Federal de Santa Catarina. Florianópolis, Brazil, 2012.

Cirilo, J. A.: Public Water Resources Policy for the semi-arid Region. *Estudos Avançados*, 22 (63), Revista USP, Universidade de São Paulo, SP, 2008.

Cunha, A.P.M.A.; Zeri, M.; Deusará Leal, K.; Costa, L.; Cuartas, L.A.; Marengo, J.A.; Tomasella, J.; Vieira, R.M.; Barbosa, A.A.; Cunningham, C.; Cal Garcia, J.V.; Broedel, E.; Alvalá, R.; Ribeiro-Neto, G.: Extreme Drought Events over Brazil from 2011 to 2019. *Atmosphere*, 10, 642, 2019.

Cunha, A. P. M. A., Alvalá, Regina C. S., Nobre, C. A., Carvalho, M. A.: Monitoring vegetative drought dynamics in the Brazilian Semiarid Region. *Agricultural and Forest Meteorology*. Vol 214-215, 494-505 p. 2015.

Cunha, A. P. M. A. ; Alvalá, R. C. dos S. ; Cuartas, L. A. ; Marengo, J. A. ; Marchezini, V. ; Leal, K. R. D. ; Tomasella, J. ; Saito, S. M. ; Zeri, M. ; Munoz, V. A. ; Ribeiro-Neto, G. ; Seluchi, M. E. ; Cunningham, C. ; Costa, L. C. O. ; Zhang, R. ; Moraes, O. L. L. . Brazilian Experience on the Development of Drought monitoring and Impact Assessment Systems. United Nations Office for Disaster Risk Reduction - UNDRR, 2019 (Contributing paper to Global Assessment Report on Disaster Risk Reduction - GAR 2019).

D' Souza, R., Fernandes, M.F., Barbosa, M.: Vulnerabilidades, semi-aridez e desertificação: cenários de riscos no Cariri Paraibano. *OKARA: Geografia em debate*, v.2, n.2, p. 190-202, 2008.

Duffy, P. B., Brando, P., Asner, G. P., and Field, C. B.: Projections of future meteorological drought and wet periods in the Amazon. *Proceedings of the National Academy of Sciences*, 112(43), 13172–13177. <https://doi.org/10.1073/pnas.1421010112>, 2015

Formatado: Português (Brasil)

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

EM-DAT: The Emergency Events Database Université Catholique de Louvain (UCL) - CRED, D. Guha-Sapir. Retrieved from: www.emdat.be, Brussels, Belgium, 2018.

Florinsky, I.V. and Pankratov, A. N.: A universal spectral analytical method for digital terrain modeling, *Int J Geogr Inf Sci*, 30:12, 2506-2528, Doi: 10.1080/13658816.2016.1188932, 2016.

Guevara, M., Olmedo, G. F., Stell, E., Yigini, Y., Aguilar Duarte, Y., Arellano Hernández, C., Arévalo, G. E., Arroyo-Cruz, C. E., Bolivar, A., Bunning, S., Bustamante Cañas, N., Cruz-Gaistardo, C. O., Davila, F., Dell Acqua, M., Encina, A., Figueredo Tacona, H., Fontes, F., Hernández Herrera, J. A., Ibelle Navarro, A. R., Loayza, V., Manueles, A. M., Mendoza Jara, F., Olivera, C., Osorio Hermosilla, R., Pereira, G., Prieto, P., Alexis Ramos, I., Rey Brina, J. C., Rivera, R., Rodríguez-Rodríguez, J., Roopnarine, R., Rosales Ibarra, A., Rosales Riveiro, K. A., Schulz, G. A., Spence, A., Vasques, G. M., Vargas, R. R., and Vargas, R.: No Silver Bullet for Digital Soil Mapping: Country-specific Soil Organic Carbon Estimates across Latin America, *Soil Discuss.*, <https://doi.org/10.5194/soil-2017-40>, in review, 2018.

Guevara, M. and Vargas, R. Downscaling satellite soil moisture using geomorphometry and machine learning. *PLoS one*, 14(9), 2019.

Hiemstra, P. H., Pebesma, E. J., Twenhöfel, C. J. W. and Heuvelink, G. B. M.: Real-time automatic interpolation of ambient gamma dose rates from the Dutch radioactivity monitoring network. *Comput. Geosci.*, 35(8), 1711–1721, doi:10.1016/j.cageo.2008.10.011, 2009.

Holden, J.: *Water Resources: An Integral Approach*. Routledge. New York, NY, 2014.

IBGE - Instituto Brasileiro de Geografia e Estatística: Mapa de Biomas e de Vegetação. Retrieved from <https://www2.ibge.gov.br/home/presidencia/noticias/21052004biomashtml.shtm> (Accessed August 20, 2018), 2004.

IBGE - Instituto Brasileiro de Geografia e Estatística: Pesquisas. Retrieved from <https://cidades.ibge.gov.br/pesquisas>, 2017.

IBGE - Instituto Brasileiro de Geografia e Estatística: Biomas e sistema costeiro-marinho do Brasil. Rio de Janeiro, 2019.

Formatado: Português (Brasil)

INPE – National Institute of Spatial Research: INPE Nordeste mapeia desmatamento da Caatinga. Retrieved from http://www.inpe.br/noticias/noticia.php?Cod_Noticia=3895 (Accessed April 17, 2018), 2018.

Ioris, A. A. R., Irigaray, C. T., and Girard, P.: Institutional responses to climate change: opportunities and barriers for adaptation in the Pantanal and the Upper Paraguay River Basin. *Climatic Change*, 127(1), 139–151. <https://doi.org/10.1007/s10584-014-1134-z>, 2014.

Kouadio, Y. K., Servain, J., Machado, L. A. T., Lentini, C. A. D.: Heavy rainfall episodes in the eastern northeast Brazil linked to large-scale ocean-atmosphere conditions in the tropical Atlantic. *Adv Meteorol*, doi:<http://dx.doi.org/10.1155/2012/369567>, 2012.

Kuppel, S., Houspanossian, J., Noretto, M. D., Jobbágy, E. G.: What does it take to flood the Pampas?: Lessons from a decade of strong hydrological fluctuations: Floods and the water cycle in the Pampas. *Water Resour Res*, 51(4), 2937–2950. <https://doi.org/10.1002/2015WR016966>, 2015.

Leal, I. R., Da Silva, J. M. C., Tabarelli, M., Lacher, T. E. (2005) Changing the Course of Biodiversity Conservation in the Caatinga of Northeastern Brazil. *Conserv Biol*, 19(3), 701–706. <https://doi.org/10.1111/j.1523-1739.2005.00703.x>

Legates, D. R., Mahmood, R., Levina, D. F., DeLiberty, T. L., Quiring, S. M., Houser, C., and Nelson, F. E.: Soil moisture: A central and unifying theme in physical geography. *Prog Phys Geog*, 35(1), 65–86. <https://doi.org/10.1177/0309133310386514>, 2011.

Liu, Y. Y., Parinussa, R. M., Dorigo, W. A., De Jeu, R. A. M., Wagner, W., van Dijk, A. I. J. M., McCabe, M. F. and Evans, J. P.: Developing an improved soil moisture dataset by blending passive and active microwave satellite-based retrievals. *Hydrol. Earth Syst. Sci.*, 15(2), 425–436, doi:10.5194/hess-15-425-2011, 2011.

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

Llamas, R.M., Guevara, M., Rorabaugh, D., Taufer, M. and Vargas, R. Spatial Gap-Filling of ESA CCI Satellite-Derived Soil Moisture Based on Geostatistical Techniques and Multiple Regression. *Remote Sensing*, 12(4), p.665, 2020.

Loon, A. F. V., Gleeson, T., Clark, J., Baldassarre, G. D., Teuling, A. J., Tallaksen, L. M., ... Wanders, N.: Drought in the Anthropocene. *Nat Geosci*, 9, 3, 2016.

Magalhães, A.: Life and drought in Brazil. Drought in Brazil - Proactive Management and Policy, in: Wilhite, D. (Ed.). Drought and Water Crisis. CRC Press. Boca Raton, FL, pp. 1 - 19, 2016.

Marengo, J., Alves, L., Alvala, R., Cunha, A., Brito, S., Moraes, O.: Climatic characteristics of the 2010-2016 drought in the semiarid Northeast Brazil region. *Anais da Academia Brasileira de Ciências (Annals of the Brazilian Academy of Sciences)*, doi:10.1590/0001-3765201720170206, 2017.

Formatado: Português (Brasil)

Marengo, J., Tomasella, J., Alves, L., Soares, W., Rodriguez, D.: The drought of 2010 in the context of historical droughts in the Amazon region. *Geophys Res Lett*, Vol. 38, L12703, doi:10.1029/2011GL047436, 2010.

McColl, K. A., Alemohammad, S. H., Akbar, R., Konings, A. G., Yueh, S. and Entekhabi, D.: The global distribution and dynamics of surface soil moisture. *Nat. Geosci.*, 10(2), 100–104, doi:10.1038/ngeo2868, 2017.

Formatado: Português (Brasil)

Medeiros, R. M.: Análise Hidroclimático do Município de Cabaceiras, PB. *Revista Brasileira de Geografia Física*, 17, 2012.

Ministry of National Integration of Brazil: Reconhecimentos Realizados e Reconhecimentos Vigentes. Retrieved from: <http://www.mi.gov.br/web/guest/reconhecimentos-realizados> (Accessed 24 March 2018), 2018.

Mishra, A. K., Singh, V. P.: A review of drought concepts. *J Hydrol*, 391(1- 2), 202-216. doi:10.1016/j.jhydrol.2010.07.012, 2010.

Moraes, C., Pereira, G., Cardozo, F.: Avaliação da precipitação e sua influência sobre as áreas inundadas no Pantanal. *Anais XVI Simpósio Brasileiro de Sensoriamento Remoto - SBSR, INPE, Foz do Iguaçu, PR, Brasil, 2013.*

Formatado: Português (Brasil)

Morim, R. L., Kolker, E., Liu, G., Liu, X., Cheng, S.: In praise of open research measures. *Nature* (498) 170, <https://doi.org/10.1038/498170b>, 2013.

Formatado: Português (Brasil)

Nascimento, S., Alves, J.: Ecoclimatologia do Cariri Paraibano. *Revista Geográfica Acadêmica*, v.2, vol.3, 28-41, 2008.

National Secretary of Civil Defense and Protection of Brazil: Relatório de Gestão: Exercício 2016. Ministério da Integração Nacional. Brasília-DF, Brazil, 2017.

Novick, K. A., Ficklin, D. L., Stoy, P. C., Williams, C. A., Bohrer, G., Oishi, A. C., Phillips, R. P.: The increasing importance of atmospheric demand for ecosystem water and carbon fluxes. *Nat Clim Change*, 6(11), 1023–1027. <https://doi.org/10.1038/nclimate3114>, 2016

NWS - National Weather Service: Drought: Public Fact Sheet. National Oceanic and Atmospheric Administration, 2008.

Overbeck, G. E., Vélez-Martin, E., Scarano, F. R., Lewinsohn, T. M., Fonseca, C. R., Meyer, S. T., Pillar, V. D.: Conservation in Brazil needs to include non-forest ecosystems. *Diversity Distrib*, 21(12), 1455–1460. <https://doi.org/10.1111/ddi.12380>, 2015.

Pontes, A. The Caatinga Biome. Percentual de água por Bacia Hidrográfica varia de 18.6% a 96.7%. Das 15 bacias apenas quatro estão em situação de segurança hídrica. Blog do Jose Carneiro, <http://joseliocarneiro.blogspot.com/2012/11/percentual-de-agua-por-bacia.html>, 2012.

Formatado: Português (Brasil)

Reuter, H.I. & Hengl, T.: Global Soil Information Facilities-Component Worldgrids.org. EGU General Assembly Conference Abstracts. Retrieved 9 September, 2018 from https://www.researchgate.net/publication/233540147_Global_Soil_Information_Facilities-Component_Worldgrids_org, 2012.

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

Roesch, L. F., Vieira, F., Pereira, V., Schünemann, A. L., Teixeira, I., Senna, A. J., and Stefenon, V. M.: The Brazilian Pampa: A Fragile Biome. *Diversity*, 1(2), 182–198. <https://doi.org/10.3390/d1020182>, 2009.

Rossato, L., Marengo, J. A., Angelis, C. F. de, Pires, L. B. M., Mendiondo, E. M.: Impact of soil moisture over Palmer Drought Severity Index and its future projections in Brazil. *RBRH*, 22(0). <https://doi.org/10.1590/2318-0331.0117160045>, 2017.

Formatado: Português (Brasil)

Santos, M. G., Oliveira, M. T., Figueiredo, K. V., Falcão, H. M., Arruda, E. C. P., Almeida-Cortez, J., ... Antonino, A. C. D.: Caatinga, the Brazilian dry tropical forest: can it tolerate climate changes? *Theor Exp Plant Phys*, 26(1), 83–99. <https://doi.org/10.1007/s40626-014-0008-0>, 2014.

Formatado: Português (Brasil)

Santos, S., Silva, L. G.: Mapeamento por imagens de sensoriamento remoto evidencia o bioma Pampa brasileiro sob ameaça. *Boletim de Geografia*, 29(2). <https://doi.org/10.4025/bolgeogr.v29i2.12366>, 2012.

SECOM- The Secretariat for Social Communication of the Presidency of Brazil: Biodiversity in Brazil. Secretariat for Social Communication of the Presidency of the Federative Republic of Brazil. United Nations Conference on Biological Diversity (COP11). Hyderabad, India, 2012.

Sen, P. K.: Estimates of the Regression Coefficient Based on Kendall's Tau. *J. Am. Stat. Assoc.*, 63(324), 1379, doi:10.2307/2285891, 1968

Sheffield, J., Wood, E. F.: Global Trends and Variability in Soil Moisture and Drought Characteristics, 1950–2000, from Observation-Driven Simulations of the Terrestrial Hydrologic Cycle. *J Climate*, 21(3), 432–458, 2008.

Siegel, A. F.: Robust Regression Using Repeated Medians. *Biometrika*, 69(1), 242, doi:10.2307/2335877, 1982.

Smith, K.: *Hydrological Hazards. Environmental Hazards - Assessing Risk and Reducing Disaster*, Routledge, 337-370. New York: NY, 2013.

The Nature Conservancy: The Atlantic Forest harbors a range of biological diversity similar to that of the Amazon. The Nature Conservancy. Retrieved from: <https://www.nature.org/ourinitiatives/regions/latinamerica/brazil/placesweprotect/atlantic-forest.xml>, 2015.

Theil, H.: A Rank-Invariant Method of Linear and Polynomial Regression Analysis, in Henri Theil's Contributions to Economics and Econometrics, in: Raj, B. and Koert, J. (Eds) vol. 23, pp. 345–381, Springer Netherlands, Dordrecht. Retrieved from: http://www.springerlink.com/index/10.1007/978-94-011-2546-8_20, 1992.

Tomasella J, et al. (2018) Desertification trends in the Northeast of Brazil over the period 2000–2016, *Int J Appl Earth Obs Geoinformation*, 73. 197–206.

Travassos, I. S., De Souza, B. I.: Os negócios da lenha: indústria, desmatamento e desertificação no Cariri paraibano. *GEOUSP: Espaço e Tempo (Online)*, 18(2), 329. <https://doi.org/10.11606/issn.2179-0892.geosp.2014.84536>, 2014

Formatado: Português (Brasil)

Vargas, R. How a hurricane disturbance influences extreme CO₂ fluxes and variance in a tropical forest. *Environmental Research Letters*, 7(3), p.035704, 2012.

Vargas, R.; Collins, S.L.; Thomey, M.L.; Johnson, J.E.; Brown, R.F.; Natvig, D.O.; Friggens, M.T.: Precipitation variability and fire influence the temporal dynamics of soil CO₂ efflux in an arid grassland. *Global Change Biol*, 18, 1401–1411, 2012.

Vargas, R., Sánchez-Cañete, P., Serrano-Ortiz, P., Curiel Yuste, J., Domingo, F., López-Ballesteros, A. and Oyonarte, C., 2018. Hot-moments of soil CO₂ efflux in a water-limited grassland. *Soil Systems*, 2(3), p.47, 2018.

Villarreal, S., Vargas, R., Yopez, E.A., Acosta, J.S., Castro, A., Escoto-Rodriguez, M., Lopez, E., Martínez-Osuna, J., Rodriguez, J.C., Smith, S.V. and Vivoni, E.R. Contrasting precipitation seasonality influences evapotranspiration dynamics in water-limited shrublands. *Journal of Geophysical Research: Biogeosciences*, 121(2), pp.494-508, 2016.

Formatado: Português (Brasil)

Formatado: Cor da fonte: Preto

Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita

Zeri, M., S. Alvalá, R., Carneiro, R., Cunha-Zeri, G., Costa, J., Rossato Spatafora, L., ... Marengo, J.: Tools for Communicating Agricultural Drought over the Brazilian Semiarid Using the Soil Moisture Index. *Water*, 10(10), 1421. <https://doi.org/10.3390/w10101421>, 2018.

Formatado: Cor da fonte: Preto
Formatado: Normal, Borda: Superior: (Sem borda), Inferior: (Sem borda), Esquerda: (Sem borda), Direita: (Sem borda), Entre : (Sem borda), Tabulações: 7,96 cm, Centralizado + 15,92 cm, Direita