

An analysis of temporal scaling behaviour of extreme rainfall in Germany based on radar precipitation QPE data

Judith Marie Pöschmann¹, Dongkyun Kim², Rico Kronenberg¹, and Christian Bernhofer¹

¹Department of Hydrosociences, Institute of Hydrology and Meteorology, Technische Universität Dresden, 01069 Dresden, Germany

²Department of Civil and Environmental Engineering, Hongik University, Wausanro 94, Mapo-gu, Seoul, 04066, Korea

Correspondence: Judith Pöschmann (Judith.Poeschmann@tu-dresden.de)

Abstract. We investigated the depth–duration relationship of maximum rainfall over all Germany based on 16 years of radar derived Quantitative Precipitation Estimates (namely, RADKLIM–YW, German Meteorological Service) with a space–time resolution of 1 km² and 5 min. Contrary to the long–term historic records that identified a smooth power law scaling behaviour between the maximum rainfall depth and duration, our analysis revealed three distinct scaling regimes of which boundaries are approximately 1 hour and 1 day. Few extraordinary events dominated a wide range of durations and deviate to the usual power law. Furthermore, the shape of the depth–duration relationship varied with the sample size of randomly selected radar pixels. A smooth scaling behaviour was identified when the sample size was small (e.g. 10 to 100), but the original three distinct scaling regimes became more apparent as the sample size increases (e.g. 1,000 to 10,000). Lastly, a pixel wise classification of the depth–duration relationship of the maximum rainfall at all individual pixels in Germany revealed three distinguishable types of scaling behaviour, clearly determined by the temporal structure of the extreme rainfall events at a pixel. Thus, the relationship might change with longer time series and can be improved once available.

1 Introduction

Extreme rainfall poses significant threats to natural and anthropogenic systems (Papalexiou et al., 2016). The frequency and magnitude of extreme rainfall are expected to increase in the future (Blanchet et al., 2016; Gado et al., 2017; García-Marín et al., 2012; Ghanmi et al., 2016; Lee et al., 2016; Madsen et al., 2009; Marra and Morin, 2015; Marra et al., 2017; Overeem et al., 2009; Yang et al., 2016) especially at sub–daily timescales (Barbero et al., 2017; Fadhel et al., 2017; Guerreiro et al., 2018; Westra et al., 2013, 2014) leading potentially to more flash floods (Dao et al., 2020), riverine floods, and landslides. A thorough understanding on magnitude, duration, and frequency of extreme rainfall is thus necessary for efficient design, planning, and management of these systems, with many requiring (sub–)hourly information especially.

It is difficult to identify and investigate extremes and record rainfall events, they occur rarely, and the spatiotemporal resolutions and coverage information are generally limited. Lengfeld et al. (2020) analysed the problems with rain gauge observations and concluded that more than 50 % of the extreme rainfall events observed were missed, especially in data with higher temporal resolutions. Remotely sensed precipitation products with high spatiotemporal resolution such as the ones provided by radar, satellite or microwave link networks may solve this issue. Weather radar systems are considered appropriate to capture

25 the spatial variability of extreme rainfalls, including events with limited spatial extent (Borga et al., 2008). While their high
spatiotemporal resolution is superior to many other rainfall products, most of the currently available radar QPE (quantitative
precipitation estimates) data sets do not cover very long periods (Lengfeld et al., 2020). Radar products also have well-known
uncertainties, like variation of reflectivity with height, relating radar reflectivity to precipitation rates, attenuation, clutter, and
beam blocking. Therefore, reprocessing these data sets is necessary in order to achieve homogeneous and consistent products
30 for spatiotemporal evaluation of rainfall characteristics.

Probable maximum precipitation (PMP) is one way to define extreme rainfall. It is defined as the "theoretically greatest depth
of precipitation for a given duration that is physically possible over a particular drainage basin at a particular time of year"
(American Meteorological Society, 2020). One of the methods to estimate the PMP is the maximum rainfall envelope curve
method, that plots the depth (y)–duration (x) relationship of the record rainfall events observed across a large geographical
35 boundary (e.g. entire country or globe) on the log–log plane. The PMP is then derived as a straight line on the plot representing
the upper boundary of the envelope containing all depth–duration relationships. This maximum rainfall envelope curve method
was first proposed by Jennings (1950), who showed that the depth of the extreme rainfall events observed across the globe is a
power function of their duration. Jennings discovered that this unique scaling behaviour holds at rainfall durations between 1
min through 24 months. Paulhus (1965) showed that the same power law relationship holds after the addition of a new world
40 rainfall record observed at the island of La Réunion at the duration between 9 h and 8 d. The envelope for these extreme values
can be expressed as:

$$P = \alpha D^\beta \tag{1}$$

where P is the maximum precipitation (in mm) occurring in duration D (in h), the coefficient α (425 in Paulhus, 1965)
represents the value at 1 hour of the depth–duration relationship plotted on the log–log plane, and the exponent β (0.47 in
45 Paulhus, 1965) is the parameter characterizing the scaling behaviour of the depth–duration relationship. The Spanish study of
Gonzalez and Bech (2017) updated the global envelope's slope to 0.51, showing a remarkable stability. Multiple exponents
describing the scaling property of rainfall extremes have been retrieved at various regions around the world (Commonwealth
of Australia, 2019; Gonzalez and Bech, 2017). Figure 1 shows the maximum rainfall–duration relationship identified by some
of these studies. All relations reveal power law relationships, with exponents ranging from around 0.5 (Spanish and global
50 estimate) and 0.2 (German) over a wide range of scales.

Several studies examined the validity of this universal scaling exponent. Galmarini et al. (2004) showed, based on rainfall
records observed at several stations in Canada, Australia, and La Réunion, that the single exponent scaling laws exist only for
single stations experiencing extremely high precipitation and that the deviation from a scaling law is caused by the intermittency
associated with a substantial number of zero precipitation intervals in data. They also showed that the scaling exponent β tend
55 to stay around 0.5 based on the stochastic simulation assuming a point rainfall process composed of the Weibull distributed
rainfall depth and a given temporal autocorrelation structure. Zhang et al. (2013) showed that the scaling exponent varies
around 0.5, if the vertical moisture flux and rainfall can be modelled by a censored (or truncated) first–order autoregressive




Figure 1. Scaling relationships of extreme (record) precipitation values for different durations based on worldwide data (World Meteorological Organization, 1994; NWS, 2017; Gonzalez and Bech, 2017), Spanish rain gauge data (Gonzalez and Bech, 2017), and a regional analysis of Eastern Germany (Dyck and Peschke, 1995)

process AR(1). However, these works showed the scaling behaviour of maximum rainfall at a single point location, and did not investigate maxima observed at different spatial locations.

60 One of the main obstacles to identify the "true" scaling behaviour of maximum rainfall is that the most rainfall is measured from sparse ground gauge networks (Dyck and Peschke, 1995; Papalexiou et al., 2016). Breña-Naranjo et al. (2015) used a satellite based rainfall product to identify the scaling behaviour of the maximum rainfall across the globe. They showed that the maximum of the areal rainfall averaged over the $\sim 20 \text{ km} \times \sim 20 \text{ km}$ data grid has the scaling exponent of ~ 0.43 which is similar to that of Jennings (1950). However, the coarse spatial resolution of the satellite data easily misses the small scale
65 rainfall variability that is closely associated with extreme values, thus the extremes found in the satellite data are lower than expected (Cristiano et al., 2017; Fabry, 1996; Gires et al., 2014; Kim et al., 2019; Peleg et al., 2013, 2018).

In this study, we analyse the rainfall depth–duration relationship for all Germany based on 16 years of RADKLIM–YW, a reprocessed QPE radar product with 1 km–5 min space–time resolution. We want to answer the following questions regarding the scaling behaviour of the maximum rainfall: (1) Does the depth–duration relationship of German extreme rainfall show
70 scale invariant behaviour? If so or if not, what is the primary reason? (2) Does this relationship vary with regard to the spatial sampling rate? (3) Does it provide any clue to modify the relationship currently applied in practice based on sparse gauge

networks? The answers to these questions would be especially intriguing because few studies have so far investigated the scaling behaviour of maximum rainfall based on a rainfall data set with such a high spatiotemporal resolution, long recording period, and large spatial extent as this study did.

75 2 Data and methods

2.1 Data description

The German National Meteorological Service (DWD) runs a radar network (currently 17 C-band radars) for almost two decades and is providing different rainfall data products. Full coverage of Germany was not reached until today, however, all neighbouring countries contribute to the rainfall information and the network extension continues on an ongoing basis. One
80 QPE from German radar data is RADOLAN (German: RADar OnLine ANeichung) (Winterrath et al., 2012), which combines ground information of fallen precipitation (rain gauge data) with radar data. Since the quality enhancement of RADOLAN is ongoing without post-correcting previous data, the so-called radar climatology project of the DWD, RADolanKLIMatologie (RADKLIM, Winterrath et al., 2017) has consistently reanalysed the complete radar archive set since 2001 for improved homogeneity despite the originally different processing algorithms. Compared to RADOLAN, RADKLIM has implemented
85 additional algorithms leading to consistently fewer radar artefacts, improved representation of orography as well as efficient correction of range-dependent path-integrated attenuation at longer time scales (Kreklow et al., 2019). Whereas RADOLAN is not well suited for climatological applications with aggregated precipitation statistics, RADKLIM is a promising data set for these climatological applications. The RADKLIM product is available in the following two versions with around 392 128 filled pixels within the German border: 1) RADKLIM-RW is an hourly precipitation product resulting from radar based precipitation
90 estimates that are calibrated with ground stations (Winterrath et al., 2018a), which was validated by several studies, such as Lengfeld et al. (2019) and 2) RADKLIM-YW (Winterrath et al., 2018b) is a 5 min product resulting from a correction/factoring of DWD's 5 min product RADOLAN-RY (rainfall estimate after basic quality correction and refined z-R-relationship) with the help of RADKLIM-RW on a sequential hourly base. The RADKLIM-YW version 2017.002 was used in this study because it has the high temporal resolution necessary for the analysis. This release is in its third version and covers the years 2001 to
95 2018. Due to comparison reasons with another study at our institute, only years 2001 to 2016 have been used for this study. The YW product covers the area composed of 1,100 x 900 pixels with the spatial resolution of 1 km (improved compared to former version of RADOLAN). Remaining weaknesses of RADKLIM (as outlined in Kreklow et al., 2019) are a greater number of missing values compared to RADOLAN as well as an underestimation of high intensity rainfall because of spatial averaging and rainfall-induced attenuation of the radar beam.

100 The data is available as one layer for each time step. Since not all raster pixels are with values (only around half of the values lay within the borders of Germany), the spatial data was converted to time series for quicker processing. The data contain missing values (NaN) of the following two types: 1) NaNs due to changes and ongoing radar network extension. This mainly affects areas near the border of Eastern, Northern, and Southern Germany. Data in some areas is only available from 2014 onwards. 2) Some locations/raster pixels have NaNs potentially due to malfunction of the radar or general (radar) errors.




Figure 2. Spatial distribution of the proportion of NaNs (in %) for each pixel of the QPE RADKLIM–YW from 2001–2016. The German boundary is obtained from the GADM Global Administrative Database (Hijmans et al., 2018)

105 Figure 2 shows the proportion of the NaNs of the time series developed for each of the pixels. The visible cones display the
 110 individual radar coverage, and the overlapping areas of the radar cones have a better data coverage than the areas without overlapping.

It is hard to handle NaNs in highly episodic geophysical events such as rainfall. We chose to not do any data interpolation, since the consequence of imputing potentially too high extreme values is more severe and uncertain for our study than missing
 110 extreme values.

2.2 Depth–duration relationships

Maximum rainfall values for each duration τ between 2001–2016 were calculated with rolling sums applied over moving windows using the R–package Rcpp–Roll (Ushey, 2018). Time windows of up to 3 days were chosen for the analysis, with special

focus on the sub-hourly and sub-daily durations. The records may include non-rainfall data and thus do not imply continuous precipitation for the period considered. Values were not aggregated spatially, since this usually reduces the maximum intensity values (Cristiano et al., 2018).

First, the extreme values for each pixel and duration $M_{\max}^{\tau, pixel}$ are calculated. Afterwards, the overall maxima for all Germany for each τ ($M_{\max}^{(\tau)}$) is extracted from these calculated extreme values. Based on these results, the depth-duration relationships can be developed for each pixel as well as for all Germany.

120 2.3 K-means clustering of depth-duration-relationships

The depth-duration relationships ($M_{\max}^{\tau, pixel}$ vs τ) for each pixel derived from Sect. 2.2 are individually clustered with the K-means clustering algorithm (Scott and Knott, 1974). "Erroneous" pixels (=having NaNs as resulting maxima) were excluded from the cluster process in order to avoid disturbances. The data was rescaled to make the characteristics more comparable with each other. If the number of clusters is not predefined, it can be identified by drawing an elbow chart. For different numbers of clusters K the measure of the variability of the observations within each cluster (Total within-cluster sum of squares, y-axis) is calculated and the curve should bend like an elbow at the optimal value. Since the algorithm did not suggest a number of clusters, we chose six clusters for a sufficiently detailed analysis since it gave consistent results when repeating the automatic algorithm for several times (each time the algorithm clusters slightly differently).

3 Results and discussion

130 3.1 Scaling behaviour of all Germany

Figure 3 shows the maximum depth-duration relationship of all Germany that was derived from the QPE radar data (dots). The same relationship based on ground gauge network (empty triangles) and the global precipitation extremes (filled triangles) are shown for reference. The rain gauge based values clearly follow a scaling relationship with a slope that is different in comparison to world extremes. Radar based maxima for the shorter duration from 2001 to 2016 do not cover all sub-daily extremes but exceed observed ones from the 1 d durations as well as for one sub-daily value. In Fig. 3, a "plateau" is visible between around 35 min up to 18 h, indicating a "one event" effect at 35 min, potentially from an extreme rainfall event in this period. Overall, a scaling behaviour can be observed at sub-hourly durations with a scaling component of around 0.65 even though the maxima are observed rather randomly across all Germany as indicated by the map showing the location of the maximum rainfall. This result implies that even though the location of extreme rainfall is different, the maximum rainfall may exhibit smooth scaling behaviour if the rainfall generation mechanism is similar. As mentioned in the data quality description, it is possible that these sub-hourly values do not represent the true extremes across Germany for 2001–2016, since radar-based measurements at fine timescales are highly sensitive to the effects of averaging. Between 25 min and 16 h, maximum values are calculated for a location at the border of Hesse and Bavaria in August 25th 2006. An extreme event around September 30th,

Table 1. Rainfall records for different duration from RADKLIM–YW for 2001–2016 with corresponding locations.

Duration	Start Date	Start Time (CET)	Precipitation Sum (mm)	Location (WGS84)
5 min	2009–07–04	14:10	40.9	48.50° N, 9.35° E
10 min	2006–07–07	9:30	80.8	51.22° N, 8.77° E
15 min	2010–07–12	23:05	105.6	52.80° N, 12.39° E
20 min	2002–07–30	17:15	127.3	48.82° N, 9.58° E
25 min–16 h	2006–08–25	05:25–13:25	141.1–230.7	50.21° N, 9.20° E
18 h–1 d	2003–09–29	09:05–15:05	258.9–327.5	52.53° N, 13.53° E
1.5–2 d	2003–09–28	14:20–21:35	471.7–503.7	52.53° N, 13.53° E
3 d	2001–04–08	06:50	525.9	53.68° N, 10.00° E

Maxima of 25 min–16 h as well as from 18 h–2 d correspond to the same location and date and are thus summarized.

2003 around Berlin comprised the maximum depth–duration relationship at between 18 h and 2 d durations. Weak scaling
145 behaviour existed in the regime at 18 h and 3 d durations with the scaling exponent of 0.20.

All locations of maxima and the corresponding dates of occurrence are provided in Table 1.

3.2 Scaling behaviour of all Germany for high–quantile rainfall

High rainfall values obtained from radar data are associated with especially great uncertainty. Thus, we also investigated
150 the scaling behaviour of high–quantile rainfall values. Figure 4 shows the maximum depth–duration relationship of several
quantiles: 0.99999, 0.9999, 0.999, and 0.99. The “three phase regime” from radar maximum values remains relatively stable,
however, the “single event” effect between 50 min and 1 d is smoothed out, because the degree of inflections in the curve
becomes weaker. Lower quantiles thus show a smoother curve rather than the “three phase regime”.

Figure 5 shows the location the high quantile rainfall. The colour of the circles represents the different rainfall durations. It
shows that at the highest considered quantile (0.99999) multiple maxima appear at similar locations, potentially referring to
155 the same rainfall events, whereas for lower quantiles (e.g., 0.9999 to 0.99), maxima are more spread over Germany and the
visible points increase in number. This suggests the reduction of the influence of one single rainfall event on the depth–duration
relationship causing inflection in the curve.

Additionally, locations of such high quantile maxima (e.g., 0.99 quantile in Fig. 5) seem to occur predominantly in the wider
Alpine region in South Germany. This suggests that natural rainfall mechanisms are dominating the scaling relationship, such as
160 regional characteristics and meteorological conditions (e.g. orographic lifting or leewards effects). Naturally, one would assume
that this heterogeneity in meteorological conditions and rainfall generating mechanisms will reflect regional characteristics and
will exhibit some irregular scaling behaviour. Contrary to this conjecture, the curves in Fig. 4 (99.9 % and 99 %) show a quite
smooth scaling behaviour.




Figure 3. Overview of maximum rainfall records in Germany. Chart: Maximum depth–duration relationship of rainfall records based on QPE RADKLIM–YW (data of this study) (blue dots), and as reference the relationships based on the German ground network (Rudolf and Rapp, 2003; DWA, 2015; DWD, 2020) (non-filled triangles) and the global precipitation extremes (World Meteorological Organization, 1994; NWS, 2017). Map: Locations of rainfall maxima (based on QPE RADKLIM–YW) for the considered duration.




Figure 4. Depth–duration relationships of rainfall values of all Germany based on QPE RADKLIM–YW for 2001–2016 from maximum values down to the 3,921st greatest per duration.

Rainfall Maxima




Figure 5. Locations of the 0.99999, 0.9999, 0.999, and 0.99 quantile rainfall with varying durations from 5 min to 3 d. Point colours represent the corresponding rainfall duration, similar for each quantile. Different numbers of data points in panels a–d result from several data points being at the same location.




Figure 6. Spatial distribution of the maximum rainfall values retrieved from QPE RADKLIM–YW (2001–2016) for different durations (5 min to 3 days).

3.3 Spatial distribution of maximum rainfall

165 Figure 6 shows the spatial distribution of 5 min, 30 min, 1 h, 6 h, 1 d, and 3 d maximum rainfall over Germany. The red and yellow spots that are spatially distributed in Fig. 6 (a) suggest that 5 min extreme rainfall can happen at any place in Germany. Note that extreme rain occurred also outside the Alpine region at the southern edge of Germany, which suggests that fine-scale extreme rainfall is not necessarily governed by topography. The influence of fine-scale intense rainfall persists until the hourly timescale as implied by the red and yellow hotspots that are located at similar places in the maps of 5 min, 30 min, and 1 h.

170 The distribution of maxima significantly changes for the duration of 6 h, and an interesting pattern emerges in the map of 1 d and 3 d duration. These maxima seem to be dominated by single events or single heavy rainfall occurrences. This is especially evident in the 2002 flooding in Saxony (mid eastern edge) with unprecedented long and heavy rainfall as well as a singular rainfall event in 2014 (narrow aisle in the Northwestern area) clearly visible in the maps.




Figure 7. Depth–duration relationships of rain records for single pixels at rain gauge locations within state capitals of German federal states

3.4 Scaling behaviour at a single point

175 Figure 7 shows the maximum rainfall–duration relationship of the radar pixels at the major cities of Germany with a single power law (blue) as reference to see the differences better. Except for Hamburg and Stuttgart, most cities exhibit slight (Hannover, Kiel, Magdeburg, Potsdam, Schwerin, Wiesbaden) to considerable (the remaining cities) deviation from a single power law behaviour. This significant deviation is similar to what was identified by Galmarini et al. (2004) who found that the inflection of the curve is inevitable because of the small (or zero) rainfall observations around a maximum rainfall event.

180 Furthermore, Galmarini et al. (2004) and Zhang et al. (2013) both showed that the maximum rainfall–duration relationship at a given point location follows a smooth and simple power law if the rainfall process can be modelled with a set of simple stochastic processes. Our results imply that natural rainfall processes might significantly deviate from this rather simple assumption, also the model framework is also based on very few time series of very different lengths and resolutions.

3.5 Classification of maximum depth–duration relationship

185 The maximum depth–duration relationships for all pixels within Germany were clustered since Fig. 7 indicated that they might show similar shapes. The K–means clustering algorithm classified the depth–duration relationship into six categories revealing different curve characteristics regarding the curve shapes. Figure 8 shows a categorical map of Germany representing each category with an individual colour. Additionally, depth–duration relationships at 100 randomly chosen grid points from each category are shown with the regression line from category 5 as reference.

190 Pixels belonging to category 1 have the highest rainfall intensities over all scales until 1 d and show a strong inflection at around 1 h similar to the scaling curve for all Germany (Fig. 3). The behaviour of the curve between 5 min and 1 h is associated with strong convective rainfall events of around 1 h within the corresponding pixel. Thus, these events are responsible for the high slope at the beginning of the curve. Some curves also show another small inflection between 12 h and 1 d that might correspond to an inter storm arrival time over which another large event contributes to the positive slope of the curve at duration
195 from 1 d and longer, or simply contributes to the general high intensity of the whole event. Category 1 pixels can be identified as yellow hotspots in Fig. 8 that occur predominantly as smaller areas in the midst of category 2 (red) and partly in category 3 (light pink) pixel clusters. Category 2 pixels (red) have a similar curve shape as those in category 1 and always occur together with category 1 pixels. The curve inflection begins around 30 min and the slope up to 3 d is a little steeper than the slope of category 1. This implies that those locations experienced strong convective patterns of a slightly shorter duration, but potentially
200 longer event durations in general. Most likely, category 1 (event centre) and category 2 (event boundary) pixels experience local convective events, which are forming in the summer months on warm days with a moist atmosphere. Categories 3 and 4 can be generally associated to large scale events dominated by regional weather patterns. The three largest clusters in the map can be identified as intense frontal rainfall in August 2002 (Saxony, large cluster in Eastern Germany), heavy downpours over Münster in July 2014 (narrow path in the Northwestern part) and orographic rainfall in the Alpine region of southern
205 Germany. In category 3, curves show steep slopes of up to one day that abruptly end with super–daily duration. This category is contributing to the scales between 12 h and 3 d for the curve of all Germany (Fig. 3). The steep slope at sub–daily duration is because the pixels experienced intense convective storms, however, there was lower intensity at this duration than categories 1 and 2. Yet, for daily–scale duration they can experience significant amounts of rainfall. Both categories 4 and 5, which compose around 50 % of all pixels, show rough power law behaviour over all scales. Category 4 (dark blue) pixels are mainly at the outer
210 borders of the described larger events as well as adjacent to pixels of categories 1 and 2. Thus, the curves have steep slopes because the corresponding pixels experienced great rainfall. Most pixels belong to category 5 (36 %), showing the smoothest scaling behaviour of all categories. Based on the data set, these regions/locations never have been hit by any ‘extreme’ extreme event that could have altered the power law behaviour of the depth–duration relationship. The locations of these pixels indicate no spatial pattern and can be seen as a kind of background colour of the map. The last category 6 contains similar characteristics
215 from categories 1 to 3 and comprises another 30 % of all pixels. This category represents pixels experiencing common types of convective events with short heavy rainfall sequences on the sub–hourly scale, indicated by a relatively steep slope until 1 h compared to categories 1 and 2. However, these pixels also experience longer rainfall sequences, thus showing an almost




Figure 8. Resulting clusters of the maximum depth–duration relationships of rainfall for all pixels. The left panel shows the spatial distribution of the groups, distinguishable by colour. The corresponding curve shapes of 100 randomly selected radar pixels from each group are displayed on the right side with the same colours as the map.

“three phase regime” as the overall curve for Germany with lower values. The found clusters can be further summarized into three classes: Pixels that have experienced very heavy rainfall on a sub-hourly scale (categories 1 and 2) exhibiting steep slopes at sub-hourly scale and mild slope for longer duration. The second class experienced heavy rainfall sequences of up to 1 d (categories 3 and 4). The third type shows power law behaviour over all scales and can be mainly found in category 5. Category 6 simultaneously shows characteristics of classes 1 and 2.

3.6 Sensitivity of scaling behaviour to ground gauge network density

An important message from Sect. (3.4) and Sect. (3.5) is that the depth-duration relationship at a given point varies location by location based on the occurred rainstorms. This implies that the maximum depth-duration relationship over the entire study area, which is fundamentally the process of the superposition of these various relationships and the picking up of the very maximum values at each duration, may vary with regard to density and spatial formation of ground gauge networks (compare to previous section). For this reason, we investigated how the depth-duration relationship would vary with regard to a different number of sampling pixels. Figure 9 shows the result corresponding to the pixel sample size of 10, 100, 1,000, and 10,000. For each of the cases, 30 ensembles of random pixel sampling were performed. For each of the plots, the maximum depth-duration relationship based on all radar pixels ($n = 392,128$) was shown for reference. Clear and smooth scaling behaviours are identified when the pixel sample size is 10 and 100, but the smooth scaling behaviours are lost when including more major rainfall events that formed the original maximum depth-duration relationship. This emphasizes that the number of rain gauges in a network is extremely relevant in order to adequately capture rainfall extremes. It is of note that most scaling relationships of the past including Jennings (1950) were based on the measurements of the ground gauge network. The station density was used as “best we could get” and was not tested against a smaller set of stations, as obviously including all reliable extremes improves the relationship. This might work for the Jennings curve as a global scale (space and time) can make up for a limit in spatial resolution. However, regional scaling suffers from the limited spatial extent, which cannot be completely balanced by a denser network or radar data.

4 Conclusions

A thorough understanding on the scaling behaviour of the depth-duration relationship of extreme precipitation has been limited because its spatiotemporal variability cannot be fully captured by a measurement network composed of limited number of ground gauges. This study tried to overcome this limitation by using the radar Quantitative Precipitation Estimates (QPE) rainfall product RADKLIM-YW. The radar QPE enabled clear identification and explanation of the characteristics of the different scaling regimes of extreme rainfall depth-duration relationships. The maximum depth-duration relationship derived from radar data did not show clear scaling behaviour compared to one based on gauge data from longer time series, but exhibited a “three phase regime” with a high slope at the duration smaller than 1 h, a plateau at the duration between 1 h and 1 d, and a low slope at the duration greater than 1 day. The relationship was developed based on only a few extreme rainfall events, which dominated the shape of the curve and this changed when examining quantiles of pixel maxima. The depth-duration relationship




Figure 9. Dependency of maximum depth–duration relationship characteristic on underlying pixel sample size. The maximum rainfall values are derived from (a) 10, (b) 100, (c) 1,000, and (d) 10,000 random pixels from all considered pixels ($n=392,128$) within Germany. For each sample size, 30 ensembles are displayed and compared to the overall maximum curve from Fig. 3 and 4 (yellow top line in Figure).

250 of lower quantile rainfall (e.g. 99 percentile) showed a smooth scaling behaviour and the rainfall events contributing to the curve
 sparsely occurred at various locations of Germany. This implies that the modest extreme rainfall events are less sensitive to the
 random effects of a limited period (under sampling) and may even share common atmospheric conditions of rainfall generation
 regardless of pixel location in a limited region like Germany. The rainfall depth–duration relationship at a single radar pixel
 did not show clear power law behaviour either. The shape of the curve was governed by the temporal structure of the extreme
 255 rainfall events at the pixel location. The point wise clustering of depth–duration relationships revealed three classes of scaling
 behaviour: a) Linear scaling over all durations, as well as inflections at b) 1 hour and c) 1 day, which shows the influence of
 small convective pixels as well as large scale weather patterns on the depth–duration relationship. The scaling behaviour thus
 can be significantly different for each pixel because the rainfall characteristics for each pixel are very different as well. Given
 that the extreme rainfall depth–duration relationship over a region is a process of overlapping the relationships observed at

260 various pixel locations and picking up the highest rainfall values at each duration, the result implies that the depth–duration relationship of extreme rainfall can significantly deviate from power law behaviour. With longer available time series of radar in the future, the deviation can be further investigated and tested. Also, the known issue of rainfall extreme underestimation by RADKLIM–YW and the potential impact on the results need further evaluation.

265 *Code and data availability.* The data sets used in this study are freely available to download in ASCII and BINARY format and are published under the following DOI: 10.5676/DWD/RADKLIM_YW_V2017.002 (Winterrath et al., 2018b). Analysis was conducted in R (R Core Team, 2019) and Matlab with freely available R–packages ggplot (Wickham, 2016), RasterVis (Perpiñán and Hijmans, 2019), Rcpp–Roll (Ushey, 2018), and fst (Klik, 2019)

270 *Author contributions.* JP and DK conceptualized research, developed model code and methodology. JP carried out the analysis together with DK and RK. DK provided the computing facilities in his lab at Hongik University in South Korea. JP and DK prepared the manuscript with contribution from all co-authors. CB contributed the scientific idea, contributed to its refinement, to the discussion within the authors’ group and to the writing.

Competing interests. There are no competing interests involved in this study.

275 *Acknowledgements.* The authors sincerely acknowledge the financial support by TU Dresden’s Institutional Strategy, which is funded by the Excellence Initiative of the German Federal and State Governments. Professor Kim’s contribution was supported by the Korea Environment Industry & Technology Institute (KEITI) through Water Management Research Program, funded by Korea Ministry of Environment (MOE) (Project No. 127557). We also thank Dr. Tanja Winterrath from the German Weather Service (DWD) for giving further insight on DWD’s data processing. We thank four anonymous reviewers for their helpful comments and remarks to improve our manuscript.

References

- American Meteorological Society: Glossary of Meteorology, American Meteorology Society, <http://glossary.ametsoc.org/wiki/>, 2020.
- 280 Barbero, R., Fowler, H. J., Lenderink, G., and Blenkinsop, S.: Is the intensification of precipitation extremes with global warming better detected at hourly than daily resolutions?, *Geophysical Research Letters*, 44, 974–983, <https://doi.org/10.1002/2016gl071917>, 2017.
- Blanchet, J., Ceresetti, D., Molinié, G., and Creutin, J.-D.: A regional GEV scale-invariant framework for Intensity–Duration–Frequency analysis, *J. Hydrol.*, 540, 82–95, <https://doi.org/10.1016/j.jhydrol.2016.06.007>, 2016.
- Borga, M., Gaume, E., Creutin, J. D., and Marchi, L.: Surveying flash floods: gauging the ungauged extremes, *Hydrological Processes*, 22, 3883–3885, <https://doi.org/10.1002/hyp.7111>, 2008.
- 285 Breña-Naranjo, J. A., Pedrozo-Acuña, A., and Rico-Ramirez, M. A.: World’s greatest rainfall intensities observed by satellites, *Atmos. Sci. Lett.*, 16, 420–424, <https://doi.org/10.1002/asl2.546>, 2015.
- Commonwealth of Australia: Australia’s Record Rainfall, online, <http://www.bom.gov.au/water/designRainfalls/rainfallEvents/ausRecordRainfall.shtml>, 2019.
- 290 Cristiano, E., ten Veldhuis, M.-C., and van de Giesen, N.: Spatial and temporal variability of rainfall and their effects on hydrological response in urban areas – a review, *Hydrol. Earth Syst. Sci.*, 21, 3859–3878, <https://doi.org/10.5194/hess-21-3859-2017>, 2017.
- Cristiano, E., ten Veldhuis, M.-C., Gaitan, S., Ochoa-Rodriguez, S., and van de Giesen, N.: Critical scales to explain urban hydrological response: an application in Cranbrook, London, *Hydrol. Earth Syst. Sci.*, 22, <https://doi.org/10.5194/hess-22-2425-2018>, 2018.
- Dao, D. A., Kim, D., Kim, S., and Park, J.: Determination of flood-inducing rainfall and runoff for highly urbanized area based on high-resolution radar-gauge composite rainfall data and flooded area GIS data, *J. Hydrol.*, 584, 124–140, <https://doi.org/10.1016/j.jhydrol.2020.124704>, 2020.
- 295 DWA: Klimawandel erfordert wassersensible Stadtentwicklung, *Korrespondenz Wasserwirtschaft*, 8, 2015.
- DWD: Nationaler Klimareport, 4. korrigierte Auflage, www.dwd.de/nationalerklimateport, 2020.
- Dyck, S. and Peschke, G.: *Grundlagen der Hydrologie*, Verlag für Bauwesen, <https://books.google.de/books?id=JYBZJgAACAAJ>, 1995.
- 300 Fabry, F.: On the determination of scale ranges for precipitation fields, *J. Geophys. Res. Atmos.*, 101, 12 819–12 826, <https://doi.org/10.1029/96JD00718>, <http://doi.wiley.com/10.1029/96JD00718>, 1996.
- Fadhel, S., Rico-Ramirez, M. A., and Han, D.: Uncertainty of Intensity–Duration–Frequency (IDF) curves due to varied climate baseline periods, *J. Hydrol.*, 547, 600–612, <https://doi.org/10.1016/j.jhydrol.2017.02.013>, https://ac.els-cdn.com/S0022169417300926/1-s2.0-S0022169417300926-main.pdf?_tid=cf511a0c-fad4-11e7-a06d-00000aac361&acdnat=1516117938_9f0e0520549b5f62fb5ed0c6bde1bf1bhttp://www.sciencedirect.com/science/article/pii/S0022169417300926http://files/3593/Fa, 2017.
- 305 Gado, T. A., Hsu, K., and Sorooshian, S.: Rainfall frequency analysis for ungauged sites using satellite precipitation products, *J. Hydrol.*, 554, 646–655, <https://doi.org/10.1016/j.jhydrol.2017.09.043>, <https://doi.org/10.1016/j.jhydrol.2017.09.043>, 2017.
- Galmarini, S., Steyn, D. G., and Ainslie, B.: The scaling law relating world point-precipitation records to duration, *Int. J. Climatol.*, 24, 533–546, <https://doi.org/10.1002/joc.1022>, 2004.
- 310 García-Marín, A. P., Ayuso-Muñoz, J. L., Jiménez-Hornero, F. J., and Estévez, J.: Selecting the best IDF model by using the multifractal approach, *Hydrol. Process.*, 27, 433–443, <https://doi.org/10.1002/hyp.9272>, 2012.
- Ghanmi, H., Bargaoui, Z., and Mallet, C.: Estimation of intensity-duration-frequency relationships according to the property of scale invariance and regionalization analysis in a Mediterranean coastal area, *J. Hydrol.*, 541, 38–49, <https://doi.org/10.1016/j.jhydrol.2016.07.002>, 2016.

- 315 Gires, A., Tchiguirinskaia, I., Schertzer, D., Schellart, A., Berne, A., and Lovejoy, S.: Influence of small scale rainfall variability on standard comparison tools between radar and rain gauge data, *Atmos. Res.*, 138, 125–138, <https://doi.org/10.1016/j.atmosres.2013.11.008>, 2014.
- Gonzalez, S. and Bech, J.: Extreme point rainfall temporal scaling: a long term (1805–2014) regional and seasonal analysis in Spain, *Int. J. Climatol.*, 37, 5068–5079, <https://doi.org/10.1002/joc.5144>, 2017.
- Guerreiro, S. B., Fowler, H. J., Barbero, R., Westra, S., Lenderink, G., Blenkinsop, S., Lewis, E., and Li, X.-F.: Detection of continental-scale intensification of hourly rainfall extremes, *Nature Climate Change*, 8, 803–807, <https://doi.org/10.1038/s41558-018-0245-3>, 2018.
- 320 Hijmans, R., Garcia, N., and Weiczorek, J.: GADM: database of global administrative areas (<http://gadm.org> version 3.6), 2018.
- Jennings, A. H.: WORLD ' S GREATEST OBSERVED POINT RAINFALLS, *Mon. Weather Rev.*, 1950.
- Kim, J., Lee, J., Kim, D., and Kang, B.: The role of rainfall spatial variability in estimating areal reduction factors, *J. Hydrol.*, 568, 416–426, <https://doi.org/10.1016/j.jhydrol.2018.11.014>, 2019.
- 325 Klik, M.: fst: Lightning Fast Serialization of Data Frames for R, <https://CRAN.R-project.org/package=fst>, r package version 0.9.0 — For new features, see the 'Changelog' file (package source), 2019.
- Kreklow, J., Tetzlaff, B., Kuhnt, G., and Burkhard, B.: A Rainfall Data Intercomparison Dataset of RADKLIM, RADOLAN, and Rain Gauge Data for Germany, *Data*, 4, 118, <https://doi.org/10.3390/data4030118>, 2019.
- Lee, J., Ahn, J., Choi, E., and Kim, D.: Mesoscale Spatial Variability of Linear Trend of Precipitation Statistics in Korean Peninsula, *Adv. Meteorol.*, 2016, 1–15, <https://doi.org/10.1155/2016/3809719>, 2016.
- 330 Lengfeld, K., Winterrath, T., Junghänel, T., Hafer, M., and Becker, A.: Characteristic spatial extent of hourly and daily precipitation events in Germany derived from 16 years of radar data, *Meteorol. Z.*, pp. 363–378, <https://doi.org/10.1127/metz/2019/0964>, 2019.
- Lengfeld, K., Kirstetter, P.-E., Fowler, H. J., Yu, J., Becker, A., Flamig, Z., and Gourley, J.: Use of radar data for characterizing extreme precipitation at fine scales and short durations, *Environ. Res. Lett.*, 15, <https://doi.org/10.1088/1748-9326/ab98b4>, 2020.
- 335 Madsen, H., Arnbjerg-Nielsen, K., and Mikkelsen, P. S.: Update of regional intensity–duration–frequency curves in Denmark: Tendency towards increased storm intensities, *Atmos. Res.*, 92, 343–349, <https://doi.org/10.1016/j.atmosres.2009.01.013>, 2009.
- Marra, F. and Morin, E.: Use of radar QPE for the derivation of Intensity–Duration–Frequency curves in a range of climatic regimes, *J. Hydrol.*, 531, 427–440, <https://doi.org/10.1016/j.jhydrol.2015.08.064>, 2015.
- Marra, F., Morin, E., Peleg, N., Mei, Y., and Anagnostou, E. N.: Intensity-duration-frequency curves from remote sensing rainfall estimates: comparing satellite and weather radar over the eastern Mediterranean, *Hydrol. Earth Syst. Sci.*, 21, 2389–2404, <https://doi.org/10.5194/hess-21-2389-2017>, 2017.
- 340 NWS: World record point precipitation measurements, http://www.nws.noaa.gov/oh/hdsc/record_precip/record_precip_world.html, 2017.
- Overeem, A., Buishand, T. A., and Holleman, I.: Extreme rainfall analysis and estimation of depth-duration-frequency curves using weather radar, *Water Resour. Res.*, 45, <https://doi.org/10.1029/2009wr007869>, 2009.
- 345 Papalexiou, S. M., Dialynas, Y. G., and Grimaldi, S.: Hershfield factor revisited: Correcting annual maximum precipitation, *J. Hydrol.*, 542, 884–895, <https://doi.org/10.1016/j.jhydrol.2016.09.058>, 2016.
- Paulhus, J. L. H.: INDIAN OCEAN AND TAIWAN RAINFALLS SET NEW RECORDS, *Mon. Weather Rev.*, 93, 331–335, 1965.
- Peleg, N., Ben-Asher, M., and Morin, E.: Radar subpixel-scale rainfall variability and uncertainty: lessons learned from observations of a dense rain-gauge network, *Hydrol. Earth Syst. Sci.*, 17, 2195–2208, <https://doi.org/10.5194/hess-17-2195-2013>, 2013.
- 350 Peleg, N., Marra, F., Fatichi, S., Paschalis, A., Molnar, P., and Burlando, P.: Spatial variability of extreme rainfall at radar subpixel scale, *J. Hydrol.*, 556, 922–933, <https://doi.org/10.1016/j.jhydrol.2016.05.033>, 2018.

- Perpiñán, O. and Hijmans, R.: rasterVis, <http://oscarperpinan.github.io/rastervis/>, r package version 0.46 — For new features, see the 'Changelog' file (in the package source), 2019.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, 2019.
- 355 Rudolf, B. and Rapp, J.: Das Jahrhunderthochwasser der Elbe: Synoptische Wetterentwicklung und klimatologische Aspekte, Abdruck aus klimastatusbericht 2002, DWD, <https://pdfs.semanticscholar.org/fdfc/a0eb2c7ac37d2d80ddd2700b3f710a7fed79.pdf>, 2003.
- Scott, A. J. and Knott, M.: A Cluster Analysis Method for Grouping Means in the Analysis of Variance, *Biometrics*, 30, 507–512, <https://doi.org/10.2307/2529204>, 1974.
- 360 Ushey, K.: RcppRoll: Efficient Rolling / Windowed Operations, <https://CRAN.R-project.org/package=RcppRoll>, r package version 0.3.0 — For new features, see the 'Changelog' file (in the package source), 2018.
- Westra, S., Alexander, L. V., and Zwiers, F. W.: Global increasing trends in annual maximum daily precipitation, *J. Clim.*, 26, 3904–3918, <https://doi.org/10.1175/JCLI-D-12-00502.1>, 2013.
- Westra, S., Fowler, H. J., Evans, J. P., Alexander, L. V., Berg, P., Johnson, F., Kendon, E. J., Lenderink, G., and Roberts, N. M.: Future changes
365 to the intensity and frequency of short-duration extreme rainfall, *Rev. Geophys.*, 52, 522–555, <https://doi.org/10.1002/2014rg000464>, 2014.
- Wickham, H.: ggplot2: Elegant Graphics for Data Analysis, Springer-Verlag New York, <https://ggplot2.tidyverse.org>, 2016.
- Winterrath, T., Rosenow, W., and Weigl, E.: On the DWD Quantitative Precipitation Analysis and Nowcasting System for Real-Time Application in German Flood Risk Management, in: *Weather Radar and Hydrology*, Proceedings of a symposium held in Exeter, UK, April
370 2011, IAHS Publ. 351, pp. 323–329, 2012.
- Winterrath, T., Brendel, C., Hafer, M., Junghänel, T., Klameth, A., Walawender, E., Weigl, E., and Becker, A.: Erstellung einer radargestützten Niederschlagsklimatologie, 2017.
- Winterrath, T., Brendel, C., Hafer, M., Junghänel, T., Klameth, A., Lengfeld, K., Walawender, E., Weigl, E., and Becker, A.: Radar climatology (RADKLIM) version 2017.002: Reprocessed gauge-adjusted radar data, one-hour precipitation sums (RW),
375 https://doi.org/10.5676/dwd/radklm_rw_v2017.002, 2018a.
- Winterrath, T., Brendel, C., Hafer, M., Junghänel, T., Klameth, A., Lengfeld, K., Walawender, E., Weigl, E., and Becker, A.: Radar climatology (RADKLIM) version 2017.002: Reprocessed quasi gauge-adjusted radar data, 5-minute precipitation sums (YW), https://doi.org/10.5676/dwd/radklm_yw_v2017.002, 2018b.
- World Meteorological Organization: Guide to hydrological practices, 1994.
- 380 Yang, Z.-Y., Pourghasemi, H. R., and Lee, Y.-H.: Fractal analysis of rainfall-induced landslide and debris flow spread distribution in the Chenyulan Creek Basin, Taiwan, *J. Earth Sci.*, 27, 151–159, <https://doi.org/10.1007/s12583-016-0633-4>, 2016.
- Zhang, H., Fraedrich, K., Zhu, X., Blender, R., and Zhang, L.: World's Greatest Observed Point Rainfalls: Jennings (1950) Scaling Law, *J. Hydrometeorol.*, 14, 1952–1957, <https://doi.org/10.1175/JHM-D-13-074.1>, 2013.