

The authors would like to thank the reviewer for his thoughtful and useful comments on our paper. We have considered all the suggestions. Our point-by-point responses (R) to comments and questions (Q) are detailed below:

Q1:

The paper under review is devoted to the experimental study of nonlinear interactions in the focused wave packet by using the wavelet bi-spectral analysis. This experiment continues a series of studies done by these authors with the investigation the nonlinear dispersive mechanism of the rogue wave formation. In their experiments, the authors used the wave trains with various spectra (Pierson-Moskowitz and JONSWAP) in intermediate depth. Strong nonlinear effects are observed on the sloping beach as it is expected. Nonlinear energy transfer in the high-frequency region is analyzed. In fact, such processes have been actively studied earlier, and, perhaps, the novel moment here is the demonstration of nonlinear effects through the bi-spectral analysis. *To add to this, I would like the authors to formulate the obtained results in Conclusion better underlying their difference from the known results.*

R1:

We would like to thank the reviewer for his remark. We tried to formulate and synthesise the obtained results in order to distinguish similarities and differences from the known results. Here is the conclusion; the added text is in red. Redundant sentences have been deleted as well.

An experimental approach is proposed for determining the non-linear wave-wave interactions, which accompany the propagation of large amplitude wave trains, that might cause damage to coastal zones, marine structures and navigation vessels. We investigate seven focused wave trains derived from JONSWAP ($\gamma = 3.3$ or 7) and Pierson-Moskowitz spectra propagating from intermediate water depth to the inner surf zone. The results presented in this study extend the parameter range of observations of triad interactions. ~~The experimental conditions were selected based on two parameters: the wave steepness and the spectrum type. Focusing waves were generated in a physical wave flume by modifying the wave spectrum and steepness.~~ The present data were collected in intermediate water with a $k_p h_0$ varying between 0.92 and 0.79 . A typical wave train consists of a large number of waves interacting with one another. Wavelet-based bicoherence is used to investigate the phase coupling between frequency components of short time series. Some consequences of non-linear transfer are briefly discussed; in particular the role played by non-linear interactions in shaping the high frequency part of the spectrum, the relative contribution of each harmonic and the downshifting of the peak spectrum demonstrated in previous studies. *Note that our experimental study is different from previous experiments (Dong et al., 2008; Ma et al., 2010) regarding the slope geometry and most importantly, the use of three different spectral types.*

Along the flat bottom ($4\text{ m} < x < 9.5\text{ m}$), one might assume that the influence of triad interactions is very weak for the three considered spectra. The bispectral analysis of the data shows that as the waves propagate along the flat bottom, the magnitude of the bicoherence increases slightly (between 0% and 20% of its initial value). Moreover, this is foreseeable because the spectrum and the wave train shape do not substantially change along the flat bottom and a small amount of energy is transferred from the peak region to high frequency components. ~~The spectra remain approximately unimodal and do not present a clear second or third harmonic and non-linear interactions concern only the peak and low frequency regions. On the flat bottom, the wave train is still dispersive and consequently, bound interactions are responsible for the slight growth of $b^2(f_p, f_p)$ and $b^2(0.5f_p, 0.5f_p)$.~~

When the wave train reaches the slope ($9.5 \text{ m} < x < x_b$), wave-wave interactions among high order harmonics increase rapidly and reach the maximum degree in the breaking/focus location. In line with previous studies (Elsayed, 2006; Dong et al., 2008; Ma et al., 2010), strong nonlinear interactions were predominantly observed in the shallower region. The analysis showed a gradual broadening of the bicoherence spectrum, which is in accordance with previous studies who demonstrated that the energy is transferred mainly to high frequencies regions (Tian et al. 2011; Abroug et al., 2020). This is partly due to significant spectral transformations which are more important during the shoaling process. Particularly, this analysis showed a considerable contribution of 2nd and 3rd harmonics for unidirectional steep wave trains and the spectral components at the second harmonic $2f_p$ have increased substantially (6 times its initial value). The bispectral analysis results show that the wave non-linearity S_0 plays an important role in the increasing trend of phase coupling, which is more important for wave trains having strong non-linearities. This last finding agrees well with the conclusions made by Ma et al. (2010).

An innovative aspect of this paper is presenting wavelet-based bispectral analysis for highly non-linear intermediate water waves with different spectral types. If we compare the three spectra, we can see that all nonlinear interactions on the flat bottom ($x < 9.5 \text{ m}$) are weak ($b^2 < 0.15$) in the case of wide spectrum wave trains (Tests 2 and 3 Fig. 11). However, in the case of narrower spectra, more frequencies (e.g. f_p , $2f_p$ and $3f_p$) are implicated in the focusing process (Tests 4, 5, 6 and 7 Fig. 11) and the corresponding phase coupling is higher ($b^2 > 0.2$). This finding is in agreement with the stable behavior of wide spectrum wave trains, which was demonstrated experimentally in Abroug et al. (2019) and Stansberg (1994). In intermediate water depth ($0.79 < k_p h < 0.92$), wide spectrum harmonics (f_p , $2f_p$, $3f_p \dots$) are less implicated in the focusing process compared to narrow spectrum harmonics. In shallow water regions ($9.5 \text{ m} < x < x_b$) and after breaking ($x_b < x$), the spatial evolution of the phase coupling is qualitatively similar for the three spectra.

The results obtained in this study show important features in wave-wave interactions during the propagation of focused waves. This study strengthens the usefulness of wavelet-based analysis in detecting features that are hidden in a Fourier-based analysis, and in explaining a number of phenomena, such as the process leading to wave breaking and the energy transfer between wave components. Nevertheless, in order to confirm the use of wavelet-based bicoherence for more realistic 3D studies with structures, efforts should be made to expand this study for example by investigating greater water depths, higher steepness and wider spectra. Furthermore, the observed evolution of bicoherence for focused waves should be compared to that of waves with similar steepness and bandwidth but with initial random distribution of phase. In other words, efforts should be made to identify and quantify the phase coupling differences between focusing wave trains and non-focusing waves. Information concerning the phase coherence can be obtained by calculating the biphasic parameter (β (a1, a2), Ma et al., 2010). It would be interesting to quantitatively measure the deviation of biphasic values between primary waves/higher harmonics and to analyse their spatial evolution through different spectra to distinguish differences. Finally, a detailed study of how bound energy at harmonics would be influenced by quadruplet interactions should be performed.

Q2: Equations (2) and (3) are written inaccurately. Function (3) does not contain tau and the parameter a.

R2:

We would like to thank the reviewer for this comment. I added Eq 4, which contains the two parameters tau and a, in order to make Eq 3 more explicit (Line 130). Equation 4 has been added between the continuous wavelet transform $WT(a, \tau)$ function and the Morlet wavelet function.

$$\psi_{a,\tau}(t) = |a|^{-0.5} \psi\left(\frac{t-\tau}{a}\right)$$

Q3:

For the wave focusing, it is necessary to vary the local frequency on the specific law for intermediate depth. I do not understand which formula for the local frequency versus time has been used. Perhaps, by using the optimal law, the focusing can occur on the flat bottom. If there is no specific focusing, there is an interference. Moreover, it should be reflected in the title.

R3:

It is an interesting comment and maybe the methodology of generation needs more clarifications from the authors. Here is the explanation, which was added to the manuscript.

Line (85-89)

The linear NewWave theory (Tromans et al. (1991)), which is able to generate targeted waves at a prescribed location and time by combining sinusoidal components of different frequencies, is used as input for the generated focused wave trains. This theory was validated at deep water locations, at intermediate water depth locations (Taylor and Williams (2004)) and at coastal regions (Whittaker et al. (2016); for $kh < 0.5$). In NewWave theory, the expected shape of a wave train is the autocorrelation function (Fourier Transform of the spectral density).

Line (100-111)

Using linear NewWave theory, the free surface elevation of a wave train at a distance x from the wavemaker can be written as follows:

$$\eta(x, t) = \sum_{i=1}^N a_i \cos[k_i(x - x_0) - \omega_i(t - t_0)] \quad (1)$$

$$a_i = A_0 \frac{S(f_i)\Delta f}{\sum_{i=1}^N S(f_i)\Delta f} \quad (2)$$

where a_i (Eq. (2)) is the amplitude of each component, i varies from 1 to N (number of waves), x_0 and t_0 denote respectively the predefined focal location and focal time, $k_i = \omega_i / g \tanh(k_i h)$ is the wavenumber, $\omega_i = 2\pi f_i$ is the angular frequency, h is the water depth, A_0 represents the theoretical linear crest amplitude of the wave train, $S(f_i)$ is the spectral density and $\Delta f = \frac{f_{max} - f_{min}}{N-1}$ is the frequency step. The wave group is generated with a given linear focus position ($x = 12$ m from the wave maker) based on linear focusing in a constant water depth. **So to answer your question, yes, by modifying $x < 9.5$ m in the EDL Software, we can obtain a focusing on the flat bottom.**

JONSWAP and Pierson-Moskowitz are the two spectra used to represent the sea state. All generated waves are crested focused waves, i.e. the phase angle of the wave group within its envelope at the focus position is equal to zero.

The authors would like to thank the reviewer for his thoughtful and useful comments on our paper. We have considered all the suggestions. Our point-by-point responses (R) to comments and questions (Q) are detailed below:

Q1:

This paper present a nonlinear phase coupling analysis of focusing wave groups propagating over a slope. Focusing wave groups with three different spectra , Gaussian, P-M, JONSWAP were generated in a wave flume. ***But, only three wave groups are discussed.*** Actually, the topic of this study is not new. Spectral and bi-spectral analysis of irregular waves over shoal have been presented previously. ***I think the new aspect of this paper is presenting bispectral analysis for waves with different spectral types.*** My comments are outlined as follow.

R1:

We would like to thank the reviewer for the comment. One wave train from each spectrum was selected to be detailed in the text in order to avoid, as much as possible, unduly redundant results and text. Each selected wave train has the strongest steepness in its spectrum type. An important objective of this study was to investigate strong steepness wave trains. The spatial evolution of bicoherence of the other four wave trains are summarised in the last figure (Figure 11) in order to investigate the impact of the steepness on the spatial evolution of the phase coupling.

A new paragraph has been added in the conclusion (Line 336-344) in order to summarise the new aspect of this paper, which is as mentioned by the reviewer “presenting bispectral analysis for waves with different spectral types”.

Q2:

Page 2, lines 47-47. The authors argues that "It is important to mention here that resonant interactions are not easily achieved in unidirectional wave train propagation since the resonant conditions cannot not be satisfied in a small area. ", I think it is better to specify that this is only true in shallow water. As for deep or intermediate water depth, resonant interactions is very strong in unidirectional wave trains.

R2:

We completely agree with the reviewer. We have added ‘shallow water regions’ to this sentence. Line 47.

Q3:

The reference Guohai et al. 2008 should be Dong et al. (2008).

R3:

I have corrected this error throughout the manuscript (Lines 35, 51, 62, 142, 143, 151, 154, 308 and 327).

Q4:

Fig. 3, There is a mistake in caption: JONSWAP (gamma = 7) (Test 7)

R4:

I have corrected this error and changed $\gamma = 3.3$ to $\gamma = 7$ (Line 199).

Q5:

Where is the incipient breaking points and spatial range should be presented in the text.

R5:

I would like to thank the reviewer for this comment. The presence of the incipient breaking points and spatial range in the text is crucial for a better understanding and have been added in lines: 186, 186, 187, 202, 237, 247, 259, 284 and 285.

Q6

The authors should concentrate on the analysis of difference biphas coupling for different spectra?

R6

I agree with the reviewer. Information concerning the phase coherence can be obtained by calculating the biphas parameter ($\beta(a_1, a_2)$). It will be interesting to quantitatively measure the deviation of biphas values between primary waves/higher harmonics and to analyse their spatial evolution through different spectra to distinguish differences. This text was added to the manuscript (conclusion and perspectives) (Line 353-355)

In this study, we wanted to exclusively study non-linear wave interactions through the spatial evolution of the bicoherence. In our future work, we will continue to examine this issue on selected high nonlinear wave groups by investigating accurately the spatial evolution of the biphas. It would also be useful to compare it to the results found in Ma et al. (2010).

Laboratory study of non-linear wave-wave interactions of extreme focused waves in the nearshore zone

Iskander Abroug ^{1,2}, Nizar Abcha ¹, Armelle Jarno ², François Marin ²

¹ Normandie Université, UNICAEN, UNIROUEN, CNRS, UMR 6143 M2C, 14000 Caen, France.

5 ² Normandie Université, UNILEHAVRE, CNRS, UMR 6294 LOMC, 76600 Le Havre, France.

Correspondence to: Iskander Abroug (iskander.abroug@unicaen.fr)

Abstract. Extreme waves play a crucial role in marine inundation hazards and coastal erosion. In this article, we experimentally study nonlinear wave-wave interactions of large amplitude focused wave groups propagating in a two-dimensional wave flume over a mild slope ($\beta = 1:25$). The influence of the frequency spectrum and the steepness on the nonlinear interactions of focused waves are examined. The generated wave trains correspond to Pierson-Moskowitz and JONSWAP ($\gamma = 3.3$ or $\gamma = 7$) spectra. Subsequently, we experimentally approach this problem by the use of a bispectral analysis applied on short time series, via the wavelet-based bicoherence parameter, which identifies and quantifies the phase coupling resulting from non-resonant or bound triad interactions with the peak frequency. The bispectral analysis shows that the number of frequency components involved in the focusing process and the resulting phase coupling increases gradually and approaches 1 just prior to breaking, accordingly with the spectrum broadening and the energy increase in high frequency components. Downstream breaking, the values of phase coupling between the peak frequency and its higher harmonics decrease drastically and the bicoherence spectrum becomes less structured.

1 Introduction

Extreme wave propagation is a highly nonlinear process observed in both open seas and coastal regions. The main physical mechanisms which may lead to an extreme wave event are illustrated in Kharif and Pelinovsky, 2003; Kharif et al., 2009; Didenkulova et al., 2010; Onorato et al., 2013. Extreme waves may occur in deep or shallow water, in energetic storm sea state or previously calm sea state. In our opinion, spatio-temporal wave focusing is one of the most important mechanisms in the extreme wave formation for shallow and deep water (Kharif and Pelinovsky, 2003). The spatio-temporal wave focusing is a classical mechanism giving rise to an important wave energy concentration in a small region. If the wave height of the focusing group exceeds 2.2 times its significant wave height, it can be defined as a rogue or freak wave (Dysthe et al., 2008). For this reason, spatio-temporal wave focusing is often employed in laboratory wave flumes with a wide variation of water depth (Merkoune et al., 2013), spectrum type (Tian et al., 2011; Xu et al., 2019; Abroug et al., 2019; 2020) and wavelength to depth ratio, in order to better understand the generation process, the dynamic behavior and the hydrodynamic loads on ocean structures in extreme sea conditions.

30 Over the past years, several studies have attempted to quantify the spatial evolution of spectral energy of unidirectional wave groups in experimental wave flumes using a classic Fourier analysis (Tian et al., 2011; Liang et al., 2017; Abroug et al., 2020). The frequency spectrum only gives the distribution of energy in the frequency domain; however, information about the phase coupling between different wave components is unknown. Consequently, higher order spectrum techniques should be adopted. A powerful tool to investigate the highly nonlinear process is the wavelet-based bispectral technique, which has been used in

35 several works to study the non-linear interactions and quadratic phase coupling between wave components (Dong et al., 2008; Ma et al., 2010). The need to detect and quantify second order non-linear interactions can be found in many disciplines, such as, geophysics (Grinsted et al., 2004), plasma physics (Milligan et al., 1995), fault diagnosis (Li et al., 2014), health-related areas, neuroscience (Bai et al., 2017) and wave analysis (Eldeberky, 1996; Eldeberky and Madsen, 1999; Young et al., 1996; Young and Eldeberky, 1998; Becq-Girard et al., 1999; Huseni and Balaji, 2017; Zhang et al., 2019). In wave analysis, the

40 propagation of wave trains in the nearshore zone has an exceptionally high spectral and temporal resolution. The majority of previous works regarding the evolution of unidirectional wave trains in numerical and experimental wave flumes have shown that spatio-temporal focusing leads to a shape and elevation of a wave crest at focus that cannot be predicted by either linear or 2nd order wave theory. This is due to high order nonlinearities, called the bound (harmonics) and resonant nonlinearities (Vyzikas et al., 2018). On the one hand, bound nonlinearities are the result of non-linear harmonics that are

45 phase-locked to the wave train and contribute in the sharpening of free surface elevation. On the other hand, resonant interactions contribute in the redistribution of energy among different frequency components. It is important to mention here that in shallow water regions exact resonant interactions are hardly realised in unidirectional propagation because the resonant conditions cannot not be satisfied in a small area. Therefore, we investigate specifically the role of bound waves generated by non-resonant three-wave coupling.

50 Over the past few decades, various experimental studies have investigated the spatial evolution of non-linear coupling between wave components. Dong et al., 2008 studied the spatial evolution of non-linear interactions between different wave components in the shoaling and de-shoaling region by carrying out two random waves experiments based on JONSWAP spectra with varying peak waves periods and root-mean-square waves heights. They showed that the degree of quadratic phase coupling increases in the shoaling region and achieves its highest level prior to wave breaking. Ma et al., 2010 studied

55 experimentally JONSWAP wave trains propagating in intermediate water depth. Recently, nonlinear transformation of unidirectional irregular waves propagating over a complex bathymetry ($1.06 < k_p h < 2.2$; where k_p is the peak wavenumber and h denotes the water depth) was performed in Zhang et al., 2019, who studied the triad wave-wave non-linear interactions in the case of long records of JONSWAP irregular waves ($1200 T_p$, where T_p is the peak period) using a Fourier-based bispectral analysis. They found that the phase coupling is strong near the end of the slope, where second and third harmonics

60 become more important. They also noticed the appearance of low-frequency waves generated by the difference interactions during wave propagation. We must note here that the main difference between Fourier-based bicoherence and wavelet-based bicoherence is the number of degrees of freedom (Dong et al., 2008). Wavelet-based bicoherence is a suitable tool to detect

nonlinear wave-wave interactions occurring in relatively short data sequences, and can be used to analyse data collected in laboratory flumes (Elsayed, 2006).

- 65 Most of the aforementioned studies were conducted in random wave conditions based on JONSWAP spectra. To the author's knowledge, few studies have attempted to quantify the degree of phase coupling resulting from the propagation of realistic spectrum wave trains in the nearshore zone using wavelet-based bicoherence. Experiments are performed on numerous Pierson-Moskowitz and JONSWAP wave trains propagating from a constant intermediate water depth to shoaling and breaking zones.
- 70 The paper is outlined as follows. The experimental set-up and test conditions are illustrated in section 2. In section 3, a short formal description of wavelet analysis and wavelet-based bicoherence is provided. The spatial evolution of wavelet-based bicoherence is discussed in section 4. Section 5 is devoted to conclusions and perspectives.

2 Experimental setup and wave train parameters

- The following is a brief consideration of present wave trains generation; more details of the experiments can be found in
- 75 Abroug et al., 2020. The experiments were conducted in a two-dimensional wave flume of the M2C (Morphodynamique Continentale et côtière) Laboratory at Caen University, France. The flume is 22 m long, 0.8 wide and the water depth is $h_0 = 0.3$ m (Fig. 1). In this study, the relative water depth $k_p h_0 < 1.363$ is verified, which means that the modulation instability effect can be neglected (Janssen et al., 2007; Fedele et al., 2019). An Edinburgh Designs Ltd piston type wavemaker is located at one end of the flume to implement wave trains using linear wave generation signal. Wave trains are generated with almost
- 80 no reflection at the end of the flume, since measurements are performed before reflected waves travel back to the measurement location. Thus, the occurrence of resonant interactions potentially driven by reflected waves is limited and we only focus on bound waves.

- The data used in this work are issued from Abroug et al., 2020. The present study relates to seven wave train simulations based on the averaged JONSWAP spectra (i.e. with peak factor $\gamma = 3.3$ or 7) or Pierson-Moskowitz spectra with varied peak wave
- 85 periods f_p and wave steepnesses S_0 (i.e. non-linearity). The linear NewWave theory (Tromans et al. (1991)), which is able to generate targeted waves at a prescribed location and time by combining sinusoidal components of different frequencies, is used as input for the generated focused wave trains. This theory was validated at deep water locations, at intermediate water depth locations (Taylor and Williams (2004)) and at coastal regions (Whittaker et al. (2016); for $kh < 0.5$). In NewWave theory, the expected shape of a wave train is the autocorrelation function (Fourier Transform of the spectral density).

- 90 For each wave train, a large number of wave signals were recorded along the flume to accurately follow the wave evolution in space. The surface elevation is measured by two aligned wave gauges located from the longitudinal coordinate $x_{\min} = 4$ m to $x_{\max} = 14$ m, where $x = 0$ is defined as the mean position of the wavemaker. The positions of these wave gauges are clearly delineated in Fig. 1. The sampling rate is 50Hz and each record duration is 35 s with a sample interval of 0.02 s. The fast Fourier transform (FFT) was applied to each signal, resulting in 1750 frequency components over the range $[0, 3f_p]$ and with

95 a spectral resolution $\Delta f = 0.023$ Hz. The distance from the wave maker for the focusing point was set to 12 m from the wave maker.




Figure 1: Schematic experimental setup, WG1 and WG2 denote wave gauge n°1 and wave gauge n°2 respectively.

100 Using linear NewWave theory, the free surface elevation of a wave train at a distance x from the wavemaker can be written as follows:

$$\eta(x, t) = \sum_{i=1}^N a_i \cos[k_i(x - x_0) - \omega_i(t - t_0)] \quad (1)$$

$$a_i = A_0 \frac{S(f_i)\Delta f}{\sum_{i=1}^N S(f_i)\Delta f} \quad (2)$$

105

where a_i (Eq. (2)) is the amplitude of each component, i varies from 1 to N (number of waves), x_0 and t_0 denote respectively the predefined focal location and focal time, $k_i = \omega_i / g \tanh(k_i h)$ is the wavenumber, $\omega_i = 2\pi f_i$ is the angular frequency, h is the water depth, A_0 represents the theoretical linear crest amplitude of the wave train, $S(f_i)$ is the spectral density and $\Delta f = \frac{f_{\max} - f_{\min}}{N-1}$ is the frequency step. JONSWAP and Pierson-Moskowitz are the two used spectra to represent the sea state.

110 All generated waves are crested focused waves, i.e. the phase angle of the wave group within its envelope at the focus position is equal to zero.

Based on Eq. (1), the varied parameters during these experiments were the spectrum type ($S(f_i)$) and the wave steepness S_0 . The peak frequency parameter was chosen in order to have a relative depth $k_p h_0$ varying between 0.79 and 0.92 (Deep side in Table 1). Deep and shallow sides in Table 1 represent respectively the flat bottom depth ($4 \text{ m} < x < 9.5 \text{ m}$) and the shallowest studied depth ($x = 14 \text{ m}$). Five of the studied wave trains have more than one breaking and breaking locations x_b are indicated as bracketed intervals in Table 1.

115

120

Table 1. Wave trains key parameters

Test	f_p (Hz)	S_0	Spectrum type	x_b (m)	$k_p h_0$	$k_p h$
					Deep Side	Shallow Side
1	0.70	0.19	Gaussian	[11.85;12.55]	0.84	0.34
2	0.66	0.14	Pierson-Moskowitz	12.9	0.79	0.31
3	0.66	0.28	Pierson-Moskowitz	[11.09;11.82]	0.79	0.31
4	0.75	0.25	JONSWAP ($\gamma=3.3$)	[12.13;12.81]	0.92	0.37
5	0.75	0.38	JONSWAP ($\gamma=3.3$)	[10.5;11.61]	0.92	0.37
6	0.75	0.11	JONSWAP ($\gamma=7$)	13.5	0.92	0.37
7	0.75	0.23	JONSWAP ($\gamma=7$)	[12.07;12.69]	0.92	0.37

3 Wavelet-based analysis

The free surface elevation of each wave train was studied through the bispectral analysis applied on short time series, via the wavelet-based bicoherence. The detailed characteristics of the wavelet-based bicoherence can be found in Van Milligen et al., 1995 and a brief introduction of this technique is given below. The continuous wavelet transform $WT(a, \tau)$ of a time series $f(t)$ is calculated as:

$$WT(a, \tau) = \int_{-\infty}^{+\infty} f(t) \psi_{a,\tau}^* dt \quad (3)$$

$$\psi_{a,\tau}(t) = |a|^{-0.5} \psi\left(\frac{t-\tau}{a}\right) \quad (4)$$

where the asterisk denotes the complex conjugate and $\psi_{a,\tau}$ (Eq. (4)) represents the mother wavelet function dilated by a factor τ and scaled by a factor a , $a > 0$. The latter parameter can be interpreted as the frequency inverse, i.e. $f = 1/a$. The wavelet transform can be interpreted as a series of bandpass filter of the time series with a mother wavelet function. We have chosen the Morlet wavelet as a mother wavelet function because it provides information about phase and amplitude, and it is adapted for capturing coherence between harmonic components. The Morlet wavelet can be considered as a modulated Gaussian waveform and is defined as:

$$\psi(t) = \pi^{-1/4} e^{-\frac{t^2}{2}} e^{(i\omega_0 t)} \quad (5)$$

where ω_0 denotes the dimensionless frequency and t is the dimensionless time. The Morlet wavelet with $\omega_0 = 6$ is a good choice, since it ensures a good balance between time and frequency localisation (Grinsted et al., 2004; Dong et al., 2008). For the Morlet wavelet the scale a is almost equal to the Fourier period T ($T = 1.03 a$). As mentioned in Dong et al., 2008, it is convenient to write the scales a as fractional powers of two (Torrence and Compo, 1998):

145

$$a_i = a_0 2^{i\delta}, i = 0, 1, 2, \dots, M \quad (6)$$

$$M = \frac{1}{\delta} \log_2 \left(\frac{N \Delta t}{a_0} \right) \quad (7)$$

150 where a_0 is the smallest resolvable scale, M represents the largest scale and δ denotes the scale factor. The a_0 parameter should be chosen equal to $2 \times \Delta t$ (Torrence and compo, 1998; Dong et al., 2008). N and Δt represent respectively the number of points in the times series and the time sampling. The scale factor δ should be sufficiently small to provide high resolution and adequate sampling in scale. Moreover, for the Morlet wavelet, a scale factor $\delta = 0.5$ is the largest value that gives adequate sampling (Dong et al., 2008). It is for that reason that we opted for a scale factor $\delta = 0.02$, giving a total of 395 scales ranging from 0.04
155 up to 11.83 for respectively high and low frequency. The wavelet-based bispectrum (Eq. (8)) measures the phase coupling in the interval $\Delta T = 35$ s that occurs between f_1, f_2 and f_3 where the latter parameters must satisfy the frequency sum rule (Eq. (9)). Quadratic non-linear coupling occurs between f_1 and f_2 , generating a third component at the sum frequency f_3 .

The bispectrum (Eq. (8)), which is the double Fourier transform of the third-order moment, measures the extent of phase coherence due to the non-linear triad interaction between three waves that satisfy the frequency and phases matching criteria
160 (Eq. (9) and (10)). The estimation of wavelet-based bispectrum in the whole bifrequency plan can be based on its values in the interval $\psi : \{f_1 > f_2 > 0, f_1 + f_2 = f_3 = 25 \text{ Hz (Nyquist sampling frequency)}\}$.

$$B(a_1, a_2) = \int WT_x(f_1, \tau) WT_x(f_2, \tau) WT_x^*(f_3, \tau) d\tau \quad (8)$$

$$165 \quad f_3 = f_1 + f_2 \quad (9)$$

$$\varphi_3 = \varphi_1 + \varphi_2 \quad (10)$$

The wavelet-based bicoherence (Eq. (11)), which can be defined as the normalised wavelet bispectrum, is used in practise to measure the degree of phase coupling (Larsen et al., 2001) and is bounded by 0 and 1 by the Schwarz inequality. A value close
170 to unity reveals a maximum amount of coupling and a value close to zero corresponds to a random phase relation.

$$b^2(a_1, a_2) = \frac{|B(a_1, a_2)|^2}{\left[\int_{t=0}^{t=35} |WT_x(a_1, \tau) WT_x(a_2, \tau)|^2 d\tau \right] \left[\int_{t=0}^{t=35} |WT_x(a_3, \tau)|^2 d\tau \right]} \quad (11)$$

Figure 2 exhibits a simple illustration of the wavelet-based bicoherence of a narrow-banded Gaussian wave train (Test 1) recorded at $x = 4$ m from the wavemaker. The shading indicates the strength of non-linear coupling; dark red ($b^2(f_1, f_2) = 1$) being totally coupled and dark blue ($b^2(f_1, f_2) = 0$) completely uncoupled. The degree of phase coupling is represented by the colorbar indicating the sum interactions between two frequencies. In this manner, a visualisation of the non-linear activity across the wave train propagation is feasible, detecting the frequency sections of the signal that contribute the most to the non-linear activity. The two frequencies f_1 and f_2 are normalised by the peak frequency f_p . Red ($b^2(f_p, f_p)$) and yellow peaks represent the phase coupling of the primary frequency component with its harmonic. In general, a non-null bicoherence $b^2(f_1, f_2) > 0$ means that the $f_3 = f_1 + f_2$ component gains energy from the f_1 and f_2 components.




Figure 2: The wavelet-based bicoherence of a narrow-banded Gaussian wave train (Test 1) at $x = 4$ m.

4 Results and discussions

Figure 3 shows three sets of time series of three wave trains with approximately the same steepness S_0 and derived from Pierson-Moskowitz (Test 3; $x_b \in [11.09; 11.82]$), JONSWAP ($\gamma = 3.3$) (Test 4; $x_b \in [12.13; 12.81]$) and JONSWAP ($\gamma = 7$) (Test 7; $x_b \in [12.07; 12.69]$) spectra at eight different locations along the flume. This preliminary figure shows surface elevation time histories including the first measurement ($x = 4$ m), the propagation along the flat bottom, the shoaling and the breaking of the focused wave group. It should be noted here that the seven studied wave trains are crest-focused wave groups ($\Phi = 0$).

Figure 4 shows, in a log scale, the spatial evolution of the Fourier spectra of the same three wave trains (Tests 3, 4 and 7). A spatial downshift of the spectral peak (Cases 4 and 7), a steepening of the low frequency side and a widening of the high frequency side are illustrated. These spectral variations, identified and quantified in Abroug et al., 2020, concern high and low frequency components. The shift of energy is essentially due to non-linear wave-wave interactions among wave frequency components during the focalisation process. Nevertheless, we do not distinguish which wave components participate in the wave-wave interactions, nor do we distinguish the wave modes that undergo the strongest non-linear interactions. Consequently, the wavelet-based bicoherence is used herein to provide information about the non-linear triad wave interactions that cannot be easily obtained from the Fourier analysis which was used in Abroug et al., 2020.




Figure 3: Three sets of time series of Pierson-Moskowitz (Test 3), JONSWAP ($\gamma = 3.3$) (Test 4) and JONSWAP ($\gamma = 7$) (Test 7) wave trains.

Figure 5 presents the spatial evolution of the wavelet-based bicoherence of a Pierson-Moskowitz wave train (Test 3; $x_b \in [11.09; 11.82]$) along the flat bottom. This figure shows that wave-wave interactions between different modes are weak on flat bottom ($4 \text{ m} < x < 9.5 \text{ m}$; $k_p h_0 = 0.79$) and few frequency components participate in the focusing process. In the intermediate water depth region ($4 \text{ m} < x < 9.5 \text{ m}$), the sea state is almost Gaussian, and for that reason non-linear wave-wave interactions are relatively moderate. For example, $b^2(f_p, f_p) = 0.1$ and $b^2(f_p, 3f_p) = 0.065$ at $x = 4 \text{ m}$, indicate respectively a weak self-self wave interaction at the energy-frequency peak coupled with the energy at $2f_p$ and a very weak wave interaction at the peak frequency coupled with the energy at $4f_p$ (Fig. 5a). A significant bicoherence magnitude band ranging from $0.5f_p$ to f_p is observed, i.e. $b^2(0.5f_p - f_p, 0.5f_p - f_p)$, which indicates an energy transfer from low frequency components to the spectral peak. This partially explains the spatial evolution of the spectrum, namely the increase of energy in the peak region, which is

210 potentially a way of compensating the energy dissipation in the transfer region, i.e. the region between the spectral peak and high frequency regions, (Abroug et al., 2020; Liang et al., 2017). Note that the magnitude of the bicoherence is consistent with the fact that spectrum shape does not vary substantially along the flat bottom ($4 \text{ m} < x < 9.5 \text{ m}$) (Fig. 4).




Figure 4 : Spatial evolution of normalised amplitude spectra in a log scale for Tests 3, 4 and 7.

215 As the wave train approaches the toe of the slope ($x \sim 9.5 \text{ m}$), more and more wave components are involved in the nonlinear phase coupling and the bicoherence values increase progressively. For $x = 9.6 \text{ m}$, just a little over the toe of the slope, the bicoherence magnitude among primary components increases slightly, i.e. $b^2(f_p, f_p) = 0.24$ and $b^2(3f_p, f_p) = 0.15$, which is consistent with the small energy increase in the high frequency region (Abroug et al., 2020).

As the wave train propagates in the shallower region ($9.5 \text{ m} < x < x_b \in [11.09; 11.82]$), the degree of phase coupling is seen to increase rapidly (Fig. 6a). The degree of phase coupling within the peak frequency increases considerably at shallower regions compared to deeper regions. Wave energy transfers increase in high frequency region and as a result, the spectrum broadens. In the vicinity of the breaking location ($x_b \in [11.09; 11.82]$), the non-linear coupling spreads over most of the wave components. The increase of the second and third harmonic is clearly noticeable in Fig. 6b. The values of bicoherence for approximately all frequency pairs are greater than 0.13, indicating that the non-linear coupling reaches its maximum level, which means that almost all of the higher harmonics waves are involved in the propagation process.





Figure 5 : The wavelet-based bicoherence spatial evolution on the flat bottom for a Pierson-Moskowitz wave train (Test 3). (a): $x = 4 \text{ m}$; (b): $x = 5 \text{ m}$; (c): $x = 8 \text{ m}$; (d): $x = 9.6 \text{ m}$.

Downstream the breaking location ($x > x_b \in [11.09; 11.82]$), the degree of phase coupling between frequency components decreases drastically and the bicoherence becomes less structured (Fig. 6d). This result is consistent with the decreasing trend of energy in higher frequency components downstream the breaking location (Tian et al., 2011; Abroug et al., 2020).



235 **Figure 6 : The wavelet-based bicoherence spatial evolution on the sloping bottom for a Pierson-Moskowitz wave train (Test 3). (a): $x = 11$ m; (b): $x = 12$ m; (c): $x = 13$ m; (d): $x = 13.8$ m.**

In fig. 7 and 8, a JONSWAP ($\gamma = 3.3$) wave train (Test 5; $x_b \in [10.5; 11.61]$) is chosen to illustrate the spatial evolution of the wavelet-based bicoherence of a narrower wave train propagating over the flat and the sloping bottom. Wave-wave interactions evolve qualitatively in the same way compared to the case of Pierson-Moskowitz. Fig. 7a ($x = 4$ m) shows that the two dominant phase coupling peaks appear at the bifrequencies (f_p, f_p) and $(0.5f_p, 0 - 0.5f_p)$, which illustrates that the quadratic non-linear interactions only occur between the peak and low-frequency modes. Note that no other peak was found to be significant. As the wave train propagates over the shallower region ($x > 9.5$ m), new phase couplings appear at the bifrequencies $(2f_p, f_p)$, $(3f_p, f_p)$ and $(2f_p, 3f_p)$ (Fig. 8). This finding illustrates that quadratic non-linear interactions between the peak frequency, the first harmonic, the second harmonic and third harmonics result from the gradual broadening of the spectrum. It is an accordance with previous studies demonstrating that energy is mainly transferred to high frequencies during the shoaling process (Tian et al., 2011; Liang et al., 2017; Abroug et al., 2020). For this wave train (Test 5), the wavelet-based bicoherence reaches its maximum shortly after the breaking ($x_b \in [10.5; 11.61]$) at $x = 12$ m. For example, $b^2(f_p, f_p) = 0.7$, $b^2(2f_p, f_p) = 0.53$ and $b^2(2f_p, 2f_p) = 0.16$. Triad interactions lead to skewed wave profiles and can characterize the near-breaking conditions (Fig. 3 for $x > 10.6$ m).

250




Figure 7: The wavelet-based bicoherence spatial evolution on the flat bottom for JONSWAP ($\gamma = 3.3$) wave train (Test 5). (a): $x = 4$ m; (b): $x = 6$ m; (c): $x = 8$ m; (d): $x = 9.6$.

Beyond the breaking location ($x > x_b \in [10.5; 11.61]$), the bicoherence decreases sharply and becomes less structured. For example $b^2(f_p, f_p) = 0.52$, $b^2(2f_p, f_p) = 0.31$ and $b^2(2f_p, 2f_p) = 0.004$ at $x = 13.6$ m, i.e. $h = 0.13$ m. This pattern is qualitatively similar to that obtained in the case of a Pierson-Moskowitz wave train. This indicates that the increasing trend of the phase coupling is one of the more important reasons for the wave train breaking in shallow water.

Figures 9 and 10 depict the wavelet-based bicoherence spectra for the case of a JONSWAP ($\gamma = 7$) wave train (Test 7; $x_b \in [12.07; 12.69]$) at eight locations along the wave flume. No bispectral peak appears at $b^2(2f_p, f_p)$ and this may be not surprising as no clear third harmonic $3f_p$ is present in the frequency spectrum (Fig. 4). Furthermore, wavelet-based bicoherence diagrams show that the phase coupling reaches its maximum level at frequencies slightly higher than the exact harmonics ($2f_p, 3f_p \dots$). This result is consistent with the results of Ma et al., 2010 who explained this process by the slight upshift of peak values in spectrum at higher harmonics, which is readily seen in Fig. 4. The fact that clear 1st, 2nd and 3rd harmonics are not present, is possibly due to other mechanisms such as quadruplet interactions ($f_1 + f_2 = f_3 + f_4$, Elgar et al., 1995) which have a shape stabilizing impact on the spectrum and are confined to free waves. This result is consistent with the peak frequency downshift demonstrated experimentally in Stansberg, 1994 and Abroug et al., 2020, where it was interpreted as a self-stabilizing feature.





Figure 8: The wavelet-based bicoherence spatial evolution on the sloping bottom for JONSWAP ($\gamma = 3.3$) wave train (Test 5). (a): $x = 10.20$ m; (b): $x = 11.20$ m; (c): $x = 12.40$ m; (d): $x = 13.60$ m.

Figure 11 summarizes the variability in the location and intensity of the wavelet-based bicoherence between the bifrequency pairs (f_p, f_p) , $(2f_p, f_p)$, $(3f_p, f_p)$, $(4f_p, f_p)$, $(2f_p, 2f_p)$ for several tests. The two vertical solid lines and the dotted line respectively indicate the breaking region and the toe of the slope. This figure indicates that the steepness has a strong influence on the non-linear phase coupling between harmonics in intermediate water depth ($h_0 = 0.3$ m). Non-linear wave-wave interactions and their increasing trend is more important for wave trains having strong non-linearities. Beyond the wave breaking ($x > x_b$), the decreasing trend of the phase coupling between harmonics is also more significant in the case of strong steepness S_0 . This result is in accordance with the dissipation related to breaking, which is particularly noticeable when the wave steepness is high (Abroug et al., 2020).

An important similarity between different spectra, is that important wave-wave interactions are mostly limited to the first harmonics of primary waves (f_p, f_p) , $(2f_p, f_p)$. This finding is consistent with the energetic behavior of wave trains downstream the wave breaking (Abroug et al., 2020). Moreover, in the case of small and moderate wave steepness (Test 2; $x_b = 12.9$ and Test 6; $x_b = 13.5$), the phase coupling varies slightly downstream the wave breaking compared to that found prior the breaking, suggesting that a small energy transfer happens downstream the breaking location.



290 **Figure 9: The wavelet-based bicoherence spatial evolution on the flat bottom for JONSWAP ($\gamma = 7$) wave train (Test 7). (a): $x = 4$ m; (b): $x = 6.4$ m; (c): $x = 8$ m; (d): $x = 9.6$ m.**

It can be concluded that bound or non-resonant interactions play an important role in the evolution and breaking of wave trains in shallow water depth. Although the bound waves are not supposed to contribute to the energy redistribution, our experimental observations raise the question of the impact of bound interactions on dissipation and energy transfers among different
 295 frequency components.

5 Conclusions and perspectives

An experimental approach is proposed for determining the non-linear wave-wave interactions, which accompany the propagation of large amplitude wave trains, that might cause damage to coastal zones, marine structures and navigation vessels. We investigate seven focused wave trains derived from JONSWAP ($\gamma = 3.3$ or 7) and Pierson-Moskowitz spectra propagating
 300 from intermediate water depth to the inner surf zone. The results presented in this study extend the parameter range of observations of triad interactions. **The experimental conditions were selected based on two parameters: the wave steepness and the spectrum type. Focusing waves were generated in a physical wave flume by modifying the wave spectrum and steepness.** The present data were collected in intermediate water with a $k_p h_0$ varying between 0.92 and 0.79. A typical wave train consists

of a large number of waves interacting with one another. Wavelet-based bicoherence is used to investigate the phase coupling between frequency components of short time series. Some consequences of non-linear transfer are briefly discussed; in particular the role played by non-linear interactions in shaping the high frequency part of the spectrum, the relative contribution of each harmonic and the downshifting of the peak spectrum demonstrated in previous studies. **Note that our experimental study is different from previous experiments (Dong et al., 2008; Ma et al., 2010) regarding the slope geometry and most importantly, the use of three different spectral types.**

Along the flat bottom ($4 \text{ m} < x < 9.5 \text{ m}$), one might assume that the influence of triad interactions is very weak for the three considered spectra. The bispectral analysis of the data shows that as the waves propagate along the flat bottom, the magnitude of the bicoherence increases slightly (between 0% and 20% of its initial value). Moreover, this is foreseeable because the spectrum and the wave train shape do not substantially change along the flat bottom and a small amount of energy is transferred from the peak region to high frequency components. **The spectra remain approximately unimodal and do not present a clear second or third harmonic and non-linear interactions concern only the peak and low frequency regions. On the flat bottom, the wave train is still dispersive and consequently, bound interactions are responsible for the slight growth of $b^2(f_p, f_p)$ and $b^2(0.5f_p, 0.5f_p)$.**





Figure 10: The wavelet-based bicoherence spatial evolution on the sloping bottom for JONSWAP ($\gamma = 7$) wave train (Test 7). (a): $x = 10.2 \text{ m}$; (b): $x = 11 \text{ m}$; (c): $x = 12.4 \text{ m}$; (d): $x = 13.8 \text{ m}$.



325 **Figure 11: Spatial variation of the wavelet-based bicoherence among harmonics.**

When the wave train reaches the slope ($9.5 \text{ m} < x < x_b$), wave-wave interactions among high order harmonics increase rapidly and reach the maximum degree in the breaking/focus location. **In line with previous studies (Elsayed, 2006; Dong et al., 2008; Ma et al., 2010), strong nonlinear interactions were predominantly observed in the shallower region.** The analysis showed a gradual broadening of the bicoherence spectrum, which is in accordance with previous studies who demonstrated that the

330 energy is transferred mainly to high frequencies regions (Tian et al. 2011; Abroug et al., 2020). This is partly due to significant

spectral transformations which are more important during the shoaling process. Particularly, this analysis showed a considerable contribution of 2nd and 3rd harmonics for unidirectional steep wave trains and the spectral components at the second harmonic $2f_p$ have increased substantially (6 times its initial value). The bispectral analysis results show that the wave non-linearity S_0 plays an important role in the increasing trend of phase coupling, which is more important for wave trains having strong non-linearities. This last finding agrees well with the conclusions made by Ma et al. (2010).

An innovative aspect of this paper is presenting wavelet-based bispectral analysis for highly non-linear intermediate water waves with different spectral types. If we compare the three spectra, we can see that all nonlinear interactions on the flat bottom ($x < 9.5$ m) are weak ($b^2 < 0.15$) in the case of wide spectrum wave trains (Tests 2 and 3 Fig. 11). However, in the case of narrower spectra, more frequencies (e.g. f_p , $2f_p$ and $3f_p$) are implicated in the focusing process (Tests 4, 5, 6 and 7 Fig. 11) and the corresponding phase coupling is higher ($b^2 > 0.2$). This finding is in agreement with the stable behavior of wide spectrum wave trains, which was demonstrated experimentally in Abroug et al. (2019) and Stansberg (1994). In intermediate water depth ($0.79 < k_p h < 0.92$), wide spectrum harmonics (f_p , $2f_p$, $3f_p$...) are less implicated in the focusing process compared to narrow spectrum harmonics. In shallow water regions ($9.5 \text{ m} < x < x_b$) and after breaking ($x_b < x$), the spatial evolution of the phase coupling is qualitatively similar for the three spectra.

The results obtained in this study show important features in wave-wave interactions during the propagation of focused waves. This study strengthens the usefulness of wavelet-based analysis in detecting features that are hidden in a Fourier-based analysis, and in explaining a number of phenomena, such as the process leading to wave breaking and the energy transfer between wave components. Nevertheless, in order to confirm the use of wavelet-based bicoherence for more realistic 3D studies with structures, efforts should be made to expand this study for example by investigating greater water depths, higher steepness and wider spectra. Furthermore, the observed evolution of bicoherence for focused waves should be compared to that of waves with similar steepness and bandwidth but with initial random distribution of phase. In other words, efforts should be made to identify and quantify the phase coupling differences between focusing wave trains and non-focusing waves. Information concerning the phase coherence can be obtained by calculating the biphasic parameter (β (a1, a2), Ma et al., 2010). It will be interesting to quantitatively measure the deviation of biphasic values between primary waves/higher harmonics and to analyse their spatial evolution through different spectra to distinguish differences. Finally, a detailed study of how bound energy at harmonics would be influenced by quadruplet interactions should be performed.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

360 This work has been carried out within the framework of the DYNAT (Dynamique du littoral et risques NATurels) project and has received funding from the Normandy region. We thank L. Perez and D. Mouazé for providing technical supports during the experiments. The authors wish also to express their gratitude to S. Baatout for her thorough re-reading of this article.

References

- Abroug, I., Abcha, N., Dutykh, D., Jarno, A and Marin, F.: Experimental and numerical study of the propagation of focused
365 wave groups in the nearshore zone, *Phys. Lett. A*, 384, 6, 126144, doi:10.1016/j.physleta.2019.126144, 2020.
- Abroug, I., Abcha, N., Jarno, A and Marin, F.: Physical modelling of extreme waves: Gaussian wave groups and solitary waves in the nearshore zone, *Advances and Applications in Fluid Mechanics*, 23, 2, 141–159, doi:10.17654/FM023020141, 2019
- Bai, Y., Xia, X., Li, X., Wang, Y., Yang, Y., Liu, Y., Liang, Z and He, J.: Spinal cord stimulation modulates frontal delta and gamma in patients of minimally consciousness state, *Neuroscience*, 346, 247–254, doi: 10.1016/j.neuroscience.2017.01.036,
370 2017
- Becq-Girard, F., Forget, Ph and Benoit, M.: Nonlinear propagation of unidirectional wave fields over varying topography, *Coast. Eng.*, 38, 2, 91–113, doi:10.1016/S0378-3839(99)00043-5, 1999.
- Didenkulova, I and Anderson, C.: Freak waves of different types in the coastal zone of the Baltic Sea, *Nat. Hazard. Earth. Syst. Sci.*, 10, 2021–2029, doi:10.5194/nhess-10-2021-2010, 2010.
- 375 Dong, G., Yuxiang, Ma., Perlin, M., Xiaozhou, M., Bo, Y and Jianwu, X.: Experimental study of wave-wave nonlinear interactions using the wavelet-based bicoherence, *Coast. Eng.*, 55, 9, 741–752, doi:10.1016/j.coastaleng.2008.02.015, 2008.
- Dysthe, K., Krogstad, H. E and Müller, P.: Oceanic rogue waves, *Annu. Rev. Fluid Mech.*, 40, 287–310, doi: 10.1146/annurev.fluid.40.111406.102203, 2008.
- Eldeberky, Y and Madsen, P. A.: Deterministic and stochastic evolution equations for fully dispersive and weakly nonlinear
380 waves, *Coast. Eng.*, 38, 1, 1–24, doi:10.1016/S0378-3839(99)00021-6, 1999.
- Eldeberky, Y.: Nonlinear transformation of wave spectra in the nearshore zone, PhD Thesis, published as Communications on Hydraulic and Geotechnical Engineering, Delft University of Technology, Faculty of Civil Engineering, Report No. 96-4, 203, 1996.
- Elgar, S., Herbers, T. H. C., Chandran, V and Guza. R. T.: Higher-order spectral analysis of nonlinear ocean surface gravity
385 wave, *J. Geophys. Res.*, 100, 4983–4997, doi:10.1029/94JC02900, 1995.
- Elsayed, M. A. K.: A novel technique in analyzing non-linear wave-wave interaction, *Ocean. Eng.*, 33, 2, 168–180, doi:10.1016/j.oceaneng.2005.04.010, 2006.
- Fedele, F., Herterich, J., Tayfun, A and Dias, F.: Large nearshore storm waves off the Irish coast, *Scientific Reports* 9, 1, 15406, doi:10.1038/s41598-019-51706-8, 2019.

- 390 Grinsted, A., Moore, J. C and Jevrejeva, S.: Application of the cross wavelet transform and wavelet coherence to geophysical time series, *Nonlinear. Proc. Geoph.*, 11, 561–566, doi:10.5194/npg-11-561-2004, 2004.
- Huseni, GH and Balaji, R.: Wavelet transform based higher order statistical analysis of wind and wave time histories, *Journal of the institution of engineers (India)*, Ser. C , 98, 635–640, doi:10.1007/s40032-016-0287-0, 2016.
- Janssen, P. A. E. M and Onorato, M.: The intermediate water depth limit of the Zakharov equation and consequences for wave prediction, *J. Phys. Oceanogr.*, 37, 2389–2400, doi:10.1175/JPO3128.1, 2007.
- 395 Kharif, C and Pelinovsky, E.: Physical mechanisms of the rogue wave phenomenon, *Eur. J. Mech. B-Fluid*, 22, 603–635, doi:10.1016/j.euromechflu.2003.09.002, 2003.
- Kharif, C., Pelinovsky, E and Slunyaev, A.: *Rogue waves in the ocean*, Springer Verlag, berlin Helderberg, 2009.
- Larsen, Y., Hanssen, A and Pecseli, H. L.: Analysis of non-stationary mode coupling by means of wavelet-bicoherence, *Int. Conf. Acoust. Spee.*, 6, IEEE, New York, 3581–3584, doi: 10.1109/ICASSP.2001.940616, 2001.
- 400 Li, Y., Wang, X and Lin, J.: Fault diagnosis of rolling element bearing using nonlinear wavelet bicoherence features, *IEEE Conference on prognostics and health management (PHM)*, Cheney, WA, 22–25 June, 1–6, doi: 10.1109/ICPHM.2014.7036369, 2014.
- Liang, S., Zhang, Y., Sun, Z and Chang, Y.: Laboratory study on the evolution of waves parameters due to wave breaking in deep water, *Wave Motion*, 68, 31–42, doi:10.1016/j.wavemoti.2016.08.010, 2017.
- 405 Ma, Y., Dong, G., Liu, S., Zang, J., Li, J and Sun, Y.: Laboratory study of unidirectional focusing waves in intermediate depth water, *J. Eng. Mech.*, 136, 1, doi:10.1061/(ASCE)EM.1943-7889.0000076, 2010.
- Merkoune, D., Touboul, J., Abcha, N., Mouazé, D., Ezersky, A.: Focusing wave group on a current of finite depth, *Nat. Hazard. Earth. Syst. Sci.*, 13, 2941–2949, doi:10.5194/nhess-13-2941-2013, 2013.
- 410 Milligen, B. P. V., Sanchez, E., Estrada, T., Hidalgo, C., Branas, B., Carrersa, B and Garcia, L.: Wavelet bicoherence: a new turbulence analysis tool, *Phys. Plasma*, 2, 8, 3017–3032, doi:10.1063/1.871199, 1995.
- Onorato, M., Residori, S., Bortolozzo, U., Montina, A and Arecchi, F.: Rogue waves and their generating mechanisms in different physical contexts, *Phys. Rep.*, 528, 48–89, doi:10.1016/j.physrep.2013.03.001, 2013.
- Stansberg, C. T.: Effects from directionality and spectral bandwidth on nonlinear spatial modulations of deep-water surface gravity wave trains, *Coast. Eng. Proceedings of the XXIV international conference*, 23–28, Kobe, Japan, 579–593, doi:10.1061/9780784400890.044, 1994.
- Taylor, P.H., Williams, B.A.: *Wave statistics for intermediate depth water – NewWaves and symmetry*, *J. Offshore. Mech. Arct.*, 126, 54-59, doi:10.1115/1.1641796, 2004.
- Tian, Z., Perlin, M and Choi, W.: Frequency spectra evolution of two-dimensional focusing wave groups in finite water depth water, *J. Fluid. Mech.*, 688, 169–194, doi:10.1017/jfm.2011.371, 2011.
- 420 Torrence, C and Compo, G. P.: A practical guide to wavelet analysis, *Bulletin of American meteorology society*, 79, 1, 61–78, doi:10.1175/1520-0477(1998)079<0061:APGTWA>2.0.CO;2, 1998.

- Tromans, P.S., Anaturk, A.R., Hagemeijer, P.: A new model for the kinematics of large ocean waves – application as a design wave, *Proceedings of the first international offshore and polar engineering Conference*, Int. J. Offshore. Polar, 64-71, 1991.
- 425 Vyzikas, T., Stagonas, D., Buldakov, E and Greaves, D.: The evolution of free and bound waves during dispersive focusing in a numerical and physical flume, *Coast. Eng.*, 132, 95–109, doi:10.1016/j.coastaleng.2017.11.003, 2018.
- Whittaker, C.N., Raby, A.C., Fitzgerald, C.J., Taylor, P. H.: The average shape of large waves in the coastal zone, *Coast. Eng.*, 114, 253-264, 2016.
- Xu, Guochun., Hao, H., Ma, Q and Gui, Q.: An experimental study of focusing wave generation with improved wave amplitude spectra, *Water*, 11, 12, 2521, doi:10.3390/w11122521, 2019.
- 430 Young, I. R and Eldeberky, Y.: Observations of triad coupling of finite depth wind waves, *Coast. Eng.*, 33, 137–154, doi:10.1016/S0378-3839(98)00006-4, 1998.
- Zhang, J., Benoit, M., Kimmoun, O., Chabchoub, A and Hsu, H. C.: Statistics of extreme waves in coastal waters: large-scale experiments and advanced numerical simulations, *Fluids*, 4, 2, 99, doi:10.3390/fluids4020099, 2019.
- 435