

Interactive comment on “Landslide susceptibility assessment based on different machine-learning methods in Zhaoping County of eastern Guangxi” by Chunfang Kong et al.

Chunfang Kong et al.

xukai@cug.edu.cn

Received and published: 17 October 2020

[Printer-friendly version](#)

[Discussion paper](#)



Response to RC1 from Anonymous Referee #1

Interactive
comment

October 17, 2020

Dear Reviewer #1:

Thank you for your comments concerning our manuscript ID nhess-2020-251 (Landslide susceptibility assessment based on different machine-learning methods in Zhaoping County of eastern Guangxi). Those comments are all valuable and very helpful for revising and improving our paper, as well as of important guiding significance to our researches. We have studied comments carefully and have made correction which we hope meet the suggestions. Revised portion are marked in highlight in the paper. The main corrections in the paper and the responds to the reviewer's comments are as flowing.

1. Lines 26-28: Is there any literature research to support such a claim that the variation of robustness and performance of these two models can be neglected among applications in different regions?

Zhou et al. (2018) used the ML methods in the landslide susceptibility analysis of Longju in the Three Gorges Reservoir area. Its result showed that the SVM model has a better performance and a strong robustness. Hence, the SVM model

[Printer-friendly version](#)

[Discussion paper](#)



can be recommended before reaching a consensus on the model of landslide susceptibility assessment.

Deng et al. (2018) analyzed the long-term observation test of temperature and gases in the gob of 40106 fully mechanized top-coal caving face at Dafosi coal mine using random forests. At the same time, the particle swarm optimization (PSO) algorithm was employed to optimize the hyper-parameters of RF and SVM for establishing the PSO-RF and PSO-SVM prediction models with optimized parameters. The results indicated that PSO-RF and PSO-SVM models had strong generalization and robustness, and the PSO-RF model could be further applied to other energy and fuel fields.

The relevant literature is as follows:

Zhou, C., Yin, K., Cao, Y., Ahmed, B., Li, Y., Catani, F., and Pourghasemi, H.R.; Landslide susceptibility modeling applying machine learning methods: A case study from Longju in the Three Gorges Reservoir area, China, *Comput. Geosci.*, 2018, 112, 23–37.

Deng, J., Lei, C., Cao, K., Ma, L., Wang, C., and Zhai, X.; Random forest method for predicting coal spontaneous combustion in gob, *J. China Coal Soc.*, 2018, 43(10), 2800–2808.

The above literatures have been labeled in the paper; please see L45, L47-48, L338, and L398.

2. In the introduction, there is in lack of a summary of the popularity of these ML algorithms. Such a summary can help readers understand why the authors chose these ML algorithms in the current study.

In the introduction of this paper, examples are given to illustrate that more and more machine learning (ML) algorithms have been optimized and applied for landslide susceptibility assessment in different regions. These have all been

[Printer-friendly version](#)

[Discussion paper](#)



used to quantitatively predict and assess the susceptibility for landslide in different regions of the world. These studies play an important role in the susceptibility evaluation and prediction of landslide. Please see L40-58. At the same time, many comparative studies on landslide susceptibility assessment using different ML methods have been performed. Please see L59-72. These previous studies have shown that the SVM and RF have been widely proved to be useful methods in the evaluation of landslide susceptibility (Marjanović et al., 2011; Tien Bui et al., 2012; Kavzoglu et al., 2014; Trigila et al., 2015; Pham et al., 2016; Ada and San, 2018). However, few studies have focused on the optimization of SVM and RF models in landslide susceptibility prediction and evaluation and compared the optimized results. Therefore, based on previous works, the objective of the present paper is to: (1) optimize SVM and RF models by using a particle swarm optimization (PSO) algorithm; (2) analyze and evaluate the susceptibility levels of landslide by using the SVM, PSO-SVM, RF, and PSO-RF models for Zhaoping County; and (3) compare the performances of four ML models for landslide susceptibility evaluation by receiver operating characteristic (ROC) curve, statistical analysis, and field-verified methods. Please see L74-84.

3. The objectives of the introduction should be supported by the gap in the literature. The current objectives jump out from nowhere without any rationale or reasoning.

In the introduction of this paper, L40-56 illustrated that more and more machine learning (ML) algorithms have been optimized and applied for landslide susceptibility assessment in different regions. At the same time, L59-72 illustrated that many comparative studies on landslide susceptibility assessment using different ML methods have been performed. These previous studies have shown that the SVM and RF have been widely proved to be useful methods in the evaluation of landslide susceptibility. However, few studies have focused on the optimization of SVM and RF models in landslide susceptibility prediction and evaluation and

[Printer-friendly version](#)

[Discussion paper](#)



compared the optimized results. Therefore, the objective of the present paper is to: (1) determine the landslide susceptibility assessment factors by multi-source data fusion and correlation factor analysis; (2) optimize SVM and RF models by using a particle swarm optimization (PSO) algorithm; (3) analyze and evaluate the susceptibility levels of landslide by using the SVM, PSO-SVM, RF, and PSO-RF models for Zhaoping County; and (4) compare the performances of four ML models for landslide susceptibility evaluation by receiver operating characteristic (ROC) curve, statistical analysis, and field-verified methods. The results provide valuable informational support for the prediction and evaluation of landslide in Zhaoping County, Guangxi. Please see L40-86.

4. **Fig.1: Please highlight the experimental site in mainland China only and ignore the outlying islands to maximize the area of interest.**

Figure 1 show the location of the study area of Zhaoping County in China and Guangxi Province. The island is an inseparable part of China, so it should not be ignored.

5. **Lines 107-108, 117-120, 128-130: Please provide proper references.**

These materials come from the field investigation report of the geological hazard project by Guangxi Geological Survey Bureau (Huang and He, 2018), and references have been added to the corresponding positions in the article. Thank you for your careful reviews. Please see revised L107-109, L118-121, L125-129, and L130-132.

6. **Lines 124-127: Why were they chosen? Were there any prior studies to support such a decision?**

According to the field investigation report of the geological hazard project by Guangxi Geological Survey Bureau and the disaster factors correlation analysis, a total of ten factors of high correlation with landslide disaster occurrence

[Printer-friendly version](#)

[Discussion paper](#)



Interactive
comment

were selected as landslide hazard assessment factors for Zhaoping County. Related reference materials have been added in the paper; please see the revised L125-129.

7. Line 140: Please define “heavy rain”.

Heavy rain generally refers to rainfall with a daily rainfall of 25-49.9 mm (24 hours) or a rainfall of 8.1-16.0 mm per hour.

8. Line 165: Is the LULC map created based on Landsat imagery? Please provide detailed calibration and validation results so the LULC data can be used. LULC determination is not a straightforward process and can be complicated.

The LULC map in the paper comes from the manual visual interpretation results of Landsat 8 OLI image (2017/12/24, 124/043). The interpretation results are shown in Figure 2(h).

The process of interpretation is as follows: (1) Radiometric calibration and atmospheric correction for the Landsat 8 OLI image; (2) Based on the ground control point, geometric correction for the Landsat 8 OLI image, and the correction accuracy is less than 1 pixel; (3) Combined with topographic map and China's Land Use/Cover Dataset (CLUD) of 2015 by Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, the LULC of Zhaoping County was divided into five categories by manual visual interpretation: 1-cultivated land; 2-woodland; 3-grassland; 4-river and lake; 5-construction land; (4) By randomly selecting 10% samples to verify the accuracy of field investigation, it was shown that the overall classification accuracy is more than 95.53%, which meets the accuracy requirements of this paper.

The LULC map is only a representative factor reflecting the impact of human activities on the environment in Zhaoping County, and due to the limited space

[Printer-friendly version](#)

[Discussion paper](#)



of the paper, detailed calibration and verification results are not provided in the paper.

9. **Fig. 2: For each subplot, please provide numerical ranges for each “grade” based on Table 1.**

The level of each subplot in Figure 2 is one-to-one corresponding to the specific numerical range in Table 1.

For example, as shown in Figure 2(a) of the slope, 1 represents 0-7°, 2 represents 7-13°, 3 represents 13-25°, 4 represents 19-25°, 5 represents 25-34°, 6 represents 34-50°, 7 represents 50-70°, and 8 represents 70-76°.

Figure 2(b) of the aspect, 1 represents 337.5-22.5°, 2 represents 22.5-67.5°, 3 represents 67.5-112.5°, 4 represents 112.5-157.5°, 5 represents 157.5-202.5°, 6 represents 202.5-247.5°, 7 represents 247.5-292.5°, and 8 represents 292.5-337.5°.

Figure 2(c) of the plan curvature, 1 represents -25-5°, 2 represents -5-2.5°, 3 represents -2.5-1°, 4 represents -1-0°, 5 represents 0-1°, 6 represents 1-2.5°, 7 represents 2.5-5°, and 8 represents 5-28.9°.

Figure 2(d) of the annual rainfall, 1 represents 0-1980 mm, 2 represents 1980-2100 mm, 3 represents 2100-2220 mm, 4 represents 2220-2340 mm, 5 represents 2340-2460 mm, 6 represents 2460-2580 mm, 7 represents 2580-2700 mm, and 8 represents 2700-2820 mm.

Figure 2(e) of the NDVI, 1 represents 0-0.01, 2 represents 0.01-0.09, 3 represents 0.09-0.17, 4 represents 0.17-0.25, 5 represents 0.25-0.33, 6 represents 0.33-0.4, 7 represents 0.4-0.5, and 8 represents 0.5-0.71.

Figure 2(f) of the stratum lithology, 0 represents river, 1 represents Quaternary, 2 represents carbonate rock, 5 represents clastic intercalated with siliceous rocks, 6 represents clastic rock, 7 represents sandstone and shale, and 8 represents granite or basal rocks.

Interactive comment

[Printer-friendly version](#)

[Discussion paper](#)



Figure 2(g) of the tectonic complexity, 1 represents 0-1.4, 2 represents 1.4-2.7, 3 represents 2.7-3.8, 4 represents 3.8-4.9, 5 represents 4.9-6, 6 represents 6-7.3, 7 represents 7.3-8.9, and 8 represents 8.9-9.4.

Figure 2(h) of the LULC, 1 represents cultivated land, 2 represents woodland, 3 represents grassland, 4 represents river and lake, and 5 represents construction land.

Figure 2(i) of the residential density, 1 represents 0-1.2, 2 represents 1.2-2.7, 3 represents 2.7-4.5, 4 represents 4.5-6.9, 5 represents 6.9-10.1, 6 represents 10.1-14.2, 7 represents 14.2-19.7, and 8 represents 19.7-25.

Figure 2(j) of the road network density, 1 represents 0-3.2, 2 represents 3.2-4.7, 3 represents 4.7-6.1, 4 represents 6.1-7.8, 5 represents 7.8-9.7, 6 represents 9.7-11.7, 7 represents 11.7-13.9, and 8 represents 13.9-14.

10. **Fig. 3: Please provide more details for each step in the text**

Figure 3 is a Flowchart of landslide susceptibility evaluation based on ML. For details, please refer to the paper of 3.1-3.4.

11. **Tables 2-4: Please merge information of these steps into the text. Tables are used to display arrayed data.**

Tables 2-4 are the specific steps of the four ML algorithms, so, it is appropriate to put it in a table rather than merge it into the text.

12. **Fig. 4: Maybe I missed this – Is the definition of the levels hidden somewhere in the text?**

Figure 4 is the evaluation results of landslide susceptibility for four ML models in Zhaoping County, and 1 represents extremely low susceptibility, 2 represents low susceptibility, 3 represents middle susceptibility, 4 represents high susceptibility, 5 represents extremely high susceptibility.

[Printer-friendly version](#)

[Discussion paper](#)



13. **Please define robustness in this case – my definition of robustness is that the algorithm consistently delivers good results at all kinds of environments. I don't see how your analyses reflect such quality.**

Robustness in this paper refers to the stable performance of the PSO-RF and PSO-SVM models established in this paper, that is, by inputting the attribute values of each evaluation factor, the results of the susceptibility level of landslide disasters in the study area are obtained, and the results are in good agreement with the results of field investigation.

The reason why the PSO-RF and PSO-SVM models have strong robustness is that we applied the models in this paper to evaluate the landslide susceptibility in 23 other regions of Guangxi and the evaluation results agree with the results of field investigation.

14. **I don't understand the message delivered by Table 5. It looks that the accuracy is not good because the landslide points that fell into high susceptibility areas are rare. Please highlight the message delivered by this table.**

Table 5 indicates the proportions of hazards points falling into different susceptibility levels.

The first line indicates that the PSO-RF model simulates the probability of the landslide point falls into the extremely high susceptibility level is the highest, which is 0.2306%, followed by the PSO-SVM model, the third is the RF model, and the last one is the SVM model.

The second line indicates that the PSO-RF model simulates the probability of the landslide point falls into the high susceptibility level is also the highest, which is 0.0845%, followed by the PSO-SVM model, the third is the RF model, and the last one is the SVM model.

The third line indicates that the PSO-RF model simulates the probability of the landslide point falls into the middle susceptibility level is the lowest, which is



0.0117%, followed by the RF model, the third is the PSO-SVM RF model, and the last one is the SVM model.

The fourth line indicates that the PSO-RF model simulates the probability of the landslide point falls into the low susceptibility level is also the lowest, which is 0.0041%, followed by the PSO-SVM model, the third is the RF model, and the last one is the SVM model.

The fifth line indicates that the PSO-RF model simulates the probability of the landslide point falls into the extremely low susceptibility level is also the lowest, which is 0.0005%, followed by the PSO-SVM model, the third is the RF model, and the last one is the SVM model.

The above analysis shows that the proportion of landslide disaster points simulated by the PSO-RF model falling into extremely high and high-prone areas is higher than that of other models. At the same time, the proportion of landslide disaster points simulated by the PSO-RF model falling into low and extremely low-prone areas is lower than that of other models, which from another aspect shows that the PSO-RF model has the highest simulation accuracy and the best performance in comparison to other landslide models.

15. **Overall, there is a serious issue with this manuscript. This manuscript simply applied several known algorithms without interpretations. To make it publishable, interpretation of results is required. Why are certain algorithms performing better? Why are certain factors having a higher influence? What does this information mean to management and disaster prevention? These are simply some quick examples on top of my head.**

It has been discussed in the introduction that this paper is based on previous works, and many previous studies have proved that SVM and RF models have better performance in landslide susceptibility evaluation and prediction than other models. Based on this, the focus of the present paper is to optimize SVM and

[Printer-friendly version](#)

[Discussion paper](#)



RF models by using a particle swarm optimization (PSO) algorithm; analyze and evaluate the susceptibility levels of landslide by using the SVM, PSO-SVM, RF, and PSO-RF models for Zhaoping County; and compare the performances of four ML models for landslide susceptibility.

Our research denoted that the PSO algorithm has a good effect on SVM and RF models. Meanwhile, our research also demonstrated that PSO-RF model has a better prediction performance than the PSO-SVM model, which is mainly due to the large number of factors selected in this study, the PSO-RF model, a type of ensemble learning, exhibited advantages over a traditional ML method by not only accounting for different types of factors but also evaluating the relative importance of the factors in terms of landslide stability. The relevant discussion has been added to the paper, Please see L341-346.

Our research also denoted that the simulation results of the paper proved that the occurrence of landslide disasters has a strong correlation with the stratum lithology, geological tectonic complexity, precipitation, human engineering activities, and vegetation index. This is mainly due to the occurrence of landslide disasters have its internal and external factors: stratum lithology, geological tectonic complexities are the internal causes of landslide disasters; precipitation, human engineering activities, and vegetation cover are the external causes of landslide disasters. Internal causes play a major role in the occurrence of landslide, while external causes play a role in promoting the occurrence of landslide.

To sum up, the information of the landslide susceptibility levels from the results can provide method support for engineering construction, ecological environment construction, rapid economic development, disaster reduction and disaster prevention in Zhaoping County, Guangxi.

Once again, we are very grateful for your comments, and those comments are all valuable and very helpful for revising and improving our paper, as well as of important



guiding significance to our researches.

NHESSD

Interactive
comment

