

Towards an efficient storm surge and inundation forecasting system over the Bengal delta: Chasing the super-cyclone Amphan

Md Jamal Uddin Khan¹, Fabien Durand^{1,2}, Xavier Bertin³, Laurent Testut^{1,3}, Yann Krien³, A.K.M. Saiful Islam⁴, Marc Pezerat³, and Sazzad Hossain^{5,6}

¹LEGOS UMR5566, CNRS/CNES/IRD/UPS, 31400 Toulouse, France

²Laboratório de Geoquímica, Instituto de Geociências, Universidade de Brasília, Brazil

³UMR 7266 LIENSs, CNRS- La Rochelle University, 17000 La Rochelle, France

⁴IWFM, BUET, Dhaka 1000, Bangladesh

⁵Flood Forecasting and Warning Centre, BWDB, Dhaka, Bangladesh

⁶Department of Geography and Environmental Science, University of Reading, UK

Correspondence: Md Jamal Uddin Khan (jamal.khan@legos.obs-mip.fr)

Abstract. The Bay of Bengal is a well-known breeding ground to some of the deadliest cyclones in history. Despite recent advancements, the complex morphology and hydrodynamics of this large delta and the associated modelling complexity impede accurate storm surge forecasting in this highly vulnerable region. Here we present a proof of concept of a physically consistent and computationally efficient storm surge forecasting system tractable in real-time with limited resources. With a state-of-the-art wave-coupled hydrodynamic numerical modelling system, we forecast the recent super cyclone Amphan in real-time. From the available observations, we assessed the quality of our modelling framework. We affirmed the evidence of the key ingredients needed for an efficient, real-time surge and inundation forecast along this active and complex coastal region. This article shows the proof of the maturity of our framework for operational implementation, which can particularly improve the quality of localized forecast for effective decision-making over the Bengal delta shorelines, as well as over other similar cyclone-prone regions.

1 Introduction

Storm surges and associated coastal floodings are one of the most dangerous natural hazards along the world coastlines. Annually, storm surges killed on average, over 8000 people and affected 1.5 million people worldwide over the past century (Bouwer and Jonkman, 2018). Among the storm surge-prone basins, the Bay of Bengal in the northern Indian Ocean is one of the deadliest. This basin has consistently been home to the world's highest surges, experiencing in each decade an average of five surge events exceeding 5 m (Needham et al., 2015). On May 18th of 2020, a super-cyclone named Amphan was identified as the most powerful ever recorded in the Bay of Bengal, with highest sustained 3-min wind speed more than 240 km per hour (130 knots) and highest 1-min wind gusts as fast as 260 km/h (<https://www.metoc.navy.mil/jtwc/jtwc.html>). Two days later, on May 20th, Amphan made landfall in West Bengal (India), with a sustained wind speed of 112 km per hour and gusts of 190 km per hour, causing massive damage in India and Bangladesh and claiming at least 116 lives (IFRC, 2020a, b).

The cyclone activity in the Bay of Bengal is very different from the other cyclone-prone oceanic basins (Needham et al., 2015). The basin experiences a distinct bi-modal distribution of cyclonic activity - one with a peak in the pre-monsoon (April-May) and another during post-monsoon (October-November) (Bhardwaj and Singh, 2019). The Bay of Bengal is a small semi-enclosed basin (Figure 1), which accounts for only 6% of cyclones worldwide. However, more than 70% of global casualties from the cyclones and associated storm surges over the last century occurred there (Ali, 1999). The number of fatalities concentrates in the Bengal delta across Bangladesh and India (Ali, 1999; Murty et al., 1986) where more than 150 million people live below 5 m above mean sea level (MSL) (Alam and Dominey-Howes, 2014). Seo and Bakkensen (2017) noted a statistically significant correlation between storm surge height and the fatality in this region. Two of the notable cyclones that struck this delta include the 1970 cyclone Bhola and 1991 cyclone Gorky which killed about 300,000 (Frank and Husain, 1971) and 150,000 (Khalil, 1993) people respectively. In recent decades, the death toll has reduced by orders of magnitude. For example, cyclone Sidr, a category five equivalent cyclone on the Saffir-Simpson scale that made landfall in 2007, claimed 3406 lives (Paul, 2009). At the same time, the cost of material damages has significantly increased (Alam and Dominey-Howes, 2014; Emanuel, 2005; Schmidt et al., 2009). During recent cyclones, government and voluntary agencies of Bangladesh and India took coordinated effort to evacuate millions of people to safety at cyclone shelters before the cyclone landfall (Paul and Dutt, 2010). This kind of informed coordination benefited from improved communication, increased shelter infrastructure, and, most importantly, from the improvement of the numerical prediction of storm track and intensity.

Over the last decades, global weather and forecasting systems have advanced significantly. Several global models now run multiple times a day, starting from multiple initial conditions, with horizontal resolution in the order of tens of km, providing forecast with a range of hours to a week. These forecasts have proven useful during weather extremes like tropical storms (Magnusson et al., 2019). Operational hurricane forecasting systems like Hurricane Weather Research and Forecasting (HWRF) have emerged and reached a level of maturity to provide reliable cyclone forecasts several days in advance throughout the tropics (Tallapragada et al., 2014).

Storm surge forecasting has been developed and implemented around the world along with the advancement of the weather and storm forecasting (Bernier and Thompson, 2015; Daniel et al., 2009; Flowerdew et al., 2012; Lane et al., 2009; Verlaan et al., 2005). Nowadays, operational surge forecasting systems typically run on high-performance computing systems, either on a scheduled basis or triggered on-demand during an event (Loftis et al., 2019; Oliveira et al., 2020; Khalid and Ferreira, 2020). In the Bay of Bengal region, the Indian Institute of Technology-Delhi (IIT-D) storm surge model has been running operationally with a horizontal resolution of 3.7 km for one decade (Dube et al., 2009). Under the Tropical Cyclone Program (TCP) of the World Meteorological Organization (WMO), IIT-D model is also implemented to run in Bangladesh Meteorological Department (BMD). Beside the IIT-D model, BMD also experimentally uses MRI storm surge model from Japan Meteorological Agency (JMA) (<http://bmd.gov.bd/p/Storm-Surge>). Storm surge forecasts have shown their potential to better target the evacuation decision, to optimize early-engineering preparations, and to improve the efficiency of the allocation of the resources (Glahn et al., 2009; Lazo and Waldman, 2011; Munroe et al., 2018). Availability of a spatially-distributed forecast of storm surge flooding can further increase the fluidity of communication toward the public (Lazo et al., 2015). Keeping in mind the cyclonic surge hazards over the densely populated Bengal delta, having a reliable real-time operational forecast system in




Figure 1. The path of the super cyclone Amphan of May 2020, overlaid on the population density. The footprint of 18 m/s (Tropical storm), 33 m/s (Cat. 1), 43 m/s (Cat. 2), and 50 m/s (Cat. 3) wind speed area is shown with the red colour bar (Classified as in Saffir–Simpson scale). Inset shows the location of the study area.

the region would be extremely valuable and would address a societal demand (Ahsan et al., 2020). The major challenges in operating and maintaining such systems are manifold for the Bengal delta, including lack of expertise, limitations of funding resources to operate and maintain necessary infrastructure and dataset, as well as availability of reliable modelling systems in operational mode (Roy et al., 2015).

60 Storm surge modelling is, however, computationally expensive and proved to be challenging in real-time forecasting mode (Glahn et al., 2009; Murty et al., 2017). Several practical solutions exist to curb the real-time constraints of cyclone forecast, such as soft real-time computation based on looking up an extensive database of pre-computed cyclones (Condon et al., 2013; Yang et al., 2020), coarse-resolution modelling (Suh et al., 2015), and modelling without waves coupling (Murty et al., 2017). Over the past decades, unstructured-grid modelling systems are getting more and more popular due to their efficiency in
 65 resolving the topographic features and their reduced computational cost compared to structured-grid equivalents (Ji et al., 2009; Lane et al., 2009; Melton et al., 2009; Fortunato et al., 2017; Khalid and Ferreira, 2020).

The published history of storm surge modelling in the northern Bay of Bengal goes hand-to-hand with the land-falling of very severe cyclones (Das, 1972; Flather, 1994; Dube et al., 2004; Murty et al., 2014; Krien et al., 2017). From the modelling of the historical storm events, previous studies have identified several ingredients as essential for accurate modelling and forecasting
 70 of storm surge induced water level and associated flooding over the Bengal delta. The most important of these ingredients is

an accurate bathymetry and topography (Krien et al., 2016, 2017; Murty et al., 1986). Second, it is required to have a large-scale modelling domain comprising the whole Ganges-Brahmaputra-Meghna (GBM) estuarine network (Johns and Ali, 1980; Oliveira et al., 2020) at a high-enough model resolution (Krien et al., 2017; Kuroda et al., 2010). The modelling framework is required to simulate tide and surge together to account for the tide-surge interactions (As-Salek and Yasuda, 2001; Johns and Ali, 1980; Kuroda et al., 2010; Murty et al., 1986). Furthermore, an online coupling of the hydrodynamics and the short waves is also necessary to account for the wave setup (Deb and Ferreira, 2016; Krien et al., 2017). The capability of the model to simulate the wetting and drying is also necessary to model the inundation (Madsen and Jakobsen, 2004).

Fully coupled tide-surge-wave models have been recommended for operational forecasting in the Bay of Bengal for years (Johns and Ali, 1980; Murty et al., 1986; As-Salek and Yasuda, 2001; Deb and Ferreira, 2016; Krien et al., 2017). Besides the development of the wave-ocean coupled hydrodynamic modelling tool itself, the main challenge in implementing such a system sums up to the high computational requirements for the modelling systems (Murty et al., 2014, 2016). Due to this constraints, operational systems are commonly run with either only surge, or coupled tide-surge model at kilometric resolution (Dube et al., 2009). In recent years, higher resolution (100 m at the coast) unstructured grid models has been used for operational forecast over part of Indian coastlines in the Bay of Bengal, but without wave-coupling (Murty et al., 2017). These limitations in operational forecasting systems can significantly impede a proper interpretation during an actual cyclonic storm, as discussed in the next section through a set of hindcast sensitivity experiments of cyclone Amphan.

Cyclone Amphan is not only the latest event on record in the Bay of Bengal but also the costliest event that struck this shoreline, with an estimated bill over 14 billion dollars in West Bengal and Bangladesh (DhakaTribune, 2020; IFRC, 2020a). During this cyclone, we have proactively forecast storm surge in real-time with common computational resources using a high-resolution coupled modelling system forced by a combination of freely available atmospheric forecasts. The general goal of this paper is to provide a proof-of-concept of a tractable operational storm surge forecasting system over this key-region of vulnerability, to identify the key ingredients of such a system, and to provide guidance in the near-future initiatives of the operational forecasting community. First, we present the various processes governing the surge dynamics to illustrate the challenges of modelling storm surges in the Bay of Bengal in Section 2. Section 3 documents the atmospheric forecasts that are required to generate a surge forecast. In Section 4, we introduce our numerically-efficient hydrodynamics-waves coupled modelling platform and present its practical real-time computational setup in Section 5. Finally, we present the remaining modelling and forecasting issues in Section 6. Section 7 provides a conclusion to this study.

2 Storm surge and inundation processes in the Bay of Bengal

The Bay of Bengal is a macro-tidal region where peak tidal range reaches as high as 5 m over the north-eastern corner of the Bay (Krien et al., 2016; Tazkia et al., 2017). A large geographical extent of the land-sea hydraulic continuum with an intricate estuarine network complicates the water level dynamics in this part of the coastal ocean. During a cyclone, the dynamics get even more complicated due to wave-induced setup and tide-surge interaction. In Figure 2, we illustrate a simplified view of

water level components – tide, surge, wave setup – and their interactions with the topography in driving inundation during a cyclone.

105 During a cyclone, the atmospheric pressure drop and the wind stress both generate the water level surge. This atmospheric storm surge is non-linearly dependent on the astronomical tide. The dependent nature of both tide and surge implies that the surge induced by a given meteorological condition differs at different stages of the tide, particularly in the nearshore shallow zone. This departure from linearly-added tide and surge component is known as tide-surge interaction. In a macro-tidal region like the Bengal delta tide-surge interaction typically amounts to several tens of centimetres (Antony and Unnikrishnan, 2013; 110 Idier et al., 2019). Due to this interaction, the highest surge (water level – tide) does not coincide with high tide as shown from observations (Antony and Unnikrishnan, 2013) and numerical modelling (Krien et al., 2017; Antony et al., 2020). The strong dependence on tide reinforces the importance of having an accurate tidal model for the region and a reliable network of water level gauge for validation (As-Salek and Yasuda, 2001; Krien et al., 2017; Kumar et al., 2015; Murty et al., 2016).




Figure 2. Conceptual diagram of the involved processes that determine the water level evolution and its interaction with the controls determining the inland flooding.

Wave setup is another component that has a significant impact on the nearshore sea level (Idier et al., 2019; Krien et al., 115 2017; Murty et al., 2014). It manifests as an increase in the sea level occurring in the nearshore zone that accompanies the dissipation of short waves by breaking (Longuet-Higgins and Stewart, 1962). During cyclone Sidr, the modelled wave-setup was around 30 cm (Krien et al., 2017). This estimation would be potentially underestimated by up to 100% due to an early dissipation of wave energy by depth-induced breaking arising from usual parameterizations utilized in spectral wave models (Pezerat et al., 2020, 2021).

120 At the seasonal scale, the mean sea level around the Bengal delta also shows considerable evolution. The amplitude of this variation can go as high as 40 cm due to freshwater influxes during monsoon from the GBM river system and offshore-ocean steric variability (Durand et al., 2019; Tazkia et al., 2017). During the wintertime cyclone-prone season, this steric variability can induce 10-15 cm variation of mean sea level of the bay.

Except for the Sundarban mangrove forest, coastal Bangladesh is mostly embanked through a network of 139 polders (Nowreen et al., 2013). These embanked areas can be flooded in three ways - (a) by overflowing/overtopping, (b) by breaching of the embankments/control structures, and (c) by heavy rainfall inside the embanked area. The height of these embankments plays the most vital role in actuating the inundation from a storm surge (Krien et al., 2017). The performance of the embankments during the cyclone is another crucial factor, depending on the composition (typically, earth vs stones/concrete). During a cyclone, the wave action as well as overtopping/overflowing on the embankments can create breaches at a weak point, thus creating local inland flooding (Islam et al., 2011). Particularly for Bangladesh, the consequence of such storm surge induced flooding can be long-lasting as the topography inside polders is often below the mean sea level due to ground compaction (Auerbach et al., 2015).

3 Atmospheric evolution of cyclone Amphan

On May 13th of 2020, a low-pressure area was persisting over northern Bay of Bengal about 300 km east of Sri Lanka. By the end of 15th, the Joint Typhoon Warning Center (JTWC) upgraded the low-pressure system to a tropical depression. Indian Meteorological Department (IMD) also reported the same development on the next day. The tropical depression continued to move northward and intensified into the cyclonic storm Amphan by May 16th 18:00 UTC. During the following 12 hours, the intensification of the system was limited. However, starting from 12:00 UTC on May 17th, Amphan started to intensify very rapidly. In just 6 hours around 18:00 UTC, the maximum wind speed increased from 140 km/h to 215 km/h, making it an extremely severe cyclonic storm (equivalent to Category 4 on the Saffir-Simpson scale). Over the next twelve hours, Amphan continued to intensify reaching a maximum of 260 km/h wind-speed and 907 millibars central pressure, making it the most intense event on record in the Bay of Bengal. During this time, it appeared to form two distinct eye-walls, which is typical of intense cyclones. Over the next 24 hours, Amphan lost most of its strength due to the eye-wall replacement cycle in the presence of moderate vertical wind shear (30-40 km/h). The system continued to decay due to easterly shear and mid-level dry air. The vertical wind shear remained moderate during this period. The cyclone made landfall between 08:00UTC and 10:00UTC around (88.35°N,21.65°E) (Figure 1), at mid-tide. At landfall, the reported central pressure amounted to 965 millibars with a maximum wind speed of 150-160 km/h. Afterwards, over its journey inland, the system further eroded and disappeared by May 21st. The black lines in Figure 3 illustrates the evolution of Amphan from JTWC best track.

The extended range outlook by IMD published on May 7th predicted cyclogenesis to occur during May 8th to May 14th with low probability RSMC (2020). Global and regional models started to predict a significant storm to happen in the Bay of Bengal as early as May 12th, eight days before the cyclone landfall and four days before the actual formation of the tropical depression. The formation of the low-pressure system triggered the operational HWRF system of NOAA (https://www.emc.ncep.noaa.gov/gc_wmb/vxt/HWRF/) on May 14th. Similarly, the operational system of IMD (<https://nwp.imd.gov.in/hwrf/IMDHWRFv3.6>) was also triggered a day later. For the rest of the paper, we confine our analysis to the forecast disseminated by NOAA-HWRF only, relying on the Automated Tropical Cyclone Forecasting System (ATCF) text output (Miller et al., 1990). Figure 3 illustrates a few of the selected forecast cyclone tracks sequentially issued from NOAA-HWRF.




Figure 3. Temporal evolution of the successive forecasts of Amphan cyclone wind and pressure. (a) Forecast track colour-coded with the date (JTWC best track in black) (b) Wind speed forecast with each epoch, (c) Pressure forecast with each epoch. The vertical red dashed line indicates the time of landfall.

The forecasts illustrate the convergence of the location of the landfall. As early as May 17th, three days before the landfall, the forecast tracks had converged towards the observed track. One day later, on May 18th, the forecast error on the storm track location had reduced to around 50 km. The forecast wind speed and central pressure did not show as much accuracy as the forecast trajectory. The initial forecasts captured the magnitude of the intensification. However, they failed to forecast the rapid intensification that occurred during the 17-18th of May. On the other hand, the subsequent forecasts initiated during May 17th and May 18th failed to capture the rapid weakening of the system that occurred over May 19th-20th, before landfall. Overall, in terms of wind speed, the forecast evolution was accurate (within 5 m/s) only at 24 hours range.

4 Storm surge model and performance

We have developed our tide-storm surge model based on the community and open-source modelling platform SCHISM (Semi-implicit Cross-scale Hydroscience Integrated System Model) developed by Zhang et al. (2016b) - a derivative code of SELFE (Semi-implicit Eulerian-Lagrangian Finite Element) model, originally developed by Zhang and Baptista (2008). This model solves the shallow-water equations using finite-element and finite-volume schemes in an unstructured grid that can combine tri- and quad- elements. The model is applicable in baroclinic as well as barotropic ocean circulation problems, for a broad range of spatial scales from the creek scale to the ocean basins (Ye et al., 2020; Zhang et al., 2020, 2016a). It has already been shown

to have an excellent performance in reproducing shallow water processes over the Bengal shoreline and elsewhere, including the coastal tide (Krien et al., 2016), wave setup (Guérin et al., 2018) and storm surge flooding (Bertin et al., 2014; Krien et al., 2017; Fernández-Montblanc et al., 2019).

4.1 Model implementation

175 We have implemented our model on an updated bathymetry of Krien et al. (2016) with the additional inclusion of 77,000 digitized sounding points from a set of 34 nautical charts published by Bangladesh Navy (Khan et al., 2019) (See supplement Figure S1). Our bathymetric dataset is a combination of two digitized sounding datasets in the nearshore zone – one from navigational charts produced by Bangladesh Navy, and another being a bathymetry of the Hooghly estuary provided by IWAI (Inland Waterways Authority of India). Depending on the sounding points, these observations are 7 to 20 years old. The river
180 bathymetry is composed of a set of cross-sections obtained from the Bangladesh Water Development Board (BWDB), which is further interpolated at about 300 m resolution using dedicated 1D river modelling tool and GIS techniques. The south-central part of the delta is composed of a high-resolution (50 m) inland topography (See supplement Figure S2). We considered GEBCO 2014 bathymetry to complement in the deeper part of the ocean (https://www.gebco.net/data_and_products/gridded_bathymetry_data/), and SRTM digital elevation model for the rest of the inland topography as appearing in the GEBCO 2014
185 dataset (<https://www2.jpl.nasa.gov/srtm/cbanddataproduts.html>). Our unstructured model mesh was developed based on this blended bathymetry, covering the northern Bay of Bengal (11°N to 24°N) with a variable resolution based on the shallow water wave propagation and bottom slope criteria. Our model grid consists of about 600,000 nodes and 1 million elements in total. The resulting grid resolution ranges from 250 m in the estuaries and the delta, to 15 km in the deeper part of the central Bay of Bengal. The model domain and mesh are identical to those presented in Khan et al. (2019)(Figure 4).

190 To account for the effect of short waves on the mean circulation, Wind Wave Model III (WWMIII), a third-generation spectral wave model, is coupled at source code level with SCHISM in our modelling framework (Roland et al., 2012). In our configuration, WWMIII solves wave action equation on the native SCHISM grid, using a fully implicit scheme. The source terms in our model comprise the energy input due to wind (Ardhuin et al., 2010), the non-linear interaction in deep and shallow water, energy dissipation in deep and shallow water due to white capping and wave breaking, and energy dissipation due to
195 bottom friction. We run WWMIII with a 12 directional and 12 frequency bins. Every 30 minutes of SCHISM runtime, water level and currents are passed to WWM for calculating the evolution of the wave fields. Calculated wave radiation stress, total surface stress, and the wave orbital velocity are passed back to SCHISM before computing the next time step. Wave breaking is modelled, according to Battjes and Janssen (1978). As the nearshore region of Bengal delta has a very mild slope, the value for the breaking coefficient α , which controls the rate of dissipation, was reduced from 1 to 0.1 to avoid over-dissipation
200 (Pezerat et al., 2020, 2021). WWMIII was also forced along the model southern open boundary by time series of directional spectra, computed from a large-scale application of the WaveWatch3 (hereafter WW3) model (WW3DG, 2019). WW3 was implemented at the scale of the whole Indian Ocean with a spatial resolution of 0.5° and forced by the FNL reanalysis wind data at 3-hour time step (National Centers for Environmental Prediction, National Weather Service, NOAA, U.S. Department of Commerce, 2015).




Figure 4. Computational domains and model mesh for SCHISM-WWMIII, as well as model boundary conditions. White arrows on the southern boundary show the forcing with the tidal solution provided by FES2012, and on the northern boundary shows the river discharges. For a hindcast experiment, wave spectra from WW3 are imposed on the southern boundary.

205 4.2 Model Forcings

Our model is forced over the whole domain by the astronomical tidal potential for the 12 main constituents (2N2, K1, K2, M2, Mu2, N2, Nu2, O1, P1, Q1, S2, T2). At the southern boundary, we have prescribed a boundary tidal water level from 26 harmonic constituents (M2, M3, M4, M6, M8, Mf, Mm, MN4, MS4, Msf, Mu2, N2, Nu2, O1, P1, Q1, R2, S1, S2, S4, Ssa, T2, K2, K1, J1, and 2N2) extracted from FES2012 global tide model (Carrère et al., 2013). At upstream river open boundaries of Ganges and Brahmaputra, a discharge time series from BWDB is forced for the benchmark tidal simulation, and a climatologic discharge timeseries is applied for storm surge simulations during Amphan. A discharge climatology is applied at Hooghly (Mukhopadhyay et al., 2006) and Karnaphuli (Chowdhury and Al Rahim, 2012). At Meghna and Rupnarayan river open boundaries, we implemented a radiating Flather boundary condition. From a year-long tidal simulation, we found that the updated version of the bathymetry performs 2-4 times better at key locations compared to the global tidal models – which includes FES (Carrère et al., 2013), GOT (Ray, 1999), and TPXO (Egbert and Erofeeva, 2002) (see supplementary Table S1).

We derived the atmospheric wind and pressure fields for SCHISM from a blending of analytical wind field inferred from a parametric wind model (close to the cyclone track) and background atmospheric field from the Global Forecast System (GFS, <https://www.ncdc.noaa.gov/data-access/model-data/model-datasets/global-forecast-system-gfs>) reanalysis (further away), following Krien et al. (2017, 2018). We used the best-track data of JTWC provided at 6-hour time step as input for the analytical wind and pressure field. Here we choose the analytical wind profile of Emanuel and Rotunno (2011) and the analytical pressure field profile of Holland (1980). To be consistent with our forecast described in Section 5, the background wind field was generated incrementally from an accumulative merging of GFS forecasts for each 6-hour forecast window, with a 1-hour time step. The analytical and background wind fields were first temporally interpolated every 15 minutes and overlaid on the background GFS fields using a distance-varying weighting coefficient to the cyclone centre to ensure a smooth transition (Figure 7b). We took into account the asymmetry of the cyclonic wind field following Lin and Chavas (2012). The final resolution after merging the analytical wind field with the interpolated background GFS fields is 0.025° . For all storm surge simulations, a spinup time of 2 days is considered in this study.

4.3 Predictive skills

With the above-described model setup, we ran our model to calculate the water level and sea state evolution during Amphan. The simulation was done a-posteriori, once the cyclone had passed, and JTWC and GFS hindcasts being already available throughout the cyclone lifetime. Figure 5 shows comparisons between observed and modelled significant wave heights (SWH), peak wave period and water level. First, we compare the significant wave height derived from the altimetric estimate of Sentinel-3B processed by the Wave Thematic Assembly Center from Copernicus Marine Environment Monitoring Service (<https://scihub.copernicus.eu/>), which overpass on May 18th 2020 at 16:03GMT. For each altimetric data point inside our domain, we interpolated the model output using a nearest-neighbour approach, both spatially and temporally. This comparison shows that SWH is reasonably reproduced, with RMSE typically within 1 m (within 18% of the mean value). However, we observed an overestimation of SWH of about 2.5 m near the cyclone centre. One of the reasons for such difference is a limitation of analytical wind field models, as discussed by Krien et al. (2017).

We present an in situ observed time-series of SWH and mean wave period during the life cycle of Amphan in Figure 5 for BD08 (Figure 5, b-c) and BD11 (Figure 5, d-e) oceanographic buoys (the dataset is collected from INCOIS data portal at <https://incois.gov.in/portal/datainfo/mb.jsp>). The two buoys are located on either side of Amphan track, 250 km to its east / 230 km to its west, respectively. We can see that at both locations, the model reproduces the SWH well. The match is particularly good at BD11, consistently within 1 m of the observed evolution. At BD08, SWH is found to be overestimated by about 2 m at its peak, with a 12-h lead shift in time. Such difference can come from positional and timing uncertainties in JTWC best track data, from our assumption of a constant translational velocity in-between each 6-hourly JTWC time step, and finally from the intrinsic uncertainty of the analytical wind field itself, as shown on Figure 3b. Despite multiple sources of potential errors and uncertainties, our model appears capable of capturing the general features of the sea state, both significant wave height (SWH) and period, relatively well. The overall predictive skills of the model in terms of short waves are only slightly below those from




Figure 5. Comparison of simulated (in orange) and observed (in blue) water level, significant wave height (SWH) and mean wave period. (a) The map shows the along-track bias in SWH compared to the one calculated from Sentinel 3B altimeter overpass at 2020-05-18 1603Z. Bottom panel shows the modelled SWH and mean wave period (orange line) compared to buoy observations (blue dots) at BD08 (b-c) and BD11 (d-e) provided by INCOIS. Comparison between observed (blue dots) and modelled (orange line) water level at the station locations – (f) Galachipa, (g) Kuakata, (h) Tajumuddin, (i) Chittagong, and (j) Visakhapatnam. Green dashed lines in (f)-(k) indicates the modelled tidal water level. Location of the buoys and the water level gauges are shown in (a). The vertical red lines in water level plots indicate the time of landfall.

hindcast exercises typically conducted over other ocean basins using similar modelling strategies (Bertin et al., 2014; Krien
250 et al., 2018).

The real-time availability of observed water level time series in this region is relatively scanty. We were able to access only
a handful of coastal water level records, most of them located eastward of the landfall location. The right panels of Figure
5(f-j) illustrated the comparison of the recorded water level and modelled storm surge. Among these tide-gauges, Chittagong
(Figure 5j) and Visakhapatnam (Figure 5k) time series are retrieved from the UNESCO IOC Sea Level Monitoring service
255 (<http://www.ioc-sealevelmonitoring.org/map.php>) and the rest are provided by Bangladesh Inland Water Transport Authority
(BIWTA, <http://www.biwta.gov.bd/>). The datum for water level timeseries at Chittagong and Visakhapatnam is changed from
in-situ datum to MSL by removing long-term mean. The timeseries provided by BIWTA are already referenced to MSL, thus
kept unchanged. We also plotted the reconstructed tidal water level from the tidal atlas derived from our model. The overall
tidal propagation is well simulated in Galachipa and Kuakata, while at Tajumuddin the simulation tidal range is found to be
260 underestimated. The local bathymetric error and friction parameterization might be the source of the discrepancy. Tide gauge
at Tajumuddin is situated at the mouth of Meghna, where bathymetry is rapidly changing (Khan et al., 2019). However, our
model correctly reproduces the peak of the water level at all locations. The maximum recorded water level at these stations is
around 2-2.5 m above mean sea level, with a storm surge (water level – tide) of the same order of magnitude.

To spatially assess the storm surge generated by cyclone Amphan, we first take a look into the temporal maximum water
265 level in Figure 6(a). We can see that the whole coastal region experienced a high-water level ranging from 2-5 m for Amphan.
To quantify the contribution of the cyclone, we have looked in the surge component, defined as the difference between the total
modelled water level and a tide-only simulation. Figure 6(b) illustrates the temporal maximum of surge over the delta region.
The highest surge level is 5 m around the location of the landfall (88.3°E, 21.6°N). The maximum non-linear interaction
between tide and surge amounts to about 30 cm in the nearshore domain (Figure 6d).

To quantify the contribution of the wave setup, we re-calculated the storm surge without the coupling with the wave model.
270 Figure 6(c) shows the maximum of the difference between the two simulated water levels. In general, wave setup contributes
to the Amphan storm surge by about 20 cm all along the Bengal shoreline, and locally over 30 cm close to the cyclone landfall.
Nevertheless, the spatial resolution at the nearshore region employed in this study (250 m) is still too coarse for capturing the
maximum wave setup that develops along the shoreline, where a resolution of a few tens of m should be employed (Guérin
275 et al., 2018). Nonetheless, this comparison shows that wave setup is not only developed along the shoreline exposed to waves
but affects the whole delta up to far upstream, a process already observed over large estuaries (Bertin et al., 2015; Fortunato
et al., 2017).

Our findings reaffirm the necessity of a proper coupling between tide, surge and wave to forecast the water level evolution.
For Amphan, the magnitude of tide-surge interaction amounts to about 10% of the maximum total water level with spatial
280 variation. Similarly, the magnitude of wave-setup is also spatially varying with a typical amplitude of 10-15% of the maximum
total water level. This non-negligible non-linear dependency shows the importance of a fully coupled hydrodynamic-wave
modelling system for a successful forecast of the water levels in the hydrodynamic setting of the Bengal delta. Our modelling
framework showed reasonable accuracy in reproducing observed storm surge water level along the Bengal coastline. The




Figure 6. Hindcast of (a) maximum water level, (b) maximum surge (c) wave setup/setdown (d) maximum non-linear interaction between tide and surge. For (a), for the areas above mean sea level, the water level is converted to water level above the topography for consistency. The inset maps show a close-up (75 km \times 45 km) of the landfall region. The black dashed line shown in (b) is the segment for error analysis of the forecast experiment in Section 5.

accuracy is in line with the typical level of performance of similar systems applied elsewhere in the world ocean (Bertin et al.,

285 2015; Fernández-Montblanc et al., 2019; Suh et al., 2015). The hindcast experiment thus shows the viability of our model for a proof-of-concept of real-time forecasting exercise.

5 Near real-time storm surge forecasting

The current state of maturity of atmospheric real-time cyclone forecast products allows their application to real-time surge forecasting, as discussed in Section 3. As explained in Section 2, the storm surge generated from cyclones depends on the atmospheric pressure, wind, and background (typically: tidal) water level. In this section, a near real-time storm surge forecasting scheme is presented using publicly available atmospheric forcing dataset.

5.1 Forecast strategy and forcing products

During cyclone Amphan, we performed near-realtime storm surge forecasts based on our model. In our forecasts, we relied on the outputs of global models (GFS, and HWRF) and advisories (JTWC) for the estimates of the atmospheric forcing. These advisories and global models are updated every 6 hours. Similar to the GFS forecast cycle, we updated our forecasts at every 00:00, 06:00, 12:00, 18:00 UTC.

For each forecast, we derived the atmospheric forcing from a blend of JTWC, HWRF, and GFS data (Table 1). JTWC publishes storm advisories at 03:00, 09:00, 15:00, 21:00 UTC. The advisory includes an analysis of the storm intensity and position from a satellite fix 3-hours prior, followed by a forecast for the following 72 hours, with 6 to 12 hours time-steps. NOAA published their HWRF model advisory for each GFS forecast cycle at 3-hour time steps. GFS provides the wind and pressure fields in hourly timesteps at 0.25° regular spatial grid. The GFS model is initialized 6-hourly at 00:00, 06:00, 12:00, 18:00 UTC, and the data is available 3 hours after the initialized period.

Table 1. Data sources for the model forcings.

Name	Datatype	Timestep	Resolution	Source
JTWC advisory	Text	Variable (6-24 hours)	NA	https://www.metoc.navy.mil/jtwc/jtwc.html
HWRF forecast	Text	3 hours	NA	https://www.emc.ncep.noaa.gov/gc_wmb/vxt/HWRF/index.php
GFS	4D array/DAP	1 hours	0.25°	https://nomads.ncep.noaa.gov/dods/gfs_0p25_1hr
FNL	netCDF	3 hours	0.25°	https://rda.ucar.edu/datasets/ds083.3/

NA: Not applicable

The wind and pressure fields around the centre of the storm are derived analytically from a concatenated JTWC and HWRF. We used GFS data as the background wind and pressure on the outer region of the analytical fields.

305 We initiated our forecast cycles on March 16th 2020 at 06:00 UTC - utilizing the first advisory from JTWC published at 03:00 UTC, merged with the forecast from HWRF issued at 18:00 UTC of the preceding day. We took the background wind

and pressure field from GFS forecast published at 00:00 UTC. In the subsequent forecast cycles, we updated the previous track file by first replacing the past time-steps with the analysis from the latest JTWC advisory, then appending the forecasts from HWRF again. For the background wind and pressure fields from GFS, we retained the first 6 hours of forecasts from the previous cycles and updated the remainder of the time series from the latest forecast with new initialization. For each forecast cycle, we kept the starting time of our model the same, on May 16th at 00:00 UTC, and ran till one-day after the landfall for a consistent initial condition. This approach is feasible as the storms in the Bay of Bengal typically form, grow and dissipate in about a week. A graphical representation of our strategy to temporally concatenate JTWC, GFS, and HWRF forecasts over a forecast cycle is shown in Figure 7a. The spatial blending of the analytical and GFS wind and pressure field is illustrated in Figure 7b.




Figure 7. (a) Temporal combination scheme of the JTWC, GFS, and HWRF forecasts for each 6-hourly storm surge forecast epoch. (b) Spatial blending of the analytical and GFS wind and pressure field.

5.2 Real-time computation and results

Since our forecast is contingent to the JTWC advisory of three hours before, we had only a three-hour-long window to achieve the pre-processing of the forcing files, to run the model, and to complete the necessary post-processing of the model outputs. The numerical efficiency of our model made it possible to cope with this time constraint of the forecast, using a modest

320 computing resource. The computing environment we used is detailed in Table 2, which essentially amounts to a single desktop computer fitted with a high-end consumer-grade processor.

Table 2. Computing environment used during the forecast

Model simulation duration	5.5 days
Model timestep	5 minutes
WWM Coupling Timestamp	30 minutes
Processor clock speed	2.8 GHz
Number of MPI processes	20
Wall clock time (pre-processing)	10 minutes
Wall clock time (model integration)	1 hour 45 minutes
Wall clock time (post-processing)	5 minutes
Disk storage (each simulation)	8 GB
Memory usage	18 GB

Figure 8 illustrates the evolution of the maximum total water level and surge for the corresponding forecasts issued at T-60 hours, T-36 hours, and T-12 hours, where T is the time of landfall (May 20th 2020 at 08:00 to 10:00 UTC). For the forecast issued at T-60 hours, the cyclone landfall is located near 90°E, associated with the strong surge simulated over the surrounding area. As the atmospheric cyclone forecasts gradually converge towards the actual landfall location more than 100 km westward, so do our storm surge forecasts at T-36 hours and T-12 hours. The dependency of the timing of the landfall to the forecast range is notable here. At T-36 hours, the maximum water level pattern is similar to the hindcast shown in Figure 6. By T-12 hours, the magnitude of the forecast maximum water level corresponds well with the hindcast estimate.

To substantiate the gradual increase in the quality of the forecasts, we compared the maximum surge level predicted at the various forecast ranges (T-60 hours, T-36 hours, T-12 hours) with the hindcast experiment, along a line encompassing the near-shore delta (segment displayed on Figure 6). The results show that, in the T-60h forecast, the location of maximum surge appears offset eastward, by as much as 150 km. The magnitude of the maximum surge is also poorly predicted, with an overestimation of about 3 m. In the T-36h forecast, the location of the maximum surge appears largely corrected, but the magnitude of the peak remains overestimated by about 3 m. In the T-12h forecast, both the location and magnitude of the peak surge are in relatively good agreement with the hindcast. Overall, along this landfalling coastal section, the standard error of the maximum surge level amounts to 2.06 m, 1.73 m, and 0.66 m for the T-60h, T-36h, and T-12h forecast, respectively..




Figure 8. Maximum surge (a, c, e) and elevation (b, d, f) evolution for forecast initiated at (a-b) T-60 hours (2020-05-18 0000Z), (c-d) T-36 hours (2020-05-19 0000Z), (e-f) T-12 (2020-05-20 0000Z) hours before landfall. (g) Comparison of maximum surge level extracted along the section shown in white line in (a, c, e). Hindcast results (BEST) is extracted along the same line shown in Figure 6(b).

6 Discussion

6.1 Coupled tide-surge-wave dynamics

The water level dynamics over the Bay of Bengal and particularly around the Bengal delta is complex, and the various components of water level have different relative contributions depending on the location. Due to tide-surge interaction, the maximum water level can be lower or higher depending on the tidal phase in a fully non-linear tide-surge coupled simulation, compared to its linear counterpart. During Amphan the tide-surge interaction was mostly negative, except within about 10 km of the landfall location. The positive interaction near the landfall location means that the coupled tide-surge estimate was higher than the linearly added tide and surge. In this case, the typical magnitude of tide-surge interaction ranged from -30cm to +30cm. The contribution from the wave-induced setup along the shoreline increases the maximum water level estimation throughout the coast by about 10-15% during Amphan.

6.2 Performance of storm surge forecasting

Aside from the necessary inclusion of relevant physical processes – tide, surge, waves and their non-linear interactions – the performance of storm surge forecasting depends on the wind and pressure forcings. The errors in the wind forcing and in the atmospheric pressure forcing are typically the most important ones among the various sources of error in storm surge modelling (Krien et al., 2017). In our forecast environment, the source of this error is two-fold – the analytical model used to synthesize the fields from the storm parameters, and the numerical weather model used to forecast the storm wind and pressure fields to derive the storm parameters themselves. There have been attempts to find a more accurate formulation from recent satellite scatterometer missions (Krien et al., 2018). However, not one formulation works best in all radial distances from the storm centre. Despite the well-known error associated with the parametric wind field, they are widely used in applications due to their computational simplicity and lightweight data requirements (Lin and Chavas, 2012). The best way to avoid the error from the analytical wind field might be not using these formulations and rely on the full-fledged atmospheric forecasts. However, atmospheric models are costly in terms of computation. It is particularly true for cyclonic storms where high spatial resolution (typically kilometric) is required (Tallapragada et al., 2014).

From the validation of water level at a few tide gauge locations for our hindcast simulation, it appears that the modelling framework proposed here could capture the maximum surge level successfully throughout the coast. Our tide-gauge sites were limited to the eastern side of Amphan landfall location and were operational throughout the life cycle of the cyclone. However, it was not possible to confirm the ability of the model to reproduce the water level on the western side of the landfall location due to data unavailability. The proper reproduction of the maximum water level by the model gives strong confidence in the formulation as well as in the coupling strategy of our modelling framework.

6.3 Inundation

One of the potential, but challenging, outcomes of storm surge forecasting is the prediction of inundation. Particularly in the northern Bay of Bengal, the existing modelling studies do not generally consider the inundation process (Dube et al., 2004; Murty et al., 2014). Some modelling systems take advantage of simplified inundation modelling schemes to tackle this problem (Lewis et al., 2013). In our modelling framework, the inland inundation is calculated seamlessly by SCHISM, solving the same hydrodynamics over the model domain, thanks to the combination of cross-scale unstructured-grid and an efficient wetting-drying algorithm. While the recent improvements in models bathymetry and the detailed accounting of dikes and coastal defences improved the overall modelling of the inundation, the associated error remains large (Krien et al., 2017). As an update to Krien et al. (2017), in our model we have incorporated a novel dike heights dataset, bearing varying dike crests heights, for the numerous polders scattered around the Bengal delta. The assessment of the impact of the updated embankment heights is, however, hard to quantify and validate.

In the absence of any operational network of inundation monitoring, to understand and better characterize the inundation dynamics, we systematically skimmed the Bengal delta local newspapers on the day of Amphan landfall and on the few subsequent days, to achieve an as-comprehensive-as-possible mapping of the inundation extent observed in situ. We digitized the reported flooded locations through © Google Earth and categorized the inundation mechanism. We digitized 88 such locations over the Bangladeshi part of Bengal delta, as shown in Figure 9 (Khan, 2020). Over the Indian side of the delta, the news reporting of inland flooding from dike breaching was not accurate enough for us to be able to geotag the locations. We overlaid the inundated locations over the inundation predicted by the model on a false-colour image derived from Sentinel-2 satellite using © Google Earth Engine (<https://earthengine.google.com>). Three categories of inundation mechanism are typically observed – inundation by breaching of the dikes (labelled as "Embankment breach"), inundation of unprotected low land by increased water level ("Unembanked lowland"), and flooding by overflowing of the dikes ("Embankment overflow"). From the reported news, it is clear that the major part of the inundation results from the breaching of dikes. We also note two breaching instances in the south-eastern corner of our domain, very far from the cyclone landfall. On the other hand, the flooding from merely increased water level in non-diked areas is mostly concentrated along the Meghna estuary. Along the estuary, designed embankments probably do not exist to protect many populated low-lying char areas (land/island formed through accretion). We could find only one report of the occurrence of overflowing of an embankment, in Maheshwaripur of Koyra upazila (89.30°E, 22.45°N). Around (90.1°E-90.5°E, 22.4°N-22.6°N), in Barishal district, our model predicted an extended inundated area, which is known not to be polderized. While we could find some reports of flooding around that region from our newspaper survey, the widespread inundation predicted by our model in this region probably did not occur in reality due to the presence of levees and city protection embankments. Indeed, this kind of small-scale geographic information, to the best of our knowledge, is not systematically available publicly and thus could not be incorporated in our model topography, despite being of utmost importance for localized inundation forecast. The forecast flooded vs non-flooded areas show a wealth of spatial scales, demonstrating that our unstructured modelling framework has the intrinsic capability to model the small-scale hydrodynamic gradients. In our modelled forecasts, the contrast between the well-predicted water level temporal evolution, and the

400 comparatively limited predictive skills of inundated locations points out the necessity of accurate knowledge and incorporation of reliable topographic and embankment information, at the local scale. Our modelling system has the ability and appropriate resolution to ingest such topographic information efficiently due to its unstructured nature.




Figure 9. Digitized location of inundation resulting from embankment breaching (red), flooding of low-lying unembanked area (orange), and embankment overflow (cyan) overlaid on a false-colour image composite derived from B12, B11, and B4 channels of Sentinel-2 during April 2020. The grey patches are the inundated regions predicted by the model, as shown in the hindcast experiment. The polders are shown with cyan outlines.

The coastal polder management has long been a boon and a bane (Warner et al., 2018). On one side, the polders have protected the fertile areas from daily tidal flooding of brackish waters, which favoured agricultural production by reducing the soil salinity (Nowreen et al., 2013). On the other hand, these embankments have restricted freshwater and sediment supply, causing significant subsidence inside the embanked areas due to sediment compaction (Auerbach et al., 2015). Over time, the land use inside these polders has also changed from agriculture to aquaculture (namely shrimp and crab farming). Although the embankments were not built to protect from the cyclonic surges, in many instances they functioned as such. The presence of these protective structures also contributed to a false sense of security in the residents living inside (Paul and Dutt, 2010).
410 Embankments have become part and parcel of survival in this low-lying region. It has now become essential to monitor the condition and topography of the embankments for efficient management and maintenance. From a forecasting point of view, embankment breaching is extremely hard to model in the state-of-the-art hydrodynamic modelling frameworks. However, consistent periodic monitoring of the embankment conditions can improve the forecast by providing a more objective view of the associated inundation hazard.

415 **6.4 Prospects in operational forecasting**

Throughout our forecast and hindcast experiments, we have relied on freely available forecast products as the forcing for our storm surge model. These forecasts data are publicly available within 3 hours of their initialization time through online portals.

The primary input to generate the forcing is the forecast storm track distributed as a text file, which is very lightweight even for a limited internet connection. We have implemented our storm surge model on a freely available open-source framework. The
420 model code is already operationally used in regional forecasting systems and services (Fortunato et al., 2017; Oliveira et al., 2020).

As we have demonstrated, the modelling setup presented here is tractable in an operational, real-time forecasting scenario. Thanks to advanced and efficient numerics, one can deploy such a system in a consumer-grade computing environment. The computing setup we have deliberately used (see Table 2) essentially amounts to a high-end x86-64 desktop personal computer
425 (e.g. Intel® Core™i9-10900 or similar). This computing requirement takes only a portion of the available computing capacity of an institute such as the BMD (15x4 cores@2.8GHz) (Roy et al., 2015). Furthermore, the model is easily deployable in cloud computing infrastructure (Oliveira et al., 2020), which could provide additional reliability and cost-savings for an event-driven operational forecasting system.

7 Conclusions

In this study, we present a retrospective evaluation of storm surge prediction during super-cyclone Amphan using an efficient, state-of-the-art coupled hydrodynamic-wave numerical model. During cyclone Amphan, the predicted maximum water level ranges from 2 to 5 m along the coastline. Comparison with in-situ measurements revealed that the water level could be modelled with high accuracy on condition that all the relevant mechanisms are considered. Notably, we demonstrated the necessity of considering a coupled hydrodynamic tide-surge-wave modelling framework for the head Bay of Bengal, for effective fore-
435 casting. For Amphan, the contribution from tide-surge interaction, and wave-setup typically amounts to 10% and 10-15% of the total water level, respectively.

From our proactive forecast initiative during cyclone Amphan, we showed that with publicly available storm forecast products and easily accessible computing resources, it is feasible to forecast the evolution of water level throughout the vast coastline of the Bengal delta in real-time. During Amphan, a sufficiently skilful storm surge forecast was achieved as early as 36-48 hours
440 before landfall. The forecast water level with 36-hour lead time seemed quite similar to our best (hindcast) simulation, in terms of maximum water level as well as the spatial pattern. We communicated the results to Bangladesh local government authority through personal communications, as well as to the scientific community through social media.

From a secondary post-disaster news-survey, we have identified the limiting factors for location-specific inundation forecasts. The main limiting factor is due to limited and outdated topographic information of the existing coastal defences. Also,
445 dike-breaching, which was the prevalent process of inland inundation during Amphan, is not explicitly modelled in our hydrodynamic framework. These two factors call for routine monitoring of embankment topography and condition.

Cyclone and storm surge warning has always been a communication and trust issue in the Bay of Bengal (Paul and Dutt, 2010; Roy et al., 2015). It is thus necessary to communicate well-grounded storm and surge forecast to the community for coordinated and informed decision-making during a storm surge (Morss et al., 2018). The forecasting system we implemented

450 and assessed in the present case study provides the proof-of-feasibility and opens short-term operational prospects to fill a gap
in the existing disaster management tools in this part of the world.

Code and data availability. The instructions to download and install the model used in this study can be accessed freely at <https://github.com/schism-dev/schism>. The sources of the data used in this study have been described in the article. The data should be requested through the mentioned institutions or downloaded from the provided websites. The geolocation of inundation mapping from newspaper can be found at
455 <https://dx.doi.org/10.5281/zenodo.4086102>.

Author contributions. JK and FD formulated the study. JK did the numerical simulations, analysis, and wrote the first draft. XB computed the simulation for WW3 boundary used in hindcast. All co-authors contributed to this study through multiple discussions. MP provided help with the WWM formulation. SH disseminated the forecasts to local government of Bangladesh, and ASI to the scientific community during cyclone Amphan.

460 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. We acknowledge financial support from CNES (through the TOSCA project BANDINO) and Embassy of France in Bangladesh. This work was supported by the French research agency (Agence Nationale de la Recherche; ANR) under the DELTA project (ANR-17-CE03-0001). We are thankful to LIENSs Laboratory (University of La Rochelle, France) for hosting JK and LT during this study.

References

- 465 Ahsan, M. N., Khatun, A., Islam, M. S., Vink, K., Ohara, M., and Fakhruddin, B. S.: Preferences for improved early warning services among coastal communities at risk in cyclone prone south-west region of Bangladesh, *Progress in Disaster Science*, 5, 100065, <https://doi.org/10.1016/j.pdisas.2020.100065>, 2020.
- Alam, E. and Dominey-Howes, D.: A new catalogue of tropical cyclones of the northern Bay of Bengal and the distribution and effects of selected landfalling events in Bangladesh, *International Journal of Climatology*, 35, 801–835, <https://doi.org/10.1002/joc.4035>, publisher: Wiley, 2014.
- 470 Ali, A.: Climate change impacts and adaptation assessment in Bangladesh, *Climate Research*, 12, 109–116, <https://doi.org/10.3354/cr012109>, publisher: Inter-Research Science Center, 1999.
- Antony, C. and Unnikrishnan, A. S.: Observed characteristics of tide-surge interaction along the east coast of India and the head of Bay of Bengal, *Estuarine, Coastal and Shelf Science*, 131, 6–11, <https://doi.org/10.1016/j.ecss.2013.08.004>, publisher: Elsevier BV, 2013.
- 475 Antony, C., Unnikrishnan, A., Krien, Y., Murty, P., Samiksha, S., and Islam, A.: Tide–surge interaction at the head of the Bay of Bengal during Cyclone Aila, *Regional Studies in Marine Science*, 35, 101–133, <https://doi.org/10.1016/j.rsma.2020.101133>, 2020.
- Ardhuin, F., Rogers, E., Babanin, A. V., Filipot, J.-F., Magne, R., Roland, A., van der Westhuysen, A., Queffelecoul, P., Lefevre, J.-M., Aouf, L., and Collard, F.: Semiempirical Dissipation Source Functions for Ocean Waves. Part I: Definition, Calibration, and Validation, *Journal of Physical Oceanography*, 40, 1917–1941, <https://doi.org/10.1175/2010jpo4324.1>, 2010.
- 480 As-Salek, J. A. and Yasuda, T.: Tide–Surge Interaction in the Meghna Estuary: Most Severe Conditions, *Journal of Physical Oceanography*, 31, 3059–3072, [https://doi.org/10.1175/1520-0485\(2001\)031<3059:tsiitm>2.0.co;2](https://doi.org/10.1175/1520-0485(2001)031<3059:tsiitm>2.0.co;2), publisher: American Meteorological Society, 2001.
- Auerbach, L. W., Jr, S. L. G., Mondal, D. R., Wilson, C. A., Ahmed, K. R., Roy, K., Steckler, M. S., Small, C., Gilligan, J. M., and Ackerly, B. A.: Flood risk of natural and embanked landscapes on the Ganges–Brahmaputra tidal delta plain, *Nature Climate Change*, 5, 153–157, <https://doi.org/10.1038/nclimate2472>, publisher: Springer Nature, 2015.
- 485 Battjes, J. A. and Janssen, J. P. F. M.: Energy Loss and Set-Up Due to Breaking of Random Waves, in: *Coastal Engineering 1978*, pp. 569–587, American Society of Civil Engineers, Hamburg, Germany, <https://doi.org/10.1061/9780872621909.034>, <http://ascelibrary.org/doi/10.1061/9780872621909.034>, 1978.
- Bernier, N. B. and Thompson, K. R.: Deterministic and ensemble storm surge prediction for Atlantic Canada with lead times of hours to ten days, *Ocean Modelling*, 86, 114–127, <https://doi.org/10.1016/j.ocemod.2014.12.002>, publisher: Elsevier BV, 2015.
- 490 Bertin, X., Li, K., Roland, A., Zhang, Y. J., Breilh, J. F., and Chaumillon, E.: A modeling-based analysis of the flooding associated with Xynthia, central Bay of Biscay, *Coastal Engineering*, 94, 80–89, <https://doi.org/10.1016/j.coastaleng.2014.08.013>, 2014.
- Bertin, X., Li, K., Roland, A., and Bidlot, J.-R.: The contribution of short-waves in storm surges: Two case studies in the Bay of Biscay, *Continental Shelf Research*, 96, 1–15, <https://doi.org/10.1016/j.csr.2015.01.005>, 2015.
- Bhardwaj, P. and Singh, O.: Climatological characteristics of Bay of Bengal tropical cyclones: 1972–2017, *Theoretical and Applied Climatology*, 139, 615–629, <https://doi.org/10.1007/s00704-019-02989-4>, publisher: Springer Science and Business Media LLC, 2019.
- 495 Bouwer, L. M. and Jonkman, S. N.: Global mortality from storm surges is decreasing, *Environmental Research Letters*, 13, 014008, <https://doi.org/10.1088/1748-9326/aa98a3>, publisher: IOP Publishing, 2018.
- Carrère, L., Lyard, F., Cancet, M., Guillot, A., and Roblou, L.: FES 2012: a new global tidal model taking advantage of nearly 20 years of altimetry, in: *20 Years of Progress in Radar Altimetry*, vol. 710, 2013.

- 500 Chowdhury, M. A. M. and Al Rahim, M.: A proposal on new scheduling of turbine discharge at Kaptai hydro-electric power plant to avoid the wastage of water due to overflow in the dam, in: 2012 7th International Conference on Electrical and Computer Engineering, pp. 758–762, IEEE, 2012.
- Condon, A. J., Sheng, Y. P., and Paramygin, V. A.: Toward High-Resolution, Rapid, Probabilistic Forecasting of the Inundation Threat from Landfalling Hurricanes, *Monthly Weather Review*, 141, 1304–1323, <https://doi.org/10.1175/mwr-d-12-00149.1>, publisher: American Meteorological Society, 2013.
- 505 Daniel, P., Haie, B., and Aubail, X.: Operational Forecasting of Tropical Cyclones Storm Surges at Meteo-France, *Marine Geodesy*, 32, 233–242, <https://doi.org/10.1080/01490410902869649>, publisher: Informa UK Limited, 2009.
- Das, P. K.: Prediction Model for Storm Surges in the Bay of Bengal, *Nature*, 239, 211–213, <https://doi.org/10.1038/239211a0>, publisher: Springer Science and Business Media LLC, 1972.
- 510 Deb, M. and Ferreira, C. M.: Simulation of cyclone-induced storm surges in the low-lying delta of Bangladesh using coupled hydrodynamic and wave model (SWAN plus ADCIRC), *Journal of Flood Risk Management*, 11, S750–S765, <https://doi.org/10.1111/jfr3.12254>, publisher: Wiley, 2016.
- DhakaTribune: Govt: Cyclone caused damage worth Tk1,100 crore, <https://www.dhakatribune.com/bangladesh/crisis/2020/05/21/govt-estimated-damages-from-amphan-worth-1-100-crore>, 2020.
- 515 Dube, S. K., Chittibabu, P., Sinha, P. C., Rao, A. D., and Murty, T. S.: Numerical Modelling of Storm Surge in the Head Bay of Bengal Using Location Specific Model, *Natural Hazards*, 31, 437–453, <https://doi.org/10.1023/b:nhaz.0000023361.94609.4a>, publisher: Springer Nature, 2004.
- Dube, S. K., Jain, I., Rao, A. D., and Murty, T. S.: Storm surge modelling for the Bay of Bengal and Arabian Sea, *Natural Hazards*, 51, 3–27, <https://doi.org/10.1007/s11069-009-9397-9>, publisher: Springer Science and Business Media LLC, 2009.
- 520 Durand, F., Piecuch, C. G., Becker, M., Papa, F., Raju, S. V., Khan, J. U., and Ponte, R. M.: Impact of Continental Freshwater Runoff on Coastal Sea Level, *Surveys in Geophysics*, <https://doi.org/10.1007/s10712-019-09536-w>, 2019.
- Egbert, G. D. and Erofeeva, S. Y.: Efficient inverse modeling of barotropic ocean tides, *Journal of Atmospheric and Oceanic Technology*, 19, 183–204, 2002.
- Emanuel, K.: Increasing destructiveness of tropical cyclones over the past 30 years, *Nature*, 436, 686–688, <https://doi.org/10.1038/nature03906>, 2005.
- 525 Emanuel, K. and Rotunno, R.: Self-Stratification of Tropical Cyclone Outflow. Part I: Implications for Storm Structure, *Journal of the Atmospheric Sciences*, 68, 2236–2249, <https://doi.org/10.1175/jas-d-10-05024.1>, publisher: American Meteorological Society, 2011.
- Fernández-Montblanc, T., Vousdoukas, M., Ciavola, P., Voukouvalas, E., Mentaschi, L., Breyiannis, G., Feyen, L., and Salamon, P.: Towards robust pan-European storm surge forecasting, *Ocean Modelling*, 133, 129–144, <https://doi.org/10.1016/j.ocemod.2018.12.001>, <https://linkinghub.elsevier.com/retrieve/pii/S1463500318301823>, 2019.
- 530 Flather, R. A.: A Storm Surge Prediction Model for the Northern Bay of Bengal with Application to the Cyclone Disaster in April 1991, *Journal of Physical Oceanography*, 24, 172–190, [https://doi.org/10.1175/1520-0485\(1994\)024<0172:asspmf>2.0.co;2](https://doi.org/10.1175/1520-0485(1994)024<0172:asspmf>2.0.co;2), publisher: American Meteorological Society, 1994.
- Flowerdew, J., Mylne, K., Jones, C., and Tittley, H.: Extending the forecast range of the UK storm surge ensemble, *Quarterly Journal of the Royal Meteorological Society*, 139, 184–197, <https://doi.org/10.1002/qj.1950>, publisher: Wiley, 2012.
- 535 Fortunato, A. B., Oliveira, A., Rogeiro, J., Tavares da Costa, R., Gomes, J. L., Li, K., de Jesus, G., Freire, P., Rilo, A., Mendes, A., Rodrigues, M., and Azevedo, A.: Operational forecast framework applied to extreme sea levels at regional and local scales, *Journal of*

- Operational Oceanography, 10, 1–15, <https://doi.org/10.1080/1755876X.2016.1255471>, <https://www.tandfonline.com/doi/full/10.1080/1755876X.2016.1255471>, 2017.
- 540 Frank, N. L. and Husain, S.: The deadliest tropical cyclone in history?, *Bulletin of the American Meteorological Society*, 52, 438–445, publisher: American Meteorological Society, 1971.
- Glahn, B., Taylor, A., Kurkowski, N., and Shaffer, W.: The role of the SLOSH model in National Weather Service storm surge forecasting, *National Weather Digest*, 33, 3–14, 2009.
- Guérin, T., Bertin, X., Coulombier, T., and de Bakker, A.: Impacts of wave-induced circulation in the surf zone on wave setup, *Ocean Modelling*, 123, 86–97, <https://doi.org/10.1016/j.ocemod.2018.01.006>, <https://linkinghub.elsevier.com/retrieve/pii/S1463500318300246>, 2018.
- Holland, G. J.: An Analytic Model of the Wind and Pressure Profiles in Hurricanes, *Monthly Weather Review*, 108, 1212–1218, [https://doi.org/10.1175/1520-0493\(1980\)108<1212:aamotw>2.0.co;2](https://doi.org/10.1175/1520-0493(1980)108<1212:aamotw>2.0.co;2), publisher: American Meteorological Society, 1980.
- Idier, D., Bertin, X., Thompson, P., and Pickering, M. D.: Interactions Between Mean Sea Level, Tide, Surge, Waves and Flooding: Mechanisms and Contributions to Sea Level Variations at the Coast, *Surveys in Geophysics*, 40, 1603–1630, <https://doi.org/10.1007/s10712-019-09549-5>, publisher: Springer Science and Business Media LLC, 2019.
- 550 IFRC: Operation Update Report, <http://adore.ifrc.org/Download.aspx?FileId=329843>, 2020a.
- IFRC: Bangladesh: Cyclone - Final report early action, <https://reliefweb.int/sites/reliefweb.int/files/resources/EAP2018BD01fr.pdf>, 2020b.
- Islam, A. S., Bala, S. K., Hussain, M. A., Hossain, M. A., and Rahman, M. M.: Performance of Coastal Structures during Cyclone Sidr, *Natural Hazards Review*, 12, 111–116, [https://doi.org/10.1061/\(asce\)nh.1527-6996.0000031](https://doi.org/10.1061/(asce)nh.1527-6996.0000031), publisher: American Society of Civil Engineers (ASCE), 2011.
- Ji, M., Aikman, F., and Lozano, C.: Toward improved operational surge and inundation forecasts and coastal warnings, *Natural Hazards*, 53, 195–203, <https://doi.org/10.1007/s11069-009-9414-z>, publisher: Springer Science and Business Media LLC, 2009.
- Johns, B. and Ali, M. A.: The numerical modelling of storm surges in the Bay of Bengal, *Quarterly Journal of the Royal Meteorological Society*, 106, 1–18, <https://doi.org/10.1002/qj.49710644702>, publisher: Wiley, 1980.
- 560 Khalid, A. and Ferreira, C.: Advancing real-time flood prediction in large estuaries: iFLOOD a fully coupled surge-wave automated web-based guidance system, *Environmental Modelling & Software*, p. 104748, <https://doi.org/10.1016/j.envsoft.2020.104748>, publisher: Elsevier BV, 2020.
- Khalil, G. M.: The catastrophic cyclone of April 1991: Its Impact on the economy of Bangladesh, *Natural Hazards*, 8, 263–281, <https://doi.org/10.1007/bf00690911>, publisher: Springer Science and Business Media LLC, 1993.
- 565 Khan, M. J. U.: Digitized flood location dataset during cyclone Amphan from newspaper survey, <https://doi.org/10.5281/ZENODO.4086102>, 2020.
- Khan, M. J. U., Ansary, M. N., Durand, F., Testut, L., Ishaque, M., Calmant, S., Krien, Y., Islam, A. K. M. S., and Papa, F.: High-Resolution Intertidal Topography from Sentinel-2 Multi-Spectral Imagery: Synergy between Remote Sensing and Numerical Modeling, *Remote Sensing*, 11, 2888, <https://doi.org/10.3390/rs11242888>, publisher: MDPI AG, 2019.
- 570 Krien, Y., Mayet, C., Testut, L., Durand, F., Tazkia, A. R., Islam, A. K. M. S., Gopalakrishna, V. V., Becker, M., Calmant, S., Shum, C. K., Khan, Z. H., Papa, F., and Ballu, V.: Improved Bathymetric Dataset and Tidal Model for the Northern Bay of Bengal, *Marine Geodesy*, 39, 422–438, <https://doi.org/10.1080/01490419.2016.1227405>, publisher: Informa UK Limited, 2016.

- 575 Krien, Y., Testut, L., Islam, A. K. M. S., Bertin, X., Durand, F., Mayet, C., Tazkia, A. R., Becker, M., Calmant, S., Papa, F., Ballu, V., Shum, C. K., and Khan, Z. H.: Towards improved storm surge models in the northern Bay of Bengal, *Continental Shelf Research*, 135, 58–73, <https://doi.org/10.1016/j.csr.2017.01.014>, publisher: Elsevier BV, 2017.
- Krien, Y., Arnaud, G., Cécé, R., Ruf, C., Belmadani, A., Khan, J., Bernard, D., Islam, A. K. M. S., Durand, F., Testut, L., Palany, P., and Zahibo, N.: Can We Improve Parametric Cyclonic Wind Fields Using Recent Satellite Remote Sensing Data?, *Remote Sensing*, 10, 1963, <https://doi.org/10.3390/rs10121963>, publisher: MDPI AG, 2018.
- 580 Kumar, T. S., Murty, P., Kumar, M. P., Kumar, M. K., Padmanabham, J., Kumar, N. K., Shenoi, S., Mohapatra, M., Nayak, S., and Mohanty, P.: Modeling Storm Surge and its Associated Inland Inundation Extent Due to Very Severe Cyclonic Storm Phailin, *Marine Geodesy*, 38, 345–360, <https://doi.org/10.1080/01490419.2015.1053640>, 2015.
- Kuroda, T., Saito, K., Kunii, M., and Kohno, N.: Numerical Simulations of Myanmar Cyclone Nargis and the Associated Storm Surge Part I: Forecast Experiment with a Nonhydrostatic Model and Simulation of Storm Surge, *Journal of the Meteorological Society of Japan*, 88, 521–545, <https://doi.org/10.2151/jmsj.2010-315>, publisher: Meteorological Society of Japan, 2010.
- 585 Lane, E. M., Walters, R. A., Gillibrand, P. A., and Uddstrom, M.: Operational forecasting of sea level height using an unstructured grid ocean model, *Ocean Modelling*, 28, 88–96, <https://doi.org/10.1016/j.ocemod.2008.11.004>, publisher: Elsevier BV, 2009.
- Lazo, J. K. and Waldman, D. M.: Valuing improved hurricane forecasts, *Economics Letters*, 111, 43–46, <https://doi.org/10.1016/j.econlet.2010.12.012>, publisher: Elsevier BV, 2011.
- 590 Lazo, J. K., Bostrom, A., Morss, R. E., Demuth, J. L., and Lazrus, H.: Factors Affecting Hurricane Evacuation Intentions, *Risk Analysis*, 35, 1837–1857, <https://doi.org/10.1111/risa.12407>, 2015.
- Lewis, M., Bates, P., Horsburgh, K., Neal, J., and Schumann, G.: A storm surge inundation model of the northern Bay of Bengal using publicly available data, *Quarterly Journal of the Royal Meteorological Society*, 139, 358–369, <https://doi.org/10.1002/qj.2040>, publisher: Wiley, 2013.
- 595 Lin, N. and Chavas, D.: On hurricane parametric wind and applications in storm surge modeling, *Journal of Geophysical Research: Atmospheres*, 117, n/a–n/a, <https://doi.org/10.1029/2011jd017126>, publisher: American Geophysical Union (AGU), 2012.
- Loftis, J. D., Mitchell, M., Schatt, D., Forrest, D. R., Wang, H. V., Mayfield, D., and Stiles, W. A.: Validating an Operational Flood Forecast Model Using Citizen Science in Hampton Roads, VA, USA, *Journal of Marine Science and Engineering*, 7, 242, <https://doi.org/10.3390/jmse7080242>, 2019.
- 600 Longuet-Higgins, M. S. and Stewart, R. w.: Radiation stresses in water waves—a physical discussion, with applications, *Deep Sea Research and Oceanographic Abstracts*, 11, 529–562, [https://doi.org/10.1016/0011-7471\(64\)90001-4](https://doi.org/10.1016/0011-7471(64)90001-4), publisher: Elsevier BV, 1962.
- Madsen, H. and Jakobsen, F.: Cyclone induced storm surge and flood forecasting in the northern Bay of Bengal, *Coastal Engineering*, 51, 277–296, <https://doi.org/10.1016/j.coastaleng.2004.03.001>, publisher: Elsevier BV, 2004.
- 605 Magnusson, L., Bidlot, J.-R., Bonavita, M., Brown, A. R., Browne, P. A., Chiara, G. D., Dahoui, M., Lang, S. T. K., McNally, T., Mogensen, K. S., Pappenberger, F., Prates, F., Rabier, F., Richardson, D. S., Vitart, F., and Malardel, S.: ECMWF Activities for Improved Hurricane Forecasts, *Bulletin of the American Meteorological Society*, 100, 445–458, <https://doi.org/10.1175/bams-d-18-0044.1>, publisher: American Meteorological Society, 2019.
- Melton, G., Gall, M., Mitchell, J. T., and Cutter, S. L.: Hurricane Katrina storm surge delineation: implications for future storm surge forecasts and warnings, *Natural Hazards*, 54, 519–536, <https://doi.org/10.1007/s11069-009-9483-z>, publisher: Springer Science and Business Media LLC, 2009.
- 610

- Miller, R. J., Schrader, A. J., Sampson, C. R., and Tsui, T. L.: The Automated Tropical Cyclone Forecasting System (ATCF), *Weather and Forecasting*, 5, 653–660, [https://doi.org/10.1175/1520-0434\(1990\)005<0653:tatcfs>2.0.co;2](https://doi.org/10.1175/1520-0434(1990)005<0653:tatcfs>2.0.co;2), publisher: American Meteorological Society, 1990.
- 615 Morss, R. E., Cuite, C. L., Demuth, J. L., Hallman, W. K., and Shwom, R. L.: Is storm surge scary? The influence of hazard, impact, and fear-based messages and individual differences on responses to hurricane risks in the USA, *International Journal of Disaster Risk Reduction*, 30, 44–58, <https://doi.org/10.1016/j.ijdr.2018.01.023>, publisher: Elsevier BV, 2018.
- Mukhopadhyay, S., Biswas, H., De, T., and Jana, T.: Fluxes of nutrients from the tropical River Hooghly at the land–ocean boundary of Sundarbans, NE Coast of Bay of Bengal, India, *Journal of Marine Systems*, 62, 9–21, <https://doi.org/10.1016/j.jmarsys.2006.03.004>,
620 2006.
- Munroe, R., Montz, B., and Curtis, S.: Getting More out of Storm Surge Forecasts: Emergency Support Personnel Needs in North Carolina, *Weather, Climate, and Society*, 10, 813–820, <https://doi.org/10.1175/wcas-d-17-0074.1>, 2018.
- Murty, P. L. N., Sandhya, K. G., Bhaskaran, P. K., Jose, F., Gayathri, R., Nair, T. M. B., Kumar, T. S., and Sheno, S. S. C.: A coupled hydrodynamic modeling system for PHAILIN cyclone in the Bay of Bengal, *Coastal Engineering*, 93, 71–81,
625 <https://doi.org/10.1016/j.coastaleng.2014.08.006>, publisher: Elsevier BV, 2014.
- Murty, P. L. N., Bhaskaran, P. K., Gayathri, R., Sahoo, B., Kumar, T. S., and SubbaReddy, B.: Numerical study of coastal hydrodynamics using a coupled model for Hudhud cyclone in the Bay of Bengal, *Estuarine, Coastal and Shelf Science*, 183, 13–27, <https://doi.org/10.1016/j.ecss.2016.10.013>, publisher: Elsevier BV, 2016.
- Murty, P. L. N., Padmanabham, J., Kumar, T. S., Kumar, N. K., Chandra, V. R., Sheno, S. S. C., and Mohapatra, M.:
630 Real-time storm surge and inundation forecast for very severe cyclonic storm ‘Hudhud’, *Ocean Engineering*, 131, 25–35, <https://doi.org/10.1016/j.oceaneng.2016.12.026>, publisher: Elsevier BV, 2017.
- Murty, T. S., Flather, R. A., and Henry, R. F.: The storm surge problem in the bay of Bengal, *Progress in Oceanography*, 16, 195–233, [https://doi.org/10.1016/0079-6611\(86\)90039-x](https://doi.org/10.1016/0079-6611(86)90039-x), publisher: Elsevier BV, 1986.
- National Centers for Environmental Prediction, National Weather Service, NOAA, U.S. Department of Commerce: NCEP GDAS/FNL 0.25
635 Degree Global Tropospheric Analyses and Forecast Grids, Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory, Boulder CO, <https://doi.org/10.5065/D65Q4T4Z>, 2015.
- Needham, H. F., Keim, B. D., and Sathiaraj, D.: A review of tropical cyclone-generated storm surges: Global data sources, observations, and impacts, *Reviews of Geophysics*, 53, 545–591, <https://doi.org/10.1002/2014rg000477>, publisher: American Geophysical Union (AGU), 2015.
- 640 Nowreen, S., Jalal, M. R., and Khan, M. S. A.: Historical analysis of rationalizing South West coastal polders of Bangladesh, *Water Policy*, 16, 264–279, <https://doi.org/10.2166/wp.2013.172>, publisher: IWA Publishing, 2013.
- Oliveira, A., Fortunato, A., Rogeiro, J., Teixeira, J., Azevedo, A., Lavaud, L., Bertin, X., Gomes, J., David, M., Pina, J., Rodrigues, M., and Lopes, P.: OPENCoastS: An open-access service for the automatic generation of coastal forecast systems, *Environmental Modelling & Software*, 124, 104 585, <https://doi.org/10.1016/j.envsoft.2019.104585>, <https://linkinghub.elsevier.com/retrieve/pii/S1364815219304967>,
645 2020.
- Paul, B. K.: Why relatively fewer people died? The case of Bangladesh’s Cyclone Sidr, *Natural Hazards*, 50, 289–304, <https://doi.org/10.1007/s11069-008-9340-5>, 2009.
- Paul, B. K. and Dutt, S.: Hazard Warnings and Responses to Evacuation Orders: the Case of Bangladesh’s Cyclone Sidr, *Geographical Review*, 100, 336–355, <https://doi.org/10.1111/j.1931-0846.2010.00040.x>, 2010.

- 650 Pezerat, M., Martins, K., and Bertin, X.: Modelling Storm Waves in the Nearshore Area Using Spectral Models, *Journal of Coastal Research*, 95, 1240, <https://doi.org/10.2112/SI95-240.1>, <https://bioone.org/journals/journal-of-coastal-research/volume-95/issue-sp1/SI95-240.1/Modelling-Storm-Waves-in-the-Nearshore-Area-Using-Spectral-Models/10.2112/SI95-240.1.full>, 2020.
- Pezerat, M., Bertin, X., Martins, K., Mengual, B., and Hamm, L.: Simulating storm waves in the nearshore area using spectral model: Current issues and a pragmatic solution, *Ocean Modelling*, 158, 101737, <https://doi.org/10.1016/j.ocemod.2020.101737>, <http://dx.doi.org/10.1016/j.ocemod.2020.101737>, 2021.
- 655 Ray, R. D.: A global ocean tide model from TOPEX/POSEIDON altimetry: GOT99. 2, National Aeronautics and Space Administration, Goddard Space Flight Center, 1999.
- Roland, A., Zhang, Y. J., Wang, H. V., Meng, Y., Teng, Y.-C., Maderich, V., Brovchenko, I., Dutour-Sikiric, M., and Zanke, U.: A fully coupled 3D wave-current interaction model on unstructured grids, *Journal of Geophysical Research: Oceans*, 117, n/a–n/a, <https://doi.org/10.1029/2012jc007952>, publisher: American Geophysical Union (AGU), 2012.
- 660 Roy, C., Sarkar, S. K., Aberg, J., and Kovordanyi, R.: The current cyclone early warning system in Bangladesh: Providers' and receivers' views, *International Journal of Disaster Risk Reduction*, 12, 285–299, <https://doi.org/10.1016/j.ijdr.2015.02.004>, publisher: Elsevier BV, 2015.
- RSMC: Super Cyclonic Storm “AMPHAN” over the southeast Bay of Bengal (16th – 21st May 2020): Summary, Tech. rep., Regional Specialised Meteorological Centre - Tropical Cyclones, New Delhi, www.rsmcnewdelhi.imd.gov.in/images/pdf/publications/preliminary-report/amphan.pdf, 2020.
- 665 Schmidt, S., Kemfert, C., and H ppe, P.: Tropical cyclone losses in the USA and the impact of climate change — A trend analysis based on data from a new approach to adjusting storm losses, *Environmental Impact Assessment Review*, 29, 359–369, <https://doi.org/10.1016/j.eiar.2009.03.003>, <https://linkinghub.elsevier.com/retrieve/pii/S0195925509000493>, 2009.
- 670 Seo, S. N. and Bakkensen, L. A.: Is Tropical Cyclone Surge, Not Intensity, What Kills So Many People in South Asia?, *Weather, Climate, and Society*, 9, 171–181, <https://doi.org/10.1175/wcas-d-16-0059.1>, publisher: American Meteorological Society, 2017.
- Suh, S. W., Lee, H. Y., Kim, H. J., and Fleming, J. G.: An efficient early warning system for typhoon storm surge based on time-varying advisories by coupled ADCIRC and SWAN, *Ocean Dynamics*, 65, 617–646, <https://doi.org/10.1007/s10236-015-0820-3>, publisher: Springer Science and Business Media LLC, 2015.
- 675 Tallapragada, V., Kieu, C., Kwon, Y., Trahan, S., Liu, Q., Zhang, Z., and Kwon, I.-H.: Evaluation of Storm Structure from the Operational HWRP during 2012 Implementation, *Monthly Weather Review*, 142, 4308–4325, <https://doi.org/10.1175/MWR-D-13-00010.1>, <https://journals.ametsoc.org/mwr/article/142/11/4308/71708/Evaluation-of-Storm-Structure-from-the-Operational>, 2014.
- Tazkia, A. R., Krien, Y., Durand, F., Testut, L., Islam, A. S., Papa, F., and Bertin, X.: Seasonal modulation of M2 tide in the Northern Bay of Bengal, *Continental Shelf Research*, 137, 154–162, <https://doi.org/10.1016/j.csr.2016.12.008>, publisher: Elsevier BV, 2017.
- 680 Verlaan, M., Zijderveld, A., Vries, H. d., and Kroos, J.: Operational storm surge forecasting in the Netherlands: developments in the last decade, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 363, 1441–1453, <https://doi.org/10.1098/rsta.2005.1578>, publisher: The Royal Society, 2005.
- Warner, J. F., van Staveren, M. F., and van Tatenhove, J.: Cutting dikes, cutting ties? Reintroducing flood dynamics in coastal polders in Bangladesh and the Netherlands, *International Journal of Disaster Risk Reduction*, 32, 106–112, <https://doi.org/10.1016/j.ijdr.2018.03.020>, <https://linkinghub.elsevier.com/retrieve/pii/S2212420917303308>, 2018.
- 685 WW3DG, T. W. I. D. G.: User manual and system documentation of WAVEWATCH III TM, 2019.

- Yang, K., Paramygin, V. A., and Sheng, Y. P.: A Rapid Forecasting and Mapping System of Storm Surge and Coastal Flooding, *Weather and Forecasting*, 35, 1663–1681, <https://doi.org/10.1175/WAF-D-19-0150.1>, <https://journals.ametsoc.org/waf/article/35/4/1663/346026/A-Rapid-Forecasting-and-Mapping-System-of-Storm>, 2020.
- 690 Ye, F., Zhang, Y. J., Yu, H., Sun, W., Moghimi, S., Myers, E., Nunez, K., Zhang, R., Wang, H. V., Roland, A., Martins, K., Bertin, X., Du, J., and Liu, Z.: Simulating storm surge and compound flooding events with a creek-to-ocean model: Importance of baroclinic effects, *Ocean Modelling*, 145, 101–126, <https://doi.org/10.1016/j.ocemod.2019.101526>, <https://linkinghub.elsevier.com/retrieve/pii/S1463500319302173>, 2020.
- Zhang, Y. and Baptista, A. M.: SELFE: A semi-implicit Eulerian–Lagrangian finite-element model for cross-scale ocean circulation, *Ocean Modelling*, 21, 71–96, <https://doi.org/10.1016/j.ocemod.2007.11.005>, publisher: Elsevier BV, 2008.
- 695 Zhang, Y. J., Stanev, E. V., and Grashorn, S.: Unstructured-grid model for the North Sea and Baltic Sea: Validation against observations, *Ocean Modelling*, 97, 91–108, <https://doi.org/10.1016/j.ocemod.2015.11.009>, publisher: Elsevier BV, 2016a.
- Zhang, Y. J., Ye, F., Stanev, E. V., and Grashorn, S.: Seamless cross-scale modeling with SCHISM, *Ocean Modelling*, 102, 64–81, <https://doi.org/10.1016/j.ocemod.2016.05.002>, publisher: Elsevier BV, 2016b.
- 700 Zhang, Y. J., Ye, F., Yu, H., Sun, W., Moghimi, S., Myers, E., Nunez, K., Zhang, R., Wang, H., Roland, A., Du, J., and Liu, Z.: Simulating compound flooding events in a hurricane, *Ocean Dynamics*, 70, 621–640, <https://doi.org/10.1007/s10236-020-01351-x>, <http://link.springer.com/10.1007/s10236-020-01351-x>, 2020.