



A Homogeneous Earthquake Catalogue for Turkey and Surrounding Region

Onur Tan

5 İstanbul University – Cerrahpaşa, Faculty of Engineering, Dept. of Geophysics, İstanbul, Turkey

Correspondence to: Onur Tan (onur.tan@istanbul.edu.tr)

Abstract. A new earthquake catalogue for Turkey and surrounding region (32° - 41° N, 20° - 52° E) is compiled for the period 1900-2017. The earthquake parameters are obtained from the Bulletin of International Seismological Centre that is fully updated in 2020. New conversion equations between moment magnitude and the other scales (m_d , M_L , m_b , M_s and M) are determined using the General Orthogonal Regression method to build up a homogeneous catalogue, which is the essential data for seismic hazard studies. The 95% confidence intervals are estimated using the bootstrap method with 1000 samples. The equivalent moment magnitudes (M_w^*) for the entire catalogue are calculated using the magnitude relation to homogenise the catalogue. The magnitude of completeness is $2.9 M_w^*$ and 3.0 - $3.2 M_w^*$ for Turkey and Greece respectively. The final dataset is not declustered or truncated using a threshold magnitude because of motivational reasons, generating a widely usable catalogue. It contains not only M_w^* , but also the average and median of the observed magnitudes for each event. Contrary to the limited earthquake parameters in the previous catalogues, the 45 parameters of approximately 700k events occurred in a wide area from the Balkans to the Caucasus are presented.

1 Introduction

The earthquake catalogues are the first output of seismological observations. National and international catalogues are generated by several institutions around the world for understanding the seismic activity of a region. Principally, a catalogue contains the parameters such as origin time, coordinates and focal depth. Although the magnitude of an earthquake, which is a dimensionless scale of energy release, is one of the main seismological parameters, it has different scales (types) based on different seismic wave types and determining approximation (Table 1). A catalogue may not contain all magnitude scales for an event. If an earthquake catalogue is used for just showing seismicity on a map, the magnitude type may not be important because the differences among the values of scales are not too big for visualisation. However, magnitude scale information used in energy calculation is crucial for seismic hazard studies.

There are several unknowns for magnitude calculations of institutions due to used equations, seismic network structures, man-made mistakes etc. Both amplitude and distance constants in the magnitude equations are the major items. Although



30 they are specific for a region because of seismic wave attenuation in the crust and mantle, the constants calculated from the Californian earthquakes (i.e. for local magnitude by Richter, 1935; Hutton and Boore, 1987) are widely used. On the other hand, individual magnitudes are calculated at each station for an event; then they are averaged. The averaged magnitude is closely related with several factors: The number of stations, the standard deviation of the average, amplification or attenuation due to the geological structure beneath the station, the radiation pattern of the seismic waves related with the azimuthal distribution of stations. Therefore, institutions report different magnitudes for an event. Another issue picked out in this study is about moment magnitude (M_w) in catalogues. M_w is determined using waveform modelling for events ($M_w \geq 3.5-4.0$) that have a high signal-to-noise ratio. However, a few institutes report M_w for small events ($M_w < 3.0$, i.e. 25.01.1999 13:06 $M_w=1.8$ by Cyprus Geological Survey Department; 29.05.2014 01:14 $M_w=1.8$ by the Earthquake Research Center, Ataturk University). It is clear that these magnitudes are determined by using a relationship equation, but it cannot be proved this type of man-made faults. Consequently, there are more than one magnitude values for an event with known and unknown calculation errors, and only one scale for each event must be used in the studies based on the geometric data such as hazard mitigation analyses. At this point, essential of a homogenised catalogue with a common magnitude arises. In the last two decades, the studies on unifying earthquake magnitudes and generating improved catalogues are carried out for different parts of the earth (i.e. Grünthal et al., 2009; Chang et al., 2016; Manchuel et al., 2018; Rovida et al., 2020).

45

Table 1. Symbols for different magnitude scales in the priority order of magnitude saturation.

M_w	Moment magnitude
M_s	Surface wave magnitude
m_b	Body wave magnitude
M_L	Local (Richter) magnitude
m_d	Duration magnitude
M	General magnitude (unreported type)

This study focuses on the earthquakes that occurred in Turkey and surroundings. The region is one of the most geodynamically active areas on the earth and deformed among Eurasian, African and Arabian plates (Fig. 1). Both the continental collision between Arabia and Eurasia and subduction of the African Plate beneath Eurasia started in the Early/Middle Miocene (11-23 Ma). The interactions of the three plates are the major driving forces for the tectonics of the region. The plate motions result in thrust faulting in Eastern Anatolia, Caucasus and Iran, normal faulting in Western Turkey and Greece, and transform faults due to escaping to west and east (see Bozkurt, 2001 for a brief synthesis). The complex tectonic character of the region causes a high number of earthquakes with different faulting mechanism and a wide range of focal depths.




Figure 1. Simplified tectonic map of Turkey and its surroundings. DSFZ: Dardanelles Sea Fault Zone, NAFZ: North Anatolian Fault Zone. Triangles indicate direction of vergence or subduction, bars indicate down-thrown side of normal faults. White arrows indicate relative plate motions. The relief model is generated with the ETOPO1 model (Amante and Eakins, 2009).

The destructive earthquakes in Turkey and the surrounding countries along the centuries are found in the historical records. Pinar and Lahn (1952), Ergin et al. (1967, 1971), Soysal et al. (1981), Güçlü et al. (1986), Ambraseys and Finkel (1995), Ambraseys and Jackson (1998) compile the historical earthquakes in the region. Tan et al. (2008) present the historical events on a digital database and publish the first catalogue that contains the focal mechanism parameters of the earthquakes in Turkey. On the other hand, Leptokaropoulos et al. (2013) and Kadirioğlu et al. (2018) introduce homogenised catalogues for the Turkish earthquakes. The main component of homogenization is to obtain reliable magnitude conversion from one scale to moment magnitude. Several empirical relations are also proposed for the region (Papazachos et al., 1997; Ambraseys 2000; Baba et al., 2000; Kalkan et al., 2004; Ulusay et al., 2004; Özalp et al., 2010; Deniz and Yücemem, 2010).

The motivation of this study is to build a widely usable earthquake catalogue (i.e. for geophysicists, geologists, earthquake engineers) that contains homogenised moment magnitudes and the other seismological parameters. During the international seismic hazard studies of the Sinop Nuclear Power Plant that is planned to construct in northernmost of Turkey, it is clearly understood that a comprehensive homogenised earthquake catalogue for Turkey is needed for future studies. For this aim, all earthquakes occurred in a wide area are analysed with a statistical approach, and the empirical magnitude relation equations are obtained using a refined data set. Then, an extensive homogenised earthquake catalogue for Turkey and the surrounding



region is constructed. The distinguishing feature of the new homogenised catalogue is that it contains all earthquakes in a manageable format from Greece to Azerbaijan without removing aftershocks and truncating small events.

80 2. Database and processing

The Bulletin of the International Seismological Centre (ISC, 2020) is as the main database to generate a new and comprehensive homogeneous earthquake catalogue for Turkey and the surrounding region. The ISC Bulletin contains a large number of parametric data for an event that occurred anywhere on the earth. Because national and international seismological centres contribute to the bulletin, it contains not only moderate-to-large events ($M \geq 4$) but also local earthquakes with small magnitudes ($M < 4$). The most important feature of the bulletin is that an event with sufficient data is manually checked and relocated by a seismologist. Therefore, the latest event information in the database is two years behind in real-time (ISC, 2020). The bulletin also presents the event parameters reported by the contributor centres. The ISC finished rebuilding the entire database in 2020. The *ak135* seismic velocity model (Kennett et al., 1995) and a relocation procedure that is recently used by the ISC is implemented to all data. Furthermore, a large number of earthquake data from the permanent and temporary networks have been added (ISC, 2020; Storchac et al., 2017). Therefore, the latest and revised international dataset is used in this study.

The earthquake parameters in the bulletin are in the IASPEI Seismic Format (ISF, 2020). Each event has its own data block, such as origin and magnitude, contains several data types and comments. Data and comment lines have no specific flag to identify their types, and it is not possible to read the database using a simple computer program or shell-scripts. A Fortran code is written to analyse the ISF lines using the parsing subroutines provided by the ISC. Each line in the database is checked by the different parsers to identify its data type. After determining the origin and magnitude sub-blocks of an event properly, the parameters are analysed. The overall data processing is given in the flowchart in Fig. 1. In the first step, the origin data such as time, location and focal depth are searched for the "PRIME" comment that indicates the residuals is useful to prefer the hypocentre parameters. The hypocentres determined by the ISC are always prime. If there is no "PRIME" flag, the origin data is searched in the secondary hypocentres using a priority order for the institutes given in the flowchart. The parameters reported by the ISC are preferred first. If there is no information from the ISC, the availability of the origin parameters of the European-Mediterranean Seismological Centre (CSEM or EMSC) are tried (see Appendix A for the institute abbreviations). The priority of both institutes is high because they use all available data in the study area. In turn, ISK (Kandilli Observatory and Earthquake Research Institute, KOERI) and DDA (General Directorate of Disaster Affairs until September 2017; Disaster and Emergency Management Presidency - AFAD after October 2017), which are the national seismological networks in Turkey, are selected. The other institutes are used for the local events around Turkey. Besides, the earthquake information reported by ISS and GUTE is used for the pre-instrument period (1900-1964). If the origin



parameters of an event are found in any step of this query order, this event is added to the homogenised catalogue with these parameters.

After determining the origin parameters of an event in the selected area, the magnitude data sub-block analysed by the magnitude parser. The values of different magnitude scales given in Table 1 are collected. If there are two or more values for each type, average, standard deviation and median are calculated. Selecting a magnitude value from a particular institute is not preferred to overcome the problems such as unreported magnitude, the effect of network distribution calculation errors. On the other hand, we have no evidence for that an institute calculates true magnitude for an earthquake.




Figure 2 Flowchart of the ISC database processing. Ave: Average (mean), Med: Median, Sdev: Standard deviation. See Appendix A for the institute codes.



More than 7.7 million events in the ISC monthly data sets are analysed for the final catalogue. The interest area is bounded by $32^{\circ}\text{N} - 47^{\circ}\text{N}$ and $20^{\circ}\text{E} - 52^{\circ}\text{E}$ (Fig. 2). These limits cover an event that occurred 350 km away from Turkish borders to ensure the requirements of seismic hazard mitigation studies and the Turkish Regulations for nuclear power plant site studies. The study area also covers the Balkans, Black Sea, Caucasus, Syria, northern Iraq and northwest of Iran. The final catalogue contains 697,139 events occurred in the period from 1900 to the end of 2017. The instrumental period (after 1964) data is used for the statistical analyses.

The number of events (1964-2017) reported with local magnitude (M_L) is 443,939 (64% of the total) and it is the highest rate respect to the other magnitudes types (Fig. 3). About 35% of the events have duration magnitudes (m_d). Because both magnitudes types are widely determined by the national institutions, especially for local events, they are dominant in the catalogue. Body (m_b) and surface wave (M_s) magnitudes are reported for only 4.7% and 1.7% of the total events in the region, respectively. Though moment magnitude (M_w) is the most preferred magnitude scale for seismic hazard studies, only 0.8% of all events have M_w because waveform analyses are not easy and routine process. On the other hand, the catalogue contains 28,630 (4%) events with no specified magnitude types (M). This magnitude M is mostly reported until 1990, and the number of events with M dramatically decreases after this year. Besides, approximately 2% of the annual activity is reported without a magnitude (in total 41,440 events). However, the rate runs up to 6% only in 2010 and 2011 because ~5000 events without a magnitude are reported by the TIF (Georgia) for the Caucasus earthquakes. The earthquakes with no magnitude assigned are also included in the catalogue to be useful in future studies.

140




Figure 3. The catalogue area (solid border) and the earthquake in the catalogue occurred in the period 1900-2017 (dots). Yellow circles are the events with magnitude greater than 6.0.

145




Figure 4. The number of events in the final catalogue for each averaged magnitudes (1964-2017). N is the total number of event for each magnitude.



150 **Table 2.** Top: Number of magnitudes with zero and non-zero values between 1964 and 2017 in the catalogue. Bottom: The number of magnitude pairs with non-zero values.

	m_d	M_L	m_b	M_s	M_w	M
=0	452,933	251,361	662,466	683,514	690,017	666,720
≠0	242,317	443,887	32,784	11,736	5,233	28,630
	m_d, M_w	M_L, M_w	m_b, M_w	M_s, M_w	M, M_w	
≠0	2,764	4,598	3,747	3,093	312	

3. Catalogue homogenization and completeness

3.1 Refining the dataset

155 The dataset is refined in detail for regression analyses to obtain the empirical relations between the magnitudes. In the first step, the catalogue is declustered using Reasenber's (1985) second-order moment approximation because removing aftershocks is sary to determine reliable magnitude completeness. For aftershock ~~analy~~  in space, a subsequent event is searching in an area with a radius of 20 times of the circular source dimension of the preceding event considering ± 4 km hypocentre uncertainties (Kanamori and Anderson, 1975; Reasenber, 1985). The maximum interaction period for the

160 next event in a sequence is 10 days to build a temporal extension for a cluster. After declustering, the earthquakes rred after 1980 are selected because the national station networks and data analyses procedure become much more reliable in  Turkey. In the third step, completeness (M_c) for each magnitude is determined and it is found that M_c is about ~ 2.8 for m_d and M_L  for m_b and M_s . The earthquakes with averaged magnitudes ~~are~~ smaller than the M_c thresholds are excluded in the regressions. In the last step, a cut-off value is applied for high differences between magnitude pairs. There are, naturally,

165 differences among the reported magnitudes for an earthquake. Occasionally, the difference between the magnitude pairs may be as large as 2 or more magnitude units. After obtaining the distribution of the differences for each pair, the data points that are out of the 95% confidence interval ($\pm 2\sigma$) are removed using the Interquartile Range (IQR) method (Galton, 1869; MacAlister, 1879), which is one of the robust methods for outliers and can be successfully applied to seismological data (i.e. Tan et al., 2010, 2014; Tan, 2013). The cut-off values are determined as 0.85 (m_d-M_w), 0.72 (M_L-M_w), 0.68 (m_b-M_w), 1.17

170 (M_s-M_w), and 1.10 ($M-M_w$). These cut-offs overcome the scattering of the pairs. After refining the magnitude pairs in the four steps, the number of data used in regression is 2100, 3098, 1691, 881 and 228 for m_d-M_w , M_L-M_w , m_b-M_w , M_s-M_w and $M-M_w$, respectively (Fig. 4).



3.2 Regression Analyses

The relationships of the refined magnitude pairs are estimated using the general orthogonal regression (GOR). The method is
175 better estimator than the least-square (LS) approximation when both x and y variables have errors of non-negligible size
(Castellaro et al., 2006). The slope (a) and intercept (b) value of the GOR line in the form of $y = ax + b$ is given by

$$a = \frac{S_Y^2 - \eta S_X^2 + \sqrt{(S_Y^2 - \eta S_X^2)^2 + 4\eta S_{XY}^2}}{2S_{XY}} \quad (1)$$

$$b = \bar{Y} - a\bar{X} \quad (2)$$

180

where S_X^2 , S_Y^2 and S_{XY}^2 are the variance of X (independent variable), Y (dependent variable) and between X and Y ,
respectively (i.e. Castellaro et al., 2006; Das et al., 2014). \bar{X} and \bar{Y} are the average values of the variables. η is the error
variance ratio of the variables (σ_{ε_X} , σ_{ε_Y}) and defined as $\eta = (\sigma_{\varepsilon_X} / \sigma_{\varepsilon_Y})^2$. When the standard errors of the variables are not
known, η is arbitrary set to a value. In practice, $\eta = 1$ (squared Euclidean distance) gives good results (Castellaro et al.,
185 2006; Das et al., 2014). In this study, η is tested for the values from 0.5 to 2.0 to seek a better fit. The R^2 values do not
increase when η is assigned different than 1.0 and a significant improvement is not observed in the regressions. Besides, the
real errors of the magnitudes are not known; $\eta = 1$ is used. The squared Euclidean distance gives better results for all
magnitude scales. The 95% confidence intervals of the best-fit lines are determined with the bootstrap method (Efron, 1979).
Total 1,000 new regressions are calculated using 50% of the total number of data of each relation. The bootstrap samples are
190 randomly selected using the Mersenne Twister random number generator (Matsumoto and Nishimura, 1998); and the
random numbers are unique in each test to prevent multiple selections of any datum. After obtaining a large set of the
constants a and b of the linear fits, the outliers are removed with the IQR method. Then, the standard deviation (σ) of the
normally distributed dataset is calculated.

195 The GOR results are given in Table 3 and Fig. 4. Because the number of magnitude pairs is high for each relation, the data is
shown with coloured density contours in 0.1 magnitude-unit grids. It is clear that all relations are linear, and minimum misfit
regression lines are in good agreement with the data distribution. The number of pairs is generally dense between the
magnitude 0 and 5.0 and decrease for larger magnitudes. In general, the slopes of the regression lines are close to 1; and
the intercept values are negative for M_s magnitude. The relation between m_d and M_w indicates, both scale is equal at
200 $m_d = 4$ and the difference increases up to 0.4 magnitude unit at larger values. M_L values are dense between 3 and 5; and the
linear fitting line extends close to the $y=x$ line. The difference between local and moment magnitudes is about 0.25 at $M_L =$
7.0. The conversion of m_d - M_w is similar to that of M_L - M_w . The largest difference between different magnitude scales is
observed for surface and moment magnitudes. M_s is always smaller than M_w and the difference is about 0.6 at $M_s = 4.0$. Both
scales are equal at $M_s = 7.5$. The magnitude M (the real type is not known) is mostly reported in the past. There are 95 events



205 only with $M \geq 5.0$ before 1964 in the study area. Therefore, an M - M_w conversion is necessary for seismic hazard analyses using long-term seismicity data. There are few magnitude pairs ($N = 228$) and they are distributed sparsely between 3 and 7 with high standard deviation (Fig. 4).

Table 3. Equivalent moment magnitude (M_w^*) relations for different magnitude scales.

Relation	$a \pm 2\sigma$	$b \pm 2\sigma$	Number of Data	Magnitude Range	R^2
$M_w^* = a \cdot m_d + b$	$+1.125 \pm 0.025$	-0.507 ± 0.102	2,100	2.8 – 7.3	0.80
$M_w^* = a \cdot M_L + b$	$+1.053 \pm 0.015$	-0.105 ± 0.059	3,098	2.8 – 7.2	0.87
$M_w^* = a \cdot m_b + b$	$+1.042 \pm 0.015$	-0.118 ± 0.072	1,691	4.0 – 7.0	0.86
$M_w^* = a \cdot M_s + b$	$+0.838 \pm 0.035$	$+1.213 \pm 0.164$	881	4.0 – 7.7	0.80
$M_w^* = a \cdot M + b$	$+1.057 \pm 0.108$	-0.199 ± 0.551	228	3.4 – 6.9	0.69

210

3.3 Homogenization

The GOR results are implemented to all events in the study area. First M_w is searched and assigned as M_w^* if found. For the events without M_w , the first averaged magnitude with non-zero value is chosen according to the priority of saturation order in Table 1. The chosen magnitude is named M_x and used to calculate the equivalent moment magnitude (M_w^*) with the relevant equation. After applying homogenization equations to all earthquakes, the catalogue is presented with a total of 45 parameters described in Appendix B. The catalogue has three sections: Event Origin Section, Magnitude Section and Comments. There are 23 parameters in the origin section. The time, coordinates and depths with their uncertainties are given. If one of these parameters is fixed, it is marked with the "f" flag. The magnitude section contains the average with standard deviation and median of the six magnitude scales. The selected M_x value, its source magnitude scale and calculated equivalent moment magnitude (M_w^*) is presented. The ISC event ID number and the epicentre region are given in the comment section as the reference.

215
220

In the homogenised catalogue, 55% of the origin parameters are flagged as "PRIME" by the ISC. The ISC and EMSC (CSEM) origin parameters are generally reported with the prime flag (~90-98%). On the other hand, approximately half of the reported parameters (~40-45%) by the national institutes in Turkey (KOERI, AFAD/DAD) and Greece (ATH) have the flag. The catalogue contains the origin information from the national sources (Fig. 5a) in a high number of percentages. The distribution of the magnitude scales for the equivalent magnitude calculation is given in Fig. 5b. The vast majority M_w^* are obtained from M_L and m_d ; the contribution of the other magnitude scales is small.

225
230




Figure 5. Magnitude relations between M_w and the other scales. The data density in 0.1 magnitude intervals is shown with coloured counters. N is the total number of magnitude pairs. The solid line is the best linear fit of the orthogonal regression, whereas the dashed lines shows the 95% confidence interval after bootstrapping. Gray line indicates 1:1 relation.




Figure 6. Distribution of the parameters in the homogenised catalogue. (a) Source institutes for the origin parameters (b) Magnitude scales used for M_w^* calculation.

240


3.4 Completeness of the Catalogue

One of the important parameters of an earthquake catalogue is the magnitude of completeness (M_c). M_c is a threshold magnitude and indicates that earthquakes with magnitudes greater than M_c are recorded in a study area. It is determined using Gutenberg-Richter's (1954) cumulative frequency-magnitude law (GR). The GR relation is simple but powerful and formulated as $\log(N) = a - b \cdot m$, where N is the cumulative number of events with magnitudes equal to or greater than m . The other useful parameter derived from this equation is the b -value (slope). The b -value is around 1 for the tectonically active areas.

245

The instrumental period (since 1964) observation for the region shows a linear relation with $b = 0.91$ between the cumulative number of earthquakes and equivalent moment magnitude, M_w^* ; (Fig. 6). If the dataset is extended to cover the instrumental period (1900-1964), the linearity for the magnitudes between 5 and 7 due to the magnitude calculation uncertainties of the earthquakes in that time span. The M_c , the lowest intercept point of the linear fit with slope b , is 2.9 for the over all catalogue.

250



255

Figure 7. Gutenberg-Richter frequency-magnitude distribution for the homogenised catalogue. White circles represents the earthquakes occurred between 1900 and 2017, grey circles represent the earthquakes later than 1964 with the b-value of 0.91. The pre-instrumental period earthquakes cause a bend for the events with $\sim 5-7 M_w^*$. The catalogue completeness is 2.9.

260




Figure 8. Top: M_c spatial distribution map for the events after 1964. There is no data in white areas. Bottom: Temporal M_c variation as a function of year. Grey area is the $\pm\sigma$ interval estimated by bootstrapping.

265

The maximum curvature method (Wyss et al., 1999; Wiemer, 2001) is applied to investigate the spatial and temporal change of M_c for the instrumental period. Equal horizontal sampling in latitude and longitude is not used to prevent artificial elongation because the length of 1° of longitude is ~ 94 and ~ 76 km in south and north of the study area, respectively. I use 20 km grid spacing and at least 100 events larger than the completeness magnitude ($M_w^* > 2.9$) in 100 km radius for the spatial distribution of M_c . On the other hand, the temporal variation is estimated using a window with 500 events and a step of 25 events. These sampling parameters are sufficient to avoid erroneous statistical results for the b -value and M_c due to under-sampling and non-homogenous subsets (Amorese et al., 2010; Kagan, 1999, 2002, 2010; Kamer and Heimer, 2013; Shi and Bolt, 1982). The contour map  in Fig. 8 shows that the homogenised catalogue is complete down to M_w^* 3.0-3.2 in Turkey and 3.2-3.3 in Greece. M_c increases dramatically up to 4.0-4.5 in the Caucasus and its abrupt transition follows



275 the eastern border of Turkey. The regional difference reflects the observation power of the seismological networks. The
white area have very low seismic activity (see Fig. 2) and there is not sufficient data to ensure the criteria. The variation of
Mc along years indicates that the standard deviation band is narrow after 1990 and Mc is stable at about 3.0 late 2000
because the number of seismological stations increase after the devastating earthquakes ($M_w > 7$) in 1999.

4 Discussions

280 Generating an earthquake catalogue is the main issue for seismologist. An institution that operates a costly seismological
network provides the main metric information of an event from raw waveform observations. The parametric catalogues
are released in paper print before the internet and are online anymore. Although accessing catalogues is very easy via the
internet, it is difficult to obtain all available data due to some limitations of the data providers' web pages. The problems of
285 online datasets, such as absence or limited observation for the past years, a limited number of parameters, lack of parameter
uncertainties, listing limitations, useless formats in web pages etc., make difficult to use the earthquake data for a large range
of users. However, most of the researchers pay only attention to the homogenised magnitudes and the number of events.
Unfortunately, the importance of a large number of parameters and their uncertainties in a catalogue are missed, and the
given datasets less useful for the studies other than seismic hazard analyses.

290 The earthquake information for Turkey comes from two national networks operated by the KOERI and AFAD. Both
institutes have a large number of stations around Turkey and report recent events online. The date, time, depth and
magnitudes without uncertainties of events are given by the search engines of both institutions. While the KOERI lists only
50k events in a single search with a downloadable text file, the AFAD search result is given with maximum 100 events at
each window and can be downloaded in the comma-separated CSV format. Another online catalogue with the same
295 parameters is provided by the EMSC. The searched events can be downloaded in the CSV format with the limitations of 5k
lines. Among the three institutions, only the KOERI provides all available magnitude scales for an event. Additionally, the
EMSC does not provide the type of magnitude scale for an event. On the contrary, the ISC provides all available parameters
for an event determined not only by itself but also by the other institutions as mentioned in the previous chapter. The
magnitudes in the ISC event list are given in separate lines, so it is not easy to use without knowledge of the comprehensive
300 bulletin format and programming. The online bulletin search of the ISC has also output limitation with 60k events.

Besides the online catalogues, some catalogue compilations based on homogenization of magnitudes for Turkey and its
vicinity are published. Leptokaropoulos et al. (2013) statistically analyse the earthquakes in detail occurred in Western
Turkey between (1964-2010) and constructs a catalogue with an equivalent moment magnitude. They obtain conversion
305 equations for different magnitude scales reported by different institutions. The catalogue contains 9875 events with only
parameters of date, time, coordinates and focal depth. Kadirioglu et al. (2018) present a homogenised catalogue for Turkey



containing ~6573 events between 1900 and 2012. They use the same dataset and conversion equations in their previous study (Kadirioğlu et al., 2014; Kadirioğlu and Kartal, 2016). Their final catalogue is declustered and contains events only $M_w^* > 4.0$ (not observed M_w as given in the catalogue, notation mistyping). On the other hand, Kadirioğlu et al. (2014, 2018) mention that a 10 km of focal depth assigned to the events without reported depth or shallower than 1 km in the final catalogue. This is an arbitrary and unrecoverable parameter assignment and may generate artificial errors in future studies using this catalogue, especially in seismic hazard analyses.

Burton et al. (2004) generate a homogenised catalogue that contains both reported and equivalent magnitudes for earthquakes in Greece and Western Turkey using the previous conversion equations. There are ~5200 events without M_c analysis. The catalogue by Bayliss and Burton (2007) contains ~3680 homogenised events in Bulgaria and the surrounding Balkan region. It is complete down to 4.0 M_w . More recently, Makropoulos et al. (2012) present homogenised event list for calculated M_s^* and $M_w^* \sim 7350$ (not observed M_s, M_w as given in the catalogue, notation mistyping) in the excel format for Greece and western Turkey.

The common structure of the previous catalogues mentioned above and others has limited earthquake parameter as date, location, depth and M_w^* . Especially, the observed magnitudes and error/uncertainty values are not included. The source institute of the parameters is also missing. Therefore, it is impossible to trace back to the origin of the parameters, and the equivalent moment magnitude (M_w^*) cannot be recalculated using newly determined conversion equations. On the other hand, a truncated final earthquake list using a magnitude threshold not useful for the researchers who not familiar details of earthquake catalogues and want to analyse or map whole instrumental period seismic activity in a region. The homogenised catalogue overcomes the common deficiency of the previous earthquake catalogues for Turkey and surroundings.

5 Conclusions

Turkey and the surrounding area is one of the most seismically active regions on the earth. Therefore, improved earthquake catalogue studies are necessary. A new, extended and homogenised earthquake catalogue is compiled in this study. The main aim is to present an earthquake database in an easily manageable ASCII format for a broad range of researchers. The study is based on the latest ISC Bulletin the rebuilding process was finished in 2020. All parameters of the earthquakes during the period from 1900 to 2017 in an extended region from the Balkans to the Caucasus are analysed. The origin parameters and magnitude data in the IASPEI Seismic Format are systematically parsed with a Fortran algorithm.

Approximately 700k events in the study area bounded by $32^\circ - 47^\circ \text{N}$ and $20^\circ - 52^\circ \text{E}$ are compiled (Fig. 3). The equivalent moment magnitude (M_w^*), which is the mandatory parameter for the seismic hazard studies, is calculated for all events. For



For these purposes, new conversion equations for m_d , M_L , m_b , M_s and M are determined using the well-refined magnitude pairs
340 ~~using~~ in the general orthogonal regression method that is useful when the two variables have different uncertainties.
According to the values of M_w^* , the overall catalogue is complete down to $M_c = 2.9$. The spatial completeness variation
indicates $M_c = \sim 3.0-3.2$ in Turkey and Greece, ~~and as high as 4.5 in the Caucasus~~. The catalogue is not declustered or
truncated using a threshold magnitude to be useful for geophysicist, geologist and geodesist. The M_w^* values can be easily
345 recalculated and the catalogue can be declustered using different parameters by seismologists and earthquake engineers for
seismic hazard studies. The final dataset contains not only M_w^* as in the previous studies but also the average with standard
deviation and median of the observed magnitudes. The ISC event ID-number and geographic region of each event are also
given to trace an event in the bulletin. Total of 45 parameters is presented.

350



Appendix A

The ISC contributor institutes mentioned in this study is given below. The ISS and GUTE catalogues are used for pre-instrumental period events.

Code	Institute
ISC	International Seismological Centre
ISS	International Seismological Summary [for 1900 - 1964]
GUTE	Gutenberg and Richter (1954) [for 1900 - 1952]
CSEM	European-Mediterranean Seismological Centre - EMSC (France)
ISK	B.U. Kandilli Observatory and Earthquake Research Institute (Turkey)
DDA	General Directorate of Disaster Affair (Turkey), until Sep.2017
AFAD	Disaster and Emergency Management Presidency (Turkey), since Oct. 2017
ATH	National Observatory of Athens (Greece)
THE	Dept. of Geophysics, Aristotle University of Thessaloniki (Greece)
MOS	Geophysical Survey of Russian Academy of Sciences (Russia)
TEH	Tehran University (Iran)
TAB	Tabriz Seismological Observatory (Iran)
TIF	Institute of Earth Sciences/ National Seismic Monitoring Center (Georgia)
SOF	National Institute of Geophysics, Geology and Geography (Bulgaria)
BUC	National Institute for Earth Physics (Romania)
SIGU	Subbotin Institute of Geophysics, National Academy of Sciences (Ukraine)
IPER	Institute of Physics of the Earth, Academy of Sciences, Moscow (Russia)
NSSP	National Survey of Seismic Protection (Armenia)
AZER	Republican Seismic Survey Center of Azerbaijan National Academy of Sciences (Azerbaijan)
NSSC	National Syrian Seismological Center (Syria)
THR	International Institute of Earthquake Engineering and Seismology (Iran)



Appendix B

The first and second lines of the homogenised catalogue are the parameter names and column numbers, respectively. The earthquake parameters are given below.

	Column	Parameter	Column	Parameter	
Event Origin Section	1	Year	24	M (average)	
	2	Month	25	Std.Dev. of M	
	3	Day	26	M (median)	
	4	Hour	27	m_d (average)	
	5	Minute	28	Std.Dev. of m_d	
	6	Second	29	m_d (median)	
	7	Time Fix Flag	30	M_L (average)	
	8	RMS (s)	31	Std.Dev. of M_L	
	9	Latitude (°)	32	M_L (median)	
	10	Longitude (°)	33	m_b (average)	
	11	Location Fix Flag	34	Std.Dev. of m_b	
	12	Semi-major Axis of 90% ellipse (km)	35	m_b (median)	
	13	Semi-minor axis of 90% ellipse (km)	36	M_s (average)	
	14	Depth (km)	37	Std.Dev. of M_s	
	15	Depth Fix Flag	38	M_s (median)	
	16	Depth Error (km)	39	M_w (average)	
	17	Number of Stations	40	Std.Dev. of M_w	
	18	Azimuthal Gap (°)	41	M_w (median)	
	19	Closest Station	42	M_x	
Magnitude Section	20	Furthest Station	43	Source magnitude for M_x	
	21	Event Type	44	M_w^*	
	22	Institute	Comments	45	# (null)
	23	Prime Flag		46	ISC information (event ID and region)

- 360 Fixing Flags: n: Not fixed (free), f: Fixed
 Prime Flags: n: Not prime location, p: Prime location
 Event Types: de: Damaging earthquake, fe: Felt earthquake,
 ke: Known earthquake, se: Suspected earthquake,
 uk: Unknown
- 365 Unreported numerical parameters in the ISC Bulletin are given as "0.00".
 Uncalculated standard deviations are given as "-1.00".
 Unknown or blank character fields are filled with "-".



Data availability

370 The catalogue is available as the electronic material of this article.

Acknowledgments

I would like to thank Dr. Özlem KARAGÖZ TAN for her support and encouragement for this study. All maps and graphs are plotted using the Generic Mapping Tools (GMT) by Wessel and Smith (1991).

375



References

- Akkar, S., Çagnan, Z., Yenier, E., Erdogan, Ö., Sandikkaya, M. A., and Gülkan, P.: The recently compiled Turkish strong motion database: Preliminary investigation for seismological parameters, *J. Seismol.*, 14, 457–479, 2010.
- Amante, C. and Eakins, B. W.: ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis. NOAA Technical Memorandum NESDIS NGDC-24. National Geophysical Data Center, NOAA, doi: 10.7289/V5C8276M, 2009.
- Ambraseys, N. N. and Finkel, C. F.: The Seismicity of Turkey and Adjacent Areas: A Historical Review, 1500–1800. Muhittin Salih Eren, İstanbul, Turkey, 1995.
- Ambraseys, N. N. and Jackson, J. A.: Faulting associated with historical and recent earthquakes in the Eastern Mediterranean region, *Geophys. J. Int.*, 205(2), 133, 390–406, 1998.
- Ambraseys, N. N.: Reappraisal of north-Indian earthquakes at the turn of the 20th century, *Curr. Sci.*, 79, 1237–1250, 2000.
- Amorese, D., Grasso, J. R., and Rydelek, P. A.: On varying b -values with depth: results from computer-intensive tests for Southern California. *Geophys. J. Int.*, 180, 347–360, 2010.
- Baba, A. B., Papadimitiou, E. E., Papazachos, B. C., Papaioannou, C. A., and Karakostas, B. G.: Unified local magnitude scale for earthquakes of south Balkan area, *Pure Appl. Geophys.*, 157, 765 – 783, 2000.
- Bayliss, T. J. and Burton, P. W.: A new earthquake catalogue for Bulgaria and the conterminous Balkan high hazard region, *Nat. Hazards Earth Syst. Sci.*, 7, 345–359, <https://doi.org/10.5194/nhess-7-345-2007>, 2007.
- Bozkurt, E.: Neotectonics of Turkey: A synthesis, *Geodinamica Acta*, 14, 3–30, doi: 10.1016/S0985-3111 (01) 01066-X, 2001.
- Burton, P. W., Xu, Y., Qin, C., Tselentis, G., and Sokos, E.: A catalogue of seismicity in Greece and the adjacent areas for the twentieth century, *Tectonophysics*, 390, 117–127, 2004.
- Castellaro, S., Mulargia, F., and Kagan, Y. Y.: Regression problems for magnitudes, *Geophys J Int.*, 165, 913-930, doi: 10.1111/j.1365-246X.2006.02955.x, 2006.
- Chang, W., Chen, K., and Tsai, Y.B.: An updated and refined catalog of earthquakes in Taiwan (1900–2014) with homogenized M_w magnitudes, *Earth Planet Sp*, 68, 45, doi: 10.1186/s40623-016-0414-4, 2016.
- Das, R., Wason, H. R., and Sharma, M. L.: Unbiased Estimation of Moment Magnitude from Body- and Surface-Wave Magnitudes, *Bull. Seismol. Soc. Am.*, 104 (4), 1802–1811, doi: 10.1785/0120130324, 201.
- Deniz, A., and Yücemem, M. S.: Magnitude conversion problem for the Turkish earthquake data, *Nat. Hazards*, 55, 333–352, 2010.
- Efron, B.: Bootstrap methods: Another look at the jackknife, *The Annals of Statistics*, 7 (1), 1–26. doi: 10.1214/aos/1176344552, 1979
- Ergin, K., Güçlü, U., and Aksay, G.: A Catalog of Earthquakes of Turkey and Surrounding Area (1965–1970). Technical Report, İstanbul Technical University, Faculty of Mines, Institute of Physics of the Earth, no. 28, 1971.
- Ergin, K., Güçlü, U., and Uz, Z.: A Catalog of Earthquakes for Turkey and Surrounding Area (11 A.D. to 1964 A.D.). Technical Report, İstanbul Technical University, Faculty of Mines, Institute of Physics of the Earth, no. 24, 1967
- Galton, F.: Hereditary Genius, Macmillan Publishers, London, 1896.
- Grünthal, G., Wahlström, R., and Stromeyer, D.: The unified catalogue of earthquakes in central, northern, and northwestern Europe (CENEC) updated and expanded to the last millennium. *Journal of Seismology*, 13, 4, 517-541, doi: 10.1007/s10950-008-9144-9, 2009



- 415 Güçlü, U., Altınbaş, G., and Eyidoğan, H.: A Catalog of Earthquakes of Turkey and Surrounding Area (1971–1975). Technical Report, İstanbul Technical University, Faculty of Mines, no. 30, 1986
- Gutenberg, B. and Richter, C.: *Seismicity of the Earth and Associated Phenomena*, Princeton Univ. Press, Princeton, NJ, 310 pp, 1954.
- Hutton, L. K. and Boore, D. M.: The M_L scale in Southern California., *Bull. Seismol. Soc. Am.*, 77 (6), 2074–2094, 1987.
- 420 International Seismological Centre (ISC): On-line Bulletin, doi: 10.31905/D808B830, 2020.
- IASPEI Seismic Format (ISF): <http://www.isc.ac.uk/standards/isf/>, Last visit: September 2020., 2020
- Kadirioğlu, F. T. and Kartal, R. F.: The new empirical magnitude conversion relations using an improved earthquake catalogue for Turkey and its near vicinity (1900–2012), *Turkish J Earth Sci* 25(4), 300–310, 2016.
- 425 Kadirioğlu, F. T., Kartal, R. F., Kılıç, T. et al.: An improved earthquake catalogue ($M \geq 4.0$) for Turkey and its near vicinity (1900–2012), *Bull Earthquake Eng*, 16, 3317 – 3338, doi: 10.1007/s10518-016-0064-8, 2018.
- Kadirioğlu, F. T., Kartal, R. F., Kılıç, T., Kalafat, D., Duman, T. Y., Özalp, S., and Emre, Ö.: An improved earthquake catalogue ($M \geq 4.0$) for Turkey and near surrounding (1900–2012), 2nd European Conference on Earthquake Engineering and Seismology Proceedings, 411–422, 2014.
- Kagan, Y. Y.: Universality of the seismic moment-frequency relation, *Pure Appl. Geophys.*, 155, 537–573, 1999.
- 430 Kagan, Y. Y.: Seismic moment distribution revisited: I. Statistical results, *Geophys. J. Int.*, 148, 520–541, 2002.
- Kagan, Y. Y.: Earthquake size distribution: power-law with exponent?, *Tectonophysics*, 490, 103–114, 2010.
- Kamer, Y. and Hiemer, S.: Comment on “Analysis of the b-values before and after the 23 October 2011 Mw 7.2 Van–Erciş, Turkey, earthquake”, *Tectonophysics*, 608, 1448–1451, doi: 10.1016/j.tecto.2013.07.040, 2013.
- Kanamori, H., and Anderson D. L.: Theoretical basis of some empirical relations in seismology, *Bull. Seismol. Soc. Am.*, 65, 1073–1095, 1975.
- 435 Kennett, B. L. N., Engdahl, E. R., and Buland, R.: Constraints on seismic velocities in the Earth from travel times, *Geophys J Int*, 122, 108–124, 1995.
- Leptokaropoulos, K. M., Karakostas, V. G., Papadimitriou, E. E., Adamaki, A. K., Tan, O., and Inan S.: A Homogeneous Earthquake Catalog for Western Turkey and Magnitude of Completeness Determination, *Bull. seism. Soc. Am.*, 103–5, 2739–2751, 2013.
- 440 MacAlister, D.: *The Law of the Geometric Mean*, Proc. R. Soc., vol. XXIX, Harrison and Sons, London, 1879.
- Makropoulos, K., Kaviris, G., and Kouskouna, V.: An updated and extended earthquake catalogue for Greece and adjacent areas since 1900, *Nat. Hazards Earth Syst. Sci.*, 12, 1425–1430, doi: 10.5194/nhess-12-1425-2012, 2012.
- Manchuel, K., Traversa, P., Baumont, D. et al.: The French seismic CATalogue (FCAT-17). *Bull Earthquake Eng*, 16, 2227–2251, doi: 10.1007/s10518-017-0236-1, 2018.
- 445 Matsumoto, M., and Nishimura, T.: Mersenne twister: a 623-dimensionally equidistributed uniform pseudo-random number generator, *ACM Transactions on Modeling and Computer Simulation*, 8 (1), 3–30, doi: 10.1145/272991.272995, 1998.
- Papazachos, B.C., Kiratzi, A. A., and Karakostas, B. G.: Toward a homogeneous moment-magnitude determination for earthquakes in Greece and surrounding area, *Bull. Seismol. Soc. Am.*, 87, 474–483, 1997.
- 450 Pınar, N. and Lahn, E.: *Turkish Earthquake Catalog with Discriptions*, Technical Report, Turkey The Ministry of Public Works and Settlement, The General Directorate of Construction Affairs, Serial 6, no. 36, 1952.
- Reasenberg, P.: Second order moment of central California seismicity, 1969–1982., *J Geophys. Res. Solid Earth*, 90 (B7), pp.5479–5495, 1985.



- Richter, C.F.: An instrumental earthquake magnitude scale, *Bull. Seismol. Soc. Am.*, 25, 1-31, 1935.
- 455 Rovida, A., Locati, M., Camassi, R. et al.: The Italian earthquake catalogue CPTI1.5, *Bull Earthquake Eng.*, 18, 2953-2984. <https://doi.org/10.1007/s10518-020-00818-y>, 2020.
- Shi, Y. and Bolt, B.: The standard error of the magnitude–frequency b value, *Bull. Seismol. Soc. Am.*, 72, 1677–1687, 1982.
- Soysal, H., Siphioğlu, S., Kolçak, D., and Altınok, Y.: Historical Earthquake Catalogue of Turkey and Surrounding Area (2100 B.C. – 1900 A.D.). Technical Report, TÜBİTAK, No. TBAG-341, 1981.
- 460 Storchak, D.A., Harris, J., Brown, L., Lieser, K., Shumba, B., Verney, R., Di Giacomo, D., and Korger, E. I. M.: Rebuild of the Bulletin of the International Seismological Centre (ISC), part 1: 1964–1979, *Geosci. Lett.*, 4 (32). doi: 10.1186/s40562-017-0098-z, 2017.
- Tan, O.: The Dense Micro-Earthquake Activity at the Boundary Between the Anatolian and South Aegean Microplates, *J Geodyn.*, doi: 10.1016/j.jog.2012.05.005, 2013.
- 465 Tan, O., Papadimitriou, E. E., Pabuçcu, Z., Karakostas, V. G., Yörük, A., and Leptokaropoulos, K.M.: A detailed analysis of microseismicity in Samos and Kusadasi (eastern Aegean Sea) areas, *Acta Geophysica*, 62 (6), 1283-1309. doi: 10.2478/s11600-013-0194-1, 2014.
- Tan, O., Tapırdamaz, M. C., and Yörük, A.: The earthquake catalogues for Turkey, *Turkish J. Earth Sci.*, 17, 405–418. 2008
- 470 Tan, O., Tapırdamaz, M. C., Ergintav, S., İnan, S., et al.: Bala (Ankara) Earthquakes: Implications for Shallow Crustal Deformation in Central Anatolian Section of the Anatolian Platelet (Turkey), *Turkish J. Earth Sci.*, 19, doi: 10.3906/yer-0907-1, 2010.
- Ulusay, R., Tuncay, E., Sonmez, H., and Cokceoglu, C.: An attenuation relationship based on Turkish strong motion data and iso-acceleration map of Turkey, *Eng. Geol.*, 74, 265–291, 2004.
- Wessel, P. and Smith, W. H. F.: New, improved version of the Generic Mapping Tools released, *EOS* 79, 579, 1998.
- 475 Wiemer, S.: A software package to analyze seismicity: ZMAP, *Seism. Res. Lett.*, 72(3), 373-382, doi: 10.1785/gssrl.72.3.373, 2001.
- Wyss, M., Hasegawa, A., Wiemer, S., and Umino, N.: Quantitative mapping of precursory seismic quiescence before the 1989, M 7.1 off-Sanriku earthquake, Japan. *Annal. Geofis.*, 42, 851–869, 1999.