Atmospheric triggering conditions and climatic disposition of landslides in Kyrgyzstan and Tajikistan at the beginning of the 21st century

Xun Wang¹, Marco Otto¹, and Dieter Scherer¹

¹Chair of Climatology, Technische Universität Berlin, Berlin, 12165, Germany **Correspondence:** Xun Wang (xun.wang@tu-berlin.de)

Abstract. Landslide is a major natural hazard in Kyrgyzstan and Tajikistan. Knowledge about atmospheric triggering conditions and climatic disposition of landslides in Kyrgyzstan and Tajikistan is limited, even though this topic has already been investigated thoroughly in other parts of the world. In this study, the newly developed, high-resolution High Asia Refined Analysis version 2 (HAR v2) data set generated by dynamical downscaling was combined with historical landslide inventories to an-

- 5 alyze the atmospheric conditions that initialized landslides in Kyrgyzstan and Tajikistan. The results indicate the crucial role of snowmelt in landslide triggering processes since it contributes to the initialization of 40% of landslide events. Objective thresholds for rainfall, snowmelt, as well as the sum of rainfall and snowmelt (rainfall+snowmelt) were defined. Peak intensity(*I_{max}*) and accumulated amount (*Q*). Thresholds defined by rainfall+snowmelt have the best predictive performance. Mean intensity, peak intensity, and the accumulated amount of rainfall+snowmelt events yield the best predictive performance show similar.
- 10 predictive performance. Using the entire period of rainfall+snowmelt events results in better predictive performance than just considering the period up-to landslide occurrence. Mean annual exceedance maps were derived from regional thresholds of $I_{max} = 12.8 \text{ mm d}^{-1}$ and Q = 17.2 mm defined regional thresholds for rainfall+snowmelt. Mean annual exceedance maps depict climatic disposition and have added value in landslide susceptibility mapping. The results reported in this study highlight the potential of dynamical downscaling products generated by regional climate models in landslide prediction.

15 1 Introduction

Landslide is one of the most severe natural hazards in Kyrgyzstan and Tajikistan. More than 300 big landslides occurred in Kyrgyzstan from 1993 to 2010, causing 256 fatalities and direct economic losses of 2.5 million USD per year (Torgoev et al., 2012). Under global warming, wildfires, glacial retreat, and permafrost degradation are much likely to enhance slope instabilities in mountainous areas (Froude and Petley, 2018; Palmer, 2020), making these regions, including Kyrgyzstan and Tajikistan,

20 more vulnerable to climate change. Landslides are predetermined by static factors The occurrence of landslides depends on disposition and triggering events. Disposition refers to the general settings that make slopes prone to failure without actually initiating it, such as slope gradient and aspect, geology, vegetation cover, climate, etc. (Dai et al., 2002). These factors that make slopes prone to failure without actually initiating it are also referred to as dispositions (Zimmermann, 1997). Landslides

are triggered by dynamic factors, which are mainly Common triggers for landslides are extreme and prolonged rainfall, rapid

snowmelt, as well as earthquakes (Wieczorek, 1996).

The majority of landslide research in Kyrgyzstan and Tajikistan focused on characterizing landslide susceptibility, i.e., "where" landslides are prone to occur (e.g., Braun et al., 2015; Saponaro et al., 2015; Havenith et al., 2015b), and how to improve the landslide susceptibility models (Ozturk et al., 2020; Barbosa et al., 2021). But little attention is paid to the atmospheric triggering conditions, and our knowledge of "when" landslides are likely to occur is limited in this region. In addition,

- 30 most landslide susceptibility studies only took non-climatic factors into account or simply applied annual precipitation as a climatic factor. According to Segoni et al. (2018), no rainfall threshold for landslide triggering has been defined for Kyrgyzstan and Tajikistan yet, even though this topic has already been thoroughly investigated in other parts of the world with high landslide susceptibility (e.g., Berti et al., 2012; Gariano et al., 2015; Giannecchini et al., 2016; Leonarduzzi et al., 2017). The reasons are twofold. Firstly, although landslide inventories have been developed in this region, e.g., the Tien Shan Geo-
- 35 hazards Database (Havenith et al., 2015a, b) and the multi-temporal landslide inventory from Behling and Roessner (2020), there is a lack of landslide inventories with the exact date of landslide occurrence. Given the highly dynamic nature of weather phenomena, at least a daily time stamp timestamp of landslide records is required to investigate weather conditions that trigger landslides. Secondly, there is a lack of atmospheric data. The number of in-situ observation stations in Kyrgyzstan and Tajikistan decreased sharply in the 1990s due to reduced funding. There are currently eight stations in Kyrgyzstan and 26
- 40 stations in Tajikistan available from Global Surface Summary of the Day (GSOD), which is a publicly available data set. These numbers are already significantly below the recommendation of the World Meteorological Organization, even for flat areas (Ilyasov et al., 2013). Despite the sparse distribution, most GSOD stations are located in low-lying valleys and are not fully representative of the area.
- Rainfall is the most common trigger of landslide all over the world (Wieczorek, 1996). Over snow-covered regions, snowmelt
 is recognized as another common trigger of shallow landslides and debris flows (Wieczorek, 1996; Mostbauer et al., 2018). In Kyrgyzstan and Tajikistan, more than half of the annual precipitation falls in the form of snow. Snow cover duration over high mountain ranges in the Tien Shan and the Pamir is more than 200 days per year (Dietz et al., 2014). A large amount of water stored in snowpacks is released during the melting season. Snowmelt is another important source of water infiltrating into the soil that increases slope instability. Thus, in Kyrgyzstan and Tajikistan, snowmelt might also play a role in landslide
 triggering besides rainfall. But snowmelt is not as easy to be observed as rainfall and might often be neglected as a landslide
 - trigger, especially when co-occurring with rainfall.

There are two main approaches to assess rainfall threshold thresholds for landslide triggering. The first approach is physically based and requires detailed lithological, morphological, and geotechnical information of each landslide event (Guzzetti et al., 2007). Unfortunately, this level of detail is usually restricted to small areas and is not available for the whole of Kyrgyztan

55 Kyrgyzstan and Tajikistan. The second one is the empirical approach based on historical landslide and rainfall data. The majority of studies applying this approach relied on rain gauge data to analyze rainfall thresholds (e.g., Berti et al., 2012; Khan et al., 2012; Bui et al., 2013). However, rain gauge data are point measurements that cannot capture the large spatial

heterogeneity of rainfall, especially over complex terrains. Gridded products can provide continuous data in both space and time and can be used in detecting atmospheric triggering conditions of landslides.

- 60 We aim to analyze the atmospheric triggering conditions of landslides and generate climatic disposition maps that contain information on these triggering conditions in Kyrgyzstan and Tajikistan. For this purpose, we combined freely available gridded atmospheric data with historical landslide events. Atmospheric triggers for each landslides events landslide event were determined by the co-occurrence of landslide and weather events. Properties (mean intensity, peak intensity, accumulated amount) of landslide triggering events and non-landslide triggering events were compared. Objective thresholds of these properties for
- 65 different atmospheric triggers (rainfall, snowmelt, and the sum of rainfall and snowmelt) were defined so that they can best separate the atmospheric conditions that resulted and did not result in landslides. Finally, we applied the thresholds with the best predictive performance to generate maps of mean annual exceedance. In this way, we can transform the weather-scale triggering conditions into climate-scale disposition dispositions (hereafter referred to as "climatic disposition").

The objective of this study is threefold: (1) investigate the role of snowmelt in landslide triggering processes; (2) find 70 appropriate quantities of atmospheric triggers for assessing landslide hazards; (3) characterize climatic disposition in terms of rainfall and snowmelt over Kyrgyzstan and Tajikistan.

The paper is organized as follows: we describe the data and methods used in this study in the following section. Results are presented in section 3 and discussed in 4. Conclusions are drawn in section 5.

2 Data and method

75 2.1 Data

80

85

2.1.1 Landslide catalog

Landslide events used in this study come from two sources: the Global Landslide Catalog (GLC) (Kirschbaum et al., 2010, 2015) and the Global Fatal Landslide Database (GFLD) (Froude and Petley, 2018). GLC has been compiled by NASA since 2007 and contains all types of mass movements triggered mostly by rainfall. The sources of the GLC are mainly media reports, disaster databases, and scientific reports. The GFLD only includes landslide events that caused fatality obtained from media reports. It currently covers the period from 2004 to 2017. Both the GLC and the GFLD provide dates of landslide events.

We selected landslide events triggered by atmospheric factors in Kyrgyzstan and Tajikistan from 2007-2018 from the GLC and 2004-2017 from the GFLD. Then we merged these two data sets and deleted duplicated events that occurred on the same day and came from the same source link, resulting in 96 landslide events for Kyrgyzstan and Tajikistan from 2004 to 2018 (Fig. 1).

2.1.2 Atmospheric data

Rainfall and snowmelt data are extracted from the HAR v2. The HAR v2 is a newly developed regional atmospheric data set. It was generated by dynamical downscaling of the ERA5 reanalysis data using the Weather Research and Forecasting model

- (WRF). It is the only gridded atmospheric data set over High Mountain Asia provides atmospheric data with high resolution and
 accuracy (Hamm et al., 2020) over High Mountain Asia (Hamm et al., 2020; Wang et al., 2021). Detailed modeling strategies of the HAR v2 are described in Wang et al. (2021). The HAR v2 has a grid spacing of 10 km and is available in hourly, daily, monthly and yearly aggregations. Daily products were used in this study to determine the climatic trigger of each landslide event (section 2.2.1) and to define thresholds for landslide triggering (section 2.2.2). Rainfall was calculated as the difference between total precipitation and snowfall. Snowmelt is not a standard output of the WRF and was calculated using the Surface
- 95 Energy Balance (SEB). The SEB in the HAR v2 is resolved by the Noah Land Surface Model (LSM) (Tewari et al., 2004):

$$H_m = R_n - H_s - H_l - H_g \tag{1}$$

where R_n , H_s , H_l and H_g are net radiation, sensible heat flux, latent heat flux, and ground heat flux in W m⁻², respectively. These four variables are directly available in the HAR v2. H_m is the heat flux for melting and refreezing in W m⁻². $H_m > 0$ indicates melting process, while $H_m < 0$ refers to refreezing process. When $H_m > 0$, snowmelt h_m (kg m⁻² s⁻¹) is calculated as:

$$h_m = H_m / \lambda_m \tag{2}$$

where λ_m is the latent heat of fusion. When the calculated h_m is greater than snow water equivalent, then h_m is set to be equal to snow water equivalent.

2.2 Methods

100

105 2.2.1 Determine the atmospheric trigger of landslide events

Atmospheric The atmospheric trigger of a landslide event is determined by the co-occurrence of the landslide event with rainfall and snowmelt event. If a landslide event only occurred within or one day after a rainfall (snowmelt) event, then this landslide event is defined as rainfall (snowmelt) triggered. If there are both a rainfall event and a snowmelt event on the day or one day before the landslide occurrence day, then the atmospheric trigger of this landslide event is mixed.

To define a rainfall (snowmelt) event, the daily time series of rainfall(snowmelt) were extracted from the grid cells where landslides occurred. For each time series, an independent rainfall (snowmelt) event is defined as a series of consecutive days in which more than $0.2 \,\mathrm{mm}\,\mathrm{d}^{-1}$ of rainfall (snowmelt) is simulated. The value of $0.2 \,\mathrm{mm}\,\mathrm{d}^{-1}$ is chosen because it is the traditional precision of daily precipitation measurement (Jarraud, 2008) and can be applied to separate dry and wet conditions (Rodwell et al., 2010).

115 2.2.2 Threshold model for atmospheric triggers

The threshold model developed in this study contains three steps: (1) define landslide triggering events and non-triggering events; (2) define the thresholds for rainfall, snowmelt, and the sum of rainfall and snowmelt (hereafter referred to as rainfall+snowmelt) based on maximizing the predictive performance using 2×2 contingency tables; (3) validate and assess the



Figure 1. Landslide events from 2004-2018 extracted from the Global Landslide Catalog (GLC) GLC (white points) and the Global Fatal Landslide Database (GFLD) GFLD (black popints). Background contour is topography from Digital Elevation Model Digital Elevation Model (DEM) data from Shuttle Radar Topographic Mission (SRTM).

uncertainties of the defined thresholds. The methods for the first two steps were adopted from Leonarduzzi et al. (2017). Only the landslide events, the climatic triggers of which could be determined, were used for threshold modeling.

120

The first step is to define landslide triggering events and non-triggering events for rainfall, snowmelt, and rainfall+snowmelt. Here, we take rainfall as an example to describe the procedure. First, the method used in section 2.2.1 is applied to define rainfall events for each time series extracted from grid cells where landslides occurred. Next, if a landslide event occurred during or one day after a rainfall event, then this rainfall event is classified as a landslide triggering event (LTE). Given the

- 125 uncertainty in time stamps timestamps of landslide events, the day after is also considered as a temporal relaxation. Otherwise, if a rainfall event is not associated with any landslide events, it is classified as a non-landslide triggering event (NLTE). For each rainfall event, we calculated three event properties: mean intensity I_{mean} , maximum intensity I_{max} , and the accumulated amount of rainfall for the entire event Q. For triggering events, we also calculated these three properties by only considering the period up to the day of the landslide occurrence (hereafter referred to as UTL, meaning Up-To-Landslide). Note that, not all
- 130 the landslide events co-occurred with a rainfall event. For these events, we set I_{mean} , I_{max} , and Q to zero. The same procedure for defining LTEs and NLTEs was conducted for snowmelt and rainfall+snowmelt as well.

The second step is to define thresholds of rainfall, snowmelt, and rainfall+snowmelt for entire events and UTL events, using I_{mean} , I_{max} , and Q. No single threshold can perfectly separate LTEs from NLTEs since their distributions overlap. We applied 2×2 contingency tables to select the threshold that yields the best predictive performance. Using a certain threshold

135

140

as a binary classifier, LTEs and NLTEs were categorized into true positive (TP), true negative (TN), false positive (FP), and false negative (FN). The Peirce Skill Score (PSS) (Hanssen and Kuipers, 1965) was applied as the measure of the predictive performance because it is trail-independent, which means it is unbiased even when the numbers of LTEs and NLTEs are not equally presented (Woodcock, 1976). The PSS is also known as the Hanssen-Kuiper skill score and the true skill statistic. It is calculated as the difference between Hit Rate (HR) and False Alarm Rate (FAR):

$$PSS = HR - FAR \tag{3}$$

$$HR = \frac{TP}{TP + FN} \tag{4}$$

$$FAR = \frac{FP}{FP + TN} \tag{5}$$

We chose the threshold that maximizes the PSS. We also computed the Euclidean distance (*d*) to the optimal point (HR=1, FAR=0), which is another commonly used skill score in this application (e.g., Gariano et al., 2015; Piciullo et al., 2017;
Postance et al., 2018; Zhuo et al., 2019). Additionally, the receiver operating characteristic (ROC) curve was used to determine the general predictive power of a certain predictor by calculating the area under the ROC curve (AUC) (Fawcett, 2006).

The last step is to validate the threshold model and assess uncertainty. For the calibration of thresholds, all landslide event samples were utilized, and corresponding statistic measures were calculated, i.e., the threshold model was trained and tested on the same data set. To test the model's predictive ability on an unseen data set, we performed k-fold cross-validation. Landslide events were randomly split into k folds with k=8. Then for each unique fold, the fold was taken as the testing set, and the remaining k-1 folds were taken as the training set. Mean values of thresholds, the corresponding statistic measures, as well as their uncertainties represented by standard deviations were reported.

2.2.3 Mean annual exceedance

Mean annual exceedance (\overline{N}_{th}) is calculated for each HAR v2 grid cell. It is defined as the number of events that exceed a 155 certain threshold over a certain period (N_{th}) divided by the total number of years (N_a) :

$$\overline{N}_{th} = \frac{N_{th}}{N_a} \tag{6}$$

The unit of \overline{N}_{th} is the number of events per year. Mean annual exceedance transforms weather-scale triggering conditions to climate-scale disposition. It depicts where are landslides-landslides are likely to occur from the climatic aspect.

3 Results

160 3.1 The role of snowmelt in landslide triggering

Fig. Figure 2 shows the climatology of seasonal rainfall, snowmelt, and rainfall+snowmelt resolved by the HAR v2. We define seasons as commonly done in meteorology, spanning three months each: winter (December-February, DJF), spring (March-May, MAM), summer (June-August, JJA), and autumn (September-November, SON). A high amount of rainfall concentrates in the western foothill of the Fergana Range, the northern foothill of the Turkestan Range, and the Tajik Basin in

165 spring and shifts northeastwards into the Tien Shan in summer. Snowmelt occurs in spring over most high elevated areas. In summer, while most regions are snowmelt-free, the Pamir plateau still experiences a high amount of continuous snowmelt, which is in line with the results by Dietz et al. (2014) using remote sensing data.

Atmospheric triggers for each landslide events event are determined using the method described in section 2.2.1, and the results are shown in Fig. 3. Table A1 lists all 96 events and the climatic triggers detected by the HAR v2. Figure A1 shows

- 170 the temporal process of rainfall and snowmelt for selected landslide cases. Nine landslide events did not occur within any rainfall event, snowmelt event, or rainfall+snowmelt event. These This mismatch between landslide information and weather information stems from the uncertainties in landslide locations and timing, as well as the uncertainties from rainfall and snowmelt simulated in the HAR v2 (detailed discussion in section 4.1). These nine events are referred to as "not detected" (white points in Fig. 3) and are excluded. The remaining 87 landslide events were used for further analysis. Landslide events
- 175 that were only triggered by rainfall mainly cluster in Tajik Basin and the northeastern rim of the Fergana Basin, where the contribution of rainfall to the annual sum of rainfall and snowmelt is high (Fig. 3).

The annual cycles of rainfall, snowmelt, and rainfall+snowmelt are compared with monthly landslide occurrences in Fig. 4. The study region experiences a peak of landslide activity in April and May, which corresponds with the peak of rainfall+snowmelt. While rainfall is the dominant trigger of landslides, snowmelt contributes to triggering 40% of landslide events

180 (35 out of 87). There are 29% of landslide events (25 out of 87) that are attributed to the combined effect of rainfall and snowmelt. Most snowmelt-contributing events occurred in April when snowmelt amount is the highest. March and June have almost the same amount of rainfall+snowmelt. However, there are more landslide occurrences in June. This could be resulted from still frozen soil in March, which stabilizes the slope. As shown in Fig. 4a, both soil temperature at the top soil layer (0-0.1m) and air temperature at 2 m are still below zero in March.



Figure 2. Seasonal rainfall, snowmelt, and rainfall+snowmelt from the High Asia Refined Analysis version 2 (HAR v2) HAR v2 from 2004-2018. Black pointscircles: seasonal landslide events from Global Landslide Catalog (GLC) GLC and Global Fatal Landslide Database (GFLD) GFLD. Topographic shading is based on Digital Elevation Model DEM data from Shuttle Radar Topographic Mission SRTM. DJF: December-February; MAM: March-May; JJA: June-August; SON: September-November.



Figure 3. Contribution (%) of snowmelt to annual sum of rainfall and snowmelt (background contour) and climatic triggers of 96 landslide events extracted from Global Landslide Catalog (GLC) GLC and Global Fatal Landslide Database (GFLD) GFLD (points). Topographic shading is based on Digital Elevation Model DEM data from Shuttle Radar Topographic Mission SRTM.



Figure 4. (a) Mean monthly soil temperature at the top soil layer (0-0.1m) and air temperature at 2m averaged over Kyrgyzstan and Tajikistan extracted from the High Asia Refined Analysis version $2 (HAR v_2) HAR v_2$; (b) mean monthly rainfall and snowmelt averaged over Kyrgyzstan and Tajikistan extracted from the HAR v2; (c) mean monthly landslide occurrences in Kyrgyzstan and Tajikistan from 2004-2018.

185 3.2 Thresholds of elimatic atmospheric triggers for landslide triggering landslides in Kyrgyzstan and Tajikistan

Statistics of different properties of LTEs and NLTEs for rainfall, snowmelt, and rainfall+snowmelt are presented in Fig. 5 in the form of eumulative distribution function (CDF)empirical cumulative distribution function (eCDF). Rainfall and snowmelt have a high percentage of events with $I_{mean} = 0$, $I_{max} = 0$, and Q = 0 because. This is because, for landslide events that cannot be detected by only rainfall (orange points in Fig. 3) or only snowmelt (blue points in Fig. 3), I_{mean} , I_{max} , and Q of

- 190 rainfall for these events were all set to zero. The same procedure was conducted for events that cannot be detected by only snowmelt (blue points in Fig. 3). It can be seen in Fig. 5 that LTEs , for both entire events and UTL events , have stronger I_{mean} and I_{max} , as well as larger Q compared to NLTEs. Besides, snowmelt events have much higher Q but lower I_{mean} and I_{max} than rainfall events, indicating that snowmelt events are in general prolonged and not as intense as rainfall events. Overall, the HAR v2 combined with landslide inventories from GLC and GFLD can distinguish LTEs from NLTEs well and has potential in landslide threshold modeling.
- 195 in landslide threshold modeling.

We calibrated thresholds of I_{mean} , I_{max} , and Q using rainfall, snowmelt, and rainfall+snowmelt as predictors. The procedure was conducted for both entire events and UTL events. Predictive performance is better when using the entire period (Table 1) than just using the UTL period (Table ??1), which was also concluded by Leonarduzzi et al. (2017). One of the reasons is that by considering a longer period, I_{mean} , I_{max} , and especially Q of LTEs generally increase, making it easier to distinguish

- 200 LTEs from NLTEs. This can also be seen from the CDFeCDFs in Fig. 5. In CDF the eCDF space, the threshold defined by maximizing PPS_PSS is the point on the x-axis, where the vertical distance between the LTE curve and the NLTE curve is the largest. CDFeCDFs of UTL events are closer to the NLTE curve than CDFeCDFs of the entire events. Therefore, the maximum PSSs of UTL events are smaller (Fig. 5).Here, only results for entire events are presented since using The better performance by considering the entire period leads to better predictive performance and was mostly used in previous
- 205 studies (Leonarduzzi et al., 2017)could also indicate that there exists some uncertainty of landslide timing reported in GLC and GFLD. It can be seen from Table 1 that rainfall+snowmelt has the best predictive performance . For rainfall+snowmelt entire event, the best performance is achieved by I_{max} with a threshold value of 12.8 mmd⁻¹, and by Q with a threshold value of 17.2 mm. The PPS and d for for both entire events and UTL events. The predictive performance indicating by d, PSS, and AUC of the three event properties (I_{mean} , I_{max} , and $Q_{are very}$) are quite similar, but these two predictors have different
- 210 advantages. Using using I_{max} as a predictor leads to a lower FAR but also a lower HR when compared with Q and I_{mean}. K-fold cross-validation results for entire events and UTL events are presented in Table ?? A2 and Table A3. Cross-validation reduces the sample size and makes the results more sensitive to outliers. The validation results are in line with the conclusions drawn by calibration: (1) among all the predictors, rainfall+snowmelt has the best predictive performance for both entire events and UTL events; (2) for rainfall+snowmelt entire event, I_{max} and Q are more suitable than I_{mean} predictive performance is
- 215 better when using the entire period than just using the UTL period; (3) calibrated thresholds for predictive performance of I_{mean} , I_{max} , and Q are robust since they fall into the uncertainty range of the thresholds obtained by cross-validation for rainfall+snowmelt are quite similar, but I_{max} has a lower FAR and also a lower HR.



Figure 5. Cumulative distribution function (CDF) cCDF curves of mean intensity (I_{mean}), maximum intensity (I_{max}), and accumulated amount (Q) of non-landslide triggering event (NLTE)NLTE, landslide-triggering entire event (LTE entire), and landslide-triggering up-tolandslide event (LTE UTL) for rainfall, snowmelt, and rainfall+snowmelt during the period of 2004–2018. Grey dashed lines represent the thresholds for UTL event defined in Table 1.

3.3 Mean annual exceedance

220

Using the thresholds defined in section 3.2 for rainfall+snowmelt entire event, Fig. UTL events, Figure 6 presents the annual number of rainfall+snowmelt events that exceed the thresholds of $I_{mean} = 5.05 \text{ mm d}^{-1}$, $I_{max} = 12.8 \text{ mm d}^{-1} 14.05 \text{ mm d}^{-1}$, and $Q = 17.2 \text{ mm} \cdot 15.65 \text{ mm}$ (hereafter referred to as $I_{mean,th}$, $I_{max,th}$, and Q_{th}). Here, only the results for UTL events are presented since the defined thresholds of entire events and UTL events for rainfall+snowmelt are very similar and only deviate within 10%, although their predictive performance is different (Table 1).

predictor	property	threshold	HR	FAR	d	PSS	AUC
rainfall	I_{mean}	3.60	0.62	0.35	0.51	0.27	0.62
(entire event)	I_{max}	11.20	0.49	0.18	0.54	0.32	0.65
	Q	16.95	0.52	0.18	0.52	0.34	0.67
snowmelt	I_{mean}	7.05	0.23	0.06	0.77	0.17	0.31
(entire event)	I_{max}	13.45	0.24	0.04	0.76	0.20	0.32
	Q	119.60	0.24	0.03	0.76	0.21	0.33
rainfall+snowmelt	I_{mean}	4.95	0.71	0.25	0.38	0.46	0.78
(entire event)	I_{max}	12.80	0.67	0.15	0.37	0.51	0.81
	Q	17.15	0.74	0.23	0.35	0.50	0.81
rainfall	Imean	3.05	0.60	0.40	0.57	0.20	0.59
(UTL event)	$\underbrace{I_{max}}$	12.40	0.34	0.16	0.67	<u>0.19</u>	0.58
	${\cal Q}$	<u>9.25</u>	0.52	$\underbrace{0.31}_{\ldots}$	$\underbrace{0.57}_{\infty}$	0.21	0.59
snowmelt	Imean	7.40	0.22	$\underbrace{0.05}_{\sim\sim\sim}$	$\underbrace{0.78}_{\sim\sim}$	0.17	0.31
(UTL event)	$\underbrace{I_{max}}$	12.80	0.24	$\underbrace{0.05}_{\sim\sim\sim}$	$\underbrace{0.76}_{\infty}$	$\underbrace{0.19}_{\infty}$	0.32
	${\cal Q}$	<u>98.30</u>	0.24	0.04	$\underbrace{0.76}_{\infty}$	0.20	0.32
rainfall+snowmelt	Imean	5.05	0.68	0.25	0.41	0.43	0.76
(UTL event)	Imax	14.05	<u>0.59</u>	$\underbrace{0.14}$	0.44	0.45	<u>0.77</u>
	Q_{-}	15.65	0.66	0.25	0.43	$\underbrace{0.40}_{\sim\sim\sim}$	<u>0.76</u>

Table 1. Calibrated thresholds of $I_{mean} \pmod{d^{-1}}$, $I_{max} \pmod{d^{-1}}$, and $Q \pmod{d}$ for entire events and UTL events of rainfall, snowmelt, and the sum of rainfall and snowmelt (rainfall+snowmelt), and corresponding performance statistics.

- Locations with higher mean annual exceedance over $I_{max,th}$ indicate a higher chance of having rainfall+snowmelt event events with high intensity, such as the Fergana Range and the northeastern Tajik Basin. These two regions have a high contribution of rainfall to annual rainfall+snowmelt (Fig. 3), and rainfall events tend to have stronger intensity than snowmelt events (Fig. 5). Locations with high mean annual exceedance over Q_{th} but low exceedance over $I_{max,th}$, including the Pamir Plateau and the Tien Shan, indicate that prolonged events instead of short and intense events are more frequent. The mean annual exceedance map-maps of Q_{th} corresponds and $I_{mean,th}$ correspond better with the landslide occurrences since it takes they encompass both extreme events and prolonged events into account. Landslide events reported from the GLC and the GFLD are generally located in areas with high exceedance over Q_{th} and $I_{mean,th}$. However, the mean annual exceedance map-maps of Q_{th} also have more areas with false alarms, i.e., areas with high mean annual exceedance but no landslide occurrence. In remote areas, such as the Tien Shan, high false alarm-alarms could be due to the fact that landslides extracted from median media reports are generally under-reported in remote regions. This is discussed in details-detail in sec-235 tion 4.1. In contrast, the mean annual exceedance map of $I_{max,th}$ misses more landslide events but has less false alarm area
 - when compared to the exceedance map maps of Q_{th} and $I_{mean,th}$.



Figure 6. Mean annual exceedance (number of events per year) of (a) $I_{mean} = 5.05 \text{ mmd}^{-1}$ (b) $I_{max} = 12.8 \text{ mmd}^{-1}$ 14.05 mmd $^{-1}$; and (bc) $Q = 17.2 \text{ mm} \cdot 15.65 \text{ mm}$ for the sum of rainfall and rainfall+snowmelt UTL events. Black pointscircles: landslide events from Global Landslide Catalog (GLC) GLC and Global Fatal Landslide Database (GFLD) GFLD. Topographic shading is based on Digital Elevation Model DEM data from Shuttle Radar Topographic Mission SRTM.

4 Discussion

4.1 Sources of uncertainties uncertainty

The uncertainty of the results depends on the accuracy of the data and the method applied to analyze the data. Our approach

- 240 is purely empirical-based, which allows us to investigate broader areas without knowing the detailed surface characteristics of each landslide event. However, slope instability often results from numerous factors. The interaction between non-climatic characteristics and atmospheric triggers is also responsible for the initiation of landslides (Berti et al., 2012; Jia et al., 2020), which can not be captured by empirical methods. This is the reason why not all rainfall+snowmelt events that exceed $I_{mean,tb}$, $I_{max,tb}$, and Q_{tb} triggered landslides (Fig. 6), even though the number of landslides is underestimated.
- 245 Uncertainty in landslide inventories and atmospheric data is a very common issue in studies investigating thresholds for landslide triggering. These two sources of uncertainty have been comprehensively discussed and quantified (e.g., Nikolopoulos et al., 2014 . Uncertainty in these two data sources generally results in an underestimation of rainfall thresholds, leading to a higher false alarm rate (Nikolopoulos et al., 2014, 2015; Marra et al., 2016; Peres et al., 2018). In the following subsections, we discuss the uncertainty stemming from the landslide inventories (GLC and GFLD) and the rainfall and snowmelt in the HAR v2.

250 4.1.1 Uncertainty of landslide inventories

Uncertainties of the GLC and GFLD are comprehensively discussed in Kirschbaum et al. (2010), Kirschbaum et al. (2015), and Froude and Petley (2018). The first major problem of these two data sets is that they underestimate the total number of landslides. This is because these two data sets' primary sources are media reports, which are biased towards events with human casualties (Carrara et al., 2003). The second issue is that the spatial distribution of landslides is biased towards populated areas. In our study area, landslide events also tend to cluster in areas with high population density, e.g., the eastern rim of the Fergana Basin and the Tajik Basin. Landslide number over remote areas is much likely to be under-reported. Last but not leastIn addition, there is large uncertainty in landslide location because most media reports do not contain the exact location where landslides were initiated, but rather just the name of the village, road, or city affected by landslides. An example in our case is

the landslide event in the Issyk-Kul Basin (Fig. 1), the location of which is in a flat area, and the location accuracy provided by

- the GLC is "exact". This landslide event's initial zone must be different from the reported location and somewhere nearby with slopes. We also failed to determine the climatic trigger of this landslide event using the HAR v2. Last but not least, landslide timing was also reported with a certain degree of uncertainty. Although it is more typical that a landslide was reported after its actual occurrence (positive errors), negative errors are also possible depending on the interpretation of historical landslide information from an analyst (Peres et al., 2018). Our results show that using the entire weather event period leads to a better
- 265 predictive performance than just using the UTL period (Table 1). This could be an indication of negative errors in the landslide timing.

Despite these known limitations, the GLC and the GFLD still provide the lower boundary of landslide number and are proven to be valuable in global and regional landslide studies (Kirschbaum and Stanley, 2018; Jia et al., 2020; Stanley et al., 2020). Although the landslide number is known to be incomplete, our results show that they can still present the seasonal distribution of landslide occurrence reasonably well (Fig. 4). This was also concluded by Kirschbaum et al. (2015), who stated that the reason for the unbiased seasonal distribution of landslide occurrence is that the compilation method depends on media alerts, which is consistent throughout the year. Additionally, even though location uncertainty exists, we could determine atmospheric triggers of 91% of landslide events (87 out of 96). The reason could be that landslide-triggering rainfall and snowmelt events generally have a large spatial extend (Leonarduzzi et al., 2017).

275 Another source of uncertainty comes from the rainfall and snowmelt data simulated in the HAR v2. Compared to

4.1.2 Uncertainty of atmospheric data

Extracting weather data that can represent the exact weather condition at landslide sites is always a challenge in studies investigating rainfall thresholds for landslide triggering. Rain gauges are the main source of rainfall information (Segoni et al., 2018), and it is very seldom that landslide initial locations are gauged. Due to the highly heterogeneous spatial distribution of

- 280 precipitation, especially over complex terrains, there exists great uncertainty when rainfall is not directly measured from landslide initial points. Additionally, Marra et al. (2016) found that the initial points of shallow landslides and debris flows generally correspond to the local peak of rainfall. Rain depth decreases with the distance, causing an underestimation when rainfall is measured away from the landslide initial point. Traditionally, the nearest gauge is used to represent the weather condition at the landslide site, which sometimes can be kilometers away. Nikolopoulos et al. (2015) examined other more complicated interpolation methods, such as inverse distance weighting and ordinary kriging, and concluded that these methods
- did not bring any particular added value to the simplest nearest neighbor method.

Using gridded data can avoid this allocation problem (Leonarduzzi et al., 2017). But uncertainties still exist since gridded data only represent the grid-mean value but not the "true" weather condition at landslide sites. Nevertheless, it is still essential that the gridded data used in our study can accurately represent the grid-mean value. The WRF model configurations of the

- 290 HAR v2, such as the forcing strategy, physical parameterization schemes, were carefully chosen to ensure its quality (Wang et al., 2021) . Several studies (Pritchard et al., 2019; Li et al., 2020) indicate the high accuracy and quality of the old version of the High Asia Refined Analysis (HAR) (Maussion et al., 2014), HAR v2 can simulate precipitation over. Wang et al. (2021) compared the performance of the two versions of the HAR against in-situ observations from 57 GSOD stations over the High Mountain Asia as accurately as the old version. Simulation of in terms of daily precipitation and air temperature at 2 mis better
- 295 than. It was concluded that compared to the old versiondue to the snow depth correction approach (Wang et al., 2021). HAR v2 generally produces slightly higher precipitation amounts with a mean bias of 0.36 mmd⁻¹. Furthermore, Hamm et al. (2020) compared the HAR v2 with other gridded precipitation data sets in different spatial resolutions, including reanalysis data and satellite-based precipitation retrieval, over a rugged terrain of the central Himalaya and the southwestern Tibetan Plateau. It was concluded that the HAR v2 is the only product that can resolve orographic precipitation, which is a funda-
- 300 mental process over complex terrain. Furthermore, the HAR v2 can capture more extreme precipitation events than coarser productsSimulation of air temperature at 2 m in the HAR v2 is better than the old version due to the snow depth correction approach (Wang et al., 2021). Snowmelt in the HAR v2 is resolved by the Noah LSM, which only considers a single layer of snowpack (Koren et al., 1999). Several studies found uncertainty of the Noah LSM in reproducing the snow-related pro-

cess, e.g., the overestimation of snow albedo (e.g., Chen et al., 2014; Minder et al., 2016; Tomasi et al., 2017). Nevertheless,

305 the snow-related process is the major weakness of LSMs and needs further improvement in the future (Chen et al., 2014). Despite the uncertainty in the HAR v2, our results show that it can distinguish LTE s from NLTEs very well, 5), suggesting that the HAR v2 and dynamical downscaling products, in general, are suitable for extracting weather conditions corresponding to landslides.

Our approach is purely empirical-based, which allows us to investigate broader areas without knowing the detailed surface

310 characteristics of each landslide event. However, slope instability often results from numerous factors. The interaction between non-climatic characteristics and atmospheric triggers is also responsible for the initiation of landslides (Berti et al., 2012; Jia et al., 2020)
 , which can not be captured by empirical methods. This is the reason why not all rainfall+snowmelt events that exceed Imax, th and Qth triggered landslides (Fig. 6), even though the number of landslides is underestimated.

4.1.3 Impact of spatial resolution of atmospheric data

- 315 Previous studies have shown that the spatial resolutions of gridded rainfall data have impacts on identifying landslide triggering thresholds (Marra et al., 2017; Nikolopoulos et al., 2017). To investigate the influence of spatial resolution of rainfall+snowmelt data on the event properties of landslide triggering weather events and the triggering thresholds, we resampled the rainfall+snowmelt data from HAR v2 to lower resolutions (20 km, 30 km, and 40 km). Then, we repeated the procedure described in section 2.2.2 to determine the event properties of LTE UTL events and their associated thresholds. The results are presented in Fig. 7.
- 320 There are nine "not detected" events when using the original HAR v2 10 km data (Fig. 3), which means the rainfall+snowmelt amounts at these landslide grid points are near zero ($\leq 0.2 \,\mathrm{mm} \,\mathrm{d}^{-1}$) at the day and one day before landslide occurrence. By lowering the spatial resolution, more events can be detected. This implies the uncertainty in the reported landslide location since resampling of rainfall+snowmelt encompasses rainfall+snowmelt information from nearby grid points. In general, I_{mean} and I_{max} decrease with the increase of grid size, which is in line with the findings of Hamm et al. (2020) that higher resolved
- 325 products generally capture more extreme events than coarser products. I_{mean} and I_{max} thresholds defined by coarser products are also generally lower. The impact of grid size on Q is the opposite: larger grid size leads to higher Q and threshold value. This is closely associated with the increase of event duration with the increase of grid spacing, resulting from the fact that the resampling process can blend several localized events temporally together. However, lowering the spatial resolution does not lead to worse predictive performance. This, on the one hand, implies again that lower resolution can partly compensate for the
- 330 uncertainty in landslide locations. On the other hand, it indicates that although landslide initiation itself is a highly localized phenomenon, the weather processes that ensure sufficient water input into the system and trigger landslides can be clearly identified at the meso-scale (Prenner et al., 2018).

Based on the above analysis, it can be expected that a convection-permitting scale (<10 km) downscaling simulation would provide a more realistic representation of weather events that initialized landslides. Compared to such a high-resolution

335 simulation, the HAR v2 10 km data would underestimate the intensity and overestimate the duration of landslide triggering rainfall+snowmelt events. Moreover, the 10 km resolution of the HAR v2 is not able to explicitly resolve convection processes. Convection-permitting scale simulations show improvement over simulations applying cumulus parameterization schemes in several aspects, such as more accurate reproduction of the timing of precipitation peaks (Ou et al., 2020; Zhou et al., 2021).

However, a finer resolution has a lower tolerance to the uncertainty in the landslide location. The potential of a kilometer-scale
 simulation cannot be realized if the landslide location uncertainty is larger than the gird size. Thus, for our study region, future studies should not only focus on acquiring high-resolution and high-quality atmospheric data, but also on developing landslide inventories with higher location accuracy.



Figure 7. Boxplots demonstrating the impact of spatial resolution of atmospheric data on I_{mean} , I_{max} , Q, and duration of LTE UTL events, as well as the

4.2 Climatic disposition

345

In probabilistic risk analysis (e.g., Scherer et al., 2013), the risk that a system experiences an adverse effect caused by a hazardous process is given as the product of hazard and vulnerability. Vulnerability itself depends on exposure and sensitivity. Adverse effects only occur when the elements at risk are exposed to a hazardous event. Thus, risk is a function of hazard, exposure, and sensitivity. Applying this risk concept to our case, the adverse effect is landslide triggered by rainfall+snowmelt, and the hazardous process is rainfall+snowmelt events that exceed $I_{max,th}$ and Q_{th} the defined thresholds. The risk that a location experiences landslide triggered by rainfall+snowmelt depends on two factors: (a) how frequent a location is exposed to rainfall+snowmelt events that exceed $I_{mean,th}$, $I_{max,th}$, and Q_{th} , and (b) how sensitive slope instability can be triggered at this location. Climatic disposition represented by mean annual exceedance is actually factor (a) and comprises both aspects of hazard and exposure. Sensitivity is non-climatic landslide susceptibility that is only controlled by terrestrial characteristics. Thus, to assess landslide susceptibility, both climatic and non-climatic aspects need to be included.





Figure 8. Non-climatic landslide susceptibility map computed using slope, geology, fault zones, road networks, and forest loss developed by Stanley and Kirschbaum (2017). Black circles: landslide events from GLC and GFLD. Topographic shading is based on DEM data from SRTM.

355

The majority of landslide susceptibility studies only considered non-climatic factors. We compared our mean annul exceedance maps with a non-climatic landslide susceptibility map developed by Stanley and Kirschbaum (2017) at a resolution of approximately 1 km (Fig. 8). This non-climatic susceptibility map was generated using a heuristic fuzzy approach, in which slope, faults, geology, forest loss, and road networks were taken into account. This map is chosen because it covers the whole of Kyrgyzstan and Tajikistan. Even though the non-climatic susceptibility map and our mean annual exceedance maps were generated by totally different methods, they share some similarities. They both show higher values over areas with steep slopes and lower values in intermontane basins and valleys. This is because topographic relief is considered the best first-order rain-360 fall predictor (Bookhagen and Strecker, 2008). The non-climatic susceptibility map includes information on topography, and topography is explicitly resolved during dynamical downscaling. Mean annual exceedance maps not only display these localscale features caused by topography but also comprise general atmospheric circulation processes. Discrepancies between the landslide susceptibility map and our exceedance maps, e.g., over the Pamir Plateau and the southwestern Tajik Basin, suggest 365 that for landslide susceptibility mapping, both climatic and non-climatic aspects need to be considered.



Figure 9. Annual sum of rainfall and snowmelt averaged over 2014-2018 from HAR v2. Black circles: landslide events from GLC and GFLD . Topographic shading is based on Digital Elevation Model data from SRTM.

In addition, some landslide susceptibility studies took climate into account, but they often simply applied averaged annual precipitation (e.g., Shahabi et al., 2014; Havenith et al., 2015b; Wang et al., 2015). Averaged annual precipitation only shows the climatological conditions in general. Mean annual exceedance is derived from weather-scale triggering conditions, and therefore, it also contains information on extreme processes. In our case, for instance, the mean annual rainfall+snowmelt map does not correspond well with landslide occurrences, especially in the Tajik Basin and the northeastern rim of the Fergana Basin (Fig. 9). But these landslide events are captured better in both mean annual exceedance maps (Fig. 6). This indicates the added value of climatic disposition derived from triggering conditions.

4.3 Thresholds for different landslide size

GLC provides six categorized landslide sizes. Landslide events in Kyrgyzstan and Tajikistan fall into the following categories: (1) small: small landslide affecting one hill slope or small area; (2) medium: moderately sized landslide that could be either a 375 single event or multiple landslides within an area, and involves a large volume of material;(3) large: large landslide or series of landslides that occur in one general area but cover a wide area; (4) unknown (Kirschbaum et al., 2015). GFLD does not contain information about landslide size. Therefore, for landslide events from GFLD, we set the landslide size as "unknown". Table 2 presents the calibrated thresholds and corresponding statistical scores for these categories for UTL events. Using entire events leads to similar results (not presented here).

380

370

Interestingly, the thresholds for landslides with small sizes are higher than other categories and have the best predictive performance. All these 5 small-sized landslide events are snowmelt contributed events that occurred from March to May. The worse predictive performance for landslides with larger sizes could indicate that for those events, the triggering mechanism is much more complicated than small-sized events, and other non-atmospheric factors might also play a role. However, the **Table 2.** Calibrated thresholds of I_{mean} (mmd⁻¹), I_{max} (mmd⁻¹), and Q (mm) for UTL events of the sum of rainfall and snowmelt (rainfall+snowmelt), and corresponding performance statistics for different categories of landslide size. n refers to the number of landslides in each category.

landslide size	property	threshold	HR	FAR	$\overset{d}{\sim}$	PSS A	<u>UC</u>
small	Imean	9.85	1.00	0.07	0.07	0.93 (0.97
<u>(n=5)</u>	I_{max}	21.55	1.00	0.07	0.07	0.93	0.97
	${Q}$	124.25	1.00	0.04	0.04	<u>0.96</u> <u>(</u>) <u>.98</u>
medium	Inean	4.80	0.63	0.25	0.44	0.39	0.71
(<u>n=41</u>)	Imax_	14.05	0.49	0.12	0.53	0.37 (0.73
	${Q}$	9.65	<u>0.73</u>	0.35	0.44	0.38	0.72
large	Inean	8.10	0.55	0.11	0.47	0.44	0.72
(<u>n=11</u>)	Imax	21.75	0.45	0.05	0.55	0.40	0.73
	${Q}$	2.85	1.00	0.63	0.63	0.37 (0.73
unknown	Imean	5.25	0.77	0.26	0.35	0.51 (0.80
(<u>n=30</u>)	Imax	13.25	<u>0.73</u>	0.17	0.32	0.57 (0.81
	${Q}$	16.90	0.77	0.25	0.34	0.51	0.79

385 sample size of small-sized landslide events is too small to draw a robust conclusion. The number of small-sized landslides is expected to be under-reported since media reports are biased towards events with more severe impacts.

5 Conclusions

In this study, we combined gridded atmospheric data from the HAR v2 with 87 landslide records extracted from the GLC and the GFLD to analyze rainfall and snowmelt conditions that triggered landslides in Kyrgyzstan and Tajikistan. Thresholds for landslide triggering were determined for different event properties for rainfall, snowmelt, and rainfall+snowmelt. Mean annual exceedance maps were generated based on the defined thresholds.

Monthly landslide counts in Kyrgyzstan and Tajikistan correspond well with the monthly distribution of rainfall+snowmelt. An exception is March when soil temperature at the top soil layer (0-0.1m) and air temperature at 2m are both below zero. Investigation of the relationship between landslides and soil temperature could be a topic for future studies. Snowmelt plays a crucial role in landslide triggering in Kyrgyzstan and Tajikistan since it contributes to the triggering of 40% of landslide events.

By including snowmelt as an additional trigger, the skill of landslide prediction was significantly improved. I_{mean} , I_{max} , and Q of rainfall+snowmelt entire events have the best-have similar predictive performance. Thresholds of I_{mean} =5.05 mm d⁻¹, I_{max} =12.8 mm d⁻¹ (HR=0.67, FAR=0.15)-14.05 mm d⁻¹, and Q =17.2 mm (HR=0.74, FAR=0.25)-15.65 mm for UTL events were defined for landslide triggering in Kyrgyzstan and Tajikistan. Using the entire period of weather events leads to similar threshold values but better predictive performance. This could indicate uncertainty in landslide timing. Mean annual

390

395

400

exceedance maps derived from these thresholds depict climatic disposition and have added value in landslide susceptibility mapping.

Our study also The majority of previous studies applied rainfall estimates from in-situ gauges or satellite retrievals. Our study demonstrates the potential of the Regional Climate Model (RCM) in landslide prediction. Dynamical downscaling prod-

- 405 ucts generated by RCMs can provide physically consistent, high-resolution data that is extremely valuable for data scare areas. Given the global applicability of the dynamical downscaling methodand global coverage of the GLC and the GFLD , our approach can also be applied in other regions, as long as the number and quality of landslide records is sufficient. The GLC and the GFLD suffer from under-reporting problem over remote areas, which suggests the need for combination of different methods in landslide detection in remote regions are sufficient. Even though a higher-resolved downscaling product can
- 410 reproduce landslide-triggering weather events more realistically, it has a lower tolerance to the uncertainty in landslide locations and does not necessarily lead to better predictive performance. This calls for the need of developing landslide inventories with higher location accuracy in Kyrgyzstan and Tajikistan.

Appendix A



Figure A1. Event-based temporal process of rainfall and snowmelt for selected landslide events with landslide triggers defined as (a) "rainfall", (b) "snowmelt"; (c) "mixed"; and (d) "not detected" according to the method described in section 2.2.1.

Table A1: Landslide events in Kyrgyzstan and Tajikistan extracted from Global Landslide Catalog (GLC) GLC and Global Fatal Landslide Database (GFLD) GFLD from 2004 to 2018. Column "trigger" indicates the trigger of landslide events detected by the High Asia Refined Analysis version 2 (HAR v2)HAR v2.

Event date	Source	Longitude	Latitude	Country	Trigger
2004-04-17	GFLD	73.0420	40.3428	Kyrgyzstan	mixed
2004-05-22	GFLD	69.2172	39.8106	Tajikistan	rainfall
2004-06-14	GFLD	70.8718	39.8734	Kyrgyzstan	rainfall
2004-11-17	GFLD	70.0802	38.8324	Tajikistan	mixed

2005-03-13	GFLD	69.0502	40.0141	Tajikistan	mixed
2005-04-09	GFLD	69.2656	38.3801	Tajikistan	mixed
2007-03-25	GLC	70.1951	39.0071	Tajikistan	mixed
2007-04-01	GLC	72.5920	37.5760	Tajikistan	mixed
2007-04-05	GLC	71.6110	36.7270	Tajikistan	snowmelt
2007-04-17	GLC	71.6849	41.5552	Kyrgyzstan	rainfall
2007-04-17	GLC	68.2140	38.5330	Tajikistan	rainfall
2007-04-22	GLC	73.1416	40.8870	Kyrgyzstan	rainfall
2007-06-05	GFLD	69.1633	37.8276	Tajikistan	rainfall
2007-07-21	GLC	73.0000	38.0000	Tajikistan	mixed
2007-07-22	GLC	70.4400	40.7500	Tajikistan	not detected
2007-07-22	GFLD	71.0363	38.5289	Tajikistan	rainfall
2009-04-16	GFLD	71.9767	41.6184	Kyrgyzstan	rainfall
2009-04-21	GLC	68.7882	37.8515	Tajikistan	rainfall
2009-05-05	GFLD	70.1529	38.1701	Tajikistan	rainfall
2009-05-07	GFLD	69.7741	38.6726	Tajikistan	rainfall
2009-05-11	GFLD	71.0363	38.5289	Tajikistan	snowmelt
2009-05-14	GLC	68.6900	37.9867	Tajikistan	rainfall
2009-05-16	GFLD	71.0363	38.5289	Tajikistan	snowmelt
2009-05-20	GFLD	69.3199	38.7221	Tajikistan	rainfall
2010-03-13	GFLD	69.0502	40.0141	Tajikistan	snowmelt
2010-05-07	GLC	69.8054	37.9148	Tajikistan	rainfall
2010-05-07	GFLD	70.0994	37.8560	Tajikistan	rainfall
2010-06-03	GLC	72.9227	39.9854	Kyrgyzstan	mixed
2011-05-11	GLC	72.8282	41.4088	Kyrgyzstan	rainfall
2011-06-12	GLC	69.1238	38.2644	Tajikistan	rainfall
2011-06-12	GLC	69.5667	39.9342	Kyrgyzstan	rainfall
2012-05-12	GLC	70.8159	40.0538	Kyrgyzstan	rainfall
2012-05-13	GFLD	70.8718	39.8734	Kyrgyzstan	rainfall
2013-06-28	GLC	72.0106	41.6518	Kyrgyzstan	rainfall
2014-04-12	GLC	69.0971	37.9107	Tajikistan	rainfall
2014-04-12	GFLD	70.0994	37.8560	Tajikistan	rainfall
2014-04-16	GFLD	68.6749	38.0710	Tajikistan	rainfall
2014-04-26	GFLD	68.7626	38.5685	Tajikistan	rainfall
2015-04-03	GFLD	69.4222	38.5428	Tajikistan	rainfall

2015-05-08	GLC	70.0162	38.0991	Tajikistan	rainfall
2015-05-24	GLC	72.9053	40.8986	Kyrgyzstan	rainfall
2015-05-24	GFLD	73.2559	41.1036	Kyrgyzstan	rainfall
2015-07-10	GLC	70.4275	39.0712	Tajikistan	not detected
2015-07-16	GLC	71.7041	37.5773	Tajikistan	rainfall
2015-07-21	GFLD	71.7929	38.4071	Tajikistan	rainfall
2016-04-26	GLC	72.9071	40.8894	Kyrgyzstan	not detected
2016-05-09	GLC	68.5748	39.3160	Tajikistan	mixed
2016-05-15	GLC	72.9293	41.3431	Kyrgyzstan	rainfall
2016-05-23	GLC	72.7907	40.5304	Kyrgyzstan	rainfall
2016-05-27	GLC	69.8266	39.8751	Kyrgyzstan	rainfall
2016-05-28	GLC	71.5577	40.0150	Kyrgyzstan	mixed
2016-06-16	GLC	72.3374	41.4850	Kyrgyzstan	rainfall
2016-06-20	GLC	73.5233	40.1293	Kyrgyzstan	rainfall
2016-06-27	GLC	74.4438	41.7246	Kyrgyzstan	rainfall
2016-06-29	GLC	73.1415	41.7649	Kyrgyzstan	not detected
2016-07-29	GLC	69.5597	39.9377	Kyrgyzstan	rainfall
2016-08-16	GLC	78.3019	42.6831	Kyrgyzstan	not detected
2016-08-18	GLC	70.5626	39.9790	Tajikistan	rainfall
2017-01-04	GLC	71.9999	39.6699	Kyrgyzstan	snowmelt
2017-01-26	GLC	72.8834	40.8960	Kyrgyzstan	not detected
2017-03-26	GFLD	73.5725	40.8316	Kyrgyzstan	mixed
2017-04-07	GLC	73.6257	40.7733	Kyrgyzstan	snowmelt
2017-04-09	GLC	73.5335	40.8320	Kyrgyzstan	snowmelt
2017-04-10	GLC	69.5091	39.9095	Kyrgyzstan	mixed
2017-04-11	GLC	72.8601	41.2047	Kyrgyzstan	mixed
2017-04-14	GFLD	73.5725	40.8316	Kyrgyzstan	mixed
2017-04-16	GLC	73.2668	40.6430	Kyrgyzstan	snowmelt
2017-04-16	GLC	73.6000	40.7836	Kyrgyzstan	snowmelt
2017-04-17	GLC	73.6047	40.8044	Kyrgyzstan	mixed
2017-04-18	GLC	71.4973	37.3628	Tajikistan	mixed
2017-04-18	GLC	72.9069	40.8838	Kyrgyzstan	rainfall
2017-04-22	GLC	73.3402	40.8663	Kyrgyzstan	mixed
2017-04-23	GLC	71.5074	39.3410	Tajikistan	snowmelt
2017-04-23	GLC	72.8835	41.1610	Kyrgyzstan	rainfall

2017-04-23	GFLD	72.9801	41.2790	Kyrgyzstan	mixed
2017-04-29	GLC	73.4724	40.8864	Kyrgyzstan	mixed
2017-04-29	GFLD	73.2203	40.1325	Kyrgyzstan	mixed
2017-04-30	GLC	72.4381	41.2550	Kyrgyzstan	rainfall
2017-04-30	GLC	73.5310	40.0774	Kyrgyzstan	mixed
2017-05-10	GLC	74.4847	42.5635	Kyrgyzstan	mixed
2017-05-11	GLC	73.3497	40.5560	Kyrgyzstan	rainfall
2017-05-16	GLC	71.0302	41.7545	Kyrgyzstan	rainfall
2017-05-17	GLC	72.6771	41.6014	Kyrgyzstan	rainfall
2017-05-28	GLC	71.2755	39.1978	Tajikistan	mixed
2017-06-19	GLC	72.9814	39.6978	Kyrgyzstan	mixed
2017-06-19	GLC	71.7318	40.0439	Kyrgyzstan	rainfall
2017-06-26	GLC	67.8173	39.5267	Tajikistan	rainfall
2017-06-28	GLC	68.5480	39.3951	Tajikistan	not detected
2017-06-29	GLC	72.7303	41.0321	Kyrgyzstan	rainfall
2017-06-29	GLC	72.4521	41.2557	Kyrgyzstan	rainfall
2017-07-03	GLC	70.3650	39.0219	Tajikistan	rainfall
2017-07-03	GLC	68.4838	39.1172	Tajikistan	not detected
2017-07-04	GLC	69.5279	39.8102	Kyrgyzstan	not detected
2018-05-13	GLC	69.5445	39.8526	Kyrgyzstan	rainfall
2018-05-16	GLC	69.1773	37.2642	Tajikistan	rainfall
2018-05-21	GLC	72.1386	40.2437	Kyrgyzstan	mixed

Table A2. K-fold validation results. Mean values and standard deviations (in parentheses) for thresholds of I_{mean} (mmd⁻¹), I_{max} (mmd⁻¹), and Q (mm) for entire events of rainfall, snowmelt, and the sum of rainfall and snowmelt (rainfall+snowmelt), and corresponding performance statistics.

predictor	property	threshold	HR	FAR	d_{\sim}	PSS	AUC
rainfall	Т	3.76	0.56	0.33	0.56	0.23	0.62
	I mean	(0.33)	(0.14)	(0.03)	(0.10)	(0.13)	(0.01)
	Imax~	11.06	0.46	0.18	0.57	0.28	0.65
		(0.66)	(0.16)	(0.02)	(0.15)	(0.15)	(0.01)
	0	12.31	0.53	0.25	0.55	0.27	0.67
	×.	(3.88)	(0.16)	(0.07)	(0.10)	(0.10)	(0.01)
snowmelt	T	7.06	0.22	0.06	0.78	0.16	0.31
	Incan	(0.02)	(0.14)	(0.01)	(0.14)	(0.14)	(0.02)
	T	13.61	0.23	0.04	0.77	0.19	0.32
	~max~	(0.44)	(0.13)	(0.01)	(0.13)	(0.12)	(0.01)
	0	122.38	0.23	0.03	0.77	0.20	0.33
	×.	(7.93)	(0.13)	(0.01)	(0.13)	(0.12)	(0.01)
rainfall±snowmelt	T	4.96	0.70	0.25	0.40	0.45	0.78
	Imean	(0.02)	(0.13)	(0.02)	(0.08)	(0.14)	(0.01)
	I	12.93	0.65	0.15	0.39	0.49	0.81
		(0.37)	(0.15)	(0.01)	(0.13)	(0.15)	(0.01)
	0	<u>17.20</u>	0.71	0.23	0.38	0.48	0.81
	×	(0.14)	(0.15)	(0.02)	(0.10)	(0.13)	(0.01)

Table A3. K-fold validation results. Mean values and standard deviations (in parentheses) for thresholds of I_{mean} (mmd⁻¹), I_{max} (mmd⁻¹), and Q (mm) for UTL events of rainfall, snowmelt, and the sum of rainfall and snowmelt (rainfall+snowmelt), and corresponding performance statistics.

predictor	property	threshold	HR	FAR	d_{\sim}	PSS	AUC
rainfall	Т	<u>4.04</u>	0.45	0.33	0.66	0.12	<u>0.59</u>
	Imean	(1.47)	(0.13)	(0.10)	(0.08)	(0.08)	(0.01)
	T	10.94	0.34	$\underbrace{0.18}_{\overset{}}\overset{}\overset{}\overset{}\overset{}\overset{}\overset{}$	0.68	0.16	0.58
	$I_{\underline{max}}$	(1.47)	(0.06)	(0.04)	(0.05)	(0.06)	(0.01)
	0	10.21	0.46	0.29	0.62	0.17	<u>0.59</u>
	×	(2.22)	(0.09)	(0.04)	(0.09)	(0.11)	(0.01)
snowmelt	Т	<u>7.14</u>	<u>0.21</u>	$\underbrace{0.06}_{\sim\sim\sim}$	<u>0.79</u>	0.15	<u>0.31</u>
~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	Inean	(0.26)	(0.10)	(0.02)	(0.10)	(0.09)	(0.02)
		12.88	0.23	$\underbrace{0.05}_{\sim\sim\sim}$	0.77	<u>0.18</u>	0.32
		(0.23)	(0.12)	(0.01)	(0.12)	(0.11)	(0.02)
		<u>99.95</u>	0.22	$\underbrace{0.04}_{\sim\sim\sim}$	<u>0.78</u>	0.18	0.32
	×	(4.67)	(0.13)	(0.01)	(0.13)	(0.13)	(0.02)
snowmelt	T	5.35	0.61	0.23	$\underbrace{0.47}_{\sim\sim\sim}$	0.38	$\underbrace{0.76}_{\sim\sim\sim}$
	Imean	(0.85)	(0.22)	(0.04)	(0.17)	(0.18)	(0.01)
	T	13.54	0.56	0.14	$\underbrace{0.47}_{\sim\sim\sim\sim}$	0.42	$\underbrace{0.77}_{\sim\sim\sim}$
	1 max	(0.56)	(0.15)	(0.01)	(0.14)	(0.14)	(0.01)
	0	15.83	0.63	0.25	0.45	0.38	0.76
	¥	(0.44)	(0.13)	(0.02)	(0.10)	(0.12)	(0.01)

*Code and data availability.* The landslide data and atmospheric data used in this study are freely available from the following links:

- 415
- Global Landslide Catalog (GLC): https://maps.nccs.nasa.gov/arcgis/home/item.html?id=eec7aee8d2e040c7b8d3ee5fd0e0d7b9
  - Global Fatal Landslide Database (GFLD): https://blogs.agu.org/landslideblog/2019/06/18/global-fatal-landslide-database-1/
  - High Asia Refined Analysis version 2 (HAR v2): https://www.klima.tu-berlin.de/HARv2

The source code used in this study is freely available upon request.

*Author contributions.* All authors were involved in study conceptualization and writing of the manuscript. XW collected the data, carried out the analyses, and produced the visualizations.

Competing interests. The authors declare that they have no conflict of interest

Acknowledgements. This work was supported by the German Federal Ministry of Education and Research (BMBF) under the framework of the "Climatic and Tectonic Natural Hazards in Central Asia (CaTeNA)" project (Grant Number FKZ 03G0878G).

### References

460

- 425 Barbosa, N., Andreani, L., Gloaguen, R., and Ratschbacher, L.: Window-Based Morphometric Indices as Predictive Variables for Landslide Susceptibility Models, Remote Sensing, 13, 451, 2021.
  - Behling, R. and Roessner, S.: Multi-temporal landslide inventory for a study area in Southern Kyrgyzstan derived from RapidEye satellite time series data (2009-2013), 2020.

Berti, M., Martina, M., Franceschini, S., Pignone, S., Simoni, A., and Pizziolo, M.: Probabilistic rainfall thresholds for landslide occurrence

430 using a Bayesian approach, Journal of Geophysical Research: Earth Surface, 117, 2012.

- Bookhagen, B. and Strecker, M. R.: Orographic barriers, high-resolution TRMM rainfall, and relief variations along the eastern Andes, Geophysical Research Letters, 35, 2008.
  - Braun, A., Fernandez-Steeger, T., Havenith, H.-B., and Torgoev, A.: Landslide Susceptibility Mapping with Data Mining Methods—a Case Study from Maily-Say, Kyrgyzstan, in: Engineering Geology for Society and Territory-Volume 2, pp. 995–998, Springer, 2015.
- 435 Bui, D. T., Pradhan, B., Lofman, O., Revhaug, I., and Dick, Ø. B.: Regional prediction of landslide hazard using probability analysis of intense rainfall in the Hoa Binh province, Vietnam, Natural hazards, 66, 707–730, 2013.
  - Carrara, A., Crosta, G., and Frattini, P.: Geomorphological and historical data in assessing landslide hazard, Earth Surface Processes and Landforms: The Journal of the British Geomorphological Research Group, 28, 1125–1142, 2003.

Chen, F., Barlage, M., Tewari, M., Rasmussen, R., Jin, J., Lettenmaier, D., Livneh, B., Lin, C., Miguez-Macho, G., Niu, G.-Y., et al.: Modeling

- 440 seasonal snowpack evolution in the complex terrain and forested Colorado Headwaters region: A model intercomparison study, Journal of Geophysical Research: Atmospheres, 119, 13–795, 2014.
  - Dai, F., Lee, C., and Ngai, Y. Y.: Landslide risk assessment and management: an overview, Engineering geology, 64, 65–87, 2002.
  - Dietz, A. J., Conrad, C., Kuenzer, C., Gesell, G., and Dech, S.: Identifying changing snow cover characteristics in central Asia between 1986 and 2014 from remote sensing data, Remote Sensing, 6, 12752–12775, 2014.

445 Fawcett, T.: An introduction to ROC analysis, Pattern recognition letters, 27, 861–874, 2006.

Froude, M. J. and Petley, D. N.: Global fatal landslide occurrence from 2004 to 2016, Natural Hazards and Earth System Sciences, 18, 2161–2181, 2018.

Gariano, S. L., Brunetti, M. T., Iovine, G., Melillo, M., Peruccacci, S., Terranova, O., Vennari, C., and Guzzetti, F.: Calibration and validation of rainfall thresholds for shallow landslide forecasting in Sicily, southern Italy, Geomorphology, 228, 653–665, 2015.

- 450 Giannecchini, R., Galanti, Y., Avanzi, G. D., and Barsanti, M.: Probabilistic rainfall thresholds for triggering debris flows in a human-modified landscape, Geomorphology, 257, 94–107, 2016.
  - Guzzetti, F., Peruccacci, S., Rossi, M., and Stark, C. P.: Rainfall thresholds for the initiation of landslides in central and southern Europe, Meteorology and atmospheric physics, 98, 239–267, 2007.

Hamm, A., Arndt, A., Kolbe, C., Wang, X., Thies, B., Boyko, O., Reggiani, P., Scherer, D., Bendix, J., and Schneider, C.: Intercomparison of

455 Gridded Precipitation Datasets over a Sub-Region of the Central Himalaya and the Southwestern Tibetan Plateau, Water, 12, 3271, 2020. Hanssen, A. and Kuipers, W.: ON THE RELATIONSHIP BETWEEN THE FREQUENCY OF RAIN AND VARIOUS METEOROLOG-ICAL PARAMETERS.(WITH REFERENCE TO THE PROBLEM OF OBJECTIVE FORECASTING)., Koninklijk Nederlands Meteorologisch Instituut, 1965.

Havenith, H.-B., Strom, A., Torgoev, I., Torgoev, A., Lamair, L., Ischuk, A., and Abdrakhmatov, K.: Tien Shan geohazards database: Earthquakes and landslides, Geomorphology, 249, 16–31, 2015a.

- Havenith, H.-B., Torgoev, A., Schlögel, R., Braun, A., Torgoev, I., and Ischuk, A.: Tien Shan geohazards database: Landslide susceptibility analysis, Geomorphology, 249, 32–43, 2015b.
- Ilyasov, S., Zabenko, O., Gaydamak, N., Kirilenko, A., Myrsaliev, N., Shevchenko, V., and Penkina, L.: Climate profile of the Kyrgyz Republic, The State Agency for Environmental Protection and Forestry under the Government of the Kyrgyz Republic and The United Nations Development Programme: Bishkek, Kyrgyzstan, 2013.
- Jarraud, M.: Guide to meteorological instruments and methods of observation (WMO-No. 8), World Meteorological Organisation: Geneva, Switzerland, 29, 2008.

465

- Jia, G., Tang, Q., and Xu, X.: Evaluating the performances of satellite-based rainfall data for global rainfall-induced landslide warnings, Landslides, 17, 283–299, 2020.
- 470 Khan, Y. A., Lateh, H., Baten, M. A., and Kamil, A. A.: Critical antecedent rainfall conditions for shallow landslides in Chittagong City of Bangladesh, Environmental Earth Sciences, 67, 97–106, 2012.
  - Kirschbaum, D. and Stanley, T.: Satellite-based assessment of rainfall-triggered landslide hazard for situational awareness, Earth's future, 6, 505–523, 2018.
  - Kirschbaum, D., Stanley, T., and Zhou, Y.: Spatial and temporal analysis of a global landslide catalog, Geomorphology, 249, 4–15, 2015.
- 475 Kirschbaum, D. B., Adler, R., Hong, Y., Hill, S., and Lerner-Lam, A.: A global landslide catalog for hazard applications: method, results, and limitations, Natural Hazards, 52, 561–575, 2010.
  - Koren, V., Schaake, J., Mitchell, K., Duan, Q.-Y., Chen, F., and Baker, J.: A parameterization of snowpack and frozen ground intended for NCEP weather and climate models, Journal of Geophysical Research: Atmospheres, 104, 19569–19585, 1999.
- Leonarduzzi, E., Molnar, P., and McArdell, B. W.: Predictive performance of rainfall thresholds for shallow landslides in Switzerland from
   gridded daily data, Water Resources Research, 53, 6612–6625, 2017.
  - Li, D., Yang, K., Tang, W., Li, X., Zhou, X., and Guo, D.: Characterizing precipitation in high altitudes of the western Tibetan plateau with a focus on major glacier areas, International Journal of Climatology, 40, 5114–5127, 2020.
  - Marra, F.: Rainfall thresholds for landslide occurrence: systematic underestimation using coarse temporal resolution data, Natural Hazards, 95, 883–890, 2019.
- 485 Marra, F., Nikolopoulos, E., Creutin, J., and Borga, M.: Space-time organization of debris flows-triggering rainfall and its effect on the identification of the rainfall threshold relationship, Journal of Hydrology, 541, 246–255, 2016.
  - Marra, F., Destro, E., Nikolopoulos, E. I., Zoccatelli, D., Creutin, J. D., Guzzetti, F., and Borga, M.: Impact of rainfall spatial aggregation on the identification of debris flow occurrence thresholds, Hydrology and Earth System Sciences, 21, 4525–4532, 2017.
- Maussion, F., Scherer, D., Mölg, T., Collier, E., Curio, J., and Finkelnburg, R.: Precipitation seasonality and variability over the Tibetan
  Plateau as resolved by the High Asia Reanalysis, Journal of Climate, 27, 1910–1927, 2014.
  - Minder, J. R., Letcher, T. W., and Skiles, S. M.: An evaluation of high-resolution regional climate model simulations of snow cover and albedo over the Rocky Mountains, with implications for the simulated snow-albedo feedback, Journal of Geophysical Research: Atmospheres, 121, 9069–9088, 2016.
- Mostbauer, K., Kaitna, R., Prenner, D., and Hrachowitz, M.: The temporally varying roles of rainfall, snowmelt and soil moisture for debris flow initiation in a snow-dominated system, Hydrol. Earth Syst. Sci, 22, 3493–3513, 2018.
  - Nikolopoulos, E., Borga, M., Creutin, J., and Marra, F.: Estimation of debris flow triggering rainfall: Influence of rain gauge density and interpolation methods, Geomorphology, 243, 40–50, 2015.

Nikolopoulos, E., Destro, E., Maggioni, V., Marra, F., and Borga, M.: Satellite rainfall estimates for debris flow prediction: an evaluation based on rainfall accumulation–duration thresholds, Journal of Hydrometeorology, 18, 2207–2214, 2017.

- 500 Nikolopoulos, E. I., Crema, S., Marchi, L., Marra, F., Guzzetti, F., and Borga, M.: Impact of uncertainty in rainfall estimation on the identification of rainfall thresholds for debris flow occurrence, Geomorphology, 221, 286–297, 2014.
  - Ou, T., Chen, D., Chen, X., Lin, C., Yang, K., Lai, H.-W., and Zhang, F.: Simulation of summer precipitation diurnal cycles over the Tibetan Plateau at the gray-zone grid spacing for cumulus parameterization, Climate Dynamics, 54, 3525–3539, 2020.
- Ozturk, U., Pittore, M., Behling, R., Roessner, S., Andreani, L., and Korup, O.: How robust are landslide susceptibility estimates?, Landslides, pp. 1–15, 2020.
  - Palmer, J.: A slippery slope: Could climate change lead to more landslides?, Eos, 101, https://doi.org/10.1029/2020EO151418, published on 23 November 2020, 2020.
  - Peres, D. J., Cancelliere, A., Greco, R., and Bogaard, T. A.: Influence of uncertain identification of triggering rainfall on the assessment of landslide early warning thresholds, Natural Hazards and Earth System Sciences, 18, 633–646, 2018.
- 510 Piciullo, L., Gariano, S. L., Melillo, M., Brunetti, M. T., Peruccacci, S., Guzzetti, F., and Calvello, M.: Definition and performance of a threshold-based regional early warning model for rainfall-induced landslides, Landslides, 14, 995–1008, 2017.
  - Postance, B., Hillier, J., Dijkstra, T., and Dixon, N.: Comparing threshold definition techniques for rainfall-induced landslides: A national assessment using radar rainfall, Earth Surface Processes and Landforms, 43, 553–560, 2018.

Prenner, D., Kaitna, R., Mostbauer, K., and Hrachowitz, M.: The value of using multiple hydrometeorological variables to predict temporal
debris flow susceptibility in an alpine environment, Water Resources Research, 54, 6822–6843, 2018.

- Pritchard, D. M., Forsythe, N., Fowler, H. J., O'Donnell, G. M., and Li, X.-F.: Evaluation of Upper Indus near-surface climate representation by WRF in the high Asia refined analysis, Journal of Hydrometeorology, 20, 467–487, 2019.
  - Rodwell, M. J., Richardson, D. S., Hewson, T. D., and Haiden, T.: A new equitable score suitable for verifying precipitation in numerical weather prediction, Quarterly Journal of the Royal Meteorological Society, 136, 1344–1363, 2010.
- 520 Rossi, M., Luciani, S., Valigi, D., Kirschbaum, D., Brunetti, M., Peruccacci, S., and Guzzetti, F.: Statistical approaches for the definition of landslide rainfall thresholds and their uncertainty using rain gauge and satellite data, Geomorphology, 285, 16–27, 2017.
  - Saponaro, A., Pilz, M., Wieland, M., Bindi, D., Moldobekov, B., and Parolai, S.: Landslide susceptibility analysis in data-scarce regions: the case of Kyrgyzstan, Bulletin of Engineering Geology and the Environment, 74, 1117–1136, 2015.

Scherer, D., Fehrenbach, U., Lakes, T., Lauf, S., Meier, F., and Schuster, C.: Quantification of heat-stress related mortality hazard, vulnera bility and risk in Berlin, Germany, DIE ERDE–Journal of the Geographical Society of Berlin, 144, 238–259, 2013.

- Segoni, S., Piciullo, L., and Gariano, S. L.: A review of the recent literature on rainfall thresholds for landslide occurrence, Landslides, 15, 1483–1501, 2018.
  - Shahabi, H., Khezri, S., Ahmad, B. B., and Hashim, M.: Landslide susceptibility mapping at central Zab basin, Iran: A comparison between analytical hierarchy process, frequency ratio and logistic regression models, Catena, 115, 55–70, 2014.
- 530 Stanley, T. and Kirschbaum, D. B.: A heuristic approach to global landslide susceptibility mapping, Natural hazards, 87, 145–164, 2017. Stanley, T., Kirschbaum, D. B., Pascale, S., and Kapnick, S.: Extreme Precipitation in the Himalayan Landslide Hotspot, in: Satellite Precipitation Measurement, pp. 1087–1111, Springer, 2020.
  - Tewari, M., Chen, F., Wang, W., Dudhia, J., LeMone, M., Mitchell, K., Ek, M., Gayno, G., Wegiel, J., and Cuenca, R.: Implementation and verification of the unified NOAH land surface model in the WRF model, in: 20th conference on weather analysis and forecasting/16th
- 535 conference on numerical weather prediction, vol. 1115, pp. 2165–2170, American Meteorological Society Seattle, WA, 2004.

Tomasi, E., Giovannini, L., Zardi, D., and de Franceschi, M.: Optimization of Noah and Noah_MP WRF land surface schemes in snowmelting conditions over complex terrain, Monthly Weather Review, 145, 4727–4745, 2017.

Torgoev, I., Alioshin, Y. G., and Torgoev, A.: Monitoring landslides in Kyrgyzstan., FOG-Freiberg Online Geoscience, 33, 2012.

Wang, Q., Wang, D., Huang, Y., Wang, Z., Zhang, L., Guo, Q., Chen, W., Chen, W., and Sang, M.: Landslide susceptibility mapping based
on selected optimal combination of landslide predisposing factors in a large catchment, Sustainability, 7, 16653–16669, 2015.

- Wang, X., Tolksdorf, V., Otto, M., and Scherer, D.: WRF-based Dynamical Downscaling of ERA5 Reanalysis Data for High Mountain Asia: Towards a New Version of the High Asia Refined Analysis, International Journal of Climatology, 41, 743–762, 2021.
  - Wieczorek, G. F.: Landslides: investigation and mitigation. Chapter 4-Landslide triggering mechanisms, Transportation Research Board Special Report, 1996.
- 545 Woodcock, F.: The evaluation of yes/no forecasts for scientific and administrative purposes, Monthly Weather Review, 104, 1209–1214, 1976.
  - Zhou, X., Yang, K., Ouyang, L., Wang, Y., Jiang, Y., Li, X., Chen, D., and Prein, A.: Added value of kilometer-scale modeling over the third pole region: a CORDEX-CPTP pilot study, Climate Dynamics, pp. 1–15, 2021.
- Zhuo, L., Dai, Q., Han, D., Chen, N., and Zhao, B.: Assessment of simulated soil moisture from WRF Noah, Noah-MP, and CLM land surface schemes for landslide hazard application, Hydrology and Earth System Sciences, 23, 4199–4218, 2019.

Zimmermann, M.: Murganggefahr und Klimaänderung-ein GIS-basierter Ansatz, vdf Hochschulverlag AG, 1997.