

Attribution of the Australian bushfire risk to anthropogenic climate change

Geert Jan van Oldenborgh¹, Folmer Krikken¹, Sophie Lewis², Nicholas J. Leach³, Flavio Lehner^{4,5,6}, Kate R. Saunders⁷, Michiel van Weele¹, Karsten Haustein⁸, Sihan Li^{8,9}, David Wallom⁹, Sarah Sparrow⁹, Julie Arrighi^{10,11}, Roop P. Singh¹⁰, Maarten K. van Aalst^{10,12,13}, Sjoukje Y. Philip¹, Robert Vautard¹⁴, and Friederike E. L. Otto⁸

¹Royal Netherlands Meteorological Institute (KNMI), De Bilt, Netherlands

²University of New South Wales, Canberra, ACT, Australia

³Atmospheric, Oceanic and Planetary Physics, Department of Physics, University of Oxford, Oxford, U.K.

⁴Department of Earth and Atmospheric Sciences, Cornell University, Ithaca, USA

⁵Climate and Global Dynamics Laboratory, National Center for Atmospheric Research, Boulder, USA

⁶Institute for Atmospheric and Climate Science, ETH Zürich, Zürich, Switzerland

⁷Delft Institute of Applied Mathematics, Delft University of Technology, Delft, Netherlands

⁸Environmental Change Institute, University of Oxford, Oxford, U.K.

⁹Oxford e-Research Centre, University of Oxford, Oxford, U.K.

¹⁰Red Cross Red Crescent Climate Centre, The Hague, Netherlands

¹¹Global Disaster Preparedness Center, Washington DC, USA

¹²Faculty of Geo-information Science and Earth Observation, University of Twente, Enschede, Netherlands

¹³International Research Institute for Climate and Society, Columbia University, New York, USA

¹⁴Institut Pierre-Simon Laplace, France

Correspondence: G. J. van Oldenborgh (oldenborgh@knmi.nl)

Abstract. Disastrous bushfires during the last months of 2019 and January 2020 affected Australia, raising the question to what extent the risk of these fires was exacerbated by anthropogenic climate change. To answer the question for southeastern Australia, where fires were particularly severe, affecting people and ecosystems, we use a physically-based index of fire weather, the Fire Weather Index, long-term observations of heat and drought, and eleven large ensembles of state-of-the-art climate models. We find large trends in the Fire Weather Index in the ERA5 reanalysis since 1979, and a smaller but significant increase by at least 30% in the models. Therefore, we find that climate change has induced a higher weather-induced risk of such an extreme fire season. This trend is mainly driven by the increase of temperature extremes. In agreement with previous analyses we find that heat extremes have become more likely by at least a factor of two due to the long-term warming trend. However, current climate models overestimate variability and tend to underestimate the long-term trend in these extremes, so the true change in the likelihood of extreme heat could be larger, suggesting that the attribution of the increased fire weather risk is a conservative estimate. We do not find an attributable trend in either extreme annual drought or the driest month of the fire season September–February. The observations, however, show a weak drying trend in the annual mean. For the 2019/20 season more than half of the July–December drought was driven by record excursions of the Indian Ocean dipole and Southern Annular Mode, factors which are included in the analysis here. The study reveals the complexity of the 2019/20 bushfire event, with some, but not all drivers showing an imprint of anthropogenic climate change. Finally, the study concludes with a

qualitative review of various vulnerability and exposure factors that each play a role, along with the hazard, in increasing or decreasing the overall impact of the bushfires.

1 Introduction

2019 was the warmest and driest year in Australia since standardized temperature and rainfall observations began (in 1910 and 1900), following two already dry years in large parts of the country. These conditions, driven partly by a strong positive Indian Ocean Dipole from the middle of the year onwards and a large-amplitude negative excursion of the Southern Annular Mode, led to weather conditions conducive to bushfires across the continent and so the annual bushfires were more widespread and intense and started earlier in the season than usual ([Bureau of Meteorology, Annual Climate Statement 2019](#)). The bushfire activity across the states of Queensland (QLD), New South Wales (NSW), Victoria (VIC), South Australia (SA), Western Australia (WA) and in the Australian Capital Territory (ACT) was unprecedented in terms of the area burnt in densely populated regions.

In addition to the unprecedented nature of this event, its impacts to date have been disastrous ([Reliefweb Australia: bushfires, 2020](#)). There have been at least 34 fatalities as a direct result of the bushfires and the resulting smoke caused hazardous air quality adversely affected millions of residents in cities in these regions. About 5,900 buildings have been destroyed. There are estimates that between 0.5 and 1.5 billion wild animals lost their lives, along with tens of thousands of livestock. The bushfires are having an economic impact (including substantial insurance claims, e.g., [PERILS press release](#)), as well an immediate and long term health impact to the people exposed to smoke and dealing with the psychological impacts of the fires ([Finlay et al., 2012](#)).

It has at times been difficult for emergency services to protect or evacuate some communities due to the pace at which the bushfires have spread, sometimes forcing residents to flee to beaches and lakes to await rescue. Interruptions of the supply of power, fuel and food supplies have been reported and road closures have been common. This has resulted in total isolation of some communities, or only accessible by air or sea when smoke conditions allow ([Reliefweb Australia: bushfires, 2020](#)).

It is well-established that wildfire smoke exposure is associated with respiratory morbidity ([Reid et al., 2016](#)). Additionally, fine particulate matter in smoke may act as a triggering factor for acute coronary events (such as heart attack-related deaths) as found for previous fires in southeast Australia ([Haikerwal et al., 2015](#)). As noted by [Johnston and Bowman \(2014\)](#), increased bushfire-related risks in a warming climate have significant implications for the health sector, including given measurable increases in illness, hospital admissions, and deaths associated with severe smoke events.

Based on the recovery of areas following previous major fires, such as Black Saturday in Victoria in 2009, these impacts are likely to affect people, ecosystems and the region for a substantial period to come.

The satellite image in Fig. 1 shows the severity of the fires between October, illustrating two regions with particularly severe events in the South West and South East. We focus our analysis on the South East due to the affected population centres and the concomitant drought in this region. The grass fires in the non-forested areas have completely different characteristics and are not considered here.




Figure 1. Moderate Resolution Imaging Spectroradiometer (MODIS) active fire data (Collection 6, near real time and standard products) showing the severity of bushfires from 1 October 2019 to 10 January 2020 with the most severe fires being depicted in red. The image also shows the forested areas in blue. The polygon shows the area analysed in this article.

Wildfires in general are one of the most complex weather-related extreme events (Sanderson and Fisher, 2020) with their occurrence depending on many factors including the weather conditions conducive to fire at the time of the event and also on the availability of fuel, which in turn depends on rainfall, temperature and humidity in the weeks, months and sometimes even years preceding the actual fire event. In addition, ignition sources and type of vegetation play an important role. The types of vegetation depend on the long-term climatology, but do not vary on the annual and shorter time scales we consider, and the dry thunderstorms providing a large fraction of the ignition sources are too small to analyse with climate models. In this analysis we therefore only consider the influence of weather and climate on the fire risk, excluding ignition sources, types of vegetation, and weather caused by the fires such as pyrocumunimbus development. There is no unified definition of what fire weather consists of as the relative importance of different factors depends on the climatology of the region. For instance, fires in grasslands in semi-arid regions behave very differently than those in temperate forests. There are a few key meteorological variables that are important: temperature, precipitation, humidity and wind (speed as well as direction). Fire danger indices are derived from these variables either using physical models or empirical relationships between these variables and fire occurrence, including observed factors such as the rate of spread of fires and measurements of fuel moisture content with different sets of weather conditions.

Southeast Australia experiences a temperate climate and on the eastern seaboard hot summers are interspersed with intense rainfall events, often linked with ‘east-coast lows’ (Pepler et al., 2014). Bushfire activity historically commences in the Austral spring (September-November) in the north and summer (December-February) in the south (Clarke et al., 2011). In Australia the Forest Fire Danger Index (FFDI, McArthur, 1966, 1967; Noble et al., 1980) is commonly used for indicating dangerous weather conditions for bushfires, including for issuing operational forecasts during the 2019/20 summer. The index is based on temperature, humidity and wind speed on a given day as well as a drought-factor which is based on antecedent temperature and rainfall.

Bushfire weather risk, as characterised by the FFDI, has increased across much of Australia in recent decades (Clarke et al., 2013; Dowdy, 2018; Harris and Lucas, 2019). Similar, increasing trends in fire weather conditions over southern Australia have been identified in other studies, both for the FFDI (e.g., Dowdy, 2018) and for indices representing pyroconvective processes (Dowdy and Pepler, 2018). These observed trends over southeast Australia are broadly consistent with the projected impacts of climate change (e.g., Clarke et al., 2011; Dowdy et al., 2019). For individual fire events, studies have shown that it can be difficult to separate the influence of anthropogenic climate change from that of natural variability (e.g., Hope et al., 2019; Lewis et al., 2019).

An alternative index is the physically based Canadian Fire Weather Index (FWI) that also includes the influence of wind on the fuel availability (Dowdy, 2018). The latter is achieved by modelling fuel moisture on three different depths including the influence of humidity and wind speed on the upper fuel layer (Krikken et al., 2019). While the FWI was originally developed specifically for the Canadian forests, the physical basis of the models allows it to be used for many different climatic regions of the world (e.g., Camia and Amatulli, 2009; Dimitrakopoulos et al., 2011), and has been shown to provide a good indication of the occurrence of previous extreme fire events in the Southeastern Australian climate (Dowdy et al., 2009). A study on the emergence of the fire weather anthropogenic signal from noise indicated that this is expected around 2040 for Southern

Australia (Abatzoglou et al., 2019) using the FWI. In this study we also consider the Monthly Severity Rating (MSR), which is derived from the FWI and reflects better how difficult a fire is to suppress (Shabbar et al., 2011). A more detailed analysis of the FWI in the context of bushfires in southeastern Australia is given in section 2.1.

As the fire risk indices depend on heat and drought and these were also extreme in 2019/2020, we also consider these factors separately. Previous attribution studies on Australian extreme heat at regional scales have generally indicated an influence from anthropogenic climate change. The ‘angry summer’ of 2012/2013—which until 2018/2019 was the hottest summer on record—was found to be at least 5 times more likely to occur due to human influence (Lewis and Karoly, 2013). The frequency and intensity of heatwaves during this summer were also found to increase (Perkins et al., 2014). Other attribution assessments that found an attributable influence on extreme Australian heat include the May 2014 heatwave (Perkins and Gibson, 2015), the record October heat in 2015 (Hope et al., 2016), and extreme Brisbane heat during November 2014 (King et al., 2015a). However, at small spatial scales, human influence on extreme heat is sometimes less clear, as in Melbourne in January 2014 (Black et al., 2015). It is worth noting that Lewis et al. (2019) found that the temperature component of the extreme 2018 Queensland fire weather had an anthropogenic influence, while no clear influence was detected on the February 2017 extreme fire weather over Eastern Australia (Hope et al., 2019). We are not aware of any extreme event attribution studies on Australian drought.

Thus, while it is clear that climate change does play an important role in heat and fire weather risk overall, assessing the magnitude of this risk and the interplay with local factors has been difficult. Nevertheless it is crucial to prioritise adaptation and resilience measures to reduce the potential impacts of rising risks.

We perform the analysis of possible connections between the fire weather risk and anthropogenic climate change in three steps. First, we assess the trends in extreme temperature and conduct an attribution study using the annual maximum of the seven-day moving average of daily maximum temperatures corresponding to the time scale chosen for the FWI (Section 3). Second, we undertake the same analysis but for meteorological drought (i.e., defined purely as a lack of rainfall) in two time windows, the annual precipitation as well as the driest month within the fire season, which is September–February in our study area (Section 4). The latter again roughly corresponds to the time scale on which precipitation deficits factor into the FWI, namely 52 days. Third, and most importantly, we conduct an attribution study on the FWI and MSR as indices of the probability of bushfires due to the weather (Section 5). These three attribution studies follow the same protocol used in previous assessments: heat waves: Kew et al. (2019); low precipitation: Otto et al. (2018b); Fire Weather Index: Krikken et al. (2019). The full and generalized event attribution protocol has recently been documented in Philip et al. (2020). In order to condense the lengthy analysis, we provide short overviews of the heat and drought analysis in the main paper, with extensive results in the Supplementary Material, and focus primarily on the FWI and MSR analysis. We also provide a short analysis and discussion of other large scale drivers that were of potential importance during 2019/20, such as El Niño Southern Oscillation (ENSO), the Indian Ocean Dipole (IOD) or the Southern Annular Mode (SAM), in Section 6 and a detailed analysis in the Supplementary Material, section S3. Finally, we briefly discuss non-climate factors, such as exposure and vulnerability, that have contributed to the impacts of the extreme fire season of 2019/20.

2 Data and methods

2.1 General event definition

Since we are investigating several different indicator or driver variables of fire risk, different event definitions are developed for different variables. The details of those definitions are given at the beginning of the respective sections on temperature, precipitation, and fire weather indices (Sections 3–5). General parameters of the event definition are given here.

The fire season (September–February) serves as the general event time window and the region with the most intense fires in 2019/20 in southeastern Australia serves as the general event spatial domain; specifically the land area in the polygon 29 °S, 155 °E; 29 °S, 150 °E; 40 °S, 144 °E and 40 °S, 155 °E (as shown in Fig. 1), which corresponds to the area between the Great Dividing Range and the coast.

The primary way we investigate the connection between anthropogenic climate change and the likelihood and intensity of dangerous bushfire conditions is through the FWI. The FWI provides a reasonable proxy for the burnt area in the extended summer months, with the strongest relationship observed from November to February. Fig. 2 shows both the Spearman rank correlation and the Pearson correlation of FWI with log-transformed burnt area. The 95% confidence intervals are also shown. Given the similarity in the correlation coefficients (r) within their confidence intervals, the log-linear relationship appears to explain equal variability (r^2) to that of the ranks.

To capture spatial variations in the start of the fire season at a given location within the event domain, we take for most quantities first the maximum per grid point over the fire season (September–February) and next the spatial average over the general event domain. This way the events do not need to be simultaneous at separate grid points within the region. We therefore investigate the question how anthropogenic climate change influences the chances of an intense bushfire season, rather than focusing on a single episode of intense bushfires.

In most years only very small areas are burnt, but the observational record also includes events with extremely large areas. Given this, we checked if the burnt area observations were heavy-tailed (Pasquale, 2013). We found that monthly burnt area was not Pareto-distributed and instead is reasonably approximated using a log-normal distribution. This supports using the log-transformation and extrapolating this relationship to the 2019/20 fire season. Temporal detrending of the observations did not alter these conclusions.

2.2 Observational data

The observational data used in this study are described in Sections S1, S2 and 5.3 for heat, drought and the fire weather index respectively, including justifications for including or excluding certain datasets for certain research questions. For the Global Mean Surface temperature (GMST) we use GISTEMP surface temperature (Hansen et al., 2010).




Figure 2. Left: correlation between the logarithm of area burnt ($10 \log(\text{km}^2)$, MODIS, Collection 6) in the event domain and the 7-day maximum Fire Weather Index for each month of the year. The correlations are based on the years 1997 to 2018 and the 95% two-sided confidence interval is based on bootstrapping those years. The horizontal line denotes the 5% significance critical value for a one-sided test of the null hypothesis that the correlation is zero against the alternative hypothesis that the correlation is positive. Right: scatterplot and regression line of the values for each month of the fire season (September–February). The grey lines denote the regression lines for the individual months, the green line for all months in the fire season.

2.3 Model and experiment descriptions

Attributing observed trends to anthropogenic climate change can only be done with physical climate models as they allow isolating different drivers. For this purpose we included as large a set of ocean-atmosphere coupled and atmosphere-only (i.e., sea surface temperature (SST) prescribed) climate model ensembles as we could find within the time constraints of this study in order to obtain estimates of both the uncertainty due to natural variability and the model uncertainty. A selection of large ensembles of climate models from the Coupled Model Intercomparison Project Phase 5 (CMIP5) has been used: CanESM2, CESM1-CAM5, CSIRO Mk3.6.0, EC-Earth 2.3, GFDL CM3, GFDL ESM2M and MPI ESM. In addition, the HadGem3-A N216 attribution model developed in the EUropean CLimate and weather Events: Interpretation and Attribution (EUCLEIA) project, the weather@home distributed attribution project model and the Atmospheric Seasonal Forecasts of the Twentieth Century (ASF-20C) seasonal hindcast ensemble have been used. These last three models are uncoupled and forced with observed historical SSTs and estimates of SSTs as they might have been in a counterfactual world without anthropogenic climate change. Finally, we used the coupled IPSL-CM6A-LR low-resolution CMIP6 ensemble. The GFDL-CM3 and MPI-ESM models that did not have daily data were not used for the extreme heat analysis. For the FWI analysis, which requires daily

name	context	resolution	members	time	reference
ASF-20C	seasonal hindcasts	T255L91 (0.71°)	51	1901–2010	Weisheimer et al. (2017)
CanESM2	CMIP5	2.8°	50	1950–2099	Kirchmeier-Young et al. (2017)
CESM1-CAM5	CMIP5	1°	40	1920–2100	Kay et al. (2015)
CSIRO-Mk3-6-0	CMIP5	1.9°	30	1850–2100	Jeffrey et al. (2013)
EC-Earth	CMIP5	T159 (1.1°)	16	1860–2100	Hazeleger et al. (2010)
GFDL-CM3	CMIP5	2.0°	20	1920–2100	Sun et al. (2018)
GFDL-ESM2M	CMIP5	2.0°	30	1950–2100	Rodgers et al. (2015)
HadGEM3-A	attribution	N216 (0.6°)	15	1960–2015	Ciavarella et al. (2018)
IPSL-CM6A-LR	CMIP6	2.5×1.5°	32	1950–2019	Boucher et al. (2020)
MPI-ESM	CMIP5	1.9°	100	1850–2099	Maher et al. (2019)
weather@home	attribution	N96 (1.8°)	1520 × 2	1987–2017	Guillod et al. (2017)

Table 1. List of climate model ensembles used.

data of relative humidity (RH), temperature, precipitation and wind speed, the list of models used is shortened to CanESM2, CESM1-CAM5, EC-Earth, IPSL-CM6A-LR and weather@home (HadAM3P).

2.4 Statistical methods

The methods employed in this analysis have been used previously for high and low temperatures (van Oldenborgh et al., 2015; King et al., 2015b; van Oldenborgh et al., 2018; Philip et al., 2018a; Kew et al., 2019), extreme precipitation (Schaller et al., 2014; Siswanto et al., 2015; Vautard et al., 2015; Eden et al., 2016; van Oldenborgh et al., 2016; van der Wiel et al., 2017; van Oldenborgh et al., 2017; Eden et al., 2018; Otto et al., 2018a; Philip et al., 2018b), drought (King et al., 2016; Martins et al., 2018; Otto et al., 2018b; Philip et al., 2018c; Uhe et al., 2018) and forest fire weather (Krikken et al., 2019). A paper describing the methods in detail was recently published as Philip et al. (2020).

Changes in the frequency of extreme events are calculated by fitting the data to a statistical distribution. In this study the highest temperature extremes and fire risk-related variables (FWI, MSR) of the fire season are assumed to follow a Generalised Extreme Value (GEV) distribution, which is the distribution that block maxima converge to (Coles, 2001). While our event definition is not exactly block maxima, the GEV fits the data well (see below for more details). The low annual mean precipitation values and lowest monthly precipitation of the fire season are fitted using a Generalized Pareto Distribution (GPD), which describes the exceedance below a low threshold and also allows the specification of a threshold that ensures the PDF is zero for negative precipitation.

The GEV distribution is:

$$P(x) = \exp \left[- \left(1 + \xi \frac{x - \mu}{\sigma} \right)^{-1/\xi} \right], \quad (1)$$

where x the variable of interest, e.g., temperature or precipitation. Here, μ is the location parameter, $\sigma > 0$ is the scale parameter, and ξ is the shape parameter. The shape parameter determines the tail behaviour: a negative shape parameter gives an upper bound to the distribution, for $\xi \geq 1$ the tail is so fat that the mean is infinite. The scale parameter corresponds to the variability in the tail.

The GPD gives a 2-parameter description of the tail of the distribution above a threshold, where the low tail of precipitation is first converted to a high tail by multiplying the variable by -1 . The GPD is then described by:

$$H(u - x) = 1 - \left(1 - \frac{\xi x}{\sigma}\right)^{(-1/\xi)}, \quad (2)$$

with x the temperature or precipitation, u the threshold, σ the scale parameter, and ξ the shape parameter determining the tail behaviour. For the low extremes of precipitation, the fit is constrained to have zero probability below zero precipitation ($\xi < 0, \sigma < u\xi$). Calculations were conducted on the lowest 20% and 30% of the data, which provide a first-order estimate of the influence of using more or less extreme events. We cannot use less data as the maximizations of the likelihood function do not converge anymore, and using more than 30% would not qualify as the ‘lower tail’.

Drought, or low precipitation, is particularly difficult to model using the existing extreme value framework (Cooley et al., 2018). While minima can be modelled by multiplying by -1 (Coles, 2001), the applicability of the underlying extreme value theory assumptions still needs to be validated. In the case of low precipitation, year-on-year autocorrelations are a concern. In southeastern Australia, these serial autocorrelations are approximately $r \approx 0.2$, so although non-zero, do not dominate the drought characteristics. Despite these theoretical limitations, in practice the diagnostic plots show that the Generalised Pareto models are able to describe the data reasonably well. In particular, they respect that precipitation is non-negative. In general this is a difficult problem, and the statistical extremes community is still developing the solutions necessary for modelling drought events (Naveau et al., 2016).

To calculate a trend in transient data, some parameters in these statistical models are made a function of the 4-yr smoothed global mean surface temperature (GMST) anomaly, T' . This smoothing is the shortest that on the one hand reduces the ENSO component of GMST, which is not externally forced and therefore not relevant for the trend, but on the other hand retains as much of the forced variability as possible (Haustein et al., 2019). A longer smoothing time scale would create problems with extrapolation in the highly relevant last few years of the instrumental record. The covariate-dependent function can be inverted and the distribution evaluated for a given year, e.g., a year in the past (with $T' = T'_0$) or the current year ($T' = T'_1$). This provides estimates and confidence intervals of the probabilities for an event at least as extreme as the observed one in these two years, p_0 and p_1 , or expressed as return periods $\tau_0 = 1/p_0$ and $\tau_1 = 1/p_1$. The change in probability between two such years is called the Probability Ratio (PR): $PR = p_1/p_0 = \tau_0/\tau_1$. We also estimate the changes in intensity (including uncertainties) ΔT for temperature, ΔP for drought and ΔFWI .

For extreme temperature we assume that the distribution shifts with GMST: $\mu = \mu_0 + \alpha T'$ or $u = u_0 + \alpha T'$, and $\sigma = \sigma_0$ with α the trend that is fitted together with μ_0 and σ_0 . The shape parameter ξ is also assumed constant. For drought and FWI-related variables we instead make the assumption that the distribution *scales* with GMST, the scaling approximation (Tebaldi and

210 [Arblaster, 2014](#)). In a GEV fit this gives:

$$\begin{aligned}\mu &= \mu_0 \exp(\alpha T' / \mu_0), \\ \sigma &= \sigma_0 \exp(\alpha T' / \mu_0),\end{aligned}\tag{3}$$

and in a GPD fit

$$\begin{aligned}u &= u_0 \exp(\alpha T' / u_0), \\ 215 \quad \sigma &= \sigma_0 \exp(\alpha T' / \mu_0),\end{aligned}$$

with fit parameters σ_0, α and ξ . The threshold u_0 is determined with an iterative procedure and the shape parameter ξ is again assumed constant. The exponential dependence on the covariate is in this case just a convenient way to ensure a distribution that is zero for negative precipitation and has no theoretical justification. For the small trends in this analysis it is similar to a linear dependence.

220 The validity of the other assumption, that the scale parameter or dispersion parameter are constant, is tested by computation of the significance of deviation of a constant of running (relative) variability plots of the observations and model data ([Philip et al., 2020](#)). The analysis of model data is more sensitive to variations of these parameters over time due to the large number of ensemble members but of course assumes the effect of external forcing on the variability is modelled correctly.

For all fits we also estimate 95% uncertainty ranges using a non-parametric bootstrap procedure, in which 1000 derived time series, generated from the original one by selecting random data points with replacement, are analysed in exactly the same way. The 2.5 and 97.5 percentile of the 1000 output parameters (defined as $100i/1001$ with i the rank) are taken as the 95% uncertainty range. For some models with prescribed SSTs or initial conditions (in the case of the seasonal forecast ensemble) the ensemble members are found to not be statistically independent, defined here by a correlation coefficient $r > 1/e$ with $e \approx 2.7182$. In those cases the same procedure is followed except that all dependent time series are entered together in the
230 bootstrapped sample, analogous to the method recommended in [Coles \(2001\)](#) to account for temporal dependencies.

When using a GEV to model tail behaviour, note that taking the spatial average of the annual maxima, does not have the same statistical justification as taking the annual maximum of the spatial average ([Coles, 2001](#)). Given this, the impact of the order of operations in the event definition was examined. For the temperature extremes, we compared the time series where we first take the annual maximum and next the spatial average, to the definition with the order reversed, which can be approximated
235 with a GEV. The Pearson correlation was $r = 0.95$, which is likely due to strong spatial dependence and the concentration of heatwaves at the peak of the seasonal cycle. Therefore, in practice, an approximation with a GEV is not entirely unsuitable for temperature, but caution should be exercised. For the FWI and MSR, the order of operations does make a clear difference. Indeed, we find that the whole distribution is not described well by a GEV for one climate model used (CanESM2). For that model we take block maxima over 5-ensemble member blocks, effectively looking only at the most extreme events. For this
240 part of the distribution the GEV fit agrees with the data points in the return time plot, as expected from taking block maxima.

We evaluate all climate models on the fitting parameters by determining whether the model-derived parameters fall within the uncertainty range of observation-derived parameters. We allow for a mean bias correction, i.e., we only check the scale and

shape parameters σ and ξ . Model biases are accounted for by evaluating the model at the same return time as the value found in the observational analysis. This was found to give better results than applying an additive or multiplicative bias correction to the position parameter μ as it also corrects to first order for biases in the other parameters, especially when the distribution has an upper or lower bound ($\xi < 0$), which is the case in all the cases here.

Finally, estimates of the PR and change in observations and all climate models that pass the evaluation test are combined to give a synthesized attribution statement. First, the observations and reanalyses were combined by averaging the best estimate, lower and upper bounds, as the natural variability is strongly correlated as they are largely based on the same observations (except for the long reanalyses). The difference is added as representation uncertainty (white extensions on light blue bars in Figs. S6, S12, S13 and 6).

Second, the model results were combined by computing a weighted average (using inverse model total variances), as the natural variability in the models, in contrast to the observations, is uncorrelated:

$$\bar{X} = \sum_i X_i / \sigma_{X_i}^2 / \sum_i 1 / \sigma_{X_i}^2, \quad (4)$$

with σ_{X_i} the estimated uncertainty in model i and X_i either the temperature or the logarithm of precipitation, FWI or MSR. The sums are over the N_{mod} models. Using this we can compare the spread expected from the natural variability with the observed spread of the model results using a χ^2 statistics

$$\chi^2 = \sum_i (X_i - \bar{X})^2 / \sigma_{X_i}^2, \quad (5)$$

If $\chi^2/\text{dof} \leq 1$, with dof the number of degrees of freedom, here $N - 1$, the spread of the results is compatible with the uncertainty estimated from the fits due to variability within the climate model and the results can be taken to be independent estimates of X and the weighted average used. However, if $\chi^2/\text{dof} > 1$ the model spread is larger than expected from variability due to sampling of weather noise alone, so a model spread term was added to each model in addition to the weighted average (white extensions on the light red bars, Figures S5, S6), to account for systematic model errors. This term is defined by requiring that $\chi^2/\text{dof} = 1$. The total uncertainty of the models is shown as a bright red bar in these figures. This total uncertainty consists of a weighted mean using the uncorrelated natural variability plus an independent model spread term added to the uncertainty if $\chi^2/\text{dof} > 1$, which we do not divide by $\sqrt{N - 1}$, i.e., we do not assume that by adding more models to the ensemble the model uncertainty decreases. This procedure is similar to the one employed by Ribes et al. (2020).

Finally, observations and models are synthesised into a single mean and uncertainty range. This can only be done when they appear to be compatible. We show two combinations. The first one is computed by neglecting model uncertainties beyond the model spread. The optimal combination is then the weighted average of models and observations, shown as a magenta bar. However, the total model uncertainty is unknown and can be larger than the model spread. We therefore also show the more conservative estimate of an unweighted average of observations and models with a white box in the synthesis plots.

3 Extreme heat

The key takeaways from the attribution analysis of trends in extreme heat are summarized here, while the details are given in the supplementary material, section S1.

Taking advantage of the longer observational record for temperature than for other variables, we analyse the highest 7-day mean maximum temperatures of the year, averaged over the event domain (Fig. 1), from 1910 (the beginning of standardised temperature observations) to 2019.

Observations show that a heatwave as rare as observed in 2019/20 would have been 1 to 2 °C cooler at the beginning of the 20th century (Fig. S6). Similarly, a heatwave of this intensity would have been less likely by a factor of about 10 in the climate around 1900 (Fig. S6). While climate models consistently simulate increasing temperature trends over this time period, they all have some limitations for simulating heat extremes: the variability of 7-day mean maximum temperature is generally too high and the long-term trend is only 1 °C (Fig. S5 and Fig. S6). We can therefore only conclude that anthropogenic climate change has made a hot week like the one in December 2019 more likely by at least a factor of two, but cannot give a best estimate or upper bound due to the model deficiencies limiting our confidence in the exact magnitude of the anthropogenic influence.

The reasons for the apparent model deficiencies in simulating trends and variability in extreme temperatures are not fully understood. In section S3 we show that the temperature variability explained by the Indian Ocean Dipole (IOD) and Southern Annual Mode (SAM) is too small to explain these mismatches as problems in the model representation of these modes of variability. The literature suggests that shortcomings in the coupling to land and vegetation (e.g., Fischer et al., 2007; Kala et al., 2016) and in parametrisation of irrigation (e.g., Thiery et al., 2017; Mathur and AchutaRao, 2019) in the exchange of heat and moisture with the atmosphere, and also in the representation of the boundary layers (e.g., Miralles et al., 2014) are more likely to be the cause of the problems. Given the larger trend in observations than in the models we suspect that climate models underestimate the trend in extreme temperatures due to climate change, although in principle the difference could also be due to a non-climatic driver that affects the trend in observations. The combination of a weaker trend and higher variability in models compared to observations yields an increase in the likelihood of such an event that is much higher in observations than in models.

4 Meteorological drought

The key takeaways from the attribution analysis of trends in low precipitation are summarized here, while the details are given in the supplementary material, section S2. The conclusions below are shown in Fig. S12 for annual mean drought and Fig. S13 for the driest month of the year.

Observations show non-significant trends towards more dry extremes like the record 2019 annual mean and a non-significant trend towards fewer dry months like December 2019 in the fire season (Fig. S12 and Fig. S13). All ten climate models we considered simulate the statistical properties of the observations well (Fig. S10 and Fig. S11). Collectively they show no trend in dry extremes of annual mean precipitation nor in the driest month of the fire season (September–February). We conclude

305 that there is no evidence for an attributable trend in either kind of meteorological drought extremes like the ones observed in 2019.

5 Fire risk indices

5.1 The fire weather of 2019/20

As discussed in the introduction, the fire risk as described by fire weather indices was extreme in the study domain in the
310 2019/20 fire season. The domain was chosen to encompass these fires and therefore the 2019/20 event cannot be included in the statistical analysis.

5.2 Temporal Event Definition

We choose two event definitions in order to represent two important aspects of the event, namely the intensity and the duration. For intensity, we first select the maximum FWI of a 7-day moving average over the fire season (September–February) for every
315 grid point over the study region, after which we compute the spatial average, hereafter FWI7x-SM (seasonal maximum). The 7-day time scale was chosen based on a good correlation with the area burnt (see Fig. 2) and good correspondence with area burnt in other forest fire attributions studies (Krikken et al., 2019).

For duration, we consider the monthly severity rating (MSR). The MSR is the monthly averaged value of the daily severity rating (DSR), which in turn is a transformation of the FWI ($DSR = 0.0272FWI^{1.71}$). The DSR reflects better how difficult
320 a fire is to suppress, while the MSR is a common metric for assessing fire weather on monthly time scales (Van Wagner, 1970). For this study, we select the maximum value of the MSR during the fire season over the study area (MSR-SM). In contrast to the FWI7x-SM, we first apply a spatial average of the study area and then select the maximum value per fire season. This event definition focuses more on changes in extreme fire weather for longer time scales and larger integrated areas than FWI7x-SM. Note that neither of the two event definitions includes ignition sources or small-scale meteorological factors such
325 as pyrocumunimbus development that could enhance the fires.

5.3 Observational analysis: return time and trend

For the observational analysis we use the fifth generation European Centre for Medium-Range Weather Forecasts (ECMWF) Atmospheric Reanalysis (ERA5) dataset 1979–January 2020 (Hersbach et al., 2019). This reanalysis dataset is heavily constrained by observations, thus providing one of the best estimates of the actual state of the atmosphere for all the variables
330 needed to compute the FWI over the study area. Other reanalyses did not yet include the full 2019/20 event at the time of the analysis.

Fig. 3 shows the time series of the highest 7-day-mean FWI averaged over the study area. Both for the FWI7x-SM and MSR-SM the event is the highest over the 1979–2020 time period. Note that for the MSR-SM, the value is considerably more extreme than for the FWI7x-SM. The GEV-fits (Fig. 3, right) illustrate this further, with return times in excess of 1000 years.




Figure 3. Left: time series with 10-yr running mean of the area-average of the highest 7-day mean Fire Weather Index in September–February (top) and Monthly Severity Rating September–February maximum (bottom). Right: stationary GEV fit to these data, the dots represent the ordered years and the grey bands the 95% uncertainty ranges.




Figure 4. Fit of a GEV that scales with the smoothed GSMT (Eqs 1,3) of the highest 7-day mean FWI computed from the ERA5 reanalysis, averaged over the index region. Top: observations (blue symbols), location parameter μ (thick line and uncertainties in 1900 (extrapolated) and 2019/20) and the 6 and 40 yr return values (thin lines). The purple square denotes the 2019/20 value, which is not included in the fit. Bottom: return time plot with fits for the climates of 1900 (blue lines with 95% confidence interval) and 2019 (red lines), the purple line denotes the 2019/20 event. The observations are plotted twice, shifted down to the climate of 1900 (blue stars) and up to the climate of 2019 (red pluses) using the fitted dependence on smoothed global mean temperature so that they can be compared with the fits for those years.

335 A fit allowing for scaling with the smoothed GMST gives a significant trend in the FWI7x-SM (Fig. 4). This fit gives a return
time for the 2019/20 fire season of about 31 yr (4 to 500 yr) in the current climate and more than 800 yr extrapolated to the
climate of 1900. This corresponds to an infinite PR, with a lower bound of four. The return time for MSR-SM is undefined and
is thus estimated to be 100 years. For the climate model analysis we thus use return times of 31 years for the FWI7x-SM and
100 years for MSR-SM to determine the event thresholds in individual climate models.

340 5.4 Model evaluation

We use four climate models with large ensembles, leaving out CESM1-CAM5 because of its failure to represent heat extremes
(see section 3). This is fewer than for the drought and heat analysis, because the FWI requires four daily input variables,




Figure 5. Model verification for FWI (top) MSR (bottom). The left figures show the dispersion parameter σ/μ and the right figures the shape parameter ξ . The bars denote the 95% uncertainty ranges.

which are not available for all models. In contrast to the heat extremes and drought analyses, the fits to the model output use as covariate the model GMST. We also take as reference climates based on GMST rather than years. The years at which the climate is evaluated are taken from the 1.1 °C temperature increase for the present-day climate and the 2 °C increase for the future reference climate and not 2019 and 2060. As the fits are invariant under a scaling of the covariate this does not make much difference.

First the models are evaluated on how well they represent the extremes of FWI7x-SM and MSR-SM. This is quantified by the dispersion parameter σ/μ and shape parameter ξ of the GEV fit for the present-day climate. We do not check the position parameter μ , assuming a multiplicative bias correction can be applied.

Fig. 5 gives an overview of these parameters. Preferably, we would like the parameters to lie within the observational uncertainty from ERA5. For the dispersion parameter CanESM2 and weather@home fall within the observational uncertainty of the FWI. The other two models (EC-Earth and IPSL CM6) show too much variability relative to the mean. The same holds for the shape parameter. This implies that it is difficult to draw strong conclusions from the model data, given that they do not accurately represent the extremes of FWI7x-SM. In particular, the models with too much variability will underestimate the probability ratios. We continue with all four models, but keep these problems in mind.

The MSR is simulated better: all model dispersion and shape parameters lie within the large observational uncertainties, although they largely disagree with one another on the dispersion parameter.

5.5 Multi-model attribution and synthesis

The model results are summarized by their PR, i.e., how more or less likely such an event will be for present or future climate, relative to the early 20th century.




Figure 6. Top: PR for a FWI as high as observed in 2019/20 or higher, (top) from 1920/21 to 2019/20 and (bottom) from 1900 to a climate globally 2 °C warmer than 1920. The last row is the weighted average of all models, the spread of which is consistent with only natural variability. Bottom: same for a 2 °C climate (GMST change from late 19th century).




Figure 7. As Fig. 6 but for the Monthly Severity Rating (MSR).

Figs. 6 and 7 show the change in probability for both the FWI7x-SM, and the MSR-SM from 1920 to 2019 (denoting the 2019/20 fire season). For the FWI7x-SM, all models agree on an increased probability for such an event in the present climate relative to the early 20th century, although the trend is not significant at $p < 0.05$ two-sided for one of the models, CanESM2.

365 As the spread of the models is compatible with natural variability ($\chi^2/\text{dof} < 1$) we take a weighted average across the models to synthesize them (Fig. 6). This shows that such an event has become about 80% more likely in the models, with a lower bound of 30%. Note that all models severely underestimate the increased risk compared to ERA5, which has a lower bound PR of a factor four relative to 1920 (extrapolated), above the upper end of the model average. Note that the ERA5 value is probably biased high as the positive contribution of trend towards a drier climate over 1979–2019 is not present over 1900–2019, see

370 sections S2 and 5.6.

For a future climate of 2 °C warming above pre-industrial we find that such events become about eight times more likely in the models, with a lower bound of about four times more likely. Note that the estimate of future climate is only based on two climate models, CanESM2 and EC-Earth, due to the absence of future data for the others.

For the MSR-SM the models on average show about a doubling of probability for the present climate relative to the early

375 20th century (Fig. 7). However, this trend is not significant as the lower bound is 0.8, i.e., a decreased probability is also possible within the two-sided 95% uncertainty range. In the fit to the ERA5 data we include 2019, as otherwise the probability of the event occurring in the current climate would be zero, contrary to the fact that it did occur. This fit shows much higher probability ratios, with a lower bound of a factor nine. As there is no overlap with the model results we cannot combine the model and observational results, but only give a conservative lower bound as an observations-model synthesis result. For a

380 future climate relative to the early 20th century climate the models show an increase in probability of about four times, with a lower bound of two.

5.6 Interpretation

The underestimation of the observed trend in fire weather indices in all models and the tendency for too much variability in some models is reminiscent of the extreme temperature results in sections 3 and S1.

385 In order to better understand which input variables cause the long-term increase in FWI7x-SM and thus the contribution to the 2019/20 FWI7x-SM value, we study the input variables to FWI7x-SM separately for each model as well as observations. For precipitation we use the cumulative precipitation (90 days) prior to each FWI7x-SM value. We calculate the change from the early 20th century to present day in each input variable to estimate its long-term change, which we then subtract from that variable's observed value in 2019/20. We then recalculate FWI7x-SM, but using each detrended individual input variables

390 in turn. Each of these newly calculated FWI7x-SM values thus illustrates the influence of the long-term trend in a particular input variable onto the observed 2019/20 FWI7x-SM value. This procedure is applied to models and ERA5. In the models, the ensemble mean change is used to estimate an individual variable's long-term trend, whereas in ERA5 a regression of each variable onto GMST is used to estimate its early 20th century value. The results of this analysis for the 2019/20 FWI7x-SM value are shown in Fig. 9.




Figure 8. ERA5 volumetric soil water from multiple levels. The data represents the spatial average over the study area.

395 The sum of the contributions from individual input variables to the 2019/20 FWI7x-SM anomaly match the effect of changing all variables at the same time, so can be considered linearly additive Fig. 9.. The underestimation of the extreme temperature trends in the climate models carries over into this analysis, such that the temperature contribution to the observed 2019/20 value is underestimated. Despite this underestimation, temperature emerges as the most important variable in EC-Earth and weather@home, as it explains roughly half of the increase in FWI. For IPSL, the simulated temperature increase explains about 400 a third of the FWI7x-SM increase, together with wind and RH. CanESM2 behaves differently, where it is mainly the decrease of RH that explains the higher FWI7x-SM. Most, but not all models analysed here therefore derive the increase in FWI7x-SM largely from the increase in temperature extremes.

In ERA5 the increase in temperature also appears to be the most important explanatory variable, followed by a decrease in RH and precipitation. As we did not find a significant trend in precipitation over longer time periods, we hypothesize this trend 405 to be due to natural variability over the short 1979–2018 period in ERA5. We explicitly verified that the dependence of the FWI7x-SM on temperature is almost linear in a range of ± 5 K around the reanalysis value (not shown). Further, volumetric soil water (Fig. 8) at multiple soil layers from ERA5 suggests that, despite the soil already being very dry in 2018 and into 2019, the 2019/20 austral spring-summer drought caused a further drying of the soil in the study area. This suggests that the late 2019 drought and high temperatures did indeed cause an additional increase in fire risk over preceding years.

410 5.7 Conclusions fire risk indices

The FWI7x-SM as computed from the ERA5 reanalysis as an approximation to the real world shows that the 2019/20 values were exceptional. They have a significant trend towards higher fire weather risk since 1979. Compared with the climate of 1900, the probability of a FWI7x-SM as high as in 2019/20 has increased by more than a factor of four. For the MSR-SM the probability has increased by more than a factor of nine.




Figure 9. Sensitivity analysis of the FWI7x-SM to changes in individual contributions from relative humidity (RH, wind, temperature and precipitation). The relative increases or decreases for the individual variables of the climate models are based on the average change in input variables between present day and the early 20th century climate (values above bars). For ERA5 the changes are based on a linear regression of the respective variable onto GMST for the years 1979 to 2018, and then extrapolated to the early 20th century. These changes are subtracted from the 2019 ERA5 data, after which the FWI is recomputed, where ΔFWI is the original FWI minus the altered FWI. In the ‘all’ experiment all input variables are changed simultaneously. ‘Sum’ is the sum of all the individual changes in FWI.

415 The four climate models investigated show that the probability of a Fire Weather Index this high has increased by at least 30% since 1900 as a result of anthropogenic climate change. As the trend in extreme temperature is a driving factor behind this increase and the climate models underestimate the observed trend in extreme temperature, the attributable increase in fire risk could be much higher. This is also reflected by a larger trend in the FWI7x-SM in the reanalysis compared to models. The MSR-SM increased by a factor of two in the models since 1900, although this increase is not significantly different from zero. As with FWI7x-SM, the attributable increase is likely higher due to the model underestimation of temperature trends and overestimation of variability in TX7x.

Projected into the future, the models project that a FWI7x-SM as high as 2019/20 would become at least four times more likely with a 2 °C temperature rise, compared with 1900. Due to the model limitations described above this could also be an underestimate.

425 6 Other drivers

The attribution statements presented in this paper are for events defined as meeting or exceeding the threshold set by the 2019/20 fire season and thus assessing the overall effect of human-induced climate change on these kinds of events. In individual years,

however, large scale climate system drivers can have a higher influence on fire risk than the trend. A detailed analysis of the influence of ENSO, the IOD and SAM is presented in section S3.

430 Besides the influence of anthropogenic climate change, the particular 2019 event was made much more severe by a record positive excursion of the Indian Ocean Dipole and a very strong negative anomaly of the Southern Annular Mode, which likely contributed substantially to the precipitation deficit. We did not find a connection of either mode to heat extremes. More quantitative estimates will require further analysis and dedicated model experiments, as the linearity of the relationship between these indices and the regional climate is not verifiable from observations alone.

435 7 Vulnerability and exposure

At least 19.4 million hectares of land have burnt as a result of the Black Summer Bush Fires of 2019/20 (CDP, 2020). This has resulted in 34 direct deaths and the destruction of 5,900 residential and public structures (IFRC, 2020). Nearly 80% of Australians reported being impacted in some way by the bushfires (The Conversation, 17 February 2020). In Sydney, Canberra and a number of other cities, towns and communities air quality levels reached hazardous levels (NY Times, 3 January 2020).
440 Over 65,000 people registered on Australian Red Cross' reunification site to look for friends and family, or to let loved ones know that they were alright (Australian Red Cross, 2020). It is estimated that over 1.5 billion animals have died nationally (Reliefweb Australia: bushfires, 2020). These impacts are not only hazard-related but also related to various vulnerability and exposure factors that each play a role in increasing or decreasing risk and impacts. Vulnerability is defined as 'The propensity or predisposition to be adversely affected. Vulnerability encompasses a variety of concepts and elements including sensitivity
445 or susceptibility to harm and lack of capacity to cope and adapt' (Agard et al., 2014). It can also be defined as 'the diminished capacity of an individual or group to anticipate, cope with, resist and recover from the impact of a natural or man-made hazard' (IFRC, 1 March 2020). Exposure is defined as 'The presence of people, livelihoods, species or ecosystems, environmental functions, services, and resources, infrastructure, or economic, social, or cultural assets in places and settings that could be adversely affected' (Agard et al., 2014).

450 Bushfires have been a part of the Australian landscape for millions of years, and are an ever-present risk for people living in rural and peri-urban areas surrounded by vegetation/bush/grasslands. In recent decades, significant bushfires occurred in 1974/75, 1983, 2002/03, and 2009, some of them including grass fires, which can have different drivers to forest fires like 2019/20. This frequent occurrence of severe bushfires, with records extending back to the 1850s, has resulted in robust preparedness and emergency management systems which serve to reduce risk and aid in swift response. Comprehensive risk
455 assessments are undertaken at local council level, and bushfire preparedness and contingency plans have been in place in most high-risk areas for decades. However, these systems were severely strained in the Black Summer Bushfires.

7.1 Excess morbidity and mortality

The time of publication is too soon for a robust estimate of excess morbidity and mortality specific to the 2019/20 Australian Bushfires. Such analysis is typically available weeks to years following the end of an event. However, the combined impacts

460 of extreme heat and air pollution can be deadly, as seen in the compounded heatwave and wildfire events in 2010 in Russia or 2015 in Indonesia (Shaposhnikov et al. (2014), Nature, 12 Aug 2010, Koplitz et al. (2016)). Those most at risk are the elderly, people with pre-existing cardiovascular, pulmonary and/or renal conditions as well as young children, as exposure to wildfire smoke can have acute respiratory effects. Health officials in New South Wales reported a 34% spike in emergency room visits for asthma and breathing problems between 30 December 2019 and 5 January 2020 (Washington Post, 12 Jan 2020). One 465 study of hospital admissions in Sydney, Australia from 1994 to 2010 found that days with extreme bushfires air pollution (as measured by PM10) resulted in 1.24% admission increase for every $10 \mu\text{gm}^{-3}$ (Morgan et al., 2010). On the other hand, it should be noted that Australia is a country with a robust healthcare system which significantly reduces vulnerability to the short and long-term consequences of smoke and extreme heat.

There is also a need for increased mental health services in the days, weeks and years following severe bushfires. As of 470 January 2020 the Australian Government announced 76 million AUD in Mental Health funding (ABC, 12 January 2020). A study of the 2009 Black Saturday bushfires found that while the majority of affected people demonstrated psychological resilience in the long-term aftermath of the fires, a significant minority of people in highly affected communities reported mental health impacts 3-4 years following the event (Bryant et al., 2014).

7.2 Early Warning

475 There is no nationally standardized system for Bushfire Warnings in Australia. However, recommendations from the Royal Commission tasked with reviewing the 2009 Victoria Bushfire (Teague et al., 2010) have helped to drive forward efforts to establish a national system. In 2014 a National Review of Warnings and Information was undertaken. It recommended the establishment of a dedicated, multi-hazard National Working Group for Public Information and Warnings. Part of the task of this group would be to ensure greater national consistency of early warning information. One outcome of this recommendation 480 is a handbook on Public Information and Warnings which has been issued to provide guidance to actors across National, State and Territory governments in issuing warning information (AIDR, 2018).

Bushfire warnings in Australia are issued by State and Territory fire authorities and generally follow the 'Prepare, Stay and Defend or Leave Early' approach. A Fire Danger Rating system is also widely applied as a way to communicate fire risks. The system, originally developed in 1967, contained five risk levels ranging from 'low-moderate', where fires can generally be 485 controlled, to 'extreme', where evacuation is recommended but home defence may be possible under certain circumstances. Following the 2009 Black Saturday bushfires a sixth 'catastrophic' level was added where evacuation is deemed the only survival option (bushfire warning levels). This guidance was adopted by all states, except in Victoria where it is called 'Code Red' (BoM, 2020). In 2017, the system was revised to update the metrics used in forecasting the most appropriate level (ABC News, 12 Dec 2017). Threat level information is provided via radio, television, social media and via signs on all major rural 490 roads. Government websites also provide information which is updated every few minutes and includes maps of fires and associated threat levels. In addition, phone calls are made house-to-house when evacuation is recommended.

While these efforts help to reduce vulnerability and exposure to the wildfires, significant barriers to early action still remain. People in bushfire areas are frequently not aware of their risk, unprepared to manage risk, wait until the final moments to

evacuate or, at times, even return to fire affected areas to defend property (Whittaker et al., 2020). This is particularly relevant in peri-urban areas which are not as frequently exposed to bushfire risks. A 2020 study of people's reactions to bushfire warnings during the 2017 bushfires in New South Wales found that people largely understood warnings, however they did not respond to the warnings before seeking and obtaining additional confirmation to avoid what they might perceive as unnecessary evacuation and associated costs.. Researchers recommend that rather than further refining messages, confirmation mechanisms need to be implemented into early warning approaches. There are further barriers to acting on warnings, such as an incorrect assessment of the defensibility of a building, attachments to pets and personal items, and a 'hero culture' around people who did defend their home. These sociological barriers to life-saving measures increase the risk of deadly impacts.

Furthermore, an inquiry into the 2009 Black Saturday bushfires found that the 'Prepare, Stay and Defend or Leave Early' approach assumed that individuals had a fire plan in-place, however many people did not. Therefore people were left in a position to make complex decisions without adequate guidance.

7.3 Controlled burning and relation to weather conditions

For this study, we did not assess vegetation cover and condition (dryness) ahead of the season in comparison with earlier years, but it is clear that fire hazard management strategies such as bush reduction, manual removal of undergrowth and controlled burning can affect fire hazard

Note that the effectiveness of such measures depends on type of vegetation, but also the specific conditions of the bushfire. For instance, prior controlled burning may be somewhat effective to suppress fire risk under average weather conditions, but much less so in cases of very high temperatures, low humidity and strong wind, i.e., when the fire risk is no longer dominated by the type, condition and quantity of fuel but by weather conditions. During the 2009 fires in Victoria, recently burnt areas (up to 5–10 years) may have reduced the intensity of the fires, but not enough to increase the chance of effective suppression given the severe weather conditions at the time (Price and Bradstock, 2012).

In addition, it should be noted that controlled burning requires a window during the cooler parts of the year when conditions allow controlled burning to take place. he Queensland Fire and Emergency Services (QFES) noted that controlled burning is highly dependent on weather conditions, and that not all planned 2019 burns had been completed, given that in some areas, it rapidly became too dry to burn safely (ABC Fact Check, 20 December 2019). Recognising the highly non-linear relationship between weather conditions through the season (and in fact across several years) and anticipatory risk management strategies, in this attribution study we have not assessed the impact of these early-season weather conditions on the ability to reduce risk, and thus on fire risk itself.

7.4 Infrastructure and land use planning

Aging electricity infrastructure may play a role in increasing the risk of bushfire outbreaks from human-related ignition (Teague et al., 2010). Electric grid fires are primarily due to elastic extension and fatigue failures, and are made increasingly worse by high wind speeds (Mitchell, 2013). A 2017 study found that fires sparked by electricity failures are more prevalent during elevated fire risk and tend to burn larger, making them worse than fires due to other causes (Miller et al., 2017). Interestingly,

a 2013 report also notes that while electricity operators have the ability to disconnect electricity grids when there is a high-risk posed to the public, only South Australia has legislation in place to protect the operator from prosecution ([Energy Networks Association, 2013](#)). All of these factors coupled together increase Australia's vulnerability to bushfire outbreaks.

530 In contrast, stringent building codes have helped to reduce vulnerability to fire risks. The Bushfire Attack Level (BAL), and associated building codes, is the guiding resource for assessing and managing risks of a building exposed to heat, embers or direct fire. The BAL is applied nationwide, however the Fire Danger Index, one of the key metrics used in calculating the site specific BAL, is under state and local level jurisdiction. The highest BAL level was established following the 2009 Black Saturday bushfires ([Country Fire Authority, 2012](#)).

535 Land use planning at a community level is also crucial in reducing bushfire risk, particularly for rural and peri-urban areas which face the highest bushfire risks. This is recognised and addressed through each state and local government planning processes, which includes ensuring accessible bushfire evacuation routes and spaces. For example, the 2009 Victorian Bushfires Royal Commission cited a need for planning which 'prioritized human life over all other policy objectives'. This led to relevant policy changes through an amendment to Victoria's Planning Provisions. The Bushfire Management Overlay, and associated
540 guidelines, are among the principle aspects of this amendment. They provide direction for approval of new construction locations as well as siting and layout requirements of approved spaces, although these guidelines do not apply to existing property which puts a limitation on their overall positive impact ([Country Fire Authority, 2012](#)).

7.5 Conclusions vulnerability and exposure

Bushfires are a natural phenomenon, but their impact is also strongly influenced by human choices. Overall, Australia is one of
545 the most prepared countries in the world to manage bushfires and thus the impacts from this season's bushfire outbreaks could have been dramatically worse if not for the systems in place. This underscores the urgent need to adapt to changing risks in all places, with a special focus on the most vulnerable, but also highlights the limits to risk reduction and preparedness.

As a result of the Black Summer bushfires, formal inquiries have been launched in Victoria ([Victoria Government, 14 Jan 2020](#)), New South Wales ([New South Wales Government, Jan 2020](#)), Queensland ([Queensland Government, 18 Dec 2019](#))
550 and South Australia ([South Australia Government, 21 Jan 2020](#)). A Federal Royal Commission has been announced with an aim to improve resilience, preparedness and response to disasters across all levels of government. The Commission will also seek to improve disaster management coordination across local government and improve relevant legal frameworks. These inquiries will shed additional light on the vulnerability and exposure elements of the Black Summer Bushfires, and hopefully help mitigate future risk.

555 8 Conclusions

We investigated changes in the risk of bushfire weather in southeastern Australia due to anthropogenic climate change, underpinned by changes in extreme heat and extreme drought. The latter have longer time series and are covered by many more

climate models, leading to more robust conclusions. The fire risk is described by the Fire Weather Index, which was shown to correlate well with the area burnt in this part of Australia.

560 The first conclusion is that current climate models struggle to represent extremes in the 7-day averaged maximum temperature which was chosen as the most impact-relevant definition of heat as well as the Fire Weather Index. They tend to overestimate variability and thus underestimate the observed trends in these variables. Both of these factors give an underestimation of the change in probability due to anthropogenic climate change (PR). We therefore do not give best fit values but only lower bounds for these variables.

565 We find that the probability of extreme heat has increased by at least a factor two. We do not find attributable trends in extreme drought, neither on the annual time scale nor for the driest month in the fire season, even when mean precipitation does have drying trends in some models. Commensurate with this we find a significant increase in the risk of fire weather as severe or worse as observed in 2019/20 by at least 30%. Both for extreme heat and fire weather we think the true change in probability is likely much higher due to the model deficiencies.

570 The fire weather of 2019 was made much more severe by record positive excursions of the Indian Ocean dipole, even when the ENSO teleconnection was removed from this. The average effect of this mode is small, but the anomaly was so large that this factor explains about one third of the anomalous drought in July–December 2019. The other factor was the Southern Annular Mode, which was also anomalously negative during this time, explaining another one third of the July–December drought. Both factors were predicted well and gave good warning of the high fire risk in late 2019. The variability due to these modes is included in our analysis, although the simulated fidelity of the modes themselves and their trends has not been assessed in detail here. It should be noted that only a small fraction of the natural variability is described by these modes.

Of course the full fire risk is also affected by non-weather factors. The bushfire warning system in place in Australia worked well, but research shows that many people do not follow the guidelines as intended. The risk of bushfires is increased due to anthropogenic factors like aging electricity infrastructure. Efforts to mitigate against that risk using controlled burning are hampered by the very high fire risk due to weather factors shrinking the window in which controlled burning can be safely executed. Overall, however, Australia is one of the most prepared countries in the world to manage bushfires and thus the impacts from this season's bushfire outbreaks could have been dramatically worse if not for the systems in place. This underscores the urgent need to adapt to changing risks in all places, and especially the most vulnerable.

580 Although we clearly identify a connection between climate change and fire weather and ascertain a lower bound, we also find, in agreement with other studies, that we need more understanding of the biases in climate models and their resolution before we can make a more quantitative statement of how strong the connection is and how it will evolve in the future. Ever more detailed attribution statements are desired by society and scientific progress in the modelling of extreme events is therefore needed.

Data availability

590 Almost all data used in the analyses can be downloaded from the KNMI Climate Explorer at https://climexp.knmi.nl/bushfires_timeseries.cgi. This also contains the scripts that we used to do all the extreme value fits. These can also be performed using the graphical user interface of the Climate Explorer.

Acknowledgements. We thank Pandora Hope, Andrew Dowdy and Mitchell Black of the Bureau of Meteorology and Sarah Perkins-Kirkpatrick of the University of New South Wales for substantial contributions to the article. G.J.v.O was supported by the ERA4CS projects SERV_FORFIRE and EUPHEME (grant #690462). The contribution of F. K. was supported by the Belmont Forum Project PREREAL (grant # 292-2015-11-30-13-43-09 to I.D.). F.L. is supported by the Regional and Global Model Analysis (RGMA) component of the Earth and Environmental System Modeling Program of the US Department of Energy's Office of Biological & Environmental Research (BER) via NSF IA 1947282, and by a Swiss NSF Ambizione Fellowship (Project PZ00P2_174128). We acknowledge the use of data and imagery from LANCE FIRMS operated by NASA's Earth Science Data and Information System (ESDIS) with funding provided by NASA Headquarters. We would like to
600 thank all the volunteers who have donated their computing time to climateprediction.net and weather@home.

References

- Abatzoglou, J. T., Williams, A. P., and Barbero, R.: Global Emergence of Anthropogenic Climate Change in Fire Weather Indices, *Geophysical Research Letters*, 46, 326–336, <https://doi.org/10.1029/2018GL080959>, 2019.
- Agard, J., Schipper, E. L. F., et al., eds.: AR5 Climate Change 2014: Impacts, Adaptation, and Vulnerability: Annex II Glossary, Cambridge University Press, New York, 2014.
- Black, M. T., Karoly, D. J., and King, A. D.: The contribution of anthropogenic forcing to the Adelaide and Melbourne, Australia, heat waves of January 2014, *Bull. Amer. Met. Soc.*, 96, S145–S148, <https://doi.org/10.1175/BAMS-D-15-00097.1>, 2015.
- Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Y., B., Bastriko, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheru, F., Cozic, A., Cugnet, D., D’Andrea, F., Davini, P., de Lavergne, C., Denvil, S., Deshayes, J., M., D., Ducharne, A., Dufresne, J.-L., Dupont, E., Ethé, C., Fairhead, L., Falletti, L., Foujols, M.-A., Gardoll, S., Gastinea, G., J., G., Grandpeix, J.-Y., Guenet, B., Guez, L., Guilyardi, E., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M., Khadre-Traoré, A., Khodri, M., Krinner, G., Lebas, N., Levvasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luyssaert, S., G., M., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meurdesoif, Y., Mignot, J., Musat, I., Ottlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thieblemont, R., Traoré, A., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N.: Presentation and evaluation of the IPSL-CM6A-LR climate model, *J. Adv. Modeling Earth Systems*, submitted, 2020.
- Bryant, R. A., Waters, E., Gibbs, L., Gallagher, H. C., Pattison, P., Lusher, D., MacDougall, C., Harms, L., Block, K., Snowdon, E., Sinnott, V., Ireton, G., Richardson, J., and Forbes, D.: Psychological outcomes following the Victorian Black Saturday bushfires, *Australian & New Zealand Journal of Psychiatry*, 48, 634–643, <https://doi.org/10.1177/0004867414534476>, PMID: 24852323, 2014.
- Camia, A. and Amatulli, G.: Weather Factors and Fire Danger in the Mediterranean, in: *Earth Observation of Wildland Fires in Mediterranean Ecosystems*, edited by Chuvieco, Springer, Berlin, Heidelberg, https://doi.org/10.1007/978-3-642-01754-4_6, 2009.
- Ciavarella, A., Christidis, N., Andrews, M., Groenendijk, M., Rostron, J., Elkington, M., Burke, C., Lott, F. C., and Stott, P. A.: Upgrade of the HadGEM3-A based attribution system to high resolution and a new validation framework for probabilistic event attribution, *Weather and Climate Extremes*, 20, 9 – 32, <https://doi.org/10.1016/j.wace.2018.03.003>, 2018.
- Clarke, H., Lucas, C., and Smith, P.: Changes in Australian fire weather between 1973 and 2010, *Int. J. Climatol.*, 33, 931–944, <https://doi.org/10.1002/joc.3480>, 2013.
- Clarke, H. G., Smith, P. L., and Pitman, A. J.: Regional signatures of future fire weather over eastern Australia from global climate models, *International Journal of Wildland Fire*, 20, 550–562, 2011.
- Coles, S.: *An Introduction to Statistical Modeling of Extreme Values*, Springer Series in Statistics, London, UK, 2001.
- Cooley, D., Hunter, B., and Smith, R.: Univariate and multivariate extremes for the environmental sciences, *Environmental Statistics*, pp. 153–180, 2018.
- Country Fire Authority: Planning for Bushfire Victoria, version 2, https://www.eastgippsland.vic.gov.au/files/content/public/planning_and_building/planning_reference_and_incorporated_documents/reference_documents/cfa_planning_conditions_guidelines_amended.pdf, 2012.
- Dimitrakopoulos, A. P., Bemmerzouk, A. M., and Mitsopoulos, I. D.: Evaluation of the Canadian fire weather index system in an eastern Mediterranean environment, *Meteorological Applications*, 18, 83–93, <https://doi.org/10.1002/met.214>, 2011.

- Dowdy, A. J.: Climatological Variability of Fire Weather in Australia, *Journal of Applied Meteorology and Climatology*, 57, 221–234, <https://doi.org/10.1175/JAMC-D-17-0167.1>, 2018.
- Dowdy, A. J. and Pepler, A.: Pyroconvection Risk in Australia: Climatological Changes in Atmospheric Stability and Surface Fire Weather
640 Conditions, *Geophys. Res. Lett.*, 45, 2005–2013, <https://doi.org/10.1002/2017GL076654>, 2018.
- Dowdy, A. J., Mills, G. A., Finkele, K., and de Groot, W.: Australian fire weather as represented by the McArthur Forest Fire Danger Index and the Canadian Forest Fire Weather Index, Technical Report 10, The Centre for Australian Weather and Climate Research, 2009.
- Dowdy, A. J., Ye, H., Pepler, A., Thatcher, M., Osbrough, S. L., Evans, J. P., Di Virgilio, G., and McCarthy, N.: Future changes in extreme weather and pyroconvection risk factors for Australian wildfires, *Scientific Reports*, 9, 10 073, [https://doi.org/10.1038/s41598-019-46362-](https://doi.org/10.1038/s41598-019-46362-x)
645 [x](https://doi.org/10.1038/s41598-019-46362-x), 2019.
- Eden, J. M., Wolter, K., Otto, F. E. L., and van Oldenborgh, G. J.: Multi-method attribution analysis of extreme precipitation in Boulder, Colorado, *Environ. Res. Lett.*, 11, 124 009, <https://doi.org/10.1088/1748-9326/11/12/124009>, 2016.
- Eden, J. M., Kew, S. F., Bellprat, O., Lenderink, G., Manola, I., Omrani, H., and van Oldenborgh, G. J.: Extreme precipitation in the Netherlands: An event attribution case study, *Weather and Climate Extremes*, 21, 90–101, <https://doi.org/10.1016/j.wace.2018.07.003>,
650 2018.
- Energy Networks Association: Bushfires + Energy Networks, <https://www.energynetworks.com.au/resources/fact-sheets/bushfires-and-energy-networks/>, 2013.
- Finlay, S. E., Moffat, A., Gazzard, R., Baker, D., and Murray, V.: Health impacts of wildfires, *PLoS Currents Disasters*, 4, e4f959 951cce2c, <https://doi.org/10.1371/4f959951cce2c>, 2012.
- Fischer, E. M., Seneviratne, S. I., Vidale, P. L., Lüthi, D., and Schär, C.: Soil Moisture–Atmosphere Interactions during the 2003 European Summer Heat Wave, *Journal of Climate*, 20, 5081–5099, <https://doi.org/10.1175/JCLI4288.1>, 2007.
- Guilod, B. P., Jones, R. G., Bowery, A., Haustein, K., Massey, N. R., Mitchell, D. M., Otto, F. E. L., Sparrow, S. N., Uhe, P., Wallom, D. C. H., Wilson, S., and Allen, M. R.: weather@home 2: validation of an improved global–regional climate modelling system, *Geoscientific Model Development*, 10, 1849–1872, <https://doi.org/10.5194/gmd-10-1849-2017>, 2017.
- Haikerwal, A., Akram, M., Del Monaco, A., Smith, K., Sim, M. R., Meyer, M., Tonkin, A. M., Abramson, M. J., and Dennekamp, M.:
660 Impact of Fine Particulate Matter (PM_{2.5}) Exposure During Wildfires on Cardiovascular Health Outcomes, *Journal of the American Heart Association*, 4, e001 653, <https://doi.org/10.1161/JAHA.114.001653>, 2015.
- Hansen, J., Ruedy, R., Sato, M., and Lo, K.: Global Surface Temperature Change, *Rev. Geophys.*, 48, RG4004, <https://doi.org/DOI10.1029/2010RG000345>, 2010.
- Harris, S. and Lucas, C.: Understanding the variability of Australian fire weather between 1973 and 2017, *PLoS ONE*, 14, e0222 328, <https://doi.org/10.1371/journal.pone.0222328>, 2019.
- Haustein, K., Otto, F. E. L., Venema, V., Jacobs, P., Cowtan, K., Hausfather, Z., Way, R. G., White, B., Subramanian, A., and Schurer, A. P.: A Limited Role for Unforced Internal Variability in Twentieth-Century Warming, *J. Climate*, 32, 4893–4917, <https://doi.org/10.1175/JCLI-D-18-0555.1>, 2019.
- Hazeleger, W., Severijns, C., Semmler, T., Stefanescu, S., Yang, S., Wang, X., Wyser, K., Dutra, E., Baldasano, J. M., Bintanja, R., Bougeault, P., Caballero, R., Ekman, A. M. L., Christensen, J. H., van den Hurk, B., Jimenez, P., Jones, C., Kallberg, P., Koenigk, T., McGrath, R., Miranda, P., Van Noije, T., Palmer, T., Parodi, J. A., Schmith, T., Selten, F., Storelvmo, T., Sterl, A., Tapamo, H., Vancoppenolle, M., Viterbo, P., and Willen, U.: EC-Earth: A Seamless Earth-System Prediction Approach in Action, *Bull. Amer. Met. Soc.*, 91, 1357–1363, <https://doi.org/DOI10.1175/2010BAMS2877.1>, 2010.

- 675 Hersbach, H., Bell, W., Berrisford, P., Horányi, A., J., M.-S., Nicolas, J., Radu, R., Schepers, D., Simmons, A., Soci, C., and Dee, D.: Global reanalysis: goodbye ERA-Interim, hello ERA5, ECMWF Newsletter, 159, 17–24, <https://doi.org/10.21957/vf291hehd7>, 2019.
- Hope, P., Wang, G., Lim, E.-P., Hendon, H. H., and Arblaster, J. M.: What caused the record-breaking heat across Australia in October 2015?, *Bull. Amer. Met. Soc.*, 97, S1–S5, <https://doi.org/10.1175/BAMS-D-16-0142.1>, 2016.
- Hope, P., Black, M. T., Lim, E.-P., Dowdy, A., Wang, G., Pepler, A. S., and Fawcett, R. J. B.: On Determining the Impact of Increasing Atmospheric CO₂ on the Record Fire Weather In Eastern Australia In February 2017, *Bull. Amer. Met. Soc.*, 100, S111–S117, <https://doi.org/10.1175/BAMS-D-18-0135.1>, 2019.
- 680 Jeffrey, S., Rotstayn, L., Collier, M., Dravitzki, S., Hamalainen, C., Moeseneder, C., Wong, K., and Syktus, J.: Australia’s CMIP5 submission using the CSIRO-Mk3. 6 model, *Aust. Meteor. Oceanogr. J.*, 63, 1–13, 2013.
- Johnston, F. and Bowman, D.: Bushfire Smoke: An Exemplar of Coupled Human and Natural Systems, *Geographical Research*, 52, 45–54, <https://doi.org/10.1111/1745-5871.12028>, 2014.
- 685 Kala, J., De Kauwe, M. G., Pitman, A. J., Medlyn, B. E., Wang, Y.-P., Lorenz, R., and Perkins-Kirkpatrick, S. E.: Impact of the representation of stomatal conductance on model projections of heatwave intensity, *Scientific Reports*, 6, 23 418, <https://doi.org/10.1038/srep23418>, 2016.
- Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J. M., Bates, S., Danabasoglu, G., Edwards, J., et al.: The Community Earth System Model (CESM) large ensemble project: A community resource for studying climate change in the presence of internal climate variability, *Bulletin of the American Meteorological Society*, 96, 1333–1349, <https://doi.org/10.1175/BAMS-D-13-00255.1>, 2015.
- 690 Kew, S. F., Philip, S. Y., van Oldenborgh, G. J., Otto, F. E., Vautard, R., and van der Schrier, G.: The exceptional summer heatwave in Southern Europe 2017, *Bull. Amer. Met. Soc.*, 100, S2–S5, <https://doi.org/10.1175/BAMS-D-18-0109.1>, 2019.
- 695 King, A. D., Black, M. T., Karoly, D. J., and Donat, M. G.: Increased likelihood of Brisbane, Australia, G20 heat event due to anthropogenic climate change, *Bull. Amer. Met. Soc.*, 96, S141–S144, <https://doi.org/10.1175/BAMS-D-15-00098.1>, 2015a.
- King, A. D., van Oldenborgh, G. J., Karoly, D. J., Lewis, S. C., and Cullen, H.: Attribution of the record high Central England temperature of 2014 to anthropogenic influences, *Environmental Research Letters*, 10, 054 002, <https://doi.org/10.1088/1748-9326/10/5/054002>, 2015b.
- King, A. D., van Oldenborgh, G. J., and Karoly, D. J.: Climate Change and El Niño increase likelihood of Indonesian heat and drought, *Bull. Amer. Met. Soc.*, 97, S113–S117, <https://doi.org/10.1175/BAMS-D-16-0164.1>, 2016.
- 700 Kirchmeier-Young, M. C., Zwiers, F. W., and Gillett, N. P.: Attribution of extreme events in Arctic sea ice extent, *Journal of Climate*, 30, 553–571, <https://doi.org/10.1175/JCLI-D-16-0412.1>, 2017.
- Kopplitz, S. N., Mickley, L. J., Marlier, M. E., Buonocore, J. J., Kim, P. S., Liu, T., Sulprizio, M. P., DeFries, R. S., Jacob, D. J., Schwartz, J., Pongsiri, M., and Myers, S. S.: Public health impacts of the severe haze in Equatorial Asia in September–October 2015: demonstration of a new framework for informing fire management strategies to reduce downwind smoke exposure, *Environ. Res. Lett.*, 11, 094 023, <https://doi.org/10.1088/1748-9326/11/9/094023>, 2016.
- 705 Krikken, F., Lehner, F., Haustein, K., Drobyshev, I., and van Oldenborgh, G. J.: Attribution of the role of climate change in the forest fires in Sweden 2018, *Natural Hazards and Earth System Sciences Discussions*, 2019, 1–24, <https://doi.org/10.5194/nhess-2019-206>, 2019.
- Lewis, S., Blake, S., Trewin, B., Black, M., Dowdy, A., Perkins-Kirkpatrick, S., King, A., and Sharples, J.: Deconstructing factors contributing to the 2018 fire weather in Queensland, Australia, *Bull. Amer. Met. Soc.*, <https://doi.org/10.1175/BAMS-D-19-0144.1>, 2019.
- 710 Lewis, S. C. and Karoly, D. J.: Anthropogenic contributions to Australia’s record summer temperatures of 2013, *Geophysical Research Letters*, 40, 3705–3709, <https://doi.org/10.1002/grl.50673>, 2013.

- Maher, N., Milinski, S., Suarez-Gutierrez, L., Botzet, M., Kornbluh, L., Takano, Y., Kröger, J., Ghosh, R., Hedemann, C., Li, C., et al.: The Max Planck Institute grand ensemble-enabling the exploration of climate system variability, *Journal of Advances in Modeling Earth Systems*, 11, 2050–2069, <https://doi.org/10.1029/2019MS001639>, 2019.
- Martins, E. S. P. R., Coelho, C. A. S., Haarsma, R. J., Otto, F. E. L., King, A. D., van Oldenborgh, G. J., Kew, S. F., Philip, S. Y., Vasconcelos Junior, F. C., and Cullen, H.: A multimethod attribution analysis of the prolonged northeast Brazil hydrometeorological drought (2012–16), *Bull. Amer. Met. Soc.*, 99, S65–S69, <https://doi.org/10.1175/BAMS-D-17-0102.1>, 2018.
- Mathur, R. and AchutaRao, K.: A modelling exploration of the sensitivity of the India’s climate to irrigation, *Climate Dynamics*, <https://doi.org/10.1007/s00382-019-05090-8>, 2019.
- McArthur, A. G.: Weather and grassland fire behaviour, Tech. Rep. Leaflet 100, Aust. For. Timb. Bur., 1966.
- McArthur, A. G.: Fire behaviour in eucalypt forest, Tech. Rep. Leaflet 107, Aust. For. Timb. Bur., 1967.
- Miller, C., Plucinski, M., Sullivan, A., Stephenson, A., Huston, C., Charman, K., Prakash, M., and Dunstall, S.: Electrically caused wildfires in Victoria, Australia are over-represented when fire danger is elevated, *Landscape and Urban Planning*, 167, 267 – 274, <https://doi.org/10.1016/j.landurbplan.2017.06.016>, 2017.
- Miralles, D. G., Teuling, A. J., van Heerwaarden, C. C., and Vilà-Guerau de Arellano, J.: Mega-heatwave temperatures due to combined soil desiccation and atmospheric heat accumulation, *Nature Geoscience*, 7, 345–349, <https://doi.org/10.1038/ngeo2141>, 2014.
- Mitchell, J. W.: Power line failures and catastrophic wildfires under extreme weather conditions, *Engineering Failure Analysis*, 35, 726–735, <https://doi.org/10.1016/j.engfailanal.2013.07.006>, special issue on ICEFA V- Part 1, 2013.
- Morgan, G., Sheppard, V., Khalaj, B., Ayyar, A., Lincoln, D., Jalaludin, B., Beard, J., Corbett, S., and Lumley, T.: Effects of Bushfire Smoke on Daily Mortality and Hospital Admissions in Sydney, Australia, *Epidemiology*, 21, 47–55, <https://doi.org/10.1097/EDE.0b013e3181c15d5a>, 2010.
- Naveau, P., Huser, R., Ribereau, P., and Hannart, A.: Modeling jointly low, moderate, and heavy rainfall intensities without a threshold selection, *Water Resources Research*, 52, 2753–2769, 2016.
- Noble, I. R., Gill, A. M., and Bary, G. A. V.: McArthur’s fire-danger meters expressed as equations, *Australian Journal of Ecology*, 5, 201–203, <https://doi.org/10.1111/j.1442-9993.1980.tb01243.x>, 1980.
- Otto, F. E. L., van der Wiel, K., van Oldenborgh, G. J., Philip, S. Y., Kew, S. F., Uhe, P., and Cullen, H.: Climate change increases the probability of heavy rains in Northern England/Southern Scotland like those of storm Desmond—a real-time event attribution revisited, *Environ. Res. Lett.*, 13, 024 006, <https://doi.org/10.1088/1748-9326/aa9663>, 2018a.
- Otto, F. E. L., Wolski, P., Lehner, F., Tebaldi, C., van Oldenborgh, G. J., Hogesteegeer, S., Singh, R., Holden, P., Fuckar, N. S., Odoulami, R., and New, M.: Anthropogenic influence on the drivers of the Western Cape drought 2015–2017, *Environ. Res. Lett.*, 13, 124 010, <https://doi.org/10.1088/1748-9326/aae9f9>, 2018b.
- Pasquale, C.: Are your data really Pareto distributed?, *Physica A: Statistical Mechanics and its Applications*, 392, 5947–5962, <https://doi.org/10.1016/j.physa.2013.07.061>, 2013.
- Pepler, A., Coutts-Smith, A., and Timbal, B.: The role of East Coast Lows on rainfall patterns and inter-annual variability across the East Coast of Australia, *Int. J. Climatol.*, 34, 1011–1021, <https://doi.org/10.1002/joc.3741>, 2014.
- Perkins, S. E. and Gibson, P. B.: Increased risk of the 2014 Australian May heatwave due to anthropogenic activity, *Bull. Amer. Met. Soc.*, 96, S154–S157, <https://doi.org/10.1175/BAMS-D-15-00074.1>, 2015.
- Perkins, S. E., Lewis, S. C., King, A. D., and Alexander, L. V.: Increased Simulated Risk of the Hot Australian Summer of 2012/13 due to Anthropogenic Activity as Measured by Heat Wave Frequency and Intensity, *Bull. Amer. Met. Soc.*, 95, S34–S37, 2014.

- Philip, S. Y., Kew, S. F., Hauser, M., Guillod, B. P., Teuling, A. J., Whan, K., Uhe, P., and Oldenborgh, G. J. v.: Western US high June 2015 temperatures and their relation to global warming and soil moisture, *Clim. Dyn.*, 50, 2587–2601, <https://doi.org/10.1007/s00382-017-3759-x>, 2018a.
- Philip, S. Y., Kew, S. F., van Oldenborgh, G. J., Aalbers, E., Otto, F. E. L., Haustein, K., Habets, F., and Singh, R.: Validation of a rapid attribution of the May/June 2016 flood-inducing precipitation in France to climate change, *Journal of Hydrometeorology*, 19, 1881–1898, <https://doi.org/10.1175/JHM-D-18-0074.1>, 2018b.
- Philip, S. Y., Kew, S. F., van Oldenborgh, G. J., Otto, F. E. L., O’Keefe, S., Haustein, K., King, A. D., Zegeye, A., Eshetu, Z., Hailemariam, K., Singh, R. K., Jjemba, E., Funk, C., and Cullen, H.: Attribution analysis of the Ethiopian drought of 2015, *J. Climate*, 31, 2465–2486, <https://doi.org/10.1175/JCLI-D-17-0274.1>, 2018c.
- Philip, S. Y., Kew, S. F., van Oldenborgh, G. J., Otto, F. E. L., Vautard, R., van der Wiel, K., King, A. D., Lott, F. C., Arrighi, J., Singh, R. P., and van Aalst, M. K.: A protocol for probabilistic extreme event attribution analyses, *Advances in Statistical Climatology, Meteorology and Oceanography*, 6, 177–203, <https://doi.org/10.5194/ascmo-6-177-2020>, 2020.
- Price, O. F. and Bradstock, R. A.: The efficacy of fuel treatment in mitigating property loss during wildfires: Insights from analysis of the severity of the catastrophic fires in 2009 in Victoria, Australia, *Journal of Environmental Management*, 113, 146 – 157, <https://doi.org/10.1016/j.jenvman.2012.08.041>, 2012.
- Reid, C. E., Brauer, M., Johnston, F. H., Jerrett, M., Balmes, J. R., and Elliott, C. T.: Critical Review of Health Impacts of Wildfire Smoke Exposure, *Environmental Health Perspectives*, 124, 1334–1343, <https://doi.org/10.1289/ehp.1409277>, 2016.
- Ribes, A., Thao, S., and Cattiaux, J.: Describing the Relationship between a Weather Event and Climate Change: A New Statistical Approach, *Journal of Climate*, 33, 6297–6314, <https://doi.org/10.1175/JCLI-D-19-0217.1>, 2020.
- Rodgers, K. B., Lin, J., and Frölicher, T. L.: Emergence of multiple ocean ecosystem drivers in a large ensemble suite with an Earth system model, *Biogeosciences*, 12, 3301–3320, <https://doi.org/10.5194/bg-12-3301-2015>, 2015.
- Sanderson, B. M. and Fisher, R. A.: A fiery wake-up call for climate science, *Nature Climate Change*, <https://doi.org/10.1038/s41558-020-0707-2>, 2020.
- Schaller, N., Otto, F. E. L., van Oldenborgh, G. J., Massey, N. R., Sparrow, S., and Allen, M. R.: The heavy precipitation event of May–June 2013 in the upper Danube and Elbe basins, *Bull. Amer. Met. Soc.*, 95, S69–S72, 2014.
- Shabbar, A., Skinner, W., and Flannigan, M. D.: Prediction of Seasonal Forest Fire Severity in Canada from Large-Scale Climate Patterns, *Journal of Applied Meteorology and Climatology*, 50, 785–799, <https://doi.org/10.1175/2010JAMC2547.1>, 2011.
- Shaposhnikov, D., Revich, B., Bellander, T., Bedada, G. B., Bottai, M., Kharkova, T., Kvasha, E., Lezina, E., Lind, T., Semutnikova, E., and Pershagen, G.: Mortality related to air pollution with the moscow heat wave and wildfire of 2010, *Epidemiology (Cambridge, Mass.)*, 25, 359–364, <https://doi.org/10.1097/EDE.000000000000090>, 2014.
- Siswanto, van Oldenborgh, G. J., van der Schrier, G., Lenderink, G., and van den Hurk, B. J. J. M.: Trends in high-daily precipitation events in Jakarta and the flooding of January 2014, *Bull. Amer. Met. Soc.*, 96, S131–S135, <https://doi.org/10.1175/BAMS-D-15-00128.1>, 2015.
- Sun, L., Alexander, M., and Deser, C.: Evolution of the global coupled climate response to Arctic sea ice loss during 1990–2090 and its contribution to climate change, *Journal of Climate*, 31, 7823–7843, <https://doi.org/10.1175/JCLI-D-18-0134.1>, 2018.
- Teague, B., McLeod, R., and Pascoe, S.: 2009 Victorian Bushfires Royal Commission Report (Summary), ISBN 978-0-9807408-1-3, http://royalcommission.vic.gov.au/finaldocuments/summary/PF/VBRC_Summary_PF.pdf, 2010.
- Tebaldi, C. and Arblaster, J. M.: Pattern scaling: Its strengths and limitations, and an update on the latest model simulations, *Climatic Change*, 122, 459–471, <https://doi.org/10.1007/s10584-013-1032-9>, 2014.

- Thiery, W., Davin, E. L., Lawrence, D. M., Hirsch, A. L., Hauser, M., and Seneviratne, S. I.: Present-day irrigation mitigates heat extremes, *Journal of Geophysical Research: Atmospheres*, 122, 1403–1422, <https://doi.org/10.1002/2016JD025740>, 2017.
- Uhe, P., Philip, S. Y., Kew, S. F., Shah, K., Kimutai, J., Mwangi, E., van Oldenborgh, G. J., Singh, R. K., Arrighi, J., Jjemba, E., Cullen, H., and Otto, F. E. L.: Attributing drivers of the 2016 Kenyan drought, *Int. J. Climatol.*, 38, e554–e568, <https://doi.org/10.1002/joc.5389>, 2018.
- van der Wiel, K., Kapnick, S. B., van Oldenborgh, G. J., Whan, K., Philip, S. Y., Vecchi, G. A., Singh, R. K., Arrighi, J., and Cullen, H.: Rapid attribution of the August 2016 flood-inducing extreme precipitation in south Louisiana to climate change, *Hydrol. Earth Syst. Sci.*, 21, 897–921, <https://doi.org/10.5194/hess-21-897-2017>, 2017.
- van Oldenborgh, G. J., Haarsma, R., De Vries, H., and Allen, M. R.: Cold Extremes in North America vs. Mild Weather in Europe: The Winter of 2013–14 in the Context of a Warming World, *Bulletin of the American Meteorological Society*, 96, 707–714, <https://doi.org/10.1175/BAMS-D-14-00036.1>, 2015.
- van Oldenborgh, G. J., Otto, F. E. L., Haustein, K., and Achuta Rao, K.: The heavy precipitation event of December 2015 in Chennai, India, *Bull. Amer. Met. Soc.*, 97, S87–S91, <https://doi.org/10.1175/BAMS-D-16-0129.1>, 2016.
- van Oldenborgh, G. J., van der Wiel, K., Sebastian, A., Singh, R. K., Arrighi, J., Otto, F. E. L., Haustein, K., Li, S., Vecchi, G. A., and Cullen, H.: Attribution of extreme rainfall from Hurricane Harvey, August 2017, *Environ. Res. Lett.*, 12, 124 009, <https://doi.org/10.1088/1748-9326/aa9ef2>, 2017.
- van Oldenborgh, G. J., Philip, S. Y., Kew, S. F., van Weele, M., Uhe, P., Otto, F. E. L., Singh, R. K., Pai, I., Cullen, H., and AchutaRao, K.: Extreme heat in India and anthropogenic climate change, *Natural Hazards and Earth System Sciences*, 18, 365–381, <https://doi.org/10.5194/nhess-18-365-2018>, 2018.
- Van Wagner, C. E.: Conversion of William’s Severity Rating for Use with the Fire Weather Index, Petawawa For. Exp. Sta. Inf. Rep. PS-X-21, Can. For. Serv., 1970.
- Vautard, R., van Oldenborgh, G. J., Thao, S., Dubuisson, B., Lenderink, G., Ribes, A., Soubeyroux, J. M., Yiou, P., and Planton, S.: Extreme fall 2014 precipitations in the Cévennes mountain range, *Bull. Amer. Meteor. Soc.*, 96, S56–S60, <https://doi.org/10.1175/BAMS-D-15-00088.1>, 2015.
- Weisheimer, A., Schaller, N., O’Reilly, C., MacLeod, D. A., and Palmer, T.: Atmospheric seasonal forecasts of the twentieth century: multi-decadal variability in predictive skill of the winter North Atlantic Oscillation (NAO) and their potential value for extreme event attribution, *Quarterly Journal of the Royal Meteorological Society*, 143, 917–926, <https://doi.org/10.1002/qj.2976>, 2017.
- Whittaker, J., Taylor, M., and Bearman, C.: Why don’t bushfire warnings work as intended? Responses to official warnings during bushfires in New South Wales, Australia, *International Journal of Disaster Risk Reduction*, 45, 101476, <https://doi.org/10.1016/j.ijdr.2020.101476>, 2020.