

Atmospheric Conditions Leading to an Exceptional Fatal Flash Flood in the Negev Desert, Israel

Uri Dayan¹, Itamar M. Lensky², Baruch Ziv³

¹Department of Geography, The Hebrew University of Jerusalem, Jerusalem, 9070227, Israel

²Department of Geography and Environment, Bar-Ilan University, Ramat-Gan, 5290002, Israel

5 ³Department of Natural Sciences, The Open University of Israel, Raanana, Israel

Correspondence to: Itamar M. Lensky (itamar.lensky@biu.ac.il)

Abstract. The study deals with an intense rainstorm that hit the Middle East between 24 and 27 April 2018 and took the lives of 13 people, 10 of them in 26 April during the deadliest flash flood in Tzafit Basin (31.0°N 35.3°E), the Negev Desert. The rainfall observed in the southern Negev was comparable to the long-term annual rainfall there, with intensities exceeding 75 years return period. The timing of the storm, at the end of the rainy season, when rain is relatively rare and spotty, raises the question what were the atmospheric conditions that made this rainstorm one of the most severe late spring storms.

The synoptic background was an upper-level cutoff low that formed south of a blocking high, which developed over Eastern Europe. The cutoff low entered the Levant near 30°N latitude, and slowed its movement from ~10 to <5 ms⁻¹, and therefore extended the duration of the storm over the region. The dynamic potential of the cutoff low, as estimated by the curvature vorticity, was the largest among the 12 late spring rainstorms that occurred during the latest 33 years. The lower levels were dominated by a cyclone centred over north western Saudi Arabia, producing north-westerly winds that advected moist air from the Mediterranean inland. During the approach of the storm, the atmosphere over Israel became unstable, with instability indices reaching values favourable for thunderstorms (e.g., CAPE = 909 J Kg⁻¹, SI = -2.7 K) and the precipitable water reaching 30 mm. The latter is explained by lower-level moisture advection from the Mediterranean and an additional contribution of mid-level moist air transport entering the region from the east. Three major rain centres were active over Israel during April 26, only one of them was orographic and the other two were triggered by instability. The build-up of the instability is explained by a negative upper-level temperature anomaly over the region, caused by a northerly flow east of a blocking high that dominated Eastern Europe.

The intensity of this storm is attributed to an amplification of a mid-latitude disturbance, which produced a cutoff low, with its implied high relative vorticity, low upper-level temperatures and slow progression. All these,

combined with the contribution of moisture supply, led to intense moist convection that prevailed over the region
30 three days successively.

1 Introduction

On April 26, 2018 an extremely intense rainstorm hit Israel. Thirteen people were killed in this storm, ten of them
by a raging flash flood in Tzafit Basin (31.0°N 35.3°E), in the northeast of the Negev Desert (denoted in Fig. 1).
The rainfall depth during this storm exceeded 20 mm over ~ half of Israel, over most of which the monthly long-
35 term mean does not exceed 10 mm (Porat et al., 2018).

Flash flood forecast and warning is a challenging task, in particular over remote arid areas, where meteorological
radar coverage is sparse. Though flash flood warning has much improved in the last few decades, it is still the most
weather-related fatal hazard globally (Montz et al. 2002), with 1,075 fatalities between 1996 and 2014 in the United
States alone (Terti et al. 2019). A significant part of flash flood victims is caused by incorrect assessment or
40 misjudging of the risks of these rapidly evolving phenomena (Sharif et al. 2015; Becker et al. 2015; Diakakis et al.
2018).

The study focuses on the Negev Desert and Judean Desert (Fig. 1), hereafter referred to as the 'study region'. The
climatic regimes of the study region span from semiarid in the north to arid in the centre and the south (south of
31.2°N, Ziv et al 2014). The majority of the annual precipitation in Israel is associated with Mediterranean cyclones,
45 while reaching its eastern part (i.e., Cyprus Lows, HMSO 1962; Saaroni et al. 2010; Zappa et al. 2015). Two-thirds
of the rainfall occur during December through February (Alpert et al. 2004; Ziv et al. 2006). During the late spring
(Apr-May), the rainfall over the northern and central parts of the Negev Desert constitutes 4 - 9% of the annual
average (5 - 10 mm). In spite of these negligible rain amounts, the number of flash flood events can not be ignored.
The flood regime in the study region was analysed by Kahana (1999), based on 37 hydrometric stations operated
50 by the Israeli Hydrological Service. He identified 59 “major floods” in which the recorded peak discharge reached
the magnitude of a 5 year recurrence interval (for the period 1947- 1994) at least in one watershed. 14% of these
major floods occurred during the late spring.


The main source of major floods in the late spring over the study region is the active "Red Sea Trough" (ARST,
Kahana et al., 2002). The RST is a low-pressure system extending from south toward the Eastern Mediterranean
55 (EM) and the Levant (Ashbel 1938; Kahana et al. 2002; Tsvieli and Zangvil 2005). The ARST is most frequent
during fall and spring (Sharon, 1978; Sharon and Kutiel, 1986; Dayan et al., 2001). The ARST is accompanied by
a pronounced upper-level trough that develops over Egypt, and at times, initiates convective storms over the Levant.

The secondary source of major floods is the Syrian Low - a derivative of the Cyprus Low. Syrian lows are Mediterranean mid-latitude cyclones that approach Syria and deepen, in contrast to the normal conditions, in which
60 Mediterranean cyclones tend to decay upon reaching Syria (Kahana et al, 2002). The implied winds over Israel are north-westerly, enriched with moisture originating from the Mediterranean that cross the Judean and the Negev Mountains perpendicular to the terrain upslope, supporting orographic rains.

The storm analysed here was associated with a Syrian low, accompanied by an upper-level closed cyclone,
65 resembling the features of a 'cutoff low' (as is shown in Sec. 3 below). A cutoff low is defined as a closed cyclone in the upper levels, overlapping with a PV maximum (e.g., Hoskins et al., 1985). Cutoff lows are considered as favourable for severe weather over the Mediterranean Basin (Porcu et al., 2007). For instance, in south Portugal cutoff lows were found to be a major source for heavy rainfall (Fragoso and Tildes Gomes, 2008). Porcu et al. (2007) identified 273 cutoff lows over the entire Mediterranean Basin during the period 1992–2001. As for the
70 Levant (30-35°N, 30-35°E), they found 6 cases (average of 0.6 y^{-1}) during the months April-September. Since the Levant is free of upper-level significant cyclonic system (and also of rain) during the summer months June-September (Kushnir et al., 2017), this result reflects, actually, the events of April – May.

This storm was severe in several aspects. One is the number of fatalities, 13, which is record breaking for Israel. Second, the north eastern part of the Negev Desert experienced rain intensities reaching 75-100-year return period,
75 resulting in discharge magnitudes of 10-50-year return period (Rinat et al., 2020). The third aspect is the rainfall totals for the storm, which reached 40-50 mm over wide parts of the study region, i.e., 10 times the monthly long-term mean (Porat et al., 2018).

The aim of this study is to analyse the storm in its seasonal context and to identify the atmospheric conditions that explain its severity. The article is organized as follows: in the next section, we present the data and methods used
80 to analyse the event. Section 3 describes the event through observational evidence. In section 4 we discuss the unique conditions that lead to the severe convection, and the sources of moisture for the rain formation in this storm. The main results are summarised in Sec. 5.



85 **Figure 1: Rainfall distribution over Israel for the period of April 25 06 UTC to April 27 06 UTC 2018, from the IMS radar and rain gauge measurements based on the INCA system integration method (Haiden et al. 2011). The black line indicates the water divide south of the Samarian Mountains.**

2 Data and Methods

The study area is the Negev Desert, south of 31.2° N, and the Judean Desert (Fig. 1). The synoptic scale data, including satellite imagery, cover the eastern and central Mediterranean Basin and South Europe. The study period, April 24-27, 2018, corresponds to a storm that influenced the EM, in particular April 26, when the Tzafit flood took place. This storm is one of 12 storms spread over 27 days ('reference storms', hereafter) with the highest surface runoff observed over the study area in the late spring during 1986 – 2018, based on the hydrometric stations of the Israeli Hydrological Service.

The rainfall data is based on the Israeli Meteorological Service (IMS) rain measuring network and radar data (Porat et al., 2018). The spatial resolution of the radar is 250 m in the radial direction and 1° in the azimuthal. This data is rectified to 1×1 km grid, and its temporal resolution is 5 min. The rain intensity was derived from the radar data (Marra and Morin, 2015).

The atmospheric processes responsible for flash flood-producing rainstorms combine synoptic and sub-synoptic scales (Bardossy and Filiz, 2005). This also holds specifically for severe convection during the spring over the Negev Desert (Dayan and Morin, 2006; Armon et al., 2018). The data for the synoptic maps, including sea level pressure (SLP) and upper-level fields and geopotential height (gph) were taken from NCEP/NCAR reanalysis 1 database, at 2.5° x 2.5° spatial resolution (Kalnay et al. 1996; Kistler et al. 2001). The data for the mesoscale maps including wind, relative and potential vorticity, temperature and relative humidity, are taken from ERA5 ECMWF reanalysis data base (C3S, 2017, <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels>) at 30 km spatial resolution. SLP maps from the two data bases for the pertinent region and period were compared for consistency and were found reasonably fit (not shown).

According to the quasi-geostrophic theory, the relative vorticity is a general measure for cyclone intensity and activity (e. g., Holton 1992). The curvature term dominates near cyclone centres and within meandering flow, whereas the shear term dominates near jets (Uccellini and Kocin, 1987). In the cases analysed here, an upper-level cyclone was found within the Levant region (or in its proximity), accompanied by a southward shift of the subtropical jet, and therefore, the curvature vorticity was considered as dominant.

Following the above, the vorticity here is approximated by a 'measure of curvature vorticity' (MCV, hereafter), which is calculated on the periphery of the upper-level (500 hPa) cyclones, based on the relation

$$\xi_c = V/R \quad , \quad (1)$$

where ξ_c is the curvature vorticity, V is the tangential wind speed, and R is the cyclone radius, measured on the outer-most closed isohypse (when derived with 15 m intervals). The tangential wind speed was calculated using the geostrophic relation, i.e.,

$$V \cong g \cdot D / (R \cdot f) , \quad (2)$$

120 where g and f are the gravitational acceleration and Coriolis parameter, respectively, and D is the depth of the cyclone, determined by the difference between its central height and that of the outer-most closed isohypse. Inserting V from Eq. (2) in Eq. (1) yields:

$$MCV = g \cdot D / (R^2 \cdot f) . \quad (3)$$

The stability indices were derived from the sounding data of Beit Dagan station, Israel (32.5°N, 34.8°E), retrieved
125 from the Department of Atmospheric Sciences, The University of Wyoming, at: <http://weather.uwyo.edu/upperair/sounding.html>. These are:

- Lifted Index (LI, Galway 1956): Temperature difference between the environment and an air parcel lifted adiabatically from 2 m above the surface to 500 hPa. Negative values indicate instability.
- Showalter stability index (SI, Showalter 1953): Similar to the lifted index (LI), but using a parcel lifted from 850
130 hPa to 500 hPa. As for the LI, negative SI values indicate instability.
- Convective Available Potential Energy (CAPE, Moncrieff and Miller 1976): The integrated energy excess of an air parcel lifted adiabatically with respect to its environmental temperature profile. Values in the order of hundreds of J Kg⁻¹ and more correspond to severe weather.

Maps of the Modified K-Index (MKI) were derived from the ERA5 ECMWF data that cover the Earth on a 30km
135 grid. The original K-Index (KI; Geer 1996), which combines instability and moisture availability, is used to predict severe thunderstorms in the US. The modified version of Haratz et al. (2010), MKI, is the version adapted for the eastern Mediterranean:

$$MKI = (T_{500} - T_{850}) \cdot RH_{850,700} + Td_{850} - (T_{700} - Td_{700}) , \quad (4)$$

140 where T and Td are temperature and dew point, respectively, RH is relative humidity and the subscripts refer to the respective pressure level (hPa). The modification gives more weight to the relative humidity at the 850 and 700 hPa levels.


In addition, we also used Precipitable Water (PW, Liu, 1986): The depth of water in the atmospheric column, if all the water in that column were precipitated as rain. The values were calculated from the soundings of Beit Dagan. METEOSAT Second Generation Water vapour imageries were retrieved from EUMETSAT data centre
145 (<https://navigator.eumetsat.int/product/EO:EUM:DAT:MSG:HRSEVIRI>).

Air back-trajectories for detecting moisture transport were retrieved from the site of NOAA HYSPLIT MODEL (Stein et al., 2015, <https://www.ready.noaa.gov/HYSPLIT.php>) Global Data Assimilation System (GDAS $1^\circ \times 1^\circ$ resolution, on pressure levels), which is used by the National Center for Environmental Prediction (NCEP) Global Forecast System (GFS) model to place observations into a gridded model space for initializing weather forecasts
150 with observed data. The 'Model vertical velocity' (derived from the gridded data) option was used.

3 Observational analysis

3.1 Synoptic evolution


During April 25–27 2018 the Levant was dominated by an upper-level closed cyclone, accompanied by a lower-level cyclone, located east of Israel (see Fig. 2).



155


Figure 2: Geopotential height (GPH) at 500 hPa (m) and SLP (hPa) projected on MODIS imagery, in 24 h intervals, for April 24–27 2018, 12UTC (a–d, respectively).

160 Prior to the storm, on April 24, 2018, the lower levels were influenced by a RST. The axis of the RST was located east of the Levant, enclosing a shallow cyclone, with a central pressure of 1008 hPa, over north western Saudi Arabia (see Fig. 2a), transporting warm and dry air from the east toward the Levant. At the 500 hPa level, a closed cyclone was approaching the region from the central Mediterranean. During April 25 the upper-level cyclone slowed its progress, from $\sim 9 \text{ ms}^{-1}$ to $\sim 4 \text{ ms}^{-1}$, while arriving to the Egyptian coast (see Fig. 3). The surface shallow cyclone deepened, attaining a central pressure of 1000 hPa, and moved northward, to Iraq, resembling the Syrian Low (Kahana et al. 2002). The winds over the south-eastern Mediterranean backed to north-westerly, hence advecting moist air from the Mediterranean inland, to central and south Israel. The rain in Israel started at April 25 and stopped at April 27 (Porat et al., 2018). The spatial distribution of the rain during the entire storm is shown in Fig. 1. The maximum rainfall was obtained east of the water divide of the Negev Mountains, and the second most intense one was found in the Jordan valley, 200 m BSL.



170
175 **Figure 3: Track of the upper-level cyclone (500 hPa GPH) during 24–27 April 2018, in 12 hours intervals. The instantaneous speed (ms^{-1}) and distance (km) spanned during each increment are denoted by s and d, respectively. For each 24 hours increment, the radius of the upper-level low (km), the maximum relative vorticity (s^{-1}), the precipitable water (PW, in mm) and 3 thermodynamic indices (LI, SI, CA) as calculated from the sounding of Beit Dagan are specified. "High", "Elevated" and "Low" values of the indices are highlighted in red, yellow and green, respectively**


During the following day, April 26, the upper-level cyclone continued its slow eastward progression (Figs. 2c and 3). The weakening of the horizontal pressure gradient at the surface in April 27 (compare Figs. 2c and 2d), in tandem with a flattening of the 500-hPa cyclone and its movement further eastward, led to the cessation of the stormy weather over Israel.



185 **Figure 4: Potential Vorticity (PVU) at 300 hPa for April 26 00 UTC, taken from ERA-5, at 30 km spatial resolution. The major feature is a closed positive anomaly (with maximum of $\text{K m}^2 \text{Kg}^{-1} \text{s}^{-1}$), corresponding to a cutoff low, centred over north-western Saudi Arabia.**

The upper-level cyclone, beside its geopotential depth (up to 100 m, Fig. 2), was accompanied by an isolated PV maximum as seen in the 300 hPa map for April 26 00 UTC (Fig. 4), indicating its being a cutoff low (Hoskins et al. 1985). Moreover, its evolution took place southeast of an upper-level blocking high that covered East Europe during April 24 and 25, as can be inferred from the PV distribution seen in Figs. 5a-c. The interaction between the blocking high and the cutoff low and its contribution to the slowing of the cutoff low while approaching the Levant are elaborated in Sec. 4 below.

195



200 **Fig. 5: PV distribution on 300 hPa, starting from April 25 2018, 12 UTC, in 6 hour intervals, covering Europe and the Mediterranean Basin, on which a conceptual sketch of a dipole type block (following Yamazaki and Itoh, 2013) is superposed. A cutoff low is seen over the south-eastern Mediterranean and a blocking high over Eastern Europe, forming a dipole. The arrows show the induced flow of each of the vortices.**

205 The synoptic-climatologic background for this storm is compared with the other rain-storms, comprising the set of 12 'reference storms', all of them occurred in the late spring and produced considerable surface runoff over the study area (see Sec. 2 above). In all of the reference storms, except one, upper-level closed cyclones were found over the EM or the Levant. The cyclones' depth varied between 10 and 130 m, with an average of 52 m, and their radii ranged from 200 to 900 km, with an average of 600 km (Table 1). The MCV for the various days belonging to the reference storms varied between 0.8 and $3.8 \times 10^5 \text{ s}^{-1}$, with an average of $2.3 \times 10^5 \text{ s}^{-1}$. The maximum value was observed during the present storm. The MCV for the present storm was $\sim 1/2$ that of the respective maximum relative vorticity (Fig. 3), presumably due to the spatial smoothing implied by the derivation method of the MCV.

210 During the approach of the cutoff low, the atmosphere over the study region became more and more unstable. Figure 3 shows the values of leading instability indices (specified in Sec. 2) derived from Beit Dagan soundings, in 24-hour increments. They reflect a gradual increase toward April 25 12 UTC, while the upper-level cyclone approached the Sinai Peninsula. At that time, the value of CAPE was 779 J Kg^{-1} , of LI: -4.9 K and of SI: -2.7 K , indicating a potential for thunderstorms. In April 26, the CAPE rose to 909 J Kg^{-1} .

220 The transition from the dry conditions that persisted at April 24 to the wet conditions that developed toward April 25 is manifested by a sharp increase in the precipitable water (PW) in Beit Dagan, from 17 mm (April 24 12 UTC) to 30 mm in April 25 12 UTC (27 mm in April 26). The moisture increased presumably due to the north-westerly wind, as evidenced by the air trajectories entering the study area at the heights of 600 and 1000 m. The trajectory

that entered the region at 1400 m originated from Iraq, entered northern Israel from the east and, after turning cyclonically, penetrated the Negev Desert from northwest (Fig. 6). The signature of the segment of the latter trajectory, extending from Iraq toward north Israel, is clearly seen as a bright band in the satellite imagery of the water vapor channel for April 25 21 UTC (12 hours prior to Tzafit flood, Fig. 7), reflecting a moisture strip.

225

Table 1. Depth, radius and the measure of curvature vorticity (MCV) of the 500 hPa cyclones for the 27 days, in which the highest surface runoff was observed over the study area in the late spring during the period 1986–2018 (the reference storms). Note that in 24 of the days an upper-level cyclone was identified.

Storm No.	Date	GPH Depth (m)	Radius (km)	MCV ($\times 10^5 \text{ s}^{-1}$)
1	19860402	70	650	2.3
1	19860403	25	400	2.1
2	19860408	45	650	1.4
2	19860409	45	900	0.8
3	19970514	no	no	
3	19970515	no	no	
4	20010404	130	900	2.2
5	20010430	15	250	3.3
5	20010501	25	350	2.8
5	20010502	90	850	1.7
5	20010503	90	750	2.2
6	20070512	50	580	2.0
7	20110404	70	900	1.2
7	20110405	20	280	3.5
8	20140507	15	330	1.9
8	20140508	50	550	2.2
9	20140511	35	500	1.9
9	20140512	10	200	3.4
10	20150416	30	350	3.3
11	20160410	no	no	
11	20160411	45	550	2.0
11	20160412	50	650	1.6
11	20160413	40	550	1.8
11	20160414	45	520	2.3
12	20180425	100	600	3.8
12	20180426	80	550	3.6
12	20180427	70	600	2.6

230




Figure 6: Sixty hours back-trajectory of the air arriving at Tzafit Basin at April 26, 2018 10 UTC at heights 600, 1000 and 1400 m.




Figure 7: Water Vapor channel 6 (6.85–7.85 μm) imagery of METEOSAT from April 25, 2018 21 UTC showing the cloud strip entering Israel from the east and the vortex over the south-eastern Mediterranean.


3.2 April 26: main rain episodes

240 In the early morning of April 26, the northern part of Israel was the only part receiving rain, mostly moderate, originating from stratified cloudiness (based on hourly radar images, not shown). After several hours of fair weather, at the noon hours, three major rain centres developed. Figure 8 shows them through two observational means. i.e., satellite IR and radar imagery, and two diagnostic tools, vorticity field and MKI, both at 30 km spatial resolution. The first rain centre, composed of several cells, was most active between 0900-1030 UTC and produced
245 the Tzafit flood. This centre yielded over 30 mm rainfall, with a maximum intensity of 180 mm/h, based on instantaneous radar measurements. This intensity has a return period of > 75 years for the pertinent region (Rinat et al. 2020). This centre can be noted in the radar image for 09 UTC (Fig. 8c).

Later on, around 12 UTC, another rain centre developed in the Jordan valley around $32.5^{\circ}\text{N } 35.5^{\circ}\text{E}$, at north Israel, in a form of a meso-scale cell, as reflected by the satellite and radar images (Figs. 8b, d). This cell yielded rainfall
250 of 30-45 mm, with a maximum of 72 mm (record breaking since start of record at 1943). The maximal 10 min averaged rate was 116 mm/h. The third rain centre started to develop at 12 UTC and lasted ~ 2 hours. This centre was aligned with the divide of the Samarian and Judean Mountains and its adjacent slopes facing the north-westerly winds (Figs. 8b, d), so it can be considered as forced by orography. This rain centre yielded 30-35 mm/h, with 10 min averaged maximum rate of 100 mm/h (in Jerusalem). The rain centres formed over Tzafit Basin and over the
255 Jordan Valley developed under highly unstable conditions, expressed by $\text{MKI} > 30^{\circ}\text{C}$, whereas the third one, over the mountain ridges, was associated with MKI around 25°C . In the late afternoon, southern Israel was still under precipitative cloudiness (not shown).

4 Discussion

Precipitation extremes in the Mediterranean Basin are usually induced by large-scale atmospheric circulation,
260 namely low-pressure systems with large pressure gradients (Jacobeit et al. 2017). Therefore, the analysis starts from the synoptic system, with which this storm was associated, namely the upper level cutoff low. This cutoff low entered the EM from northwest, deepened and crossed the Levant at 30°N latitude (Figs. 2, 3). The maximum vorticity measured in 500 hPa level was $> 7.2 \times 10^{-5} \text{ s}^{-1}$, which is comparable with the value of $5 \times 10^{-5} \text{ s}^{-1}$ at the same level found by Flokas et al. (2001) for the winter season in the Mediterranean. Apparently, this implies that the
265 synoptic factor was dominant in rain production. However, convective nature of the rain and spotty distribution was demonstrated by radar images (Figs 8c, d) and was even reflected by the rain map of the entire storm (Fig. 1).



270 **Figure 8: April 26 imageries for 09 (a,c,e,g) and 12 UTC (b,d,f,h) corresponding to the major rain centres including MSG IR images (a,b), one hour forward integration Radar [mm h^{-1}] images (c,d), relative vorticity [$\text{s}^{-1} \times 10^5$] and streamlines at 500 hPa (e,f), and MKI [$^{\circ}\text{C}$] (g,h). Panels e-f are based on ERA5 data at 30 km spatial resolution.**

Following the quasi-geostrophic approach, most convective outbreaks occur in broad south westerly flow aloft ahead of an approaching trough (Doswell, 1987). In April 26, under the north westerly flow behind the upper-level cyclonic system, its direct dynamic supporting effect on rain formation is expected to be weak, or even negative (Fig. 2c). At this sector, negative upper-level vorticity advection and lower-level cold advection are expected, both implying subsidence in the mid-levels. Inspection of the lower-level temperature and wind fields (not shown) indicate that at that stage, the cold core associated with the upper-level cyclone was centred over the Negev Desert. The lower-level winds over the study region were parallel to the isotherms, so that no temperature advection existed. Similar configuration existed with respect to the upper-level vorticity field, implying that vorticity advection also did not take place there. Moreover, the omega field at the mid-levels in the synoptic scale over the region was near zero. The above implies that the synoptic scale dynamics did not have a direct effect on the rain formation on that day.

The major synoptic factor that directly contributed to the rain formation in this storm is the wind. One implication is the onshore moisture transport, accompanied by an uplift imparted by its encounter with the coastline and later on, with the mountain ridges. The other is the upper-level cold advection, leading to thermal instability. Actually, only a small fraction of the rainfall in this storm was orographic (over Judean Mountains), whereas the majority of the rain was observed far from the coastline and beyond the water divide of the mountain ridges (Fig. 1, note the divide line). Two out of the three major rain centres were convective: the one that produced the flood in Tzafit, over 60 km inland, and the other, developed in the Jordan valley, >200 m BSL. The dominance of convective over orographic rain suggests that instability was the prominent factor in this storm.

In spite of the affinity of the Mediterranean cyclones to mid-latitude cyclones, the rain they produce highly depends on moisture supply, rather than on the presence of conveyor belts or front, due to the limited moisture sources in the Mediterranean Basin (Ziv et al. 2010). The lower-level system in the storm analysed here, can be considered as a 'Syrian low' (Kahana et al., 2002, see Sec. 1 above), which belongs to the Mediterranean cyclones. The Syrian low resembles the 'deep low to the east', which is one of the 7 types of Cyprus low defined by Alpert et al. (2004). Saaroni et al. (2010) showed that most of the rain associated with the 'deep low to the east' is distributed over the coastal regions and the western slopes of the Judean Mountains, facing the offshore north westerly winds blowing from the Mediterranean. This indicates that the Judean Mountains, extending up to 800-1000 m, block effectively the moisture, which is presumably, concentrated in the lower-levels. The heavy rains that were observed inland in this storm, and the signature of non-orographic major rain cells in the rain maps indicate that the moisture sources were not limited to the lower-levels. The presence of moisture at the mid-levels is deduced from back-trajectory

arriving at Tzafit at the time where the flood occurred (Fig. 6). This show a band of mid-level moist air that originated east of the Levant, turned cyclonically around the cyclone centre and entered the region from the west.

In the EM, rains associated with Mediterranean cyclones, as in the case studied here, are convective in nature. In the winter season, cold air originating from south Europe moves over the warmer Mediterranean water and becomes unstable before entering Israel (Shay-El and Alpert 1991). In the late spring, Europe becomes warmer, whereas the Mediterranean remains cool, due to its lagged response to the annual cycle, so that the passage of a Cyprus low over the EM does not necessarily lead to instability. In the case studied here, the instability can be attributed to a negative temperature anomaly in the upper levels (in the order of -4 K in 500 hPa, not shown) that covered the southern Levant, as a part of the cutoff low. A tendency of the rain in the Negev Desert to be convective, especially in the transition seasons, has been noted previously. Sharon (1978) pointed at a gradual increase in the proportion of localized showers in the transition from the Mediterranean part of Israel further into the more arid south. Systematic evidence for this tendency is well reflected in the low spatial correlation obtained among several rain stations for April over southern Israel (Kutiel, 1982).

The occurrence of the major rains in the noon hours of 26 April (12 LST for Tzafit and 14 LST for Beit Shean) suggests that the convection was enhanced by intense solar radiation, as implied by the date, latitude and the clear sky (Fig. 8a) during the morning hours. This effect is reflected by the MKI maps (Figs. 8g,h), showing much higher values inland compared to these over the coastal region ($>30^{\circ}\text{C}$ in Tzafit Basin compared to 13.2°C in Beit Dagan). The mesoscale features seen in the 500 hPa vorticity maps for April 26 12 UTC (Fig. 8f) indicate supportive forcing on the formation of the rain cell developed near Beit Shean. This cell was formed under a region of positive vorticity advection (by easterly winds) ahead of a positive core of $>2 \times 10^{-4} \text{ s}^{-1}$, with >100 km diameter, centred over western Jordan (32.3°N , 36.7°E). This core was one of the two vorticity centres that constituted the synoptic scale cutoff low at 12 UTC (Fig. 2). It is possible that the conic structure of the Jordan rift near Beit Shean, narrowing southward, caused the northerly lower-level winds to horizontally confluent, and consequently to ascend. The convective rains observed over the Judean Mountains at the noon hours are regarded here as initiated by orography. Despite the major direct contribution of the instability, moisture supply and topographic features to the evolution of the major rain cells, the role of the synoptic factor, namely the cutoff low, can be considered as a supportive background for these effects. The impact of cutoff lows on rainstorms over the Mediterranean has also been shown by Porcu et al. (2007) and by Fragoso and Tildes Gomes, 2008 (Sec. 1). This goes in line with the fact that 11 out of the 12 reference rainstorms that occurred over the study region in the late spring were accompanied by upper-level closed cyclones.

The slow eastward propagation of the cutoff low exacerbated the severity of the storm by extending its duration. The typical translation speed of Mediterranean cyclones is 5-10 m s⁻¹ (Alpert and Ziv, 1989). The speed of the cutoff low was 7-9 m s⁻¹ while moving along the Libyan coast, but slowed down to 3-5 m s⁻¹ while approaching the EM (Fig. 3). The series of 300 hPa PV maps (Figs. 5a-c), on which a conceptual sketch of a dipole type is superposed, consisting of a blocking high located north of the cut-off low, follows Yamazaki and Itoh (2013). These charts show a lobe of negative PV (blue) over the Balkans, north of the cutoff low, lingering over the Nile delta. The power of PV maps stems from the feature of PV anomalies (in the spatial sense) to induce circular flow out of their boundaries, i.e., cyclonic for positive and anticyclonic for negative. The negative anomaly centred over Eastern Europe exerted an anticyclonic flow over the region, which produced easterly winds south of it, including the Levant, where the positive anomaly (with which the cutoff low was associated) was located. At the same time, the positive anomaly over the southern Levant exerted a cyclonic flow around it, including easterly winds over Eastern Europe, where the negative anomaly was positioned. This is most clearly demonstrated in Fig. 5c. The outcome of the superposed effects is that both anomalies were subjected to forces acting against the mid-latitude westerlies. This explains the slow progression of the cutoff low, while reaching the Levant.

5. Summary

An intense rainstorm hit the Middle East between 24 and 27 April 2018, producing heavy flash floods in Israel that took the lives of 13 people. Ten of them lost their lives in April 26, in a flash flood that resulted from rain showers with intensities of >75 years return period. The study describes the major rain systems affecting Israel in that day, and analyses the atmospheric processes leading to their formation. The dominating synoptic feature was a cyclone in all levels that crossed Israel in its way eastward. The rain distribution for the entire storm was not evenly distributed. Three major rain centres were active during April 26. One, which caused the fatal flood, developed over a relatively flat terrain, the second developed upslope the mountains, i.e., orographic, and the third formed in the Jordan valley. The rain distribution implies that synoptic and sub-synoptic atmospheric processes acted in concert.

The synoptic background of the storm was an upper-level cutoff low that originated from south Europe and slowed its movement while approaching the Levant. This cutoff low was found to have the highest vorticity with respect to the set of 12 'reference storms' observed over the study region in the late spring during the latest 33 years. The lower levels were dominated by a cyclone, centred east of Israel. The implied winds were north-westerly, hence advecting moist air from the Mediterranean inland. During the approach of the upper-level cyclone, the atmosphere

over the study region became conditionally unstable, with instability indices reaching values indicating potential for thunderstorms. At the same time, the precipitable water increased by a factor of 2 over the study region, which triggered deep moist convection from April 25 onward. The mesoscale distribution of MKI (Modified K index) indicated that the instability was considerably higher inland than along the coastal plain. This may explain why most of the rain was observed there. In addition to the lower-level moisture advection, a mid-level band of moist air, extending from Iraq westward, curved cyclonically through Syria and the EM and entered south Israel from northwest. The combination of both moisture contributions can explain the severe rain showers over Israel observed during April 26.

The present study raises several questions for further research. One of them concerns the involvement of closed upper-level cyclones as a major factor for severe rainstorms over the Negev Desert in the late spring. The spring season in the Mediterranean is characterized by a general weakening and reduced frequency of Cyprus lows. Moreover, the late spring is the time in which the descending branch of the Hadley cell takes over the Mediterranean. In light of the above, the specific conditions that lead to intensification of Rossby waves that lead to their breaking into closed cyclones should be investigated.

The most intense showers associated with the storm studied here were observed while the study region was situated west of the upper-level cyclone, where negative vorticity advection is expected. Similar situations were noted by Kahana et al. (2002) in the framework of the 'Syrian low', which produced major floods in the Negev Desert. A similar situation was analysed by Morin et al. (2007), which resulted in a fatal flood in Israel, between 31 March and 1 April 2006.

Another issue is the need to gain a better insight on the factors initiating convection, which are not supplied by the large-scale, but rather in the smaller scales. Such factors are not well captured by the finer resolution data; therefore, their identification necessitates the use numerical simulations.

Acknowledgements:

The authors wish to thank the reviewers for their valuable comments, which considerably contributed to the quality of the paper. The authors thank Noam Halfon and Yoav Levi from the IMS for the rain map radar images and Efrat Morin and Yair Rinat from HUJI for the hydrometric data. BZ thanks the Israeli Science Foundation (ISF, grant number 1123/17).

390 Data availability. ERA5 ECMWF reanalysis data have been downloaded from the Copernicus Climate Data Store (<https://cds.climate.copernicus.eu>, Copernicus, 2020). Meteosat Second Generation water vapour imageries were retrieved from EUMETSAT data centre (<https://navigator.eumetsat.int/product/EO:EUM:DAT:MSG:HRSEVIRI>). Air back-trajectories for detecting moisture transport were retrieved from the site of NOAA HYSPLIT MODEL (Stein et al., 2015, <https://www.ready.noaa.gov/HYSPLIT.php>)

395

Author contributions. UD conceived the study, set the goals of the article, and developed its methodology. IML, selected and analyzed the data sources. BZ was engaged in the dynamic aspects of the study and their issuing calculations. All the authors contributed to the interpretation of the results and to the manuscript writing

400 Competing interests. The authors declare that they have no conflict of interest.

References

- Alpert, P., Osetinsky, I., Ziv, B., and Shafir, H.: Semi-objective classification for daily synoptic systems, Application to the Eastern Mediterranean climate change, *Int. J. Clim.*, 24, 1001-1011, DOI: 10.1002/joc.1036, 2004.
- 405 Ashbel, D.: Great floods in Sinai Peninsula, Palestine, Syria and the Syrian Desert, and the influence of the Red Sea on their formation. *Quart. J. R. Met. Soc.* 64: 635 – 639, 1938.
- Aref, H.: Integrable, chaotic, and turbulent vortex motion in two-dimensional flows. *A. Rev. Fluid Mech.* 15, 345-389, 1983.
- Armon, M. Morin, E., and Enzel, Y.: Overview of modern atmospheric patterns controlling rainfall and floods into the Dead Sea: Implications for the lake's sedimentology and paleohydrology, *Quaternary Science Reviews*, 216, 58-73 DOI: 0.1016/j.quascirev.2019.06.005, 2019.
- 410 Bardossy, A., and Filiz, F.: Identification of flood producing atmospheric circulation patterns, *J. of Hydrology*, 313(1–2), 4857, <https://doi.org/10.1016/j.jhydrol.2005.02.006>, 2005
- Becker, J.S., Taylor, H.L., Doody, B.J., Wright, K.C., Grunfest, E., and Webber, D.: A review of people's behavior in and around floodwater. *Weat., Clim. Soc.* 7, 321–332, DOI: 10.1175/WCAS-D-14-00030.1, 2015.
- 415 Campa, J., and Wernli, H.: A PV Perspective on the Vertical Structure of Mature Midlatitude Cyclones in the Northern Hemisphere, *J. Atmos. Sci.* 69, 725-740, <https://doi.org/10.1175/JAS-D-11-050.1>, 2012.
- Campins, J., Genoves, A., Picornell, M.A., and Jansa, A.: Climatology of Mediterranean cyclones using the ERA-40 dataset, *Int. J. Clim.*, 31, 11, 1596-1614, <https://doi.org/10.1002/joc.2183>, 2011.
- 420 Copernicus Climate Change Service (C3S): ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate. Copernicus Climate Change Service Climate Data Store (CDS), accessed: 25 December 2019. <https://cds.climate.copernicus.eu/cdsapp#!/home>, 2017.
- Dayan, U., and Morin, E. : Flash Flood-Producing rainstorms over the Dead Sea, Israel: A Review, in Enzel, Y., Agnon, A., and Stein, M., eds., *New Frontiers in Dead Sea paleo-environmental research: Geological Society of America Special Paper 401*, doi: 10.1130/2006/2401(04), 2006.
- 425 Dayan, U. and Sharon, D.: Meteorological Parameters for Discriminating Between Widespread and Spotty Storms in the Negev. *Israel Journal of Earth Sciences*, 29 (4), 253-256, 1980.
- Dayan U., B. Ziv, A. Margalit, E. Morin, D. Sharon,: A severe autumn storm over the middle-east: synoptic and meso-scale convection analysis, *Theor. App. Clim.*, 69, 1/2, 103-122, 2001.
- 430 Diakakis, M., Priskos, G. and Skordoulis, M.: Public perception of flood risk in flash flood prone areas of Eastern Mediterranean: the case of Attica Region in Greece. *International Journal of Disaster Risk Reduction*, 28, 404-413, <https://doi.org/10.1016/j.ijdrr.2018.03.018>, 2018.
- Flocas, H.A., Maheras, M.P., Karacostas, T.S., Patrikas, I. and Anagnostopoulos, C.: A 40-year climatological study of relative vorticity distribution over the Mediterranean, *Int. J. Climat.*, 21, 1759 – 1778, doi: 10.1002/joc.705, 2001.
- 435 Fragoso M, Tildes Gomes P.: Classification of daily abundant rainfall patterns and associated large-scale atmospheric circulation types in Southern Portugal. *Int. J. Climatol.* 28(4): 537–544, doi: 10.1002/joc.1564, 2008.

- Galateia, T., Ruin, I., Anquetin, S., and Gourley, J.J.: A situation-based analysis of flash flood fatalities in the united states, *B. Am. Meteorol. Soc.*, 333-345, <https://doi.org/10.1175/BAMS-D-15-00276.1>, 2017.
- 440 Galway, J. G.: The lifted index as a predictor of latent instability, *B. Am. Meteorol. Soc.*, 37, 10, 528-529, 1956.
- Geer, I. W.: Glossary of Weather and Climate, *Am. Meteorol. Soc.*, 131, 1996.
- Greenbaum, N., Schwartz, U., and Bergman, L.: Extreme floods and short-term hydroclimatological fluctuations in the hyper-arid Dead Sea region, Israel, *Global and Planetary Change*, 70, 1–4, 125-137, <https://doi.org/10.1016/j.gloplacha.2009.11.013>, 2010.
- 445 Haiden, T., Kann A., Wittmann C., Pistotnik G., Bica B., and Gruber C.: The Integrated Nowcasting through Comprehensive Analysis (INCA) System and Its Validation over the Eastern Alpine Region. *Wea. Forecasting*, 26, 166–183, <https://doi.org/10.1175/2010WAF2222451.1>, 2011.
- Harats, N., Ziv, B., Yair, Y., Kotroni, V., and Dayan, U.: Dynamic and thermodynamic predictors for lightning and flash floods in the Mediterranean, *Adv. Geophys.*, 23, 57-64, 2010
- 450 HMSO. *Weather in the Mediterranean I: General Meteorology*. 2nd edn. Her Majesty's Stationery Office: London, 362, 1962.
- Holton, J. R.: *An Introduction to Dynamic Meteorology*. 2d ed. Academic Press, 507 pp, 1992.
- Hoskins, B. J., McIntyre, M.E., and Robertson, A.W.: On the use and significance of isentropic potential vorticity maps, *Q. J. Roy. Meteor. Soc.*, 877-946, <https://doi.org/10.1002/qj.49711147002>, 1985.
- 455 Jacobeit, J., Homann, M., Philipp, A., and Beck, C.: Atmospheric circulation types and extreme areal precipitation in southern central Europe. *Advances in Science and Research* 14: 71-75, 2017.
- Kahana, R., Ziv, B., Enzel, Y. and Dayan, U.: Synoptic climatology of major floods in the Negev Desert, Israel. *Int. J. Climat.* 22, 867-882, <https://doi.org/10.1002/joc.766>, 2002.
- Kahana R.: *Synoptic Hydro-climatology of Major Floods in the Negev and Arava, Southern Israel*. M.Sc Thesis, Institute of Earth Sciences, The Hebrew University of Jerusalem (in Hebrew), 1999.
- 460 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-years reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–472,
- 465 [https://ui.adsabs.harvard.edu/link_gateway/1996BAMS...77..437K/doi:10.1175/1520-0477\(1996\)077%3C0437:TNYRP%3E2.0.CO;2](https://ui.adsabs.harvard.edu/link_gateway/1996BAMS...77..437K/doi:10.1175/1520-0477(1996)077%3C0437:TNYRP%3E2.0.CO;2), 1996.
- Kistler, R., Kalnay, E., Collins, W., Saha, S., White, G., Woollen, J., Chelliah, M., Ebisuzaki, W., Kanamitsu, M., Kousky, V., Van Den Dool, H., Jenne, R., and Fiorino, M.: The NCEP-NCAR 50-year Reanalysis: Monthly means CD-ROM and documentation, *B. Am. Meteorol. Soc.*, 82, 247–267, [doi:10.1175/1520-0477\(2001\)082<0247:TNNYRM>2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<0247:TNNYRM>2.3.CO;2), 2001.
- 470 Krichak, S.O., Alpert, P., and Dayan, M.: An evaluation of the role of hurricane Olga (2001) in an extreme rainy event in Israel using dynamic tropopause maps. *Meteor Atmos Physics*, 98, 1-2, 35-53, DOI 10.1007/s00703-006-0230-7, 2007.
- Kushnir, Y., Dayan U., Ziv B., Morin E., and Enzel Y.: "Climate of the Levant: phenomena and mechanisms" in *"Quaternary of the Levant: Environments, Climate Change, and Humans"*, Y. Enzel and O. Bar-Yosef Editors. Cambridge University Press, London (pp 31-44), 2017.
- 475

- Kutiel, H.: Spatial Coherence of Monthly Rainfall in Israel, *Arch. Met. Geoph. Biokl., Ser. B*, 31,353-367, 1982.
- Liu, W. T.: Statistical relation between monthly mean precipitable water and surface-level humidity over global oceans, *Mon. Weather Rev*, 114, 1591-1602, [http://dx.doi.org/10.1175/1520-0493\(1986\)114<1591:SRBMMP>2.0.CO;2](http://dx.doi.org/10.1175/1520-0493(1986)114<1591:SRBMMP>2.0.CO;2), 1986.
- 480 Marra, F., and Morin, E.: Use of radar QPE for the derivation of Intensity–Duration–Frequency curves in a range of climatic regimes, *Journal of Hydrology* 531, 427–440. <http://dx.doi.org/10.1016/j.jhydrol.2015.08.064>, 2015.
- Moncrieff, M.W. and Miller, M., J.: The dynamics and simulation of tropical cumulonimbus and squall lines, *Quart. J. R. Met. Soc.*, 102, 373-394, 1976.
- 485 Montz, B.E., Gruntfest, E.: Flash flood mitigation: Recommendations for research and applications. *Global Environmental Change Part B: Environmental Hazards*, 4(1), 15–22, [https://doi.org/10.1016/S1464-2867\(02\)00011-6](https://doi.org/10.1016/S1464-2867(02)00011-6), 2002.
- Morin, E., Harats, N., Jacoby, Y., Arbel, S., Getker, M., Arazi, A., Grodek, T., Ziv, B., and Dayan, U.: Studying the extremes: hydrometeorological investigation of a flood-causing rainstorm over Israel, *Adv. Geosci.*, 12, 107–114, <https://doi.org/10.5194/adgeo-12-107-2007>, 2007.
- 490 Palmen, E. and Newton, C.W. (Eds.): *Atmospheric Circulation Systems*. Academic Press, New York, 1969.
- Porat, A, Halfon, N and Forshpan A.: Examining the exceptionality of 25-27 April 2018, *IMS August* (in Hebrew), 2018.
- Porcù, F., Carrassi, A., Medaglia, C. M., Prodi, F., & Mugnai, A.: A study on cut-off low vertical structure and precipitation in the Mediterranean region. *Meteorology and Atmospheric Physics*, 96(1-2), 121-140, 2007.
- 495 Rinat, Y., Marra, F., Armon, M., Metzger, A., Levi, Y., Khain, P., Vadislavsky, E., Rosensaft, M., and Morin, E.: Hydrometeorological analysis and forecasting of a 3-day flash-flood-triggering desert rainstorm, *Nat. Hazards Earth Syst. Sci. Discuss.*, <https://doi.org/10.5194/nhess-2020-189>, in review, 2020.
- Saaroni, H., Halfon, N., Ziv, B., Alpert, P., and Kutiel, H.: Links between the rainfall regime in Israel and location and intensity of Cyprus lows, *Int. J. Climat.*, 30, 1014-1025, DOI: 10.1002/joc.1912, 2010.
- 500 Sharif, H., Jackson, T., Hossain, M., and Zane, D.: Analysis of flood fatalities in Texas. *Nat. Hazards Rev.* 16, 04014016, DOI: 10.1061/(ASCE)NH.1527-6996.0000145, 2015.
- Sharon, D.: Rainfall Fields in Israel and Jordan and the Effect of Cloud Seeding on Them. *J. Appl. Met.* 17, 40-48, 1978.
- 505 Shay-El, Y., and Alpert, P.: A diagnostic study of winter diabatic heating in the Mediterranean in relation to cyclones, *Quarterly Journal of the Royal Meteorological Society* 117, 715–747, <https://doi.org/10.1002/qj.49711750004>, 1991.
- Showalter, A., K.: A Stability Index for Thunderstorm Forecasting. *Bull. Amer. Meteor. Soc.*, 34, 25-252, 1953.
- 510 Stein, A.F., Draxler, R.R, Rolph, G.D., Stunder, B.J.B., Cohen, M.D., and Ngan, F.: NOAA’s HYSPLIT atmospheric transport and dispersion modeling system, *Bull. Amer. Meteor. Soc.*, 96, 2059-2077, <http://dx.doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- Terti, G., Ruin, I., Gourley, J.J., Kirstetter, P., Flamig, Z., Blanchet, J., Arthur, A. and Anquetin, S.: Toward probabilistic prediction of flash flood human impacts. *Risk Analysis*, 39, 140-161, doi: 10.1111/risa.12921, 2019.

- Uccellini, L. W., and Kocin, P. J.: The interaction of jet streak circulations during heavy snow events along the east coast of United States. *Wea. Forecasting*, 2, 289–308, 1987.
- Wei, M., Pasquero, C., and Primeau, F.: The effect of translation speed upon the intensity of tropical cyclones over the tropical ocean, *Geophysical Research Letters*, 39, L07801, doi:10.1029/2011GL050765, 2012.
- Yamazaki, A., and Itoh, H.: Vortex-vortex interactions for the maintenance of blocking: Part I: The selective absorption mechanism and a case study, *Journal of the Atmospheric Sciences*, 70, 725-742, DOI: 10.1175/JAS-D-11-0295.1, 2013.
- Zappa, G., Hawcroft, M.K., Shaffrey, L., Black, E., and Brayshaw, O.J.: Extratropical cyclones and the projected decline of winter Mediterranean precipitation in the CMIP5 models, *Climate Dynamics*, doi: 10.1007/s00382-014-2426-8, 2015.
- Ziv, B., and Paldor, N.: The Divergence field associated with time dependent jet streams, *J. Atmos. Sci.*, 56 No. 12, 1843-1857, 1999.
- Ziv, B., Dayan, U., Kushnir, Y., Roth, C., and Enzel, Y.: Regional and Global Atmospheric Patterns Governing Rainfall in the Southern Levant, *Int. J. Clim.*, 26, 55-73, 2006.
- Ziv, B., Saaroni, H., Romem, M., Heifetz, E., Harnik, N., and Baharad, A.: Analysis of conveyor belts in winter Mediterranean cyclones, *Theor. App. Clim.*, 99, 3-4, 441 – 455, 2010.
- Ziv, B., Saaroni, H., Pargament, R., Harpaz, T., and Alpert, P.: Trends in rainfall regime over Israel, 1975-2010, and their relationship to large-scale variability, *Reg Environ Chan*, 14(5), 1751-1764, 2014.
- Ziv, B., Harats, N., Morin, E. et al. Can severe rain events over the Mediterranean region be detected through simple numerical indices? *Nat Hazards* 83, 1197–1212, 2016.