



## 1 CHILDA – Czech Historical Landslide Database

2 Michal Bíl<sup>1</sup>, Pavel Raška<sup>2</sup>, Lukáš Dolák<sup>3,4</sup>, Jan Kubeček<sup>1</sup>

3 <sup>1</sup> CDV – Transport Research Centre, Brno, 636 00, Czechia

4 <sup>2</sup> Department of Geography, J. E. Purkyně University in Ústí nad Labem, Czechia

5 <sup>3</sup> Department of Geography, Masaryk University in Brno, Czechia

6 <sup>4</sup> Global Change Research Institute of the Czech Academy of Sciences, Brno, Czechia

7 *Correspondence to:* Michal Bíl ([michal.bil@cdv.cz](mailto:michal.bil@cdv.cz))

8 **Abstract:** National and regional historical landslide databases are increasingly viewed as providing empirical  
9 evidence for the geomorphic effects of ongoing environmental change and for supporting adaptive territorial  
10 planning. In this work, we present the design and current content of the Czech Historical Landslide Database  
11 (CHILDA), the first of its kind for the territory of Czechia (the Czech Republic). We outline the CHILDA system,  
12 its functionality and technical solution. The database was established by merging and extending the fragmented  
13 regional datasets for highly landslide prone areas in Czechia. Currently, the database includes 699 records (619  
14 landslides, 75 rockfalls, and 5 other movement types) encompassing the period from the oldest determined records  
15 (1132) up to 1989 which represents an important cultural, political and socioeconomic divide. Along with  
16 characterizing the content of the database, we discuss its further developments and applications.

### 17 1 Introduction

18 Historical landslide inventories and databases are among the key challenges within landslide risk reduction efforts  
19 as they fill the gap, on one hand, between the landslide occurrence in the past environments studied with the use  
20 of various proxies and the present-day landslides, for which different monitoring and mapping techniques may be  
21 used, on the other (e.g., Glade et al., 2001; Raška et al., 2015; Piacentini et al., 2018). In light of the recent global  
22 climate change, the historical landslide databases contribute a better understanding to changes in various  
23 meteorological triggers of landslides in different environmental settings (Gariano and Guzzetti, 2016; Caracciolo  
24 et al., 2017). Given the severe impacts of landslides on society (Froude and Petley, 2018), these databases also  
25 make it possible to identify changes in hot spots of landslide occurrence and the character of their impacts (Klose  
26 et al., 2016). In both these respects, the historical databases complement current landslide inventories that can be  
27 used to trace the spatial patterns in landslide occurrence and their causative factors (Van Den Eeckhaut and Hervás,  
28 2012; Herrera et al., 2018; Marc et al., 2018). Within the landslide hazard and risk assessments, the historical  
29 landslide databases provide evidence as it is generally assumed that past landslide occurrence frequencies may be  
30 used to describe the probabilities of landslide occurrence in the near future (Remondo et al., 2008; Van Den  
31 Eeckhaut et al., 2009; Wu and Yeh, 2020).

32 Connecting these directions, increasing attention has been also paid to revealing the vulnerabilities and adaptive  
33 behaviours of past societies regarding landslides (Tropeano and Turconi, 2004; Caloiero et al., 2014; Klose et al.,



34 2016; Raška, 2019; Klimeš et al., 2020). These studies argue that historical landslide databases – if approached  
35 critically – may inform current efforts for adaptive management of landslide risks (Klose et al., 2016; Raška and  
36 Dubišar, 2017). Historical landslide databases have been recently established for various countries and regions,  
37 for instance, in Italy (Guzzetti et al., 1994; Piacentini et al., 2018), Nicaragua (Devoli et al., 2007), USA (Elliott  
38 and Kirschbaum, 2007), Norway (Hermanns et al., 2013), the United Kingdom (Taylor et al., 2015), Germany  
39 (Damm and Klose, 2015), Portugal (Pereira et al., 2014), most of them covering ca. the last 150 years, but some  
40 databases also including scarce records dating back as early as the twelfth century.

41 The aim of this work is to present the Czech Historical Landslide Database (CHILDA), a project that unified and  
42 extended the fragmented existing regional databases and established an open access and concurrently updated  
43 map inventory of historical landslides in Czechia. The presented database thus further fills in the gap of missing  
44 historical databases for Central-European mid-mountain environments (Damm and Klose, 2015). Within this  
45 paper, *landsliding* is used as a generic term covering all major types of rapid mass movements (cf. Hungr et al.,  
46 2014) that are usually recorded in the documentary data. Slow slope deformations are not studied here since they  
47 usually did not cause rapid harm to society and have not been registered by past societies. For CHILDA content,  
48 we only differentiate the three following groups according to major mechanism: (a) landslides sensu stricto (also  
49 including spreading and flows), (b) rock falls (including topples) in solid bedrock, while (c) all remaining mass  
50 movements are grouped as 'others'. This rough classification is used since the documentary data often do not  
51 allow for detailed and reliable identification of the mass movement type.

52 In the following sections, we will first review the previous studies on historical landslides in Czechia with  
53 emphasis given to attempts to establish systematic historical landslide databases. We will then outline the  
54 availability of the documentary sources and present a design of the CHILDA. Finally, the current content of this  
55 database and its completeness will be presented in a comparative perspective along with discussion of its future  
56 directions. Although CHILDA is an open database, the last analysed year was set to 1989 for the purpose of this  
57 study. The year is considered an important cultural, political and socioeconomic divide in the recent history of  
58 Czechia, turning the country into a democratic regime and thus influencing production, diversity and  
59 accessibility of documentary data and the data from ongoing landslide monitoring.

## 60 **2. Landslides in Czechia**

### 61 **2.1 Landslide Predispositions in Czechia**

62

63 Despite the fact that Czechia can be generally considered a low-risk country, given the relatively low landslide  
64 frequencies and impacts (Klimeš et al., 2017), the country displays high spatial variability in landslide  
65 occurrence with some highly landslide-prone regions due to their predisposition and presence of causative  
66 factors. Among the most affected by landslides are the Outer Western Carpathians (OWC), NW Czechia (České  
67 středohoří Mts., Děčínská vrchovina) and several of the scattered spatially limited areas across the country (see  
68 Figure 1).


69 The eastern part of Czechia, OWC, is built up of Tertiary and Mesozoic flysch rocks which are particularly  
70 susceptible to landsliding. As a consequence, for example, of the 1997 landslide period as many as 3,700



71 individual landslides were mapped in that region (Krejčí et al., 2002). High numbers of reactivated landslides  
72 were also further identified during the periods of intense landsliding which followed, specifically in 2006 (Bíl  
73 and Müller, 2008), and 2010 (Pánek et al., 2011). The Registry of Slope Deformations of the Czech Geological  
74 Survey ([www.geology.cz](http://www.geology.cz)) contains in all approximately 14,500 landslides in this area of the Czech part of the  
75 OWC (7,200 km<sup>2</sup>), which was 82 % of all the landslides registered within Czechia (Bíl et al., 2016).

76 In NW Czechia, two major areas display high landslide susceptibility, namely the České středohoří Mts. (CS)  
77 and the Děčínská vrchovina Highland (DV). The lithology of CS is built up of Neogene volcanic rocks,  
78 including basalts and phonolites and their volcanoclastics, surrounded or underlain by Mesozoic weak  
79 sedimentary rocks (Cajz, 1999). In a rugged terrain, such conditions often result in landsliding accelerated during  
80 the snow-melt season and heavy rainfalls (Hroch et al., 2002; Raška et al., 2014a; CHMI, 2020). DC is formed in  
81 an uplifted and dissected Mesozoic sandstones, which are prone to rockfalls and toppling (Kalvoda and Balatka,  
82 1995; Zvelebil et al., 2005). These mainly occur in the tectonically predisposed and deeply eroded valley of the  
83 Labe/Elbe River and its tributaries. Weathering, bioturbation as well as human alterations of the terrain act as the  
84 most frequent causative factors.

85 Other parts of Czechia are not as susceptible to landsliding to the extent comparable to the above-mentioned  
86 three primary landslide areas (OWC, CS, DV). Landslide activity has been long recorded in the Neogene and  
87 Quaternary sediments of the Most basins (MB, see Figure 1), along the banks of the Ohře/Eger River and in the  
88 anthropogenic landscape at the edges of the open-pit brown coal mines (e.g., Burda et al., 2013). Another area  
89 prone to landslides is in central-eastern Czechia in the Mesozoic sandstones of the Bohemian Paradise (BP)  
90 which form steep elevations and rockfall-prone areas (e.g., Forczek, 2008). Rockfalls occur along a number of  
91 deeply incised valleys in the Bohemian Massif (e.g., along the Vltava/Moldau river valley south of Prague, VR)  
92 as well as some transportation corridors, particularly along rail tracks (as documented in a database of road and  
93 railway blockages due to natural processes, [www.rupok.cz](http://www.rupok.cz) (Bíl et al., 2017)).



94

95 *Figure 1: Delimitation of the primary areas where landsliding concentrates in Czechia. Neogene volcanic rocks*  
96 *(CS – České středohoří Mts.), Mesozoic sandstones (DV – Děčínská vrchovina Mts.), Neogene and Quaternary*  
97 *sediments (MB – Most Basin) on the west, Mesozoic sandstones in central parts of Czechia (BP – Bohemian*  
98 *Paradise sandstones) and Mesozoic and Tertiary flysch belt (OWC – Outer Western Carpathians) in the east of*  
99 *Czechia represent the most susceptible parts to landsliding. VR – concentration of rockfalls along the Vltava river*  
100 *[https://mapy.geology.cz/arcgis/rest/services/Inspire/GM2\\_5mil/MapServer](https://mapy.geology.cz/arcgis/rest/services/Inspire/GM2_5mil/MapServer). © Czech Geological Survey.*

101

## 102 2.2 Historical Landslide Research in Czechia

103

### 104 2.2.1 The Beginning of Landslide Research in Czechia

105 The first protoscientific works that also included reports on landsliding in Czechia are dated to the eighteenth  
106 century (e. g. Strnad, 1790), followed by geological and engineering geological scientific works emerging as of  
107 the end of the nineteenth century (Zahálka, 1890; Dědina, 1896; Woldřich, 1899) and at the beginning of the  
108 twentieth century (Čermák, 1912, Dědina, 1916). More systematic landslide research which described landsliding  
109 in the area of modern Czechia started, however, in the 1920s with the work of Záruba (1922, 1923, 1926, 1938).  
110 Particular attention was paid to a general description of historical landslide areas in the Pavlovské vrchy Hills  
111 situated in OWC (Jüttner 1931, 1937) as well as historical landslides themselves in the same area (Stejskal, 1931;  
112 Woldřich and Stejskal, 1934). Záruba and Myslivec (1942) documented landslides related to transportation  
113 infrastructure in OWC. Landsliding in the broader area of the city of Zlín (OWC) was described by Krejčí (1943).  
114 The first Czech modern landslide classification was published by Záruba and Mencl (1954).



115

## 116 2.2.2 Systematic Works Describing Landslide Occurrence Based on Historical Data

117 We present here an overview of works from Czechia which created at least a regional landslide chronology based  
118 on documentary data analyses. The only systematic studies of landslide occurrence, based on a range of historical  
119 sources in Czechia, were conducted by Špůrek (1967, 1972, 1985). These studies were mainly based on the  
120 investigation of articles published in national newspapers covering the territory of former Czechoslovakia (and  
121 also mentions landsliding all over the world). The recorded information includes each landslide date, location, type  
122 and amount of damage as well as the bibliographic source.

123 There have not been any other attempts to prepare an overview of dated landslides for all of Czechia which  
124 would follow up on Špůrek's work from 1970 onwards. Concerning debris avalanches, which are only located in  
125 the highest parts of Czechia, Pilous (1973) presented their overview from the Krkonoše Giant Mts. The efforts  
126 have been re-established since the 2000s and have focused on the major Czech areas prone to landsliding.

127 For the Outer Western Carpathians, Bíl et al. (2014) studied historical landsliding in an area around the village of  
128 Halenkovice (central part of OWC). They analysed documentary data, chronicles and interviewed eyewitnesses.  
129 They determined dates for 120 individual landslides. The oldest records were found in local chronicles and  
130 described landsliding in two villages (Jankovice and Košíky) in the close vicinity of Halenkovice in 1915. Bíl et  
131 al. (2020) created an overview of the chronology of landsliding in the Pavlovské vrchy Hills, an area at the  
132 Czech-Austria border belonging to the Western Carpathian Flysch Belt. They determined dates for 30 historical  
133 landslides. The first written resource relates to a landslide record dating back to 1663.

134 In NW Czechia, Raška et al. (2015) established a regional historical landslide database which was compared  
135 with the central part of OWC in respect to the data availability and content. The multihazard database for the  
136 latter half of the nineteenth century (Raška and Dubišar, 2017) allowed for an exploration of the relative direct  
137 impacts of landslides on society. Finally, Raška (2019) used the landslide database to suggest five phases in the  
138 evolution of community-based landslide risk reduction and the various approaches and mechanisms employed in  
139 landslide mitigation measures.

140 Apart from these works, several studies have been published which used historical data to understand current  
141 local landslide hazards (Krejčí et al., 2017) and which explored the availability of documentary sources (Kozák  
142 and Rybář, 2003).

## 143 3. Design of the CHILDA Database


### 144 3.1 Data availability

145 Czechia disposes of an extremely diverse and extensive range of documentary data that may be explored to build  
146 historical landslide inventories. The number of these sources have been, however, subjected to academic scrutiny  
147 mostly in historical climatology and hydrology (Kjeldsen et al., 2014; Brázdil et al., 2018a). Similarly, Bíl et al.  
148 (2020) combined different documentary and archaeological data to compile a chronology of landsliding in the  
149 Pavlovské vrchy Hills (Czechia, OWC) and described the basic historical landslides terminology. The conceptual  
150 differences in hydrometeorological and geomorphologic hazards do not allow for uncritical transposition of the



151 climatological insights into the historical landslide research and therefore call for new insights into the potential  
 152 of the documentary data (Crozier and Glade, 1999; Raška et al., 2014b).

153 The present landslide database is based on a systematic data search in documentary data (both written and  
 154 iconographic) of several types with varying content, coverage and availability (Figure 2). Most of these data are  
 155 available in local archives in a printed or hand-written form (e.g., municipal and school chronicles, official  
 156 correspondence, photos), while some were found in private collections. The primary documentary data were also  
 157 complemented by a search in secondary (published) literature (e. g. newspapers, historical synthesis, historical  
 158 research into forests) and in some areas also with oral inquiry. Within the data search, the lower time boundary for  
 159 the database has not been set. The upper (recent) boundary, for the aim of the analysis in this work, was set at  
 160 1989, representing a significant socio-economic and political turn in Czech history and thus marking the change  
 161 in public data availability as well as in approaches to scientific inquiry regarding landslides. We have allowed,  
 162 however, the database to remain open in order to allow users to add new landslides as well.



164 *Figure 2 An overview of available documentary data applicable for historical landslide research in Czechia with*  
 165 *respect to their time coverage and informational content*

166

167



168 **3.2 Database Structure**

169 Attributes related to each database record are presented in Table 1. Some of the attributes are added by users via a  
 170 form. The items with an asterisk are mandatory and the items which are not part of the input form are underlined.  
 171 They are processed automatically, within the system.

172

173 *Table 1: Structure of records in Childa.*

Name *	Meaning	Values
<u>ID</u>	Unique identifier of a landslide	Number
Type *	Determination of kind of landsliding	List: Landslide; rockfall; earthflow; debris flow; human-induced landslide
Position *	Latitude and longitude of the record inserted via a click on the map	WGS 84 coordinates
Locality *	Description of the locality	String
Accuracy *	Describes the spatial precision of landslide localization by DB user (not a precision in an original source)	List: Metres; tens of metres; hundreds of metres; kilometres
<u>MASL</u>	Height above mean sea level - Landslide highest point elevation	Metres
Count	The number of landslides related to the particular location, given its accuracy; default value 1	String
Start *	The earliest possible date for the beginning of landsliding indicated by a record	Date
End *	The latest possible date for the beginning of landsliding as indicated by a record	Date
<u>Period</u>	Shows an interval during which the landslide originated. It is computed as End – Start.	String
Causes	Description of landslide cause, more causes can be selected	List: Earthquake; lithology; flooding; precipitation; mining; snow thaw; storm; artificial cause
Magnitude	Degree of landslide damage in three categories	List: 1 - Negligible or no losses; approx. 1ha or road infrastructure; 2 - hundreds of metres, >1ha; 3 - large volumes and areas, deep seated landslides
Impact	List of losses caused by landsliding, more impacts can be selected	List: Human fatality; human injury; buildings; transport infrastructure; other infrastructure (mine, water tower, utilities, etc); landscape including old mines, etc.
Remedies	Kind of remediation if applied	String





Source *	Full citation of the source of the landslide record	String
Details	Additional information and original data availability and accessibility (e.g., museum, archive, private collection, etc.)	String
Notes	Other relevant information about landsliding	String
Photo	More than one graphics file can be attributed to a record, e.g., photo, map, a copy of a written source, etc.	Graphics file, pdf

174 \* *Mandatory attributes.*

175 *Note: The attributes which are underlined are automatically added by the system and are not part of the user*

176 *form for data entering.*

177

### 178 3.3 Web-map Application

179 The database can be accessed through a web-map application at <https://childa.cz> (Figure 3). CHILDA is

180 administrated and hosted on the CDV – Transport Research Centre servers. The software requirement is as

181 follows: PostgreSQL/PostGIS, php, php NetteFramework, HTML, CSS, JQuery.



182

183 *Figure 3: Title page of CHILDA ([www.childa.cz](http://www.childa.cz)).*

184 CHILDA users can select a background map (see Figure 4, A). There are the following possibilities: base map,

185 orthophoto, Open Street Map, Mapy.cz map and a hiking map. A municipality or other geographical feature with

186 a conventional name (mountains, etc.) can be selected via a form (B). The full-screen mode is launched when a

187 user clicks on the icon (C). Standard map control features such as full extent (D), backward and forward screen


188 (E), Google Street View (G) and zoom in/out (H) are also available here. A detailed landslide description is

189 shown in the pop-up window after clicking on the landslide map symbol (F). A new landslide record can be





190 added through a form (I). This panel contains two bookmarks. In the *Layer*, the symbology and the time filter  
191 can be seen. In the *New*, a new landslide record can be added, or an existing one can be edited. Only registered  
192 users can edit their own landslides. The edit button is visible in the pop-up window in the map or in the Database  
193 after log in. There is currently no automatic validation of data entered by users to have the app as accessible and  
194 user friendly as possible. Three roles of users are defined: non-registered users, registered users (who are  
195 allowed to edit, delete their own data) and administrators (editing, deleting all data).



196  
197 *Figure 4: Additional information on a landslide record can be obtained after clicking on a point in the map*  
198 *view. A new landslide can be added through the form (right side). A – background map, B – municipality, C –*  
199 *full-screen mode, D – full extent control, E – backward/forward screen, F - landslide description, G – Google*  
200 *Street View, H – zoom in/out, I – new landslide button. © Mapy.cz*

201

202 The landslide database is also accessible via a table (Figure. 5). Four filters are available: movement type,  
203 magnitude, only selection of records added by a particular user and time period. Full text searching is also  
204 possible. The given filter is kept when the map is selected unless it is reset by the user.



**CZECH HISTORICAL LANDSLIDE DATABASE**

ABOUT   GUIDE   MAP   DATABASE   SOURCES   TEAM   LOGIN   CZ

**DATABASE**

Select the movement's type    Select magnitude    From  1912-04-01   Fulltext   

ID	Type	Locality	Accuracy	MASL	Count	Date of origin	Causes	Magnitude	Impact	Remedies	Source	D	M	P
2043	landslide	Blilna	kilometers		vice	3/1531 - 5/1531	precipitation				Hutter, T. (1891): Die Stadt...			
2042	landslide	Usti nad Orlici	kilometers		1	4.5.1911	storm	3	infrastructure (roads), injury, death		Lidové noviny, 6.5.1911, roc....			
2011	landslide	Marsov	kilometers		1	1.1.1911					Špůrek (1972)			
2010	landslide	Potstejn	hundreds of meters		1	1.1.1909			infrastructure (roads)		Špůrek (1972)			

205

206 *Figure 5: A database table visualizes information on landslides and allows for filtering data according to a*  
 207 *number of parameters*

208 **4. Results**

209 **4.1 Landslide Records in Childa**

210 We present below an overview of data contained in CHILDA for the 1132–1989 period. The database contains  
 211 699 records, 619 of them categorised as landslides, 75 as rockfalls and five as 'other' kind of mass movements  
 212 (earth flow, rockslide or human-induced landslide). As regards temporal accuracy, 231 records were determined  
 213 exactly at single day precision, 17 records are known with a week and 88 with a month precision. In total, 363  
 214 records were only attributed to a given year. Concerning the location accuracy, 111 records were localized  
 215 precisely, 71 records with a precision of "tens of metres", 260 records to "hundreds of metres" and 478 to  
 216 kilometres (mostly between 1–2 km, exceptionally up to 5 km in the mountain terrains of Czechia).

217 Table 2 presents database completeness that was determined based on several non-mandatory fields (cause of  
 218 landsliding, magnitude, impact, etc.). The relative number [%] of particular fields always refers to all records in  
 219 CHILDA (i.e., a proportion of 100%). The average number of database completeness amounted to 33 %. An  
 220 increase, however, in database completeness in the next years, based on the incorporation of new results arising  
 221 from future research, is assumed.

222 Table 2: Database completeness on the basis of particular non-mandatory fields.

Filed name	Number of records	Relative number [%]
Cause of landsliding	120	17.2
Magnitude	181	25.9
Impact	354	50.6
Remedies	38	5.4
Source details	645	92.3
Photo	53	7.6


223

224 Our data can also be compared to the previous landslide chronology compiled by Špůrek (1972) for the area of  
 225 Czechia. Špůrek's last data on landsliding come from 1970. Our comparison consequently also ends this year. As  
 226 for 1970, CHILDA currently (April 2021) contains 667 records. In total, 359 of these records were also part of  
 227 Špůrek's overview. This indicates that our new investigation constitutes almost 50% of new records in the



228 database, i.e., records which were not previously covered by Špůrek's catalogue. A major source of Špůrek's  
229 information about landsliding was newspapers. We focused in our research, however, apart from newspapers on  
230 the primary documentary data available in archives (e.g., chronicles, memory books, official correspondence) as  
231 well. This approach implies huge potential for revealing new and unique landslide events in the future.

232 As Figure 6 indicates, the highest density of historical landslide records in Czechia in the studied period  
233 concentrates on three primary landslide areas (OWC, CS, DV). Tens of landslide records come, however, from  
234 the area of the capital Prague (VR) represented mainly by rockfalls. A higher occurrence of records is also  
235 typical for Bohemian Paradise sandstones (BP).




236

237 *Figure 6: Landslide density across Czechia.*

237

238 While landslide records come from all three core landslide areas, the rockfall records were typical for the west  
239 part of Czechia (DV area) and along the Vltava/Moldau river valley (VR) south of Prague (Figure 7).



240

241

242 *Figure 7: Landslide and rockfall distribution.*

242




243 The temporal variability of the landslide records, represented by their numbers within centuries in the whole  
 244 studied period, is shown in Table 3. The only one (and at the same time the oldest) record is known from the  
 245 twelfth century (19 January 1132, see chapter 4.2). No reports were detected between the thirteenth and fifteenth  
 246 centuries and only five and two reports were recorded in the sixteenth and seventeenth centuries, respectively.  
 247 While only 8 records were found between the twelfth and seventeenth centuries, and 30 records in the eighteenth  
 248 century, the majority of the records are evidenced in the last two centuries. Landslide records covering the  
 249 nineteenth century account for 27.8 % whereas records from the twentieth century embrace 66.8% of all reports.

250 *Table 3: The number of landslide records within centuries in Czechia between the twelfth and twentieth*  
 251 *centuries*

Century	12 <sup>th</sup>	13 <sup>th</sup>	14 <sup>th</sup>	15 <sup>th</sup>	16 <sup>th</sup>	17 <sup>th</sup>	18 <sup>th</sup>	19 <sup>th</sup>	20 <sup>th</sup> *
<i>n</i>	1	0	0	0	5	2	30	194	467

252 \* up to 1989 including

253 Due to a rapid increase in landslide records after 1770, a decadal frequency of landslide records was created  
 254 starting in 1761. The records in the period 1761–1989 represent 98% of all records in CHILDA (Figure 8). The  
 255 highest numbers of records were registered during the 1891–1900 decade (19 %) followed by 1931–1940 (14 %)  
 256 and 1961–1970 (14 %) decades. No records were detected, however, between 1791–1810. A significant rise in  
 257 the number of records is apparent since the 1891–1900 decade and the course over the rest of the twentieth  
 258 century is more or less variable. The variability has been in all probability influenced by important rainfall  
 259 and/or snow thaw events.



260

261

*Figure 8: Decadal frequency of landslide records in Czechia between 1761–1989*



262 The first most important landslide year occurred in 1770. A minimal number of landslide records to determine a  
263 particular year as the most important landslide year was set at 10 reports. From 1770, 15 such years were  
264 revealed encompassing 55% of all records on mass movements (385 out of 699) (Table 4). The highest number  
265 of landslide records was found in 1900 (61), 1941 (51), and 1939 (46), representing 15.8%, 13.2, and 11.9% of  
266 the most important 15 landslide years, respectively.

267 *Table 4: The most important 15 landslide years (1770–1989) when at least 10 records were found*

Year	1770	1898	1899	1900	1915	1919	1926	1937
<i>n</i>	17	39	13	61	10	10	32	10
Year	1938	1939	1940	1941	1965	1967	1970	
<i>n</i>	11	46	19	51	38	14	14	

268

#### 269 4.2 The Oldest Records on Landsliding

270 The issue of precise determination of dates of landslide activity is rising when we look back in history. The  
271 oldest records describing landslides in Czechia suffer from spatiotemporal inhomogeneity. CHILDA currently  
272 contains 14 reports of mass movements which took place before 1770, the first most important landslide year  
273 (see Tab. 4). It is important to mention that in 50 % of all recorded cases, citations referring to historical  
274 landslides in this study came from the Špůrek landslide catalogue (1972). Because of our efforts at maximum  
275 authenticity, we took over these citations although we were not able to study some referred citations personally  
276 in certain cases.

277 The oldest known written report describes a rockfall in Praha – Chuchle (VR area) on 19 January 1132 (Strnad,  
278 1790). More detailed information about three landslide events in spring 1531 is described by chroniclers from  
279 Litoměřice and Bílina (CS area). Landsliding was reported from the beginning of April until the middle of May  
280 and affected Radobýl Hill near Litoměřice and Holý vrch Hill near Zahořany village. Vineyards planted on the  
281 hills and slopes slid and two great parts of Radobýl Hill slipped down, including trees and plants. Similarly, a  
282 large portion of Holý vrch Hill slid at that time and a series of other landslides were observed (Smetana, 1978).  
283 In the wet spring of 1531, several landslides also occurred in the surroundings of the nearby Bílina River after a  
284 flood (Hutter, 1891).

285 A day before Christmas Eve of 1595 a landslide occurred near Vraclav – Domoradice village (Špůrek, 1967).  
286 According to Kárník et al. (1957), seismic activity preceded this event. In March 1599, extraordinary damage  
287 was described in Litoměřice (CS area) as a consequence of a great deal of snow and wet weather (Brázdil et al.,  
288 2013b). The chronicler described the situation as follows: “A piece of town wall near St. Laurentius [church] fell  
289 and collapsed [...] On 16 March [...] at the cemetery [...] a huge section slipped down too so the graves opened  
290 and the dead bodies were thrown out [...] That same year, in the month of March, extensive damage to vineyards  
291 was experienced by many [people], the walls caved in and one vineyard after another slid and all of this was  
292 happening due to great wetness”. In addition, a number of springs appeared on the surface, the cellars were full  
293 of water which had to be pumped out and one house even slipped away (Smetana, 1978).

294 The oldest landslide report from OWC, based on a written record, occurred in the Pavlovské vrchy Hills in  
295 Pavlov village in 1663 (Maca, 1994; Kryčér). In contrast to previous mass movement events, continuing



296 landsliding was also recorded on the same street in 1667 (Štefková-Vajayová, 2001), 1715, 1730, and 1763  
 297 (Maca, 1994; Kryčer). A detailed description of landslide damage before 1763 is missing, but the particular  
 298 landslide consequences in the area of the Pavlovské vrchy Hills in the following years were described by Bíl et  
 299 al. (2020).

300 A brief report of landslide occurrence comes from Dečín – Chrochvice (DV area) in 1736 (Mauder, 1931). This  
 301 event was the first one from a series of consequent landslides in 1823, 1850 and 1914 which affected an area of  
 302 16 ha and damaged buildings and local infrastructure (Špůrek, 1972). A similar lack of information relates to a  
 303 landslide in Ústí nad Labem (DV area) in 1767 (Špůrek, 1972). In contrast the sliding down of a parish cellar in  
 304 Libeznice village in 1769 is relatively well described by Třebízský (1885) who mentioned great wetness and  
 305 surfeit of water this year.


306

## 307 5. Discussion

### 308 5.1 CHILDA and Other Historical Landslide Databases

309 We presented an overview of the CHILDA database where as many as possible records on historical landsliding,  
 310 which took part in the area of modern Czechia, were collected. This database currently contains 699 records  
 311 (between 1132 and 1989) and can be compared to other similar databases which have been completed in other  
 312 countries (Figure 9).

313 It should be noted, however, that similarly to other databases CHILDA displays high asymmetry in the number  
 314 of recorded events over time. First, the sole oldest record dated to 1132 extends the span by four centuries as  
 315 further records are only dated to 1531. Second, the majority of records (93%) relates of landsliding that occurred  
 316 since 1850. In contrast, and unlike the other databases, CHILDA also records only a few increased landslide  
 317 frequencies in the pre-industrial periods, namely the 1770 landslide year (17 landslides) resulting from the  
 318 Central-European adverse climate (Raška et al., 2016), and 1817 (with five landslides) possibly influenced by  
 319 the Tambora eruption in 1815 (Brázdil et al., 2016b).



320  
 321 *Figure 9: A comparison of CHILDA and other existing historical landslide databases in terms of the number of*  
 322 *records and recurrence.*



323 *Note: dark-blue stripes mark the periods under study. In case of compound (multi-hazard) databases, light-blue*  
324 *marks complete the studied period for which geohazards other than landslides have been found, whereas dark-*  
325 *blue stands for landsliding. Dashed stripes represent an unspecified beginning (e.g., since the fifteenth century*  
326 *without explicit dating of the oldest record).*

327

## 328 **5. 2 Limitations of the CHILDA Database**

329 As with similar databases which focus on historical records and therefore depend on availability, accessibility  
330 and reliability of original sources, CHILDA also has certain limitations that may be grouped in the following  
331 kinds of uncertainties:

332 (a) Uncertainty is usually related to a description of events as landslides (mass movements in general) in the  
333 documentary records which were named differently. The terminological inconsistency, closely connected with  
334 exploiting the wide range of documentary data, lasted at least until the twentieth century and persists in media  
335 reports up until recent days. This inconsistency was caused by gaps in scientific knowledge in the first Czech  
336 modern landslide classification published in 1954 (Záruba and Mencl, 1954; compared with much earlier  
337 attempts in English listed in Cruden, 2003), and by cultural and resulting language diversity in the Czech Lands  
338 up until 1945 (i.e. Czech and German culture realms). Raška et al. (2015) analysed historical landslides in two  
339 Czech regions and found seven different terms referring to landslides and three for rockfalls. In total, five  
340 German different terms were used to describe the landslides in the Pavlovské vrchy Hills (OWC) between the  
341 middle of the seventeenth–twentieth centuries (Bíl et al., 2020).

342 (b) Moreover, documentary evidence was not continuous, sometimes only depending on concrete persons.  
343 Chroniclers were often not interested in this phenomenon as there were more dangerous ones in this area, such as  
344 floods (Brázdil et al., 2011; Elleder et al., 2020), strong winds (Brázdil et al., 2004, 2017, 2018b), episodes of  
345 drought (Brázdil et al., 2013a; Dolák et al., 2015; Řezníčková et al., 2015; Brázdil et al. 2016a, 2019), etc. There  
346 is obviously also a significant growth of available sources especially since the nineteenth century, which may  
347 distort the represented landslide frequencies over time. The availability of sources also displays a distinct  
348 geographical variation due to different historical developments in individual regions. Several smaller  
349 uncertainties related to historical landslides have thus appeared, e.g., a lack of narrative sources before 1920 and  
350 in the Czech border areas after 1945 (Bíl et al., 2020) or difficulties with recognition of landslides and scoured  
351 slopes during the floods in early documentary records. As regards the Czech border areas, chronicles were lost or  
352 carried away mainly by German inhabitants displaced after 1945. This occurred, along with other regions, in the  
353 landslide prone areas of the Pavlovské vrchy Hills or parts of NW Czechia (CS and DV). Only some of the lost  
354 information from the chronicles became available again via a historical synthesis published in the second half of  
355 the twentieth century (Bíl et al., 2020). In addition, keeping memory books was recommended by the state in  
356 1836 and chronicles were made mandatory from 1920 onward, therefore any older events are either missing in  
357 these sources or were recorded retrospectively with some risk of misinterpretation.

358 (c) Additional problems include uncertainties resulting from difficult to identify duplicities in records of different  
359 coverage (i.e., the issue of upscaling and downscaling) and from false frequency peaks caused by a combination





360 of continual (regularly published) and stationary (published once as a collection of records) data (Raška et al.,  
361 2014b). Both these issues may result in false peaks and gaps in the landslide time series. Verification of data  
362 through checking of the data reliability based on comparison among more sources, or field research is not always  
363 possible and reliable, and therefore any database must always be considered a catalogue of records rather than  
364 events themselves. In the case of the catalogue of events utilisation, these uncertainties relate, e.g., to the example  
365 of 12 records in CHILDA that describe more than one event, but the real number is not known. Such events could  
366 therefore present events with higher intensities than other events where only single landslides were reported.

367 In addition, any quantification of landslide extents and their impact is also complicated. We therefore decided to  
368 only select from a few attributes generally describing impacts. Reactivations of landslide at the same place could  
369 also not be determined exactly. Sometimes, as documented in Bíl et al. (2014) from the village of Halenkovice,  
370 villagers from landslide prone areas were used to seeing the landslides often and as a result did not pay attention  
371 to them. In contrast, rare landsliding in other areas attracted the attention of the locals. Construction work, related  
372 to the first railways across OWC, also both found and caused some landslides (e.g., Záruba, 1938). The apparent  
373 lack of documented landslides before 1920 in the Carpathians (except for the Pavlovské vrchy Hills) was attributed  
374 by Bíl et al. (2014) to dispersed settlements built primarily from wood, the majority of the unpaved roads and  
375 relatively sparse inhabitation in the area. Limited spatial accuracies of historical records often influence any  
376 reliable evaluation regarding the possible structural or anthropogenic triggers.

377 Despite all these uncertainties, documentary evidence stands as a valuable and indispensable source of data  
378 describing the occurrence and as well as the consequences of landslides in Czechia during the last five centuries.

### 379 **5. 3 Further Applications and Development of the Database**

380 Data, currently contained in CHILDA, can be further analyzed in order to delineate and explain their temporal  
381 and spatial concentration. While the spatial extent of landsliding can easily be understood as an apparent relation  
382 to bedrock, temporal distribution has been influenced by important rainfall and/or snow thaw events. Information  
383 on landslides with known dates of activities, which is the case of CHILDA records, can therefore be used in such  
384 a determination of regional rainfall or total water content triggers. Bíl et al. (2016) have already utilized, for  
385 example, the information on landslide periods, defined for an area in the central part of OWC, to determine  
386 rainfall thresholds for this area. Further applications of the data will include analyses of the long-term changes in  
387 landslide risk reduction approaches, their effectiveness and efficiency (Caloiero et al., 2014; Klose et al., 2016)  
388 framed by disaster risk reduction strategies (DRR) (Bíl et al., 2014; UNDRR, 2015). Analysis of community  
389 responses to landslide risk in individual landslide-prone areas has already been published by Raška (2019) and  
390 Klimeš et al. (2020) and allowed for an exploration of both formal (planning, DRR administration) and informal  
391 (community help, mobilization of local knowledge) mechanisms in landslide risk reduction.

### 392 **6. Conclusions**

393 We presented the online landslide database CHILDA (Czech historical landslide database) which summarizes  
394 information about landslides which took place in the area of Czechia (the Czech Republic). The database is  
395 freely accessible via the <https://childa.cz/> website, and currently includes 699 records (spanning the 1132–1989



396 period). The oldest record relates to a rockfall which took place in 1132. In total, the database doubled the  
397 number of records known from the previous historical database in Czechia. We further described in detail  
398 another eight of the oldest records (1531 to 1730) and analysed centennial and decadal frequencies of landslide  
399 records. It was demonstrated that 55 % of all recorded landslide events occurred only within 15 years of the  
400 extreme landslide incidence. Finally, the limitations of the documentary data sources have been summarized  
401 pointing at uncertainties within the database. The future research direction should focus on analysing historical  
402 landslide triggers and their thresholds, changes in spatiotemporal patterns of landslide impacts on society and on  
403 narratives of societal adaptive management to landslide risk.

#### 404 *Acknowledgements*

405 *Thanks go to Vojtěch Čícha for help with the preparation of figures, Vilém Zábanský for suggesting new data*  
406 *sources that included records on landsliding and David Livingstone for assistance with language editing.*

407 *Funding: M. Bíl and J. Kubeček worked with the financial support of the Ministry of Transport of the Czech*  
408 *Republic within the program of long-term conceptual development of research institutions on the research*  
409 *infrastructure acquired from the Operation Program Research and Development for Innovations*  
410 *(CZ.1.05/2.1.00/03.0064). P. Raška acknowledges the financial support of the project Smart City—Smart*  
411 *Region—Smart Community (CZ.02.1.01/0.0/0.0/17\_048/0007435) within the Operational Program*  
412 *Research, Development and Education of the Czech Republic. L. Dolák was supported by SustES -*  
413 *Adaptation strategies for sustainable ecosystem services and food security under adverse environmental*  
414 *conditions (CZ.02.1.01/0.0/0.0/16\_019/0000797).*

#### 415 **References**

- 416 Bíl, M., and Müller, I.: The origin of shallow landslides in Moravia (Czech Republic) in the spring of 2006,  
417 *Geomorphology*, 99, 246–253, <https://doi.org/10.1016/j.geomorph.2007.11.004>, 2008.
- 418 Bíl, M., Krejčí, O., Bílová, M., Kubeček, J., Sedoník, J., and Krejčí, V.: A Chronology of landsliding and its  
419 impacts on the village of Halenkovice, Outer Western Carpathians, *Geografie*, 119, 342–363,  
420 <https://doi.org/10.37040/geografie2014119040342>, 2014.
- 421 Bíl, M., Andrášik, R., Zahradníček, P., Kubeček, J., Sedoník, J., and Štěpánek, P.: Total water content thresholds  
422 for shallow landslides, Outer Western Carpathians, *Landslides*, 13, 337–347, [https://doi.org/10.1007/s10346-](https://doi.org/10.1007/s10346-015-0570-9)  
423 [015-0570-9](https://doi.org/10.1007/s10346-015-0570-9), 2016.
- 424 Bíl, M., Andrášik, R., Kubeček, J., Krivánková, Z., and Vodák, R.: RUPOK: An Online Landslide Risk Tool for  
425 Road Networks, *Advancing Culture of Living with Landslides, Vol 5: Landslides in Different Environments*,  
426 edited by: Mikoš, M., Vilímek, V., Yin, Y., and Sassa, K., Springer International Publishing Ag, Cham, 19–26  
427 pp., 2017.
- 428 Bíl, M., Krejčí, O., Dolák, L., Krejčí, V., Martínek, J., and Svoboda, J.: A chronology of landsliding based on  
429 archaeological and documentary data: Pavlovské vrchy Hills, Western Carpathian Flysch Belt, *Scientific*  
430 *Reports*, 10, 976, <https://doi.org/10.1038/s41598-020-57551-4>, 2020.
- 431 Brázdil, R., Dobrovolný, P., Štekl, J., Kotyza, O., Valášek, H., and Jež, J.: History of weather and climate in the  
432 Czech Lands. VI, Strong winds, 1st published ed., *History of Weather and Climate in the Czech Lands*, Masaryk  
433 University, Brno, 377 pp., 2004.
- 434 Brázdil, R., Řezníčková, L., Valášek, H., Havlíček, M., Dobrovolný, P., Soukalová, E., Řehánek, T., and  
435 Skokanová, H.: Fluctuations of floods of the River Morava (Czech Republic) in the 1691–2009 period:  
436 interactions of natural and anthropogenic factors, *Hydrological Sciences Journal-Journal Des Sciences*  
437 *Hydrologiques*, 56, 468–485, <https://doi.org/10.1080/02626667.2011.564175>, 2011.
- 438 Brázdil, R., Dobrovolný, P., Trnka, M., Kotyza, O., Řezníčková, L., Valášek, H., Zahradníček, P., and Štěpánek,  
439 P.: Droughts in the Czech Lands, 1090–2012 AD, *Climate of the Past*, 9, 1985–2002, [https://doi.org/10.5194/cp-](https://doi.org/10.5194/cp-9-1985-2013)  
440 [9-1985-2013](https://doi.org/10.5194/cp-9-1985-2013), 2013a.



- 441 Brázdil, R., Kotyza, O., Dobrovolný, P., Řezníčková, L., and Valášek, H.: Climate of the Sixteenth Century in  
442 the Czech Lands, History of Weather and Climate in the Czech Lands, Masaryk University, Brno, 286 pp.,  
443 2013b.
- 444 Brázdil, R., Raška, P., Trnka, M., Zahradníček, P., Valášek, H., Dobrovolný, P., Řezníčková, L., Tremil, P., and  
445 Stachoň, Z.: The Central European drought of 1947: causes and consequences, with particular reference to the  
446 Czech Lands, *Climate Research*, 70, 161–178, <https://doi.org/10.3354/cr01387>, 2016a.
- 447 Brázdil, R., Řezníčková, L., Valášek, H., Dolák, L., and Kotyza, O.: Climatic effects and impacts of the 1815  
448 eruption of Mount Tambora in the Czech Lands, *Climate of the Past*, 12, 1361–1374, [https://doi.org/10.5194/cp-](https://doi.org/10.5194/cp-12-1361-2016)  
449 [12-1361-2016](https://doi.org/10.5194/cp-12-1361-2016), 2016b.
- 450 Brázdil, R., Szabó, P., Dobrovolný, P., Řezníčková, L., Kotyza, O., Suchánková, S., and Valášek, H.: Windstorm  
451 of the eighteenth century in the Czech Lands: course, extent, impacts, *Theoretical and Applied Climatology*, 129,  
452 623–632, <https://doi.org/10.1007/s00704-016-1806-x>, 2017.
- 453 Brázdil, R., Kiss, A., Luterbacher, J., Nash, D. J., and Řezníčková, L.: Documentary data and the study of past  
454 droughts: a global state of the art, *Clim. Past*, 14, 1915–1960, <https://doi.org/10.5194/cp-14-1915-2018> 2018a.
- 455 Brázdil, R., Stucki, P., Szabó, P., Řezníčková, L., Dolák, L., Dobrovolný, P., Tolasz, R., Kotyza, O., Chromá,  
456 K., and Suchánková, S.: Windstorms and forest disturbances in the Czech Lands: 1801–2015, *Agricultural and*  
457 *Forest Meteorology*, 250, 47–63, <https://doi.org/10.1016/j.agrformet.2017.11.036>, 2018b.
- 458 Brázdil, R., Dobrovolný, P., Trnka, M., Řezníčková, L., Dolák, L., and Kotyza, O.: Extreme droughts and human  
459 responses to them: the Czech Lands in the pre-instrumental period, *Clim. Past*, 15, 1–24,  
460 <https://doi.org/10.5194/cp-15-1-2019>, 2019.
- 461 Burda, J., Hartvich, F., Valenta, J., Smitka, V., and Rybář, J.: Climate-induced landslide reactivation at the edge  
462 of the Most Basin (Czech Republic) – progress towards better landslide prediction, *Nat. Hazards Earth Syst. Sci.*,  
463 13, 361–374, <https://doi.org/10.5194/nhess-13-361-2013>, 2013.
- 464 Cajz, V.: The České středohoří Mts.: volcanostratigraphy and geochemistry, *Geolines*, 9, 21–28, 1999.
- 465 Caloiero, T., Pasqua, A. A., and Petrucci, O.: Damaging Hydrogeological Events: A Procedure for the  
466 Assessment of Severity Levels and an Application to Calabria (Southern Italy), *Water*, 6, 3652–3670,  
467 <https://doi.org/10.3390/w6123652> 2014.
- 468 Caracciolo, D., Arnone, E., Conti, F. L., and Noto, L. V.: Exploiting historical rainfall and landslide data in a  
469 spatial database for the derivation of critical rainfall thresholds, *Environmental Earth Sciences*, 76, 222,  
470 <https://doi.org/10.1007/s12665-017-6545-5> 2017.
- 471 Crozier, M. J., and Glade, T.: Frequency and magnitude of landsliding: Fundamental research issues, *Zeitschrift*  
472 *fur Geomorphologie, Supplementband*, 115, 141–155, 10.1127/zfgsuppl/115/1999/141, 1999.
- 473 Cruden, D. M.: The First Classification of Landslides?, *Environmental and Engineering Geoscience*, 9, 197–200,  
474 <https://doi.org/10.2113/9.3.197>, 2003.
- 475 Čermák, J.: Sesutí stráně a hrazené jezero u Mladotic: předběžná zpráva [Landslide and the landslide-dammed  
476 lake of Mladotice: a preliminary report], *Sborník České společnosti zeměvědné*, 18, 19–23, 1912.
- 477 CHILDA - Czech Historical Landslide Database: <https://childa.cz>, access: 9 April 2021, 2021.
- 478 CHMI - Czech Hydrometeorological Institute: [www.chmi.cz](http://www.chmi.cz), access: 9 April 2021, N/A.
- 479 Czech Geological Survey: [www.geology.cz](http://www.geology.cz), access: 8 February 2021, 2012a.
- 480 Czech Geological Survey Map Server:  
481 [https://mapy.geology.cz/arcgis/rest/services/Inspire/GM2\\_5mil/MapServer?f=json](https://mapy.geology.cz/arcgis/rest/services/Inspire/GM2_5mil/MapServer?f=json) access: 5 March 2021,  
482 2012b.
- 483 Damm, B., and Klose, M.: The landslide database for Germany: Closing the gap at national level,  
484 *Geomorphology*, 249, 82–93, <https://doi.org/10.1016/j.geomorph.2015.03.021>, 2015.
- 485 Dědina, V.: Sesouvání stráně na Chlomku u Ml. Boleslavě [Landsliding at Chloumek by Ml. Boleslav town],  
486 *Sborník České společnosti zeměvědné*, 2, 40–41, 1896.
- 487 Dědina, V.: Sesutí půdy pod Chlomkem u Ml. Boleslavi [Landslide of earth at Chloumek by Ml. Boleslav town],  
488 *Sborník České společnosti zeměvědné*, 22, 160–166, 1916.
- 489 Devoli, G., Morales, A., and Høeg, K.: Historical landslides in Nicaragua - collection and analysis of data,  
490 *Landslides*, 4, 5–18, <https://doi.org/10.1007/s10346-006-0048-x>, 2007.
- 491 Dolák, L., Brázdil, R., Řezníčková, L., and Valášek, H.: Selected drought impacts in South Moravia in the 18th  
492 and 20th centuries based on documentary evidence, In: Otmár Urban, Mirka Šprtová, Karel Klem. *Global*  
493 *Change: A Complex Challenge. Conference Proceedings. Global Change Research Centre, The Czech Academy*  
494 *of Sciences, v.v.i., Brno*, 30–33 pp., 2015.
- 495 Elleder, L., Krejčí, J., Racko, S., Daňhelka, J., Šírová, J., and Kašpárek, L.: Reliability check of flash-flood in  
496 Central Bohemia on May 25, 1872, *Global and Planetary Change*, 187, 103094,  
497 <https://doi.org/10.1016/j.gloplacha.2019.103094>, 2020.
- 498 Elliott, A., and Kirschbaum, M.: The preliminary landslide history database of Utah, 1850–1978, *Utah*  
499 *Geological Survey*, 514, 5, 2007.



- 500 Forczek, I.: Destruction of marginal parts of sandstone plateaus in the protected landscape area Bohemian  
501 Paradise, *Acta Geodyn. Geomater.*, 5, 267–274, 2008.
- 502 Froude, M. J., and Petley, D. N.: Global fatal landslide occurrence from 2004 to 2016, *Natural Hazards and*  
503 *Earth System Sciences*, 18, 2161–2181, <https://doi.org/10.5194/nhess-18-2161-2018>, 2018.
- 504 Gariano, S. L., and Guzzetti, F.: Landslides in a changing climate, *Earth-Sci. Rev.*, 162, 227–252,  
505 <https://doi.org/10.1016/j.earscirev.2016.08.011>, 2016.
- 506 Glade, T., Albini, P., and Frances, F.: An introduction to the use of historical data in natural hazard assessments,  
507 *Use of Historical Data in Natural Hazard Assessments*, edited by: Glade, T., Albini, P., and Frances, F.,  
508 Springer, Dordrecht, 2001.
- 509 Guzzetti, F., Cardinali, M., and Reichenbach, P.: The AVI project: A bibliographical and archive inventory of  
510 landslides and floods in Italy, *Environmental Management*, 18, 623–633, <https://doi.org/10.1007/BF02400865>,  
511 1994.
- 512 Hermanns, R., Hansen, L., Sletten, K., Böhme, M., Bunkholt, H., Dehls, J., Eilertsen, R., Fischer, L., L'Heureux,  
513 J.-S., Høgaas, F., Nordahl, B., Oppikofer, T., Rubensdotter, L., Solberg, I.-L., Stalsberg, K., and Yugsi Molina,  
514 F. X.: Systematic geological mapping for landslide understanding in the Norwegian context, in: *Landslide and*  
515 *engineered slopes: protecting society through improved understanding*, edited by: Eberhardt, E., Froese, C.,  
516 Turner, A. K., and Leroueil, S., Taylor & Francis Group, London, 265–271, 2013.
- 517 Herrera, G., Mateos, R. M., García-Davalillo, J. C., Grandjean, G., Poyiadji, E., Maftai, R., Filipciuc, T.-C.,  
518 Jemec Auflič, M., Jež, J., Podolszki, L., Trigila, A., Iadanza, C., Raetzo, H., Kociu, A., Przyłucka, M., Kułak,  
519 M., Sheehy, M., Pellicer, X. M., McKeown, C., Ryan, G., Kopačková, V., Frei, M., Kuhn, D., Hermanns, R. L.,  
520 Koulermou, N., Smith, C. A., Engdahl, M., Buxó, P., Gonzalez, M., Dashwood, C., Reeves, H., Cigna, F.,  
521 Liščák, P., Paudiš, P., Mikulénas, V., Demir, V., Raha, M., Quental, L., Sandić, C., Fusi, B., and Jensen, O. A.:  
522 *Landslide databases in the Geological Surveys of Europe*, *Landslides*, 15, 359–379,  
523 <https://doi.org/10.1007/s10346-017-0902-z>, 2018.
- 524 Hroch, Z., Kycl, P., and Šebesta, J.: Landslide hazards in North Bohemia, in: *Landslides: Proceedings of the*  
525 *First European Conference on Landslides*, edited by: Rybář, J., Stemberk, J., and Wagner, P., Swets &  
526 Zeitlinger, Lisse, 207–212, 2002.
- 527 Hungr, O., Leroueil, S., and Picarelli, L.: The Varnes classification of landslide types, an update, *Landslides*, 11,  
528 167–194, <https://doi.org/10.1007/s10346-013-0436-y>, 2014.
- 529 Hutter, T.: *Die Stadt Bilin und ihre Geschichte von der ältesten Zeit bis auf unsere Tage (1890)*, Verlag des  
530 Bürgermeisteramtes, Bilin, 168 pp., 1891.
- 531 Jüttner, K.: *Die Rutschgebiete an den Hägen der Pollauer Berge*, *Firgenwald*, 4, 108–121, 1931.
- 532 Jüttner, K.: *Die erdgeschichtliche Entwicklung des Bodens der Gemeinde Pollau, Deutsch-mähr.-schles. Heimat*,  
533 11/12, 23, 1–6, 1937.
- 534 Kalvoda, J., and Balatka, B.: Chronodynamics of the Labe River antecedence in the Děčínská vrchovina  
535 Highland, Czech Republic, *Acta Montana IRSM AS CR Series A*, 8, 43–60, 1995.
- 536 Kárník, V., Michal, E., and Molnár, A.: *Erdbebenkatalog der Tschechoslowakei bis zum Jahre 1956*, *Travaux*  
537 *Géophysiques*, 69, 411–598, 1957.
- 538 Kjeldsen, T. R., Macdonald, N., Lang, M., Mediero, L., Albuquerque, T., Bogdanowicz, E., Brázdil, R.,  
539 Castellarin, A., David, V., Fleig, A., Gul, G. O., Kriauciuniene, J., Kohnová, S., Merz, B., Nicholson, O., Roald,  
540 L. A., Salinas, J. L., Sarauskienė, D., Šraj, M., Strupczewski, W., Szolgay, J., Toumazis, A., Vanneuville, W.,  
541 Veijalainen, N., and Wilson, D.: Documentary evidence of past floods in Europe and their utility in flood  
542 frequency estimation, *Journal of Hydrology*, 517, 963–973, <https://doi.org/10.1016/j.jhydrol.2014.06.038>, 2014.
- 543 Klimeš, J., Stemberk, J., Blahut, J., Krejčí, V., Krejčí, O., Hartvich, F., and Kycl, P.: Challenges for landslide  
544 hazard and risk management in 'low-risk' regions, Czech Republic-landslide occurrences and related costs (IPL  
545 project no. 197), *Landslides*, 14, 771–780, <https://doi.org/10.1007/s10346-017-0798-7>, 2017.
- 546 Klimeš, J., Müllerová, H., Woitsch, J., Bíl, M., and Křížová, B.: Century-long history of rural community  
547 landslide risk reduction, *International Journal of Disaster Risk Reduction*, 51, 101756,  
548 <https://doi.org/10.1016/j.ijdr.2020.101756>, 2020.
- 549 Klose, M., Maurischat, P., and Damm, B.: Landslide impacts in Germany: A historical and socioeconomic  
550 perspective, *Landslides*, 13, 183–199, <https://doi.org/10.1007/s10346-015-0643-9> 2016.
- 551 Kozák, J., and Rybář, J.: Special Contribution: Pictorial Series of the Manifestations of the Dynamics of the  
552 Earth: 3. Historical Images of Landslides and Rock Falls, *Studia Geophysica et Geodaetica*, 47, 221–232,  
553 <https://doi.org/10.1023/A:1022220126463>, 2003.
- 554 Krejčí, J.: *Sesuvná území na Zlínsku [Landslide areas in the Zlín region]*, *Práce Moravské přírodovědecké*  
555 *společnosti*, 15, F156, 1943.
- 556 Krejčí, O., Krejčí, V., Kycl, P., Paleček, M., and Rybář, J.: Bohyně in Děčín district - the largest landslide area in  
557 the Czech Republic, *Geoscience Research Reports*, 50, 227–234, [10.3140/zpravy.geol.2017.38](https://doi.org/10.3140/zpravy.geol.2017.38) 2017.



- 558 Krejčí, O., Baroň, I., Bíl, M., Hubatka, F., Jurová, Z., and Kirchner, K.: Slope movements in the Flysch  
559 Carpathians of Eastern Czech Republic triggered by extreme rainfalls in 1997: a case study, *Physics and*  
560 *Chemistry of the Earth, Parts A/B/C*, 27, 1567–1576, [https://doi.org/10.1016/S1474-7065\(02\)00178-X](https://doi.org/10.1016/S1474-7065(02)00178-X), 2002.
- 561 Kryčér, R.: Dějiny obcí Bavory, Bulhary, Klentnice, Mušov, Pavlov, Perná, Sedlec, Dolní Věstonice, Horní  
562 Věstonice [Histories of the municipalities of Bavory, Bulhary, Klentnice, Mušov, Pavlov, Perná, Sedlec, Dolní  
563 Věstonice and Horní Věstonice], The State District Archive in Břeclav, (undated document).
- 564 Maca, J.: Pollauer Heimatbuch: 1334–1946: Geschichte und Schicksal einer deutschen Gemeinde in Südmähren,  
565 Selbstverlag, Wien, 551 pp., 1994.
- 566 Marc, O., Stumpf, A., Malet, J. P., Gosset, M., Uchida, T., and Chiang, S. H.: Initial insights from a global  
567 database of rainfall-induced landslide inventories: the weak influence of slope and strong influence of total storm  
568 rainfall, *Earth Surf. Dyn.*, 6, 903–922, <https://doi.org/10.5194/esurf-6-903-2018>, 2018.
- 569 Mauder, E.: Chronik von Bodenbach, Verlag der Stadtgemeinde, Tetschen-Bodenbach, 445 pp., 1931.
- 570 Pánek, T., Brázdil, R., Klimeš, J., Smolková, V., Hradecký, J., and Zahradníček, P.: Rainfall-induced landslide  
571 event of May 2010 in the eastern part of the Czech Republic, *Landslides*, 8, 507–516,  
572 <https://doi.org/10.1007/s10346-011-0268-6>, 2011.
- 573 Pereira, S., Zêzere, J. L., Quaresma, I. D., and Bateira, C.: Landslide incidence in the North of Portugal: Analysis  
574 of a historical landslide database based on press releases and technical reports, *Geomorphology*, 214, 514–525,  
575 <https://doi.org/10.1016/j.geomorph.2014.02.032>, 2014.
- 576 Piacentini, D., Troiani, F., Daniele, G., and Pizzolo, M.: Historical geospatial database for landslide analysis:  
577 the Catalogue of Landslide Occurrences in the Emilia-Romagna Region (CLOCKER), *Landslides*, 15, 811–822,  
578 <https://doi.org/10.1007/s10346-018-0962-8>, 2018.
- 579 Pilous, V.: Strukturní mury v Krkonoších – I. část, *Opera Corcontica*, 10, 15–69, 1973.
- 580 Raška, P., Hartvich, F., Cajz, V., and Adamovič, J.: Structural setting of the Certovka landslide (Ustí nad Labem,  
581 Czech Republic): morphostructural analysis and electrical resistivity tomography, *Geol. Q.*, 58, 85–98,  
582 <https://doi.org/10.7306/gq.1134>, 2014a.
- 583 Raška, P., Záborský, V., Dubišar, J., Kadlec, A., Hrbáčová, A., and Strnad, T.: Documentary proxies and  
584 interdisciplinary research on historic geomorphologic hazards: a discussion of the current state from a central  
585 European perspective, *Natural Hazards*, 70, 705–732, <https://doi.org/10.1007/s11069-013-0839-z>, 2014b.
- 586 Raška, P., Klimeš, J., and Dubišar, J.: Using Local Archive Sources to Reconstruct Historical Landslide  
587 Occurrence in Selected Urban REGions of the Czech Republic: Examples from Regions with Different Historical  
588 Development, *Land Degradation & Development*, 26, 142–157, <https://doi.org/10.1002/ldr.2192>, 2015.
- 589 Raška, P., Záborský, V., Brázdil, R., and Lamková, J.: The late Little Ice Age landslide calamity in North  
590 Bohemia: Triggers, impacts and post-landslide development reconstructed from documentary data (case study of  
591 the Koží vrch Hill landslide), *Geomorphology*, 255, 95–107, <https://doi.org/10.1016/j.geomorph.2015.12.009>,  
592 2016.
- 593 Raška, P., and Dubišar, J.: Impacts of natural hazards on an early industrial community: A case study of North  
594 Bohemia and its implications for long-term vulnerability assessment, *Moravian Geographical Reports*, 25, 13–  
595 23, <https://doi.org/10.1515/mgr-2017-0002>, 2017.
- 596 Raška, P.: Contextualizing community-based landslide risk reduction: an evolutionary perspective, *Landslides*,  
597 16, 1747–1762, <https://doi.org/10.1007/s10346-018-1099-5> 2019.
- 598 Remondo, J., Bonachea, J., and Cendrero, A.: Quantitative landslide risk assessment and mapping on the basis of  
599 recent occurrences, *Geomorphology*, 94, 496–507, <https://doi.org/10.1016/j.geomorph.2006.10.041>, 2008.
- 600 Řezníčková, L., Brázdil, R., Kotyza, O., and Valášek, H.: Documentary evidence as a source of data for studying  
601 droughts in the Czech Lands, In: *Global Change: A Complex Challenge. Conference Proceedings*. Brno: Global  
602 Change Research Centre, The Czech Academy of Sciences, v.v.i., Brno, 26–29 pp., 2015.
- 603 Smetana, J.: Nejstarší kronikářské záznamy litoměřických radních písařů [The earliest chronicle records of  
604 councillor scribes in Litoměřice], *Vlastivědný sborník Litoměřicko*, 14, 119–142, 1978.
- 605 Stejskal, J.: Svážná území na Pavlovských vrších [Landslide areas of the Pavlovské vrchy Hills], *Sborník*  
606 *československé akademie zemědělské*, 6, 55–94, 1931
- 607 Strnad, A.: Chronologisches Verzeichnis der Naturbegebenheiten im Königsreich Böhmen vom Jahre Christi  
608 633 bis 1700, Prague, 139 pp., 1790.
- 609 Špůrek, M.: Historická analýza působení klimatického sesuvného faktoru v Českém masivu [Historical analysis  
610 of the climatic effect on landsliding in the Bohemian Massif], *Archive of the Czech Geological Survey –*  
611 *Geofond Prague*, 42 pp., 1967.
- 612 Špůrek, M.: Historical catalogue of slide phenomena, *Stud Geographica, ČSAV*, Brno, 178 pp., 1972.
- 613 Špůrek, M.: Československá bibliografie svahových deformací. Dodatky 1. 1, 105 [The Czechoslovak  
614 bibliography of slope deformations, Amendments 1.1.105], MS Geofond, Prague, non-paginated pp., 1985.
- 615 Štefková-Vajayová, M.: Kronika obce Pavlov [Chronicle of Pavlov municipality], Municipal office, Pavlov, 168  
616 pp., 2001.





- 617 Taylor, F. E., Malamud, B. D., Freeborough, K., and Demeritt, D.: Enriching Great Britain's National Landslide  
618 Database by searching newspaper archives, *Geomorphology*, 249, 52–68,  
619 <https://doi.org/10.1016/j.geomorph.2015.05.019>, 2015.
- 620 Tropeano, D., and Turconi, L.: Using Historical Documents for Landslide, Debris Flow and Stream Flood  
621 Prevention. Applications in Northern Italy, *Natural Hazards*, 31, 663–679,  
622 <https://doi.org/10.1023/B:NHAZ.0000024897.71471.f2> 2004.
- 623 Třebizský, V. B.: Paměti Jiřího Václava Paroubka, někdy vikáře a faráře v Libeznicích (1740–1774) [Memoirs  
624 of Jiří Václav Paroubek, a vicar and a priest in Libeznice (1740–1774)], *Sborník historický*, 2, 73–87, 1885.
- 625 Van Den Eeckhaut, M., Muys, B., Van Loy, K., Poesen, J., and Beekman, H.: Evidence for repeated re-  
626 activation of old landslides under forest, *Earth Surface Processes and Landforms*, 34, 352–365,  
627 <https://doi.org/10.1002/esp.1727>, 2009.
- 628 Van Den Eeckhaut, M., and Hervás, J.: State of the art of national landslide databases in Europe and their  
629 potential for assessing landslide susceptibility, hazard and risk, *Geomorphology*, 139–140, 545–558,  
630 <https://doi.org/10.1016/j.geomorph.2011.12.006>, 2012.
- 631 Woldřich, J., and Stejskal, J.: Geologický profil hlíněm a svážením v cihelně u Dolních Věstonic na Moravě  
632 [Geological profile of clay-pit in brickfield by Dolní Věstonice village in Moravia], *Věda přírodní: měsíčník pro*  
633 *šíření a pěstování věd přírodních: časopis pro přírodovědeckou vlastivědu. Zvláštní otisk z časopisu Věda*  
634 *přírodní*, 15, 129–137, 1934.
- 635 Woldřich, J. N.: Sesutí u Klapého z roku 1898 [Landslide at Klapý in 1898], *Královská česká společnost nauk.*  
636 *Věstník Královské české společnosti nauk. Třída mathematicko-přírodovědecká, non-paginated*, 1899.
- 637 Wu, C. Y., and Yeh, Y. C.: A Landslide Probability Model Based on a Long-Term Landslide Inventory and  
638 Rainfall Factors, *Water*, 12, 17, <https://doi.org/10.3390/w12040937>, 2020.
- 639 Zahálka, Č.: O ssutinách čedičových a znělcových v Českém středohoří [On basaltic and phonolitic scree slopes  
640 of the České středohoří Mts.], *Vesmír*, 20, 66–67, 1890.
- 641 Záruba, Q.: Studie o sesuvných terénech na Vsatsku a Valašsku [A study on landslide terrains of the Vsatsko and  
642 Valašsko regions], *Časopis Moravského musea zemského*, 20–21, 170–180, 1922.
- 643 Záruba, Q.: Studie o sesouvání půdy na Hlučínsku [A study on landsliding in the Hlučínsko area], *Technický*  
644 *obzor*, 37, 8, 1923.
- 645 Záruba, Q.: Sesouvání půdy v oblasti českého útvaru křídového [Landsliding in the Bohemian Cretaceous  
646 Basin], *Věstník státního geologického ústavu*, 2, 226–235, 1926.
- 647 Záruba, Q.: Sesuvy v Lyském průsmyku a jejich význam pro komunikační stavby [Landslides in the Lysský  
648 průsmyk pass and their significance for the communication buildings], *Technický obzor*, 46, 1–29, 1938.
- 649 Záruba, Q., and Myslivec, A.: Sesuvy při komunikačních stavbách ve flyšovém území [Landslides at  
650 transportation constructions in flysch area], *Technický obzor*, 50, 199–230 1942.
- 651 Záruba, Q., and Mencl, V.: *Inženýrská geologie [Engineering Geology]*, Academia, Prague, 428 pp., 1954.
- 652 Zvelebil, J., Vařilová, Z., and Paluš, M.: Tools for Rock Fall Risk Integrated Management in Sandstone  
653 Landscape of the Bohemian Switzerland National Park, Czech Republic (M121), in: *Landslides*, edited by: K.,  
654 S., H. F., F., W., and G., W., Springer, Berlin, Heidelberg, 119–126, 2005.
- 655