



Data-based wildfire risk model for Mediterranean ecosystems. Study case of Concepcion Metropolitan Area in Central Chile

Edilia Jaque Castillo¹, Alfonso Fernández¹, Rodrigo Fuentes Robles², Carolina G. Ojeda³.

5

¹ Department of Geography, Universidad de Concepción, Concepción, Chile.

² Department of Forestry Sciences, Universidad de Concepción, Chile.

10

³ Facultad de Arquitectura, Diseño y Estudios Urbanos, Pontificia Universidad Católica de Chile, Santiago Chile

Correspondence to: Alfonso Fernández alfernandez@udec.cl

15

Abstract. Wildfire risk is latent in Chilean metropolitan areas characterized by the strong presence of Wildland-Urban Interfaces (WUI). The Metropolitan Area of Concepción (CMA) constitutes one of the most representative samples of that dynamic. The wildfire risk in the CMA was addressed by establishing a model of 5 categories (Near Zero, Low, Medium, High, and Very High) that represent discernible thresholds in fire occurrence, using geospatial data and satellite images describing anthropic - biophysical factors that trigger fires. Those were used to deliver a model of fire hazard using machine learning algorithms, including Principal Component Analysis and Kohonen Self-Organizing Maps in two experimental scenarios: only native forest and only forestry plantation. The model was validated using fire spots obtained from the forestry government organization. The results indicated that 12.3% of the CMA's surface area has a high and very high risk of a forest fire, 29.4% has a medium risk, and 58.3% has a low and very low risk. Lastly, the observed main drivers that have deepened this risk were discussed: first, the evident proximity between the increasing urban areas with exotic forestry plantations, and second, climate change that threatens to trigger more severe and large wildfires because of human activities.

20

25

30

Key Words: Wildfires, Wildfire Urban Interface, data-driven Modeling, Chile, Climate Change, Weather-related Hazards.

35



40 1. Introduction

In the last few decades, the world has seen an increasing trend in wildfires affecting large populations (Moritz et al., 2012), generally being attributed to atmospheric warming fueled by anthropogenic climate change (Spies et al., 2014) and extreme weather events (Stott, 2016) creating a riskier environment. However, wildfire hazard
45 is a product of interlinked socio-environmental processes including the proximity between Wildland-Urban Interface (WUI) and urban areas (Kumagai et al., 2004; Kolden and Henson, 2019; Goldman, 2018; Sarricolea et al., 2018); unregulated extractive economic activities in fire-prone landscapes (Castree, 2008; Spies et al., 2014; Freudenburg, 1992; Gago and Mezzadra, 2017); traditional cultural practices which increase the availability of flammable material -construction, forestry or agriculture- (Harari, 2013; Frene and Nuñez, 2010);
50 and the traditional practice of clearing land “slash and burn” (Shahriar et al., 2019:1). This way, the analysis of this hazard must consider biophysical factors such as altitude, slope, climate conditions, solar radiation, and the vegetation cover (Chuvieco et al., 2004; Chuvieco et al., 2011). Likewise, windy, and dry conditions with steep slope, rapidly lead to quick fire spread and burn large areas of forest within a short time (Shahriar et al., 2019:2). Identifying and managing fire hazards is part of a political agenda rather than a solely biophysical phenomenon
55 (Pyne, 2009; Doerr and Santin, 2016; Change, 2017). The experiences with fire in the underdeveloped countries are radically different from developed countries which have controlled burns, a strict forestry policy, solid territorial planning, and usually take advantage of the ecological benefits of the fire for the ecosystems and livelihoods (Hutto, 2008; González, 2005; González-Mathiesen and March, 2018; Adams, 2013).

Risk and vulnerability mapping usually identify the categories of wildfires’ likelihood that corresponds to one
60 of the most used tools in research. The use of risk categories is considered a useful method to provide understandable information for policymaking and decision making as attested by the style of the “Summary for Policymakers”, a document regularly delivered during the publication of the IPCC’s (Intergovernmental Panel on Climate Change) assessment reports and that contains many examples of categorically organized information (IPCC, 2014 and 2019). However, feeding the predictive models with precise data of land cover changes,
65 accurate meteorological data or trace the human activities that could start a wildfire in real-time remains a challenge, mainly because data are sparse or outdated, while sometimes stored in multiple agencies (Dapeng et al., 2019; Otero and Nielsen, 2017; Knowles et al., 2015).

Reducing the exposure of the population to this hazard becomes sorely needed, which requires provision of tools that support landscape planning in a way that minimizes occurrence and intensity (Gonzalez-Mathiesen
70 and March, 2018) and manage the social impacts after the disturbance (Paveglio et al. 2015). Especially in highly fragile regions such as Mediterranean ecosystems (Turco et al. 2016; Pausas et al. 2008; Darques, 2015; Diffenbaugh, 2007) existing in Spain (Vilar del Hoyo et al. 2011), Italy (Terranova et al., 2009), Australia (McGee and Russell, 2003), Portugal (Gómez-González et al., 2018), California (Koltunov et al., 2012), and Chile (de la Barrera et al., 2017).

75 Chile’s Central-South region (~35°S to ~40°S) (Figure 1) is one of the most transformed in the country, with a long history of mining, industrialization, and forest exploitation (Bustamante and Varela, 2007; Aguayo et al.,



2009). Here, intensive land use changes interact with the replacement of native land-cover for plantations, urban sprawling, and socio-environmental conflicts associated with forest property (Andersson et al., 2016, Nahuelhual et al., 2012; Altamirano et al., 2013; Heilmayr et al., 2016; McWethy et al., 2018; Cid, 2015; Schulz
80 et al., 2010) the lead to a characteristic environment prone to wildfire occurrence.

The Concepción Metropolitan Area (CMA) is a conspicuous example of wildfire activity in this region of Chile. Available studies suggest that wildfires will become more frequent and aggressive, given the changing climate conditions in the CMA (Castillo et al., 2003; CONAF, 2017-2018; Sarricolea et al., 2020; CR2, 2020) following global trends (Moritz et al., 2012). One of those changes are related with more frequent droughts (Fernández et al.,
85 et al., 2018), which are coincident with recent findings that attribute part of precipitation decrease to anthropogenic sources (Boisier et al., 2016) impacting the lives, crops, and neighborhoods of more than a million people (Gonzalez et al., 2018; de la Barrera et al., 2018, CONAF, 2018; Araya-Muñoz et al., 2017; Cid, 2015).

In this work, a model for wildfire risk mapping in the CMA (~36.7°S, Fig.1) was applied and validated. An updated categorical map at relatively high spatial resolution was delivered. This model aims to support urban
90 planning and further studies for wildfire hazards. The paper is organized as follows: Section II describes the study area, materials, and methods; section III presents and analyzes the results; section IV corresponds to the discussion, and in section V we conclude while suggesting avenues of future work.

2. Materials And Methods

95

Located in Chile's Biobío administrative region, the CMA (~36.7° and ~73° W, Figure 1) is the third-largest urban area with over 1 million total population (INE, 2021). CMA has many interconnected small urban centers (Rojas Quezada et al., 2009) which are expanding its surface every day, mostly for housing development and industry (Rojas Quezada et al., 2013). Also, has a variety of important biodiversity hotspots (Smith Ramírez,
100 2000) and wetlands (Martínez Poblete, 2014).

A Mediterranean climate with warm/dry summers and cold/wet winters characterize this region (Sarricolea et al., 2020), with an average temperature of 12.4°C while annual rainfall is 1,332 mm, with 70% concentrated between May to August (BCN, 2017). The CMA has one of the largest Wildland-Urban Interfaces (WUI) in the country (Ruiz et al., 2017).

Several economic activities were developed in the CMA and its surroundings since its foundation in the 18th century. However, today the region is mainly known for timber production and export from plantations of exotic fast-growing species (Torres et al., 2015). With more than 46,697 hectares of forestry plantations in 2016, the CMA has steadily undergone an expansion of exotic forestry plantations since 1974, when there were 44,123 hectares (INFOR, 2017).

110




Figure 1. Map of the study area. The right map highlights Chile's Biobío region in a continental context and the CMA is shown in the left map. Source: CEDEUS, 2016.

115

Previous work on wildfire hazard mapping indicates the need to include a number of spatially distributed factors that contribute to the susceptibility of the landscape, as for example slopes, orientation and the effect of insolation (You et al. 2017), and these types of models organize space into categories as a result of weighted sums of contribution factors. In this case, available research portrays wildfires as products of human activities, topographic characteristics, land cover, and climate (CONAF, 2017, de la Barrera et al. 2018; Ubeda and Sarricolea, 2016). While most approaches to map wildfires hazards have been based upon framework tested in other regions, data-driven approaches are still under-utilized. This hazard modeling takes advantage of available national databases and satellite products, ingested into machine learning algorithms to produce maps that allow spatially distributed identification and assessment of wildfire hazards. The model combines Principal Component Analysis (PCA) and Kohonen self-organizing maps (SOM) to determine locations classified in five categories: near zero, low, median, high, and very high. The following subsections present the steps to compile the analyzed database, model development, and the experiments performed.

120

125

2.1. Geodatabase Compilation

130



Input data for the modeling corresponded to a 12-variable geodatabase that included several descriptors related to wildfire spot recurrence, such as topographic features, land cover characteristics, built environment descriptors, and climatic indices (Table 1). Wildfire spot locations were utilized as reference coordinates to produce a raster that counted the number of spots within 900m pixel size. Center coordinates of each pixel are
135 the locations utilized in the compiled geodatabase. The 900m spatial resolution corresponds to a trade-off between the representation of the different input databases, which range from 30m to 5km. Locations of wildfire spots used for the geodatabase correspond to the period 2008-2019 available from the Chilean Forest Service (CONAF, Corporación Nacional Forestal). The forest fire database is constructed from information collected by CONAF brigades and private forestry companies in peri-urban and rural (forestry) areas. Forest fire detection
140 is carried out through three ways: a) Fixed terrestrial (observation towers), b) Mobile terrestrial (surveillance) and c) Aerial detection (Tapia and Castillo, 2014). Detected fires are GPS georeferenced and subsequently added to a GIS. The minimum area for a forest fire to be mapped is 10m².

Spot locations were also used to determine their distances to the closest streams, urban centers, and major roads and then were averaged at the 900m pixel size to be assigned to the corresponding location in the geodatabase.
145 These vector data, including the stream network, was retrieved from the map portal¹ of the Centre for Sustainable Urban Development (CEDEUS). Elevations of the study area were retrieved from the ASTERGDEM version 2, which is a digital elevation model produced at 30 m pixel size using stereo-correlation techniques applied to scenes from the ASTER sensor of the Terra satellite (Abrams et al., 2010). Three land cover characteristics were included in the database: a raster land cover map, a Normalized Difference Infrared
150 Index (NDII), and a Normalized Difference Vegetation Index (NDVI). The land cover map was derived from two Landsat images obtained from the platform Earth Explorer (<https://earthexplorer.usgs.gov>). The images were corrected geometrically, radiometrically and atmospherically (Chuvieco et al., 2002; Heilmayr et al., 2016). A maximum likelihood statistic of the supervised classification method (Chuvieco et al., 2002) was used to classify native forest, scrub, pasture/cropland, urban areas, exotic plantations, water bodies, bare soil and
155 burned areas. We used approximately 700 training points for each classified image, acquired through two sources, a) cadastral of the native plant resources of Chile (CONAF et al., 2015) and b) Google Earth (specifically its “time slider”) to obtain input to classify images.

NDII and NDVI data entered the geodatabase correspond to a pixel wise linear trends map for each index. All these raster maps were aggregated by simple averaging into a 900m pixel size and assigned to the nearest center
160 coordinate in the geodatabase.

Climatic descriptors included average summertime potential solar radiation, a temperature index, and a precipitation index. The ASTERGDEM was used to compute average summertime potential direct solar radiation employing the insol package within the R programming Language, package that implements algorithms presented by Corripio (2003 and references therein). The same procedure described for elevations
165 was implemented to add these data into the geodatabase.

¹ Publicly available at <http://datos.cedeus.cl/>



For temperatures, the value employed for each location was the linear trend in the number of summer (December, January, and February) days in which maximum temperatures were larger than the 90th percentile for all summers during the period 1980-2016, i.e., the linear trend in the Tx90p climate index (Klein Tank et al., 2009). For precipitation, the linear trend ingested into the geodatabase is from the number of Consecutive
170 Dry Days (CCD) in summer for the period 1980-2016, and index used in fire risk analysis (da Silva et al., 2020). The climatic data utilized for these calculations was the CR2MET product, a gridded climatology at 5km pixel size at daily to monthly frequency produced by the Chilean Center for Climate Resilience Research (CR2) covering the period 1979-2016². CR2MET was produced using a statistical downscaling of the ERA-Interim reanalysis supplemented by topographic data, land surface temperatures retrieved from satellites, and
175 instrumental observations (Alvarez-Garreton et al., 2018). The database also included linear trends of skin temperatures retrieved from the 0.05° (~5 km) daytime monthly land surface temperature product MODC11C6, version 6, derived from the Moderate Resolution Imaging Spectroradiometer (MODIS) aboard the Terra satellite (Wan et al. 2015), accessed from the GIOVANNI tool (Geospatial Interactive Online Visualization ANd aNalysis Infrastructure) at NASA's Goddard Earth Sciences Data and Information Services Center (Acker and Leptoukh, 2007). Within the geodatabase, trends in Tx90p, CDD, and skin temperatures were added to the
180 closest location falling within the respective 5 km pixel.

Table 1. Data sources of satellite products used in this study.

Source	Associated factor	Original Spatial resolution	Date
ASTERGDEM	Elevation and Summertime solar radiation	30m	Various
LANDSAT	NDII and NDVI Land cover	30m	2016
CEDEUS	Human infrastructure and activities (Highways and railways, Roads, Controlled burn areas, High voltage power lines, Camping zones)	Vector	CEDEUS 2016
Center for Climate Resilience Research (CR2)	Consecutive Dry Days (CDD) in 30 years	5km	1979-2016
CONAF (Chilean Forest Service)	Locations of wildfire spots	Vector	2008-2019

185 **2.2. Category development**

² Available at <http://www.cr2.cl/datos-productos-grillados/>



Implementing a data-driven approach allows for determination of discernible susceptibility thresholds according to the records available, which here are derived from observed spot recurrence. Thus, one of the firsts tasks in this research was the study of the 900m pixel map to determine whether there were detectable differences in spot recurrence. The categories were defined using a geometric sequence of the form:

$$C = R_{100\%} \times r \quad (1)$$

The 5 categories (C) were then computed by grouping recurrences (i.e., fires per year in each pixel) within the 2008-2019 period. $R_{100\%}$ represents the maximum value in the study area, assumed to be 100% recurrence. Thus, according to equation (1) the very high (VH) category considered recurrences from the maximum to half that recurrence, while the near zero (NZ) included the minimum of 0% recurrence ($r=0$). Thresholds for the intermediate categories high (H), media (M), and low (L) recurrence, are calculated using a ratio $r = \{0.5, 0.25, 0.125\}$, applied to $R_{100\%}$, respectively.

2.3. Model implementation, validation, and experiments

Based upon the known locations showing different recurrence categories, the modeling involved the development of a supervised classification scheme meant to determine the recurrence probability in the whole study area. To do so, two procedures were applied to the compiled geodatabase. First, a Principal Component Analysis (PCA) to reduce dimensionality from the eleven descriptors (excluding spot recurrence) to a new set of uncorrelated variables, called principal components or PCs, which maximize the explained variance while reducing redundancy among similar variables from the original database (Demšar et al., 2013). In this procedure, land cover classes were ingested using binary encoding, effectively enlarging the database to 18 descriptors. Afterwards, the PCs explaining most of the variance were used as input to a supervised classification using a Kohonen self-organizing map (SOM) algorithm (Kohonen, 1990). A SOM is a class of neural networks that reveals the structure of a dataset by competitive learning. The supervised classification was implemented as an iterative process where a random selection of locations from the recurrence categories was presented to the SOM, using the corresponding PCA output as descriptors. During a given iteration the algorithm selected 50 locations per category, classified by comparing those locations with the rest of the study area. With output of all iterations, the model calculates a simple probability to determine which category a certain location falls more often, assigning the respective value. Computation of the SOM's network size and iteration number was determined following recommendations by Kohonen (1990) and Vesanto (2000). Evaluation and validation of model output included using the MCD14ML product, a MODIS standard quality Thermal Anomalies/Fire locations database, accessed through NASA's Fire Information for Resource Management System (FIRMS).

A long-standing debate exists on whether forestry plantations enhance wildfire occurrence in this region (Úbeda and Sarricolea, 2016; Urrutia-Jalabert et al., 2018; de la Barrera et al., 2018). This presented model allows



225 testing the sensitivity of the study area to different scenarios and thus two extreme situations were compared.
A first model configuration assumed all non-urban areas as covered by native vegetation while a second
scenario considered all non-urban areas as plantation. Ingestion of these two scenarios into the SOM model was
through using the corresponding weights of the PCA to recalculate the score of each location relative to the
selected PCs.

230 3. Results: Fire Hazard in A Metropolitan Area: Analysis of Factor Maps and Model Output.

3.1. Analysis of geodatabase components

235 Spatial patterns of forest fire recurrence during the period under study was associated with the peri-urban areas
of the main cities of the CMA, since the quadrants with a recurrence of more than 20 fires were found at less
than 650 m from urban centers and highways. This suggests that the causes of these forest fires are mostly
anthropogenic. On the contrary, the areas that did not record fire outbreaks during the study period were
associated with remote locations with an average elevation of 250 m, distant 8.1 km from urban centers and 1.5
km from highways.

240 Climatic descriptors show a general increase in 0.59°C/year for Tx90, 0.51 days/year in CDD, while a decrease
of -0.52°/year in skin temperature. Tx90 shows a range of 1.8°/year and a standard deviation of 0.4°/year,
indicating that most locations have been undergoing an increase in maximum temperatures. In fact, only 7% of
the pixel's present negative linear trends, with most of them located on a buffer of about 5 to 7 km from the
South East of the main urban area of Concepción. The distribution of CDD trends, on the other hand, is
245 completely positive, with a minimum value of 0.26 days/year and maximum of 0.8 days/year. Skin temperature
is the only variable with a relatively clear spatial differentiation; while 62% of the region records a decreasing
trend, the 38% of the study area that shows the opposite behavior is on or near urban sectors of the CMA. This
satellite dataset indicates cooling as low as -4.2°/year and warming of 4.6°/year, with a standard deviation of
4.02°/year.

250 The comparison of CONAF hotspot's distances to streams, roads, and urban areas suggests a significant impact
of roads on fire occurrence, since on average hotspots are mapped at about 1.3 km from roads. In turn, streams
and urban areas are at 5.5 km and 6.6 km, respectively. Despite this pattern, the large corresponding standard
deviations of 1.1 km, 5.9 km, and 6.3 km indicates a significant spread.

255 The values of the NDVI index, which is associated with vegetation composition and structure, show values
ranging from -0.2312 to 0.4460. Negative values of this indicator are related to non-vegetation land covers/uses
such as water bodies, urban areas, or bare soils. On the other hand, the positive values are related to coverages
with low vegetation such as pastures or scrub (values close to zero) to dense vegetation such as arborescent
scrub, forest plantations or native forest (NDVI>0.25). Within the study area, 9.1% of the pixels presented
negative NDVI values (no vegetation), 35.7% corresponded to positive intermediate NDVI values (grasslands,
260 scrublands, young forest plantations), whereas 55.2% corresponded to high NDVI values (adult forest
plantations, native forest). The values of the NDII index, which is associated with the moisture content of the




265 vegetation, showed values ranging from -0.154 to 0.450. The negative values of this index are related to cover/uses without vegetation such as water bodies or urban areas. Positive values are associated with water content in the vegetation; the value of the index increases with increasing water content in the vegetation. Within the study area, 7.4% of the pixels presented negative NDII values, 42.8% corresponded to positive NDII values (but close to zero), mainly associated with shrublands and grasslands. Finally, 64.7% corresponded to medium-high NDII values, which are associated with adult forest plantations and native forest.

270 The elevation ranges in the study area fluctuated between 0 m to 910 m, with an average elevation of 212m. The highest elevations are found in the southern part of the study area and are mostly associated with the presence of forest plantations and small fragments of native forest. The western area presents the lowest terrain elevations, presenting itself as a coastal plateau where the urban areas of the CMA are present. The insolation on the ground, which directly influences the formation of fuel for the formation of forest fires, reached mean values of 887.7 W/m², with insolation ranges from 882.06 W/m² to 905.8 W/m². The highest insolation values are associated with higher elevation sectors, with north-facing terrain aspect. Insolation on the ground showed 275 little variation due to the low variability of elevation gradients in the study area. Finally, with respect to the land use/cover present in the study area, 55% of the surface corresponded to exotics forest plantations of *Pinus radiata* and *Eucalyptus globulus* species, which are associated with the highest sectors of the study area. The 7% corresponds to native forests of *Nothofagus* sp., which are present in areas of high slope (>30%). On the other hand, 15.2% corresponds to agricultural zones, which are associated with low sectors in the eastern part 280 of the study area. Finally, urban areas represent 5.3% of the total surface area and are found mainly in the coastal zone (west) of the study area.

When the data of the geodatabase is inspected according to the categories determined from spot recurrence (equation (1)), several patterns emerge (Figure 2). A first finding is that the Medium (M) to Very High (VH) 285 categories tend to present much less spread, with almost no outliers. It is also noticeable that the Near Zero (NZ) and Low (L) categories spread to about the same range, suggesting that low recurrences do not conform to a distinguishable pattern of recurrence and that instead they correspond to random events. Variables associated with vegetation characteristics, i.e., trends in NDII (t-NDII) and in NDVI (t-NDVI), tend to show that the VH category is mostly associated with negative trends, deepening the decreasing tendency from H.

290 As expected from the method utilized to calculate insolation, that variable and elevation show a similar pattern where M, H, and VH fall in regions of progressive higher values. Spot distance to Urban centers (d-Urban) appear to be within a narrow range for M, H, and VH relative to L. Although with absolute values lower than d-Urban, distances to roads (d-Road) are largely shorter than 2 km for the three highest categories, with the median for VH being marginally shifted towards 2 km relative to H and M. For the distance relative to streams (d-Stream), spot recurrence tends to be higher at a separation below 2.5 km for H and VH.




295

Figure 2. Main features of the variables in the geodatabase, grouped according to the wildfire categories: Near Zero (NZ), Low (L), Medium (M), High (H), Very High (VH).

The distribution of climatic variables according to categories shows a clear concentration within a narrow range for VH. As previously detected, the trend in CDD (t-CDD) for the CMA is completely positive, although for VH it is concentrated just below 0.5d/year. In the case of Tx90 (t-Tx90), results indicate that M, H, and VH spot recurrence have only occurred on areas with positive change, with H showing the largest interquartile range. For skin temperature trends (t-SkinT), most locations with positive trends coincide with high and very high recurrence. Figure 3 shows the land use data pertaining to each category. The most striking pattern is that 100% of VH occurs over plantations, about 70% for H, and nearly 45% for M.

305



310

Figure 3 Land use (%) classified for the CMA according to the wildfire categories: Near Zero (NZ), Low (L), Medium (M), High (H), Very High (VH) and All. “All”, represents the distribution of the land cover along the whole study area.



In addition, the progressive importance of urban land use connected with plantations as the recurrence increases, suggests that the connection between these two land uses explains most of the damaging effects of wildfires in the CMA. As already seen in the distribution of other variables, L and NZ present a similar partitioning as the whole study area (“All” in Figures 2 and 3), further attesting for a random pattern of low recurrence. This way, the analysis of input variables tends to indicate that there is a relatively consistent pattern of landscape conditions that allow for certain locations to record fires more recurrently than others.

3.2. PCA and SOM model output.

Six principal components (PCs) explain about 71% of the geodatabase variance within the study area. Since the PCA was applied to the whole CMA, these results represent the relationships including zones with zero fire spots. Although no PC within these 6 accounts for more than 20%, certain patterns emerge that suggest the procedure has been able to suppress redundancies in the database (Figure 4). The PC1 reaffirms the relationship between Elevation and Insolation and the similar behavior of d-Stream and d-Urban. Tx90 tends to be important in both PC1 and PC2, while in the latter t-NDVI and t-NDII show the highest weights, indicating important correlations between them. In turn, d-Road and t-CDD are largely influential mostly in PC3. A remarkable finding is that this PCA standardization does not show land use as a prominent variable, as plantations appear to be somewhat key in PC6 only.

330





Figure 4. Input to the SOM Model presented in the percentage of six principal components (PCs).



335

The analysis of the PCA partitioned by recurrence categories reveals that only for certain PCs M, H, and VH develop distinct signatures (Figure 5). These results also reaffirm the similarity of L with a situation of nearly zero spots. Although in almost all PCs the increase in recurrence shows a trend of a narrow range relative to the immediate previous category, there are certain remarkable exceptions, such as the slight reverse trend of VH relative to H in PC1 and the significant interquartile range for H in PC6. A noticeable finding is that VH's interquartile range seems different from the rest of the categories for PC1, PC2, and PC4.



340

Figure 5. Boxplots showing the distribution of PC's weights according to the recurrence categories. "All" represents the distribution of each PC along the whole study area.

345

SOM output compared with CONAF and MODIS data indicates that this data-driven model is skilled in predicting an increase in spot density according to the corresponding category (Table 3). Given that the model was trained with CONAF data, it is expected that a better match is found in that comparison. In effect, the model predicts increasing density according to the recurrence category. The VH's 10.98 spots per pixel is 3 times denser than H, and about 5 times relative to M. For the case of MODIS data, the model also finds a significant increase in density for VH, almost 4 times higher than M and H; however, it does not predict density differences between M and H while records a slightly larger density for L. The NZ category is well predicted by the model compared to both sources of fire spot activity: ~6 times smaller for CONAF and about half for MODIS.

350

According to the model, 55.8% of the CMA presents conditions for low recurrence of fire spots, with about 1/3 for M, and just 12.3% for high and very high recurrence. The model also predicts that spot recurrence is a phenomenon that may affect almost the whole study area (Figure 6A).

355



Table 2. Comparison of SOM model output versus CONAF and MODIS fire spot density, respectively. Density is calculated as the number of CONAF or MODIS spots that fall within a given CMA's model category, divided by the number of pixels corresponding to that category.

Category	Modeled Area covered (%)	Density (spots per pixel)	
		CONAF	MODIS
Near zero	3.3	0.13	0.45
Low	55.8	0.62	0.78
Medium	28.6	2.13	0.72
High	10.3	3.66	0.72
Very High	2.0	10.98	2.84

Note: MODIS reports 50.04% less hotspots than CONAF.

360

The native scenario tends to show more pixels in the medium category than the plantation. Also, the native scenario sees an increase in the VH category. On the other hand, plantations tend to show an increase in the L and H categories, while reducing the NZ. Although these differences are not extreme, they attest for a different dynamic depending on the prevalent land cover. Both models show clustering patterns in which very low and low values are associated with higher elevation sectors within the CMA, which in turn have the lowest insolation values. On the other hand, the high and very high values are associated with low elevations near the roads and urban areas of the CMA (Figure 6 B and C).

365

Table 3. Modeled proportion of spot recurrence within the CMA (%) according to the five categories considering the scenario in which the whole study area is assumed to be covered by native (native experiment) and exotic plantation (exotic plantation experiment). The lower section of the table compares spot counts and surface area per category relative to the full model using the original database, and the experiments.

370

Category	Native experiment (%)	Exotic plantation experiment (%)
Near zero	11,25	10,30
Low	50,73	54,22
Medium	29,27	25,58
High	6,50	8,49
Very high	2,24	1,41

Category	Spot counts			Surface (ha)		
	Full model	Exotic plantation experiment	Native experiment	Full model	Exotic plantation experiment	Native experiment
Near zero	86	296	310	6966	23976	25110
Low	1556	1501	1438	126036	121581	116478
Median	827	732	822	66987	59292	66582



High	280	239	174	22680	19359	14094
Very high	59	40	64	4779	3240	5184
TOTAL	2808	2808	2808	227448	227448	227448

375




Figure 6. Model results of wildfire recurrence on the CMA. On the left (A), model results using the original geodatabase; (B) corresponds to the scenario of only exotic plantations; (C) is the only native forest experiment.

380

4. Discussion

In Mediterranean Central Chile, land cover changes that characterize current landscape organization resulted mostly from the application of the government subsidies granted by Law Decree 701 for Forestry Development (DL701) in 1974 (Nahuelhual et al. 2012; INFOR, 2017:25). This policy favored plantations of exotic fast-growing species along the region, with staggering consequences: in 1974, the surface area of forestry plantations was 480,000 ha, during the 1990s close to 2 million ha (Aguayo et al. 2009), reaching nearly 5 million ha in 2015 (INFOR, 2017:49). This ten-fold increase in plantations motivated by public policy contrasts with the little attention paid to restoring native forests, which have historically contributed to the local population's livelihoods (Reyes and Nelson, 2014; Frene and Núñez, 2010) causing that the rural and indigenous

390



communities must compete for the use of the land against the plantations inciting environmental conflicts (INDH, 2015).

Concomitant with the plantation spread along the region, an increase in the recurrence and magnitude of fire disturbances in WUI has been observed, due to the blurred border between land covers or the substitution of certain land use for others (Goldman, 2018; Ruiz et al. 2017; Ladislao et al. 2007). According to CONAF, over 35 million hectares of vegetation are vulnerable to fires, including grasslands and shrubland (20 million), native forest (13 million) and exotic plantations (2,1 million) (Castillo et al. 2003). Of this vegetation, over 50 thousand hectares are burned annually in approximately 5,900 wildfires. Under these political and economic conditions, the land change cover seems to become a critical factor that contributes to the wildfire risk, whose conflictive evolution has built a double pressure scenario that shows no sign of changing (Ubeda and Sarricolea, 2016; Andersson et al. 2016). Likewise, the urban expansion fomented by the National Policy of Urban Development (NPUD) from 1979 has been deregulating the land use market (Brites, 2017) fomenting the urbanization of agricultural lands, wetlands, or forests (IDB-ECLAC, 2015; Vilar del Hoyo et al., 2011; Hidalgo et al. 2018).

The machine learning model developed in this work shows that more than 90% of the CMA may be subject to some degree of fire risk, with about 40% with at least a medium probability of recurrence. Wildfire hotspot density is well represented by the model, which suggests this tool could be a powerful decision-making tool for the public sector (i.e., national government, municipalities) and the private sector (universities, timber companies, real estate developers). Hotspot density is concentrated on roads (1.3 km), leaving far behind the water streams (5.5 km) and urban areas (6.6 km), consistent with the literature that assigns the major responsibility of the fire recurrence to the presence of human infrastructure and human activities (Harari, 2013; Doerr and Santin, 2016), and is consistent with CONAF's previous reports (CONAF 2017; 2018). However, anthropic factor is not the only one to count in, as Barbati et al. (2013) said, the distance from the nearest water body is determinant for short-term fire recurrence in Mediterranean countries, between other landscape factors (slope roughness, exposure, pre-fire dominant forest type). Additionally, the proximity to roads and maximum temperature dynamics, both variables severely altered by the human activities, tend to organize the randomness observed in this model. This high random component in the occurrence of events is associated with a lower wildfire hazard, which reveals that after a random appearance, the recurrence increases according to the conditions of each zone. This last idea of random distribution of low and almost zero recurrences has been around a long time, and literature reports similar results from their GIS models in Sardinia – Italy (Ricotta and Di Vito, 2014), California – USA (Minnich and Chou, 1997) and Spain (Chuvieco et al., 2011:49; Vilar del Hoyo et al. 2011). Nevertheless, is contradictory with the official data coming from CONAF, which established that “a high occurrence was recorded in the interface areas of the region, because of the repeated occurrence of forest fires during the 2015-2016 season” (CONAF, 2021) and the study of central Chile from McWethy et al. (2018) “fire activity was highly variable in any given year, with no statistically significant trend in the number of fires or mean annual area burned”.

Experiments comparing total plantation and total native scenarios, while not generating a significant change in categories, suggests that there are more areas that will be subject to fires (fewer Near Zero areas) under more plantations. The fact that category Low tends to increase, even if it is marginally, perhaps indicating that the



430 probability of prediction is reduced when coverage is mostly plantation, as the system tends to become more
random. Also, it is relevant that the model does detect a difference between scenarios of total plantation and
native forest cover, pointing that the changes in fire regime and fire occurrence relies heavily on the vegetal
cover for central Chile, which is consistent with the literature (McWethy et al., 2018; de la Barrera et al., 2018;
Úbeda and Sarricolea, 2016), press reports (CIPER, 2018), and official reports (CONAF, 2017; CONAF-BIRF,
1999). Other studies suggest the same relevance of landscape drivers for Mediterranean countries (Darques,
435 2015; Pausas et al., 2008; Turco et al., 2016). By examining the features of the model presented here, it is
possible to propose two, no necessarily exclusive possibilities that may explain the relatively weak contribution
of forest plantations to fire risk. The first is that anthropogenic activities may become more important at the
local rather than the regional scale. For instance, whereas the model shows that the urban boundary is
overwhelmingly associated with categories M to VH, if one zooms out to the whole Mediterranean Central
440 Chile, cities become small spots. The second possibility may be a “saturation effect” in the sense that plantations
now occupy such a significant surface area within the CMA that the influence is already permanent in the
current regime of fires, meaning that any data-driven treatment sees plantations as a constant and thus attributes
a small contribution. That’s the reason why the native vs plantation experiments are important, because indicates
that the plantation tends to reduce the areas with near zero recurrence relative to the native scenario, although
445 the difference is marginal, likely associated with the “saturation effect”.

Results of the model thus are relevant as they serve to accumulate and analyze historical, cartographical, and
other types of data, leading to a better understanding of controls and drivers on fire activity in the CMA at high
resolution. However, the model can be substantially improved with near real-time (NRT) information from
terrestrial platforms (e.g., vehicles, towers, cranes), airborne platforms (e.g., aircraft, unmanned aerial vehicles
450 (UAVs), helicopters) or space-borne platforms (e.g., satellites) using electromagnetic sensors (Van Ackere et
al. 2019) leading us to a truly smart metropolitan area (Costa et al., 2020). In that sense, becomes necessary to
put more effort in the future to extend the timeframe of the present study, as Chuvieco et al. (2011:54) accurately
said: “*Since fire occurrence changes in space and time, the validation of integrated indices should be done with
long time series, because short periods may bias some of the theoretical assumptions that are required to build
455 the model*”.

Future scenarios for the CMA are filled with uncertainty, especially for climate change and associated impacts.
Projections from the work of Araya-Muñoz et al. (2017) indicated that the most relevant hazards for the CMA
will be wildfires, water scarcity, and heat stress. Likewise, the droughts are becoming more recurrent (Garraud
et al., 2020; Fernández et al., 2018). As the model suggests, climatic indicators play a role in fire recurrence,
460 which allows us to infer that changes in those will lead to increases in wildfire hazard for the CMA. What is
changing fast are the climate conditions, creating riskier scenarios globally. Therefore, there is an opportunity
to improve or made mandatory the nature-based solutions, controlled burns for a social-ecological
transformation (Otero and Nielsen, 2017), the ecological restoration of soils, wetlands, and forests, REDD++,
20x20 initiative, promoting the carbon emission market for carbon sequestration (Wright et al. 2000). For
465 example, the *Pinus radiata* plantations in Chile and Australia have a potential average net annual rate of CO²



accumulation of 4.5 tons (IPCC, 1996), sequestering greenhouse gas emissions faster at a lower cost, returning the investments quickly, and mitigating some of the impacts of climate change (Pawson et al. 2013).

5. Conclusions

470

The results indicated that 12.3% of the CMA's surface area has a high and very high risk of a forest fire, 29.4% has a medium risk, and only 58.3% has a low and very low risk. This calls for reflection on the importance of spatial planning with a resilient focus on wildfires, according to the recurrence of these phenomena in these settings as they are increasingly more forced in the WUI, urban residential areas and industrial or port areas.

475

These maps and this model are of vital importance for the Chilean government emergency agencies (CONAF-ONEMI) as well as for the city governments within the CMA. They are also relevant for understanding how these phenomena affect the Mediterranean ecosystems to which the CMA belongs, and therefore should be beneficial for researchers in other latitudes working on similar ecosystems: California, Australia, Italy, Spain, Portugal, etc.

480

This study aimed to establish the relationship between the natural and the anthropogenic factors associated with infrastructure that has been proved to generate wildfires, providing valuable data about the model itself and the capacity of fire risk prediction in a holistic viewpoint. Nevertheless, it will be challenging to generate further studies of the change associated with the inclusion of meteorological data about this Mediterranean ecosystem as the indexes used to define the thresholds tend to put medium risk areas in higher ranges. Also, the CC has been an international challenge mandatory for most countries to fulfill the Paris agreements of December 2015 (COP21). Particularly, Chile has compromised, between other politics, the plantation of 100,000 ha with mostly native forests for carbon sequestration. Similarly, the 20x20 initiative and the National Restoration Landscape Program are searching for CC mitigation through soil organic carbon sequestration and the reduction of forest fire occurrences in Chile. This program aims to restore 100,000 ha of degraded native forest into forest plantations and restore 400,000 ha of degraded land for agriculture and cattle ranching through the System of Incentives for Recuperation of Degraded Soil.

485

490

Code availability: Computer code for the model available on request

Data availability: all databases used in this research are free to access from the links included in the paper.

495

Author contributions: Edilia Jaque Castillo designed the study in consultation with Rodrigo Fuentes Robles and Carolina G. Ojeda. Alfonso Fernández developed, coded, and run the model aided by all co-authors. Rodrigo Fuentes Robles processed and analyzed remote sensing data. All co-authors participated in statistical analyzes, discussion of results and writing.

500

Competing interests: The authors declare no competing interests

References



- 505 Abrams, M., Bailey, B., Tsu, H., Hato, M. 2010. The ASTER Global DEM. *Journal of American Society Photograph Remote Sensing* 20: 344–348.
- Acker, J.G. and Leptoukh, G. 2007. Online analysis enhances use of NASA Earth science data. *Eos, Transactions of American Geophysical Union*, 88: 14. <http://dx.doi.org/10.1029/2007EO020003>
- 510 Adams, M. 2013. Mega-fires, tipping points and ecosystem services: Managing forests and woodlands in an uncertain future. *Forest Ecology and Management*, 294: 250-261. <https://doi.org/10.1016/j.foreco.2012.11.039>
- Aguayo, M., Pauchard, A., Azócar, G., Parra, O. 2009. Cambio del uso del suelo en el centro sur de Chile a fines del siglo XX: Entendiendo la dinámica espacial y temporal del paisaje. *Revista Chilena de Historia Natural*, 82:361-374 <https://dx.doi.org/10.4067/S0716-078X2009000300004>
- 515 Álvarez - Garretón, C., Mendoza, P.A., Boisier, J.P. 2018. The CAMELS-CL dataset: catchment attributes and meteorology for large sample studies – Chile dataset. *Hydrology Earth System Science*, 22:5817–5846. <https://doi.org/10.5194/hess-22-5817-2018>
- Andersson, K., Lawrence, D., Zavaleta, J., Guariguata, M.R. 2016. More trees, more poverty? The socioeconomic effects of tree plantations in Chile, 2001–2011. *Environmental Management* <https://doi.org/10.1007/s00267-015-0594-x>
- 520 Araya-Muñoz, D., Metzger, M.J., Stuart, N., Meriwether, A., Wilson, W., Carvajal, D. 2017. A spatial fuzzy logic approach to urban multi-hazard impact assessment in Concepción, Chile. *Science of the Total Environment*, 576: 508–519 <http://dx.doi.org/10.1016/j.scitotenv.2016.10.077>
- Barbati, A., Corona, P., D’amato, E. Cartisano, R. 2013. Is landscape a driver of short-term wildfire recurrence? *Landscape Research*, 40(1):99-108 <https://doi.org/10.1080/01426397.2012.761681>
- 525 Biblioteca del Congreso Nacional (BCN). 2017. *Reportes comunales*. Available at: <https://reportescomunales.bcn.cl/2015/index.php/Categor%C3%A0Da:Comunas>
- Boisier, J.P., Rondanelli, R., Garreaud, R.D., Muñoz, F. 2016. Anthropogenic and natural contributions to the Southeast Pacific precipitation decline and recent megadrought in central Chile. *Geophys Research Letters*, 1–9. <https://doi.org/10.1002/2015GL067265>
- 530 Brites, W.F. 2017. La ciudad en la encrucijada neoliberal. Urbanismo mercado-céntrico y desigualdad socio espacial en América Latina. *Urbe, Revista Brasileira de Gestão Urbana*, 9(3):573-586. <https://doi.org/10.1590/2175-3369.009.003.ao14>
- Bustamante, L.P. and Varela, E.S. (2007). Crecimiento urbano y globalización: transformaciones del Área Metropolitana de Concepción, Chile, 1992-2002. *Scripta Nova. Revista Electrónica de Geografía y Ciencias Sociales*, 11(251). Available at: <http://www.ub.edu/geocrit/sn/sn-251.htm>
- 535 Castillo, M., Pedernera, P., Peña, E. 2003. Incendios forestales y medio ambiente: una síntesis global. *Revista Ambiente y Desarrollo de CIPMA*, XIX (3-4): 44-53. Disponible en: <http://www.keneamazon.net/Documents/Publications/Virtual-Library/Economia-Desarrollo/29.pdf>
- 540 Castree, N. 2008. Neoliberalising nature: the logics of deregulation and reregulation. *Environment and Planning A*, 40(1): 131-152. <https://doi.org/10.1068/a39100>



- Cid, B. 2015. Peasant economies, forestry industry and fires: socio-natural instabilities and agriculture as means of resistance. *Ambiente & Sociedad*, XVII (1): 93–114. <https://doi.org/10.1590/1809-4422ASOC720V1812015esp>.
- Center for Climate and Resilience (CR2MET). 2020. *Climate data from Chile*. [dataset]. Available at: <http://www.cr2.cl/>
- 545 CENTRO INVESTIGACIONES PERIODÍSTICAS (CIPER Chile). 2018. *Alertas que anunciaban la catástrofe se arrastran desde 2012. Mega incendios: el historial de omisiones de las autoridades que abonó la tragedia*. Por Alberto Arellano. Available at: <https://ciperchile.cl/2017/02/01/mega-incendios-el-historial-de-omisiones-de-las-autoridades-que-abono-la-tragedia/>
- 550 CONAF-BIRF. 1999. *Catastro y evaluación de los recursos vegetacionales nativos de Chile. Informe nacional con variables ambientales*. CONAF/CONAMA/BIRF, Santiago.
- CONAF. 2017. *Análisis de la afectación y severidad de los incendios forestales ocurridos en enero y febrero de 2017 sobre los usos de suelo y los ecosistemas naturales presentes entre las regiones de Coquimbo y La Araucanía de Chile*. Santiago de Chile, CONAF. Available at: http://www.conaf.cl/tormenta_de_fuego-2017/INFORME-AFECTACION-Y_SEVERIDAD-DE-INCENDIOS-FORESTALES-VERANO-2017-SOBRE-ECOSISTEMAS-VEGETACIONALES-CONAF.pdf
- 555 CONAF. 2018. *Estadísticas de incendios forestales en Chile*. Available at: <http://www.conaf.cl/incendios-forestales/incendios-forestales-en-chile/>
- CONAF 202. *Prevención de Incendios Forestales en Zonas de Interfaz de la Región del Biobío*. Disponible en: <https://www.conaf.cl/incendios-forestales/prevencion/yo-tambien-soy-forestin-campana-de-prevencion-de-incendios-forestales-2020/prevencion-de-incendios-forestales-en-zonas-de-interfaz-de-la-region-del-biobio/>
- 560 Corripio, J. G. 2003. Vectorial algebra algorithms for calculating terrain parameters from DEMs and the position of the sun for solar radiation modelling in mountainous terrain. *International Journal of Geographical Information Science*, 17(1): 1-23. <https://doi.org/10.1080/713811744>
- 565 CR2. (2020). Climatic Explorer. Center for Climate and Resilience Research (CR)2. Available at: <http://explorador.cr2.cl>
- Costa, D.G.; Vasques, F.; Portugal, P.; Aguiar, A. 2020. A Distributed Multi-Tier Emergency Alerting System Exploiting Sensors-Based Event Detection to Support Smart City Applications. *Sensors*, 20: 170. <https://doi.org/10.3390/s20010170>
- 570 Change, N.C. 2017. Spreading like wildfire. *Nature Climate Change*. Doi: 10.1038/nclimate3432
- Chuvieco, E. 2002. *Teledetección ambiental. La observación de la tierra desde el espacio*. Barcelona, Ariel.
- Chuvieco, E., Salas, J., De la Riva, J., Pérez, F., Lana-Renault, N. 2004. *Métodos para la integración de variables de riesgo: el papel de los sistemas de información geográfica. Nuevas tecnologías para la estimación del riesgo de incendios forestales*. CSIC-Instituto de Economía y Geografía, Madrid.
- 575 Chuvieco, E., Aguado, I., Yebra, M., Nieto, H., Salas, J., Martín, M.P., De La Riva, J. 2011. Development of a framework for fire risk assessment using remote sensing and geographic information system technologies. *Ecological Modelling*. <https://doi.org/10.1016/j.ecolmodel.2008.11.017>



- 580 Dapeng, L., Cova, T.J., Dennison, P.E., Wan, N., Nguyen, Q.C., Siebeneck, L.K. 2019. Why do we need a national address point database to improve wildfire public safety in the US? *International Journal of Disaster Risk Reduction*, 39: 101237 <https://doi.org/10.1016/j.ijdrr.2019.101237>
- Darques, R. 2015. Mediterranean cities under fire. A critical approach to the wildland–urban interface. *Applied Geography*, 59: 10–21. <https://doi.org/10.1016/j.apgeog.2015.02.008>
- da Silva, A.S., Justino, F., Setzer, A.W., Ávila - Díaz, A. 2020. Vegetation fire activity and the Potential Fire Index (PFIv2) performance in the last two decades (2001–2016). *International Journal of Climatology*, 77: 1–15. DOI: 10.1002/joc.6648.
- 585 de la Barrera, F. and Henríquez, C. 2017. Vegetation covers change in growing urban agglomerations in Chile. *Ecological Indicators*, <http://dx.doi.org/10.1016/j.ecolind.2017.05>
- de la Barrera, F., Barraza, F., Philomène, F., Ruíz, V., Quense, J. 2018. Megafires in Chile 2017: Monitoring multiscale environmental impacts of burned ecosystems. *Science of the Total Environment*, 637–638:1526–1536. <https://doi.org/10.1016/j.scitotenv.2018.05.119>
- 590 Demšar, U., Harris, P., Brunson, C. 2013. Principal Component Analysis on Spatial Data: An Overview. *Annals of the Association of American Geographers*, 103:106–128. <https://doi.org/10.1080/00045608.2012.689236>
- Diffenbaugh, N.S., Pal, J.S., Giorgi, F., Gao, X. 2007. Heat stress intensification in the Mediterranean climate change hotspot. *Geophysical Research Letters*, 34: L11706 <https://doi.org/10.1029/2006GL025734>
- 595 Doerr, S.H., Santin, C. 2016. Global trends in wildfire and its impacts: perceptions versus realities in a changing world. *Philosophical Transactions of Royal Society B*, 371(20150345): 1–10. <http://doi.org/http://dx.doi.org/10.1098/rstb.2015.0345>
- Fernández, A., Muñoz, A., González-Reyes, Á. 2018. Dendrohydrology and water resources management in south-central Chile: lessons from the Río Imperial streamflow reconstruction. *Hydrol. Earth Systems Science* 22:2921–2935. <https://doi.org/10.5194/hess-22-2921-2018>
- 600 Frene, C. and Núñez, M. 2010. Hacia un nuevo Modelo Forestal en Chile. *Revista Bosque Nativo*, 47: 25–35. Available at: <http://www.sendadarwin.cl/espanol/wp-content/uploads/2011/06/frene-y-nunez.pdf>
- Freudenburg, W.R. 1992. Addictive economies: Extractive industries and vulnerable localities in a changing world economy. *Rural Sociology*, 57(3): 305–332. <https://doi.org/10.1111/j.1549-0831.1992.tb00467.x>
- 605 Gago, V., Mezzadra, S. 2017. A critique of the extractive operations of capital: Toward an expanded concept of extractivism. *Rethinking Marxism*, 29(4): 574–591 DOI: 10.1080/08935696.2017.1417087
- Garraud, R.D., Boisier, J.P., Rondanelli, R., Montecinos, A., Sepúlveda, H.H., Veloso - Águila, D. 2020. The Central Chile Mega Drought (2010–2018): A climate dynamics perspective. *International Journal of Climatology*, 40: 421–439. <https://doi.org/10.1002/joc.6219>
- 610 Goldman, J.G. 2018. Living on the edge: Wildfires pose a growing risk to homes built near wilderness areas. Building houses at the edge of the wilderness increases the danger of catastrophic blazes. *Scientific American*, 318(6): 12–15. Available at: <https://www.scientificamerican.com/article/living-on-the-edge-wildfires-pose-a-growing-risk-to-homes-built-near-wilderness-areas/>



- 615 Gómez-González, S., Ojeda, F., Fernandes, P.M. 2018. Portugal and Chile: Longing for sustainable forestry while rising from the ashes. *Environmental Science and Policy*, 81: 104–107. <http://doi.org/10.1016/j.envsci.2017.11.006>
- González-Mathiesen, C. and March, A. 2018. Establishing design principles for wildfire resilient urban planning. *Planning Practice & Research*, DOI:10.1080/02697459.2018.1429787
- 620 Harari, Y. 2013. *From Animals into Gods: A brief history of Humankind*. New York: Penguin Random House.
- Heilmayr, R., Echeverría, C., Fuentes, R., Lambin, E.F. 2016. A plantation-dominated forest transition in Chile. *Applied Geography*, 75: 71-82. <https://doi.org/10.1016/j.apgeog.2016.07.014>
- Hidalgo, R., Rodríguez, L., Alvarado, V. 2018. Up the hill or over the wetland: production of nature and expanding real estate in marine and fluvial cities. The case of Valparaíso and Valdivia, Chile. *Diálogo Andino*,
625 56: 87-100 <http://dx.doi.org/10.4067/S0719-26812018000200087>
- IDB-ECLAC. (2007). Information on disaster risk management. Case study of five countries. Main technical report. Mexico City: Inter-American Development Bank (IDB) – Economic Commission for Latin American and the Caribbean (ECLAC). Available at: <https://www.cepal.org/en/publications/25845-information-disaster-risk-management-case-study-five-countries-main-technical>
- 630 Instituto Forestal (INFOR). 2017. *Anuario forestal 2017*. INFOR: Santiago de Chile. Available from: <http://wef.infor.cl/publicaciones/anuario/2017/Anuario2017.pdf>.
- Instituto Nacional De Estadísticas (INE). 2021. *Censo 2017. Síntesis de resultados*. Santiago de Chile: Ministerio de Economía, Fomento y Reconstrucción. Available at: <http://www.censo2017.cl>
- Instituto Nacional de Derechos Humanos (INDH). 2015. *Mapa de conflictos socioambientales en Chile*.
635 Santiago de Chile: INDH. Disponible en: <https://mapaconflictos.indh.cl/#/>
- Intergovernmental Panel on Climate Change (IPCC). 1996. *Guidelines for national greenhouse gas inventories, and good practice guidance reports*. Cambridge, USA: Cambridge University Press.
- IPCC 2014. *Summary for policymakers*. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and sectoral aspects. Contribution of Working Group II to the Fifth Assessment Report of the*
640 *IPCC*. Cambridge, USA: Cambridge University Press.
- Pp. 1-32. Available at: https://www.ipcc.ch/site/assets/uploads/2018/02/ar5_wgII_spm_en.pdf
- IPCC 2019. *Summary for Policymakers*. In: *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. Cambridge, USA: Cambridge University Press. Available at: <https://www.ipcc.ch/srocc/chapter/summary-for-policymakers/>
- 645 Klein Tank, A., Zwiers, F., Zhang, X. 2009. *Guidelines on analysis of extremes in a changing climate in support of informed decisions for adaptation*. *Climate Data and Monitoring WCDMP-No. 72 WMO-TD No. 1500*. Available at: http://www.wmo.int/pages/prog/wcp/wcdmp/documents/WCDMP_72_TD_1500_en_1.pdf
- Knowles, A.K., Westerveld, L., Strom, L. 2015. Inductive visualization: A humanistic alternative to GIS. *GeoHumanities*, 1(2): 233-265. DOI: 10.1080/2373566X.2015.1108831
- 650 Kohonen, T. 1990. The Self-Organizing Map. *Proc. IEEE*, 78: 1464–80.



- Kolden, C.A. and Henson, C. 2019. A Socio-Ecological approach to mitigating wildfire vulnerability in the Wildland Urban Interface: A case study from the 2017 Thomas fire. *Fire*, 2 (9): 2-19
<http://doi.org/10.3390/fire2010009>
- 655 Koltunov, A., Ustin, S.L. Prins, E.M. 2012. On timeliness and accuracy of wildfire detection by the GOES WF-ABBA algorithm over California during the 2006 fire season. *Remote Sensing of Environment*, 127:194-209
<https://doi.org/10.1016/j.rse.2012.09.001>
- Kumagai, Y., Carroll, M. S., Cohn, P. 2004. Coping with interface wildfire as a human event: lessons from the disaster/hazard's literature. *Journal of Forestry*, 102(6): 28-32. <https://doi.org/10.1093/jof/102.6.28>
- 660 Martínez Poblete, J. 2014. Land registry and condition of conservation of the wetlands coastal/marines in the region of the Biobío. *Tiempo y Espacio*, 33, 104-130.
- McGee, T.K. and Russell, S. 2003. "It's just a natural way of life..." An investigation of wildfire preparedness in rural Australia. *Environmental Hazards*, 5: 1-12 <https://doi.org/10.1016/j.hazards.2003.04.001>
- 665 McWethy, D.B., Pauchard, A., García, R.A., Holz, A., González, M.E., Veblen, T.T. 2018. Landscape drivers of recent fire activity (2001-2017) in south-central Chile. *PLoS ONE* 13(8): e0201195.
<https://doi.org/10.1371/journal.pone.0201195>
- Minnich, R.A. and Chou, Y.H. 1997. Wildland Fire Patch Dynamics in the Chaparral of Southern California and Northern Baja California. *International Journal of Wildland Fire* 7, 221-248.
<https://doi.org/10.1071/WF9970221>
- 670 Montgomery, D.R. 2007. Soil erosion and agricultural sustainability. *Proceedings of National Academy of Sciences of USA*, 104 (33): 13268-13272 <https://doi.org/10.1073/pnas.0611508104>
- Moritz, M. A., Parisien, M.-A., Batllori, E., Krawchuk, M. A., Van Dorn, J., Ganz, D. J., Hayhoe, K. 2012. Climate change and disruptions to global fire activity. *Ecosphere*, 3(6):49. <http://dx.doi.org/10.1890/ES11-00345.1>
- 675 Nahuelhual, L., Carmona, A., Lara, A. 2012. Landscape and urban planning land cover change to forest plantations: Proximate causes and implications for the landscape in south-central Chile. *Landscape and Urban Planning*. <http://dx.doi.org/10.1016/j.landurbplan.2012.04.006>.
- Otero, I. and Nielsen, J. 2017. Coexisting with wildfire? Achievements and challenges for a radical social-ecological transformation in Catalonia (Spain). *Geoforum*, 85: 234-246,
<https://doi.org/10.1016/j.geoforum.2017.07.020>
- 680 Pausas, J.G., Llovet, J., Rodrigo, A., Vallejo, R. 2008. Are wildfires a disaster in the Mediterranean basin? – A review. *International Journal of Wildland Fire*, 17: 713-723. <https://doi.org/10.1071/WF07151>
- Paveglio, T., Brenkert-Smith, H., Hall, T.E., Smith, A. 2015. Understanding social impact from wildfires: advancing means for assessment. *International Journal of Wildland Fire*, 24: 212-224.
<https://doi.org/10.1071/WF14091>
- 685 Pawson, S.M., Brin, A., Brockerhoff, E.G., Lamb, D., Payn, T.W., Paquette, A., Parrotta, J.A. 2013. Plantation forests, climate change and biodiversity. *Biodiversity Conservation* 22(5):1203–1227. Doi:10.1007/s10531-013-0458-8



- Pyne, S.J. 2009. The human geography of fire: a research agenda. *Progress in Human Geography*, 33(4): 443 - 446. <https://doi.org/10.1177/0309132508101598>
- 690 Reyes, R. and Nelson, H. 2014. A tale of two forests: why forests and forest conflicts are both growing in Chile. *International Forestry Review*, 16(4):379-388. <http://www.bioone.org/doi/full/10.1505/146554814813484121>
- Ricotta, C. and Di Vito, S. 2014. Modeling the landscape drivers of fire recurrence in Sardinia (Italy). *Environmental Management*, 53(6): 1077-1084. DOI 10.1007/s00267-014-0269-z
- 695 Rojas Quezada, C.A., Muñoz Olivera, I., García-López, M.Á. 2009. Estructura urbana y policentrismo en el Área Metropolitana de Concepción. *Revista Eure*, 35(105):47-70. <http://dx.doi.org/10.4067/S0250-71612009000200003>
- Rojas Quezada, C., Pino, J., Jaque, E. 2013. Strategic Environmental Assessment in Latin America: A methodological proposal for urban planning in the Metropolitan Area of Concepción (Chile). *Land Use Policy*, 30(1): 519-527. <https://doi.org/10.1016/j.landusepol.2012.04.018>
- 700 Ruiz, V., Munizaga, J., Burrows, A.S. 2017. Plantaciones forestales y su extensión hacia áreas urbanas en el área metropolitana de Valparaíso y su relación con el aumento de incendios forestales. *Investigaciones Geográficas*, (54): 23-40. DOI: 10.5354/0719-5370.2017.48040
- Sarricolea, P., Herrera, M.J., Meseguer-Ruiz, O. 2016. Climatic regionalization of continental Chile. *Journal of Maps*. <http://dx.doi.org/10.1080/17445647.2016.1259592>
- 705 Sarricolea, P., Serrano-Notivoli, R., Fuentealba, M., Hernández-Mora, M., de la Barrera, F., Smith, P., Meseguer-Ruiz, O. 2020. Recent wildfires in Central Chile: Detecting links between burned areas and population exposure in the wildland urban interface. *Science of the Total Environment*, 706 (1): 135894 <https://doi.org/10.1016/j.scitotenv.2019.135894>
- Schulz, J.J., Cayuela, L., Echeverría, C., Salas, J., Rey Benayas, J. M. 2010. Monitoring land cover change of the dryland forest landscape of Central Chile (1975–2008). *Applied Geography*, 30: 436–447. www.doi.org/10.1016/j.apgeog.2009.12.003
- Spies, T.A., White, E. M., Kline, J. D., Fischer, A. P., Ager, A., Bailey, J., Bolte, J., Koch, J., Platt, E., Olsen, C. S, Jacobs, D., Shindler, B., Steen-Adams, M. M., Hammer R. 2014. Examining fire-prone forest landscapes as coupled human and natural systems. *Ecology and Society*, <http://dx.doi.org/10.5751/ES-06584-190309>
- 715 Stott, P. (2016). How climate change affects extreme weather events. *Science*, 352(6293): 1517-1518. <https://science.sciencemag.org/content/352/6293/1517>
- Tapia, G. and Castillo, M. 2014. Propuesta de diseño de un sistema de torres de detección de incendios forestales: aplicación a la Región Metropolitana de Chile Central. *Bosque* (Valdivia), 35(3): 399-412. <https://dx.doi.org/10.4067/S0717-92002014000300014>
- 720 Terranova, O., Antronico, L., Coscarelli, R., Iaquina, P. 2009. Soil erosion risk scenarios in the Mediterranean environment using RUSLE and GIS: an application model for Calabria (southern Italy). *Geomorphology*, 112: 228–245 <https://doi.org/10.1016/j.geomorph.2009.06.009>
- Torres, R., Azócar, G., Rojas, J., Montecinos, A., Paredes, P. 2015. Vulnerability and resistance to neoliberal environmental changes: An assessment of agriculture and forestry in the Biobío region of Chile (1974–2014). *Geoforum*, 60: 107–122 <https://doi.org/10.1016/j.geoforum.2014.12.013>
- 725



- Turco, M., Bedia, J., Di Liberto, F., Fiorucci, P., Von Hardenberg, J., Koutsias, N. 2016. Decreasing fires in Mediterranean Europe. *PLoS ONE* 11(3): e0150663. <https://doi.org/10.1371/journal.pone.0150663>
- Úbeda, X. and Sarricolea, P. 2016. Wildfires in Chile: A review. *Global and Planetary Change*, 146: 152–161. <https://doi.org/10.1016/j.gloplacha.2016.10.004>
- 730 Urrutia-Jalabert, R., Gonzalez, M. E., Gonzalez-Reyes, A., Lara, A., Garreaud, R. 2018. Climate variability and forest fires in central and south-central Chile. *Ecosphere* 9(4): e02171. DOI: 10.1002/ecs2.2171. Available at: http://dgf.uchile.cl/rene/PUBS/Climate_Fire_CentralChile-Ecosphere.pdf
- Van Ackere, S., Verbeurgt, J., De Sloover, L., Gautama, S., De Wulf, A., De Maeyer, P. 2019. A Review of the Internet of Floods: Near Real-Time Detection of a flood event and its impact. *Water*, 11(11): 2275. <https://doi.org/10.3390/w11112275>
- 735 Vesanto, J. 2000. Neural network tool for data mining: SOM toolbox. *Proceedings of Symposium on Tool Environments and Development Methods for Intelligent Systems (TOOLMET2000)*. Oulu, Finland. 184–96
- Vilar del Hoyo, L., Martín Isabel, M.P., Martínez Vega, F.J. 2011. Logistic regression models for human-caused wildfire risk estimation: analyzing the effect of the spatial accuracy in fire occurrence data. *European Journal of Forestry Research* <https://doi.org/10.1007/s10342-011-0488-2>
- 740 Wan, Z., Hook, S., Hulley, G. 2015. MOD11C3 MODIS/Terra Land Surface Temperature/Emissivity Monthly L3 Global 0.05Deg CMG V006 [Data set]. *NASA EOSDIS Land Processes DAAC*. Accessed 2020-09-03 from <https://doi.org/10.5067/MODIS/MOD11C3.006>
- Wright, J.A., Di Nicola, A., Gaitan, E. 2000. Latin American forest plantations. Opportunities for carbon sequestration, economic development, and financial returns. *Journal of Forestry*, 12: 20-23. <https://doi.org/10.1093/jof/98.9.20>
- 745 You, W., Li, L., Wu, L., Ji, Z., Yu, J., Zhu, J., Fan, Y., He, D. 2017. Geographical information system-based forest fire risk assessment integrating national forest inventory data and analysis of its spatiotemporal variability. *Ecological Indicators*, 77: 176-184 <https://doi.org/10.1016/j.ecolind.2017.01.042>.