

# Geo-historical database of flood impacts in Alpine catchments (HIFAVa database, Arve River, France, 1850 – 2015).

Eva Boisson<sup>1</sup>, Bruno Wilhelm<sup>1</sup>, Emmanuel Garnier<sup>2</sup>, Alain Mélo<sup>3</sup>, Sandrine Anquetin<sup>1</sup>, Isabelle Ruin<sup>1</sup>.

<sup>1</sup> Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, 38000 Grenoble, France.

<sup>2</sup> Univ. Franche-Comté, CNRS, LCE, 25000 Besançon, France.

<sup>3</sup> AXALP, Annecy, France; associate member at Univ. Savoie Mont-Blanc, CNRS, EDYTEM - UMR 5204, 73370 Le Bourget du Lac, France.

Correspondance to: Eva Boisson ([eva.boisson@univ-grenoble-alpes.fr](mailto:eva.boisson@univ-grenoble-alpes.fr))

## Abstract

In France, flooding is the most common and damaging natural hazard. Global warming is expected to exacerbate flood risk and could be more pronounced in the European Alps which are experiencing a high warming rate, likely to lead to heavier rainfall events. Alpine valleys are densely populated, potentially increasing exposure and vulnerability to flood hazard. The study of historical records is highly relevant to understand long-term flood occurrence and related socio-economic impacts in relation to changes in the flood risk components (i.e. hazard, exposure and vulnerability).

To this aim we introduce the newly constituted database of *Historical Impacts of Floods in the Arve Valley* (HIFAVa) located in French Northern Alps starting in 1850. This quite unique database reports historical impacts related to impact events occurrences in a well-documented Alpine catchment that encompasses both hydrological and socio-economical diversity.

After a complete description of the database (collection, content and structure), we explore the distribution of the recorded impacts with respect to their characteristics and evolution in both time and space. The analysis reveals that small mountain streams and particularly glacial streams caused more impacts (67%) than the main river. While an increase in heavy rainfall and ice melt are expected to enhance flood hazard in small Alpine catchments, this finding calls for greater attention to flood risk assessment and management in small catchments. The analysis also reveals an increasing occurrence of impacts from 1920 onwards, for which possible factors are discussed. Further work is, however, needed to conclude on the respective contribution of the source effect, the increase in flood hazard or the exposure of goods and people.

**Keywords:** flood risk, history, socio-economic impacts, exposure, vulnerability, French Alps.

## 1. Introduction.

On the mainland French territory flood is the most common and damaging natural hazard in terms of economic cost and number of municipalities concerned (Ministère de la Transition écologique, 2020). In highland regions, these events can be caused, among others, by summer thunderstorms, rain on thaw saturated soils, rain on snow or by glacial lake outburst (Merz and Blöschl, 2003). The topography induces flood events with highly contrasted dynamics; from sudden events with large sediment transport in the upstream small catchments to multi-day events flooding large parts of the valley floor. This diversity of hydrological dynamics adds to the complexity in flood risk management. Furthermore, climate change is expected to increase extreme precipitation (Min et al., 2011) that could in turn increase flooding (Gobiet et al., 2014; Blöschl et al., 2020). This is especially the case for the European Alps where an increase in summer heavy rainfalls (Giorgi et al., 2016; Ménégoz et al., 2020) may threaten densely populated mountainous valleys, which are especially vulnerable to climate extremes (IPCC, 2019). With its long history of flooding, the densely populated Arve valley located in the Northern French Alps is indeed prone to experience increased flood risk as a result of global warming in the future.

Historical records constitute a source of reliable data to characterize past hydrological events because they contribute to give a comprehensive representation of these events and of their changes over long time scales in spite of the lack of instrumental data (Garnier and Desarthe, 2013; Barriendos et al., 2014; Wetter, 2017; Macdonald and Sangster, 2017; Wilhelm et al., 2019). They also allow to apprehend changes in flood risk since they document how societies were impacted by past flooding events. Here, we consider impacts accordingly to the IPCC (2012) definition as all types of outcomes for humans, society and ecosystems occurring in the aftermath of a physical phenomenon, i.e., any disturbance, damage, casualties or destruction described in the historical archives and related to a flood event.

The historical analysis of past events is useful for the study of catastrophe as we can hypothesize that these remarkable events are etched in the community's memory (Papagiannaki et al., 2013a). Indeed, it is because these events have impacted society that they are recorded in the historical records, i.e. have left a "social signature". Those high impact events can come close to the notion of a catastrophe, as they can lead to societal upheaval (Soanes and Stevenson, 2009) sometimes deleterious but also beneficial (behavioral change promoting prevention) (Garnier, 2017). High impact events are by nature rare, often resulting in a lack of available data (e.g. description of the event, time, extent, damages caused etc.). However, historical analysis allows a social and spatial-temporal contextualization of the data (Giacona et al., 2017), making the reconstruction (date, impacts) of major flood events possible (Barriendos et al., 2003, 2019) and attesting the social apprehension of the phenomenon (Gil-Guirado et al., 2016).

Numerous historical databases were built to document past flood occurrence and magnitude, such as the Prediflood database (Barriendos et al., 2014), and some, as the database from Thoumas (2019), allow to analyze the climatic fluctuations. In contrast to these latter databases focusing on hydrological events, some databases gather information on the socioeconomic impacts of floods such as the APAT database (Lastoria et al., 2006), the press database on natural hazards and climate change from Llasat et al. (2009), the database of high-impact weather events in Greece from Papagiannaki et al. (2013a), the EUFF database (Petrucci et al., 2019) and the SMC-Flood database (Gil-Guirado et al., 2019). Some databases stand out as the participative flood database ORRION (Giacona et al., 2019), the on-line information resources of the Chronology of British Hydrological Events (Black and Law, 2004) or the database built by the RISC-KIT project (Garnier et al., 2018) which aim at developing methods, management approaches and explore trajectories of vulnerability.

95 To our knowledge there are no similar works published about flood impacts identified in the  
96 archives over historical timescale (i.e. over longer time periods with only the last few decades  
97 covered by instrumental data) and in a mountainous catchment. There are numerous  
98 databases on flood impacts but most of them refer either to the instrumental period (Schlögl et  
99 al., 2021) or to other hazard than flooding (Zgheib et al., 2020; Papagiannaki et al., 2013b;  
100 Giacona et al., 2017) or cover a larger area (Barriendos et al., 2014; Macdonald and Sangster,  
101 2017).

102  
103 Floods, as natural hazards, are physical phenomena naturally occurring and can, when  
104 certain conditions are met, cause harm to societies. They can be interpreted as a social  
105 construction (Beck, 1992) since exposure of human activities and social vulnerability play a  
106 large role in the severity of the impacts. Flood impacts databases, constituted from historical  
107 records, can be considered as the expression of society's concerns, risk perceptions (fear,  
108 habit) and values (based on reported impacts). The recording of flood impacts, or the failure  
109 to record them, provides a subjective measure of the events that were considered worth  
110 reporting for various reasons across historical periods. Flood impacts result from the  
111 interaction between the natural phenomenon and the dynamics of exposure and vulnerability.  
112 As vulnerability we understand the inclination to damage of various exposed goods, activities  
113 or people constituting a given territory (Leone and Vinet, 2006). We consider the vulnerability  
114 as a dynamic system articulated to numerous physical and societal factors (Antoine, 2011;  
115 Terti et al., 2015). This system can evolve in time and space (Cutter, 2003). Major natural  
116 disasters, such as floods, are often displayed as unforeseeable events whereas the historical  
117 facts give evidence of the contrary (Garnier, 2016, 2019). Yet society's vulnerability may  
118 increase as past disasters are forgotten, leading to a "society of risk" (Garnier, 2019). The  
119 historical approach allows consideration and exploration of the trajectories of hazard and  
120 vulnerability in response to changes in climate, land use and flood risk management (Gil-  
121 Guirado et al., 2016).

122  
123 The present paper introduces a newly constituted database of flood impacts of the Arve  
124 River and its tributaries (Northern French Alps). The database called "*Historical Impacts of*  
125 *Floods in the Arve Valley*" (HIFAVa) covers all impacts caused by hydrological events that  
126 occurred since 1850.

127 The study of this database, probably the first one documenting flood impacts over historical  
128 time scale in a mountainous catchment, ultimately aims at analyzing the interactions between  
129 social and natural dynamics engendering these impacts. In this paper we present the dataset  
130 and first results of the impacts analysis with respect to their nature and evolution in both time  
131 and space.

132

133

## 134 **2. Study area: the Arve River.**


135

### 136 **2.1. Description of the physical setting of the Arve River.**

137 The Arve River is located in the Northern French Alps (Figure 1), flowing from the high  
138 elevations of the Mont-Blanc summit (4810m a.s.l.) to the Swiss lowlands (330m a.s.l.), where  
139 it flows into the Rhône River. The surface area of its catchment is 2164 km<sup>2</sup> with the largest  
140 part higher than 1000m a.s.l.. The main tributaries of the Arve River are the Giffre, the Borne,  
141 the Menoge and the Foron Rivers.

142 Since 1850, i.e. the start date of the studied period, almost all the current diking systems were  
143 already in place (Mougin, 1914; Gex, 1924; Peiry and Bravard, 1989; ACTHYS-Diffusion,  
144 2017) and from 1880 onward, most of the dyke construction work was completed and their  
145 nature did not significantly change after this date. As shown by the study of ACTHYS-Diffusion  
146 (2017), most of the 21<sup>st</sup> century's development of the diking systems of the Arve and the Borne

147 Rivers protecting the city of Bonneville concern the construction of weirs to fight against  
 148 generalized stream incision due to the important extraction of materials in the rivers. Some  
 149 repair works were carried out on dikes during the 20<sup>th</sup> century and new works (development of  
 150 weirs, raising of dikes) only happened at the beginning of the 2000s. Unfortunately, this study  
 151 is limited to the area of Bonneville and has not been replicated to the rest of the territory.  
 152



153 Figure 1. The Arve catchment location, topography, main hydrological network and the studied  
 154 cities (sources: IGN, 2017, 2015; SITG, 2020; GADM, 2018).  
 155

156  
 157 Due to large differences in altitude between high and lowlands, the Arve flows can be defined  
 158 by two hydrological regimes following an upstream to downstream continuum:

- 159 - The upstream part of the catchment (down to the city of Sallanches; Figure 1), has a  
 160 glacio-nival regime due to the numerous glacial tributaries flowing from the Mont Blanc  
 161 massif (Viani et al., 2018). Low flows occur in winter and early spring (December to  
 162 March) and the high flows in summer (maximum in July and August) because of the  
 163 strong contribution of ice melt (Bernard, 1900). Floods mainly occur in summer due to  
 164 the synchronicity of both ice melt and intense subdaily rain storms. In this part of the  
 165 catchment, the flood plain is narrow and the slope inclination is high.
- 166 - At lower elevations (i.e. downstream Sallanches) the regime becomes more nival  
 167 downward. Low waters mainly occur in winter and reach the highest levels between  
 168 late spring and early summer with the snowmelt. Between Sallanches and Bonneville,  
 169 floods mainly occur in summer and autumn due to the conjunction of intense rain storm,  
 170 snow melt and, in a lesser extent, ice melt contribution. Downstream Bonneville, floods  
 171 occur at any time of the year due to various hydro-meteorological interplays. The valley  
 172 floor is wide and may be affected by extended flooding.

173  
 174 **2.2. Socioeconomic setting and land use.**

175 There are 106 municipalities located in the Arve catchment, with major population  
176 growth since 1850. For instance, the population has been multiplied by a factor of three in  
177 about 170 years at Chamonix (2 304 to 8 611 inhabitants between 1848 and 2016) and by a  
178 factor of thirty-four in Annemasse (1 047 to 35 712 inhabitants between 1848 and 2017)  
179 (INSEE, 2019). These numbers also hide large seasonal variations related to tourism activities.  
180 This is particularly the case in Chamonix where the number of residents increases up to ten  
181 times in high season. Most inhabitants live in the valley floor and foothills with most of the  
182 farming, industrial and tourism activities as well as the main transportation routes and urban  
183 areas. Since 1965 and the opening of the Mont-Blanc Tunnel Highway the Arve valley is a  
184 major trans-Alpine route connecting France and Italy.

185 The socioeconomic setting of the valley follows an upstream-downstream distribution  
186 pattern. The period from 1850 to 1913 experienced great tourist development (thermal bath of  
187 Saint-Gervais-les-Bains and mountaineering in Chamonix). The economy around Chamonix is  
188 essentially based on mountain tourism. In 1921, 250 000 tourists were visiting Chamonix each  
189 year (Gex, 1924). In 2015 the lodging capacity in the valley reached about 416 400 equivalent  
190 tourist beds. This part of the valley has undergone rapid urbanization. In 1804, the discovery  
191 and exploitation of spring water for hydrotherapy in Saint-Gervais-les-Bains (Gex, 1924)  
192 fostered the construction of tourist accommodations. Around Bonneville, the valley is a densely  
193 populated corridor characterized by an old metal working industry, born from the watch  
194 manufacture and nearby hydropower resources. The smaller valleys of the Arve tributaries are  
195 sparsely populated and the economy is based on tourism and farming. Due to the  
196 attractiveness of the city of Geneva, the valley from Bonneville to the Rhône River confluence  
197 is characterized from the 1960's by a major population growth, extended industrial areas and  
198 strong urbanization. Between the town of Cluses and Geneva the valley floor is almost  
199 continuously built-up.

200

201

### 202 **3. Material and methods of the HIFAVa database.**

203

#### 204 **3.1. Collecting data from historical archives.**

205 Mélo et al. (2015) undertook historic research to develop a flood chronology and hydro-  
206 meteorological circumstances of the flooding events which happened in the Arve catchment  
207 between the 18<sup>th</sup> and the 21<sup>th</sup> century. As sources are more abundant and richer in information  
208 over the last 165 years (1850-2015), this period was defined as the studied time frame of the  
209 HIFAVa database. Only events that triggered impacts and that were mentioned in at least two  
210 sources were integrated in the database. Since 2015, data have been further collected to  
211 complete the preliminary dataset from Mélo et al. (2015).

212 Information on flood related impacts were collected from various sources. Primary sources  
213 range from handwritten archives like municipal acts to departmental archives (e.g. reports of  
214 the *Préfecture* and of town councils). Secondary and tertiary sources are respectively made of  
215 published documents (newspapers, reports, books) and pre-existing databases. The database  
216 of historical records providing a chronological and synthetic layout of the data is composed of  
217 (Figure 2 and Table A1):

#### 218 Manuscript materials:

- 219 - Most of the manuscripts are kept in the departemental archives (*Archives*  
220 *Départementales de Haute-Savoie*: ADHS) (Conseil départemental de la Haute-  
221 Savoie, n.d.) or in the municipal archives (*Archives municipales* : AM) of the towns of  
222 Chamonix, Cluses and Bonneville. The departmental and municipal archives also  
223 contain records older than the Savoy annexation by France in 1860. The records can  
224 be private or from a public institution.

#### 225 Printed materials:

- 226 - Articles from scientific journals and books used in this study have been published since
- 227 1914. They mostly correspond to analyses of the regional hydrology (Mougin, 1914;
- 228 Rousset-Mestrallet, 1986) but also focus on single hydrological events (Pardé 1931;
- 229 Rougier 2002; Goy 2002) and risk assessment (Douvinet et al., 2011).
- 230 - Open-access online municipal risk prevention plans (*Plan de Prévention des Risques* :
- 231 PPR ; and *Plan de Prévention du Risque Inondation* : PPRI) (Préfecture de la Haute-
- 232 Savoie, n.d.) as preventive regulatory documents used to delineate risk areas, often
- 233 compile historical flood events that affected municipalities.

#### 234 Newspapers:

- 235 - Most of the newspapers used are regional, but one is published at national level (*Le*
- 236 *Figaro*) and another is printed in Geneva, Switzerland (*Le Journal de Genève*). Most of
- 237 the newspapers can be found online or are kept in the departmental or municipal
- 238 archives. Newspapers describe the causes and consequences of the flood events and
- 239 sometimes provide instrumental data and illustrations. They also contain information
- 240 concerning the public authority response and past discussions.

#### 241 Other records:

- 242 - The national database of historical flooding (*Base de Données Historiques sur les*
- 243 *Inondations* : BDHI) gathering floods events considered as “remarkable” in the French
- 244 territory (Ministère de la Transition Ecologique, n.d.; Boudou, 2015) was also used.
- 245 - The database created by the department of Restoration of Mountainous Areas
- 246 (*Restauration des Terrains en Montagne* : RTM) from the public institution managing
- 247 the French public forests (*Office National des Forêts* : ONF). The ONF-RTM database
- 248 gathers transcriptions of observations of the RTM officials as well as information
- 249 collected from diverse primary sources. These data (labelled RTM in the HIFAVa
- 250 database) are freely available through the open-access and online ONF-RTM database
- 251 (RTM and ONF, 2012). Some specific ONF-RTM reports are also included in the
- 252 HIFAVa. The RTM database was built to assist the management of small tributaries.
- 253 - A movie realized in 1990 by the RTM is also mentioned as a source.
- 254 - Some records are from the *Syndicat Mixte d'Aménagement de l'Arve et de ses*
- 255 *Affluents* (SM3A), which is the institution in charge of the management of the Arve River
- 256 and its tributaries since 1994.

257

### 258 **3.2. Characteristics of the HIFAVa database.**

259 The database has been built using the free and open-source relational database

260 management system PostgreSQL and is accessible through its package pgAdmin.

261 HIFAVa contains 916 distinct flood impacts caused by 321 flood events. The primary key is

262 the impact ID. Therefore, each impact is recorded as a unique line and described through

263 various variables (Table 1 and Table A2). The river that triggered the flood is mentioned when

264 possible (94% of cases). For instance, no river name has been attributed to the impacts related

265 to overland flow in January 1979. The accuracy of the impact location varies from specific

266 addresses (house, bridge, neighborhood) to the municipality scale. When the source is not

267 accurate enough to distinguish distinct locations of several impacts, they are all referenced

268 under a unique impact ID. In other words, sometimes numerous impacts caused by a single

269 flood event are registered under distinct ID because it was possible to localize each impact

270 precisely (at the hamlet scale). Sometimes we can only localize the impacts at the municipality

271 scale, meaning that all impacts are registered under the same ID. The severity of an event can

272 not be estimated by the number of ID registered in the database. The most recent sources are

273 often highly informative, allowing impacts to be more precisely located.

274 Impacts occurring on the same day on a given river are expected to be caused by the

275 same flood event. As a result, the date is the key used to connect each impact to a flood event.

276 This “flood event” definition has been extrapolated to impacts occurring on the same day in  
 277 different catchments, assuming that two impacts occurring the same day can be caused by the  
 278 same hydrometeorological event given the moderate surface area of the Arve catchment. The  
 279 accuracy of the date is rated on a certainty scale (hour, day, month, year). Based on  
 280 information contained in the records, we distinguish when possible the hydrometeorological  
 281 events (e.g. rainfall, intense and short rainfall, melting of snow, frozen soil, glacial outburst,  
 282 wet period before the event) which caused the flood and the different flood types (e.g. river  
 283 flooding, overland flow, sediment transport) leading to the impacts.  
 284

### 285 3.3. Text mining.

286 The flood impacts of the HIFAVa were categorized through a text quantitative content  
 287 analysis with the KH Coder software (Higuchi, 2015). The description of the impacts comes  
 288 from comments contained in the records. The most frequent words were gathered in order to  
 289 determine representative categories of the database content. A category is made of several  
 290 words assigned to a coding rule. Categories were inspired by the flash flood impact severity  
 291 scale of Diakakis (2020). This analysis led to the following seven categories with example of  
 292 the assigned words:

- 293 - Transport network: e.g. “road”, “bridge”, “railway”, “street”.
- 294 - Urbanized areas and residential buildings: e.g. “house”, “town”, “basement”.
- 295 - Natural environment: e.g. “forest”, “field”, “yield”, “sediment transport”.
- 296 - Protection infrastructures and dams: “dikes” and “dam”.
- 297 - Industrial, commercial and recreational facilities: e.g. “mill”, “factories”, “golf”,  
 298 “camping”, “hotel”, “school”.
- 299 - Critical installations: “drain”, “power transformer”.
- 300 - Victims: “dead”, “injured”, “missing”, “evacuee”.
- 301

ID	Event	Sources	Start_day	Start_month	Start_year
140	58	Payot 1951 / Goy 2002 / RTM / BDHI	12	7	1892
670	263	RTM / Dauphiné libéré : 26/07/1996, 27/07/1996, 30/07/1996 and 02/08/1996	24	7	1996

Start_date	Time_unc	Hour	River	Impacts	Municipality
12/07/1892	D		Bon Nant	Thermal Bath, victims	Saint-Gervais-les-Bains
24/07/1996	H	11 pm	Arve	City center	Chamonix-Mont-Blanc

Impact_latitude	Impact_longitude	Space_unc	Count	Hydrometeo_descript	Water_level
45.8965	6.70596	A	175	No entry	
45.9257	6.87057	A	0	No entry	0,4m

Flood_description	Impacts_decription
Debris flow. 30 minutes to flow from the Tête Rousse glacier to the Arve confluence.	More than 175 victims, destruction of most of Thermal Bath, mud over the first floor.
Debris flow, logjam under bridges.	Numerous houses and the city center are under water. Fuel oil pollution due to basement flooding.

302 Table 1. Extract of the HIFAVa database showing its structure and examples of content. Refer  
 303 to the Online Resource 2 for details about the column contents.  
 304


## 305 306 307 4. Results and discussion.

308 In this section, we present the detailed content of the database, the results of the  
 309 preliminary analyses and the first key findings.  
 310

311  
 312  
 313  
 314  
 315  
 316  
 317  
 318  
 319

#### 4.1. Evolving sources over time.

During the studied period, the diversity and the quantity of sources in which mentions of impacts were found fluctuate (Figure 2). Among the existing databases used, the BDHI database for instance continuously covers the studied period but was sporadically informative since it only contains two mentions of impact events in the Arve catchment, respectively of the Bon Nant River in 1892 and of the Borne River in 1987 (Figure 2.a). By contrast, the SM3A database appears later (1979) in the studied time frame.



320  
 321  
 322  
 323  
 324  
 325  
 326  
 327  
 328  
 329  
 330  
 331

Figure 2. a) Number and b) percentage of mentions from the studied sources to document flood impacts since 1850 in the Arve Valley. In panel a, the ticks indicate each mentions of impacts and the colored areas display the cumulative number of mentions.

RTM (38% of the mentions from all sources), regional newspapers (20%) and scientific literature (13%) constitute the main sources of information on past flood impacts throughout the studied time period. Sources from the literature are particularly useful (54%) to document the period between 1850 and 1910. Then, materials from the departmental and municipality archives are abundant between 1900 and 1970, especially those from Haute-Savoie (20% of the total registered sources for this period) and Chamonix (7%). Between 1940 to 2015, the RTM represents 58% of the mentions from all sources describing the impacts (Figure 2.b).



332 One of the evolutions of the sources is the increase in newspaper articles mentioning flood  
333 impacts. Following the 1881 press freedom French law, the 1880-1889 decade marks the  
334 emergence of articles recording natural hazards, such as flood impacts (Ferenczi, 1996).  
335 However, the 1855 flood in Bonneville was already reported by the swiss newspaper, *Le*  
336 *Journal de Genève*.

337 Although few sources (e.g. the municipal archives of Sallanches) remain to be examined, we  
338 consider that most of the main sources (newspapers, existing databases, and public archives)  
339 have been analyzed. As a result, we are confident that no event that was deemed damageable  
340 by local communities was missed and we consider that we have a comprehensive view of past  
341 flood impacts since 1850 over the whole Arve catchment.


342

#### 343 **4.2. Changes in impacts over time and space.**

344 From 1850 to 1920, the number of impacts fluctuates and only four years are  
345 considered remarkable with more than 15 impacts (1852, 1878, 1895 and 1910). From 1920,  
346 years with 15 or more impacts become more frequent (1930, 1940, 1944, 1968, 1979, 1987,  
347 1990, 1996, 1997 and 2007) and the total amount of impacts per year reaches the maximum  
348 value of 54 in 1996 (Figure 3). The decennial moving average of the impacts' number highlights  
349 an overall increase over the 165 years, punctuated by periods with less frequent impacts (in  
350 1910-1923, 1950-1960 and 1975-1980).

351 The number of recorded impact events – i.e. flood events recorded in the historical sources  
352 because of the impacts they caused – stays relatively stable between 1.5 and 3 events per  
353 year on average until 1990, then it rises up to 4.5 events per year. As hydrological data are  
354 available from 1950 for the Arve River, it is not possible to assess whether the increase in  
355 impact event was a function of increased flood occurrence or changes in vulnerability or  
356 recording.

357




358

359 Figure 3. Yearly occurrence of impacts and decennial moving averages of impacts (red  
360 curve) and associated impact events (blue curve). Events discussed in the article are labeled  
361 with their year of occurrence.

362

363 When analyzing the spatial distribution of the flood impacts, we can see that they are  
364 spread over the entire catchment (Figure 4a). Chamonix and Bonneville gather, however,  
365 respectively 24% and 12,5% of total impacts recorded in the Arve catchment. These high  
366 numbers may be due to the fact that these towns are both among the most densely populated  
367 and the closest to the Arve River. The impacts caused by the Arve River floods represent 33%  
368 of all recorded impacts, and its two main tributaries, the Giffre and the Borne Rivers, have only

369 caused 8% of the recorded impacts (Figure 4b). In fact, most impacts are due to small torrential  
 370 streams (53%). Among them, almost a third are related to glacial tributaries, while these  
 371 tributaries are localized only in the uppermost part of the catchment near Chamonix. For  
 372 instance, small torrential tributaries such as the Arveyron, the Grépon (left bank tributary close  
 373 to Chamonix) or the Bon Nant caused alone more impacts than the Borne River itself.  
 374



375 Figure 4. a) Location and b) distribution of flood impacts caused by the Arve River and its  
 376 tributaries. The category "non-attributed" corresponds to the impacts for which it was not  
 377 possible to attribute a river, either because events are related to overland flows or because the  
 378 source did not mention the river (sources: IGN, 2017, 2015; SITG, 2020; GADM, 2018;  
 379 HIFAVa).  
 380

381  
 382 The Arve tributaries produced disasters characterized by numerous and major flood damage.  
 383 Among them, the 1987 Borne River flooding in its uppermost part washed away the municipal  
 384 campsite of the village of the Grand-Bornand causing 23 casualties and heavy economic  
 385 losses (Meunier, 1990). In addition, the 1892 glacial lake outburst from the Tête Rousse glacier  
 386 in the Bon Nant River (which literally translated means "Good Stream") swept away the thermal  
 387 bath of Saint-Gervais-les-Bains (Figure A1) and 33 houses causing at least 175 casualties.  
 388 The glacier was drained in 2010 and is today closely monitored to avoid such a brutal and  
 389 disastrous natural event (Garambois et al., 2016; Vincent et al., 2012).

390 All these high impacts events are due to sudden, highly-dynamic summer floods of tributaries,  
 391 often aggravated by large sediment transport. Some towns located along the Arve River – such  
 392 as Sallanches – are more prone to tributaries floods because embankments have been built  
 393 and efficiently prevent impacts from the Arve. In contrast, there are very few impacts recorded  
 394 in high altitude locations, probably due to the sparsity of human settlements.  
 395


### 396 4.3. Potential drivers of changes in the number and distribution of 397 impacts.

398 The increase in the number of impacts starting in the 1920's and well-marked from the  
 399 1960's can be explained by multiple factors, such as an increase in flood occurrence and/or  
 400 magnitude, a source effect, an increase in exposure of goods and people, a deterioration of  
 401 the diking systems, a break in the risk memory transmission, an evolution of the risk perception  
 402 or an evolution of the local political risk management. Due to the lack of available data  
 403 regarding changes in flood hazard, protection infrastructures and the risk memory, perception

404 and management, we only explore whether source effect and changes in exposure may  
405 explain the observed increase in impacts.

406 To decipher the potential source effect in the increase in impacts particularly noticeable  
407 since 1960, maps of the impacts by sources have been drawn for the periods before and after  
408 1960 (Figure 5). In addition to the noticeable increase in impacts, this date marks a strong  
409 change in the Arve valley economy, from a rather homogeneous agricultural society to an  
410 industrial society exploiting the river bed for materials extractions. After 1960, the Arve  
411 watershed also experienced a strong tourist development and a rapid demographic expansion.  
412 From 1850 to 1959, three main sources describe 64% of the impacts (literary records 28%,  
413 RTM 18% and departmental and municipal archives 18%) and for 29% the information comes  
414 from more than one source. The impacts are mainly gathered along the Arve and the Giffre  
415 Rivers, especially in the valley of Chamonix and between the towns of Cluses and Bonneville.  
416 For the second period (1960-2015), the RTM reports 65% of the impacts, and 20% come from  
417 multiple sources while departmental and municipal archives and the PPR/PPRI describe 5%  
418 each. Information coming only from literature decreases substantially (122 described impacts  
419 in the first period to 3 impacts in the second), SM3A records start in 1979 and only document  
420 the Giffre and the Bon Nant Rivers. The distribution of the impacts is much more scattered  
421 across the whole catchment than during the first period (Figure 5). The impacts are not  
422 gathered along the Arve River, since most of them result from small tributaries. Impacts  
423 described by more than one source are located in the valley of Chamonix and around  
424 Bonneville, probably because these economic and tourist centers arouse interest of many  
425 sources (newspapers, departmental and municipal archives and RTM). In addition, the strong  
426 emergence of the RTM since 1940 (Figure 2) can explain the rise in documented impacts  
427 caused by small tributaries (Figure 5). Following the floods of the Rhône, the Loire and the  
428 Garonne in 1854, the 1858 law against urban flooding places flood control at the heart of the  
429 national legislation for the first time. In the following, the RTM department was formed for the  
430 reforestation of mountains slope in order to prevent the reproduction of major floods. The  
431 department became quickly efficient and since 1860 collected numerous reports. Built for the  
432 study and management of small tributaries, the RTM database became the main source of  
433 information since 1930 for the HIFAVa database. Hence, the strong emergence of the RTM  
434 source among the others may play a role in the observed increase in impacts since 1920, even  
435 more noticeable since 1960.

436




437

438 Figure 5. Comparison of the spatial distribution of impacts by categories of sources in the Arve  
439 catchment during the periods (a) 1850-1959 and (b) 1960-2015 (sources: IGN, 2017, 2015;  
440 SITG, 2020; GADM, 2018; HIFAVa).

441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477

The rise in the number of impacts per flood may be partly explained by the fact that distinct impact types in the same location were reported and therefore referenced under distinct impact ID, while they were not differentiated in previous periods. Recent sources seem to provide more accurate information on the impacts and their locations. In older sources, impacts are most of the time documented at the city scale (21% of the impacts for the first period, and 10% for the second period). Thereby, all these impacts are stored in the database in a single line with an uncertainty code for the impact location corresponding to the municipal level. In most recent sources, impacts' locations are described more accurately allowing them to record at a resolution up to the building scale. As a result, impacts are stored in as many lines as impacts locations can be identified, with an uncertainty code for the impact's location corresponding to the building or neighborhood level (85% of the impacts for the second period). For example, in 1996 fifty-three impacts were recorded for the same event and fifty of them were located in distinct places in Chamonix. The rise in impacts can also be due to numerous impacts in different locations, as the flood of 1990 which impacted six towns in two different sub-catchments (the Arve and the Giffre catchments). However, in order to overcome the bias induced by the recording of impacts according to their location, we aggregated the impacts at the municipality level. That is to say, all the impacts reported for a given municipality that were caused by the same event (thus the same day) are recorded under the same line in the database. This results in 562 "aggregated" impacts instead of 916 impacts initially. From these data, we redrawn Figure 3 (Figure A2) comparing the moving average of impacts and associated events. We can see that in both figures the trends of increasing impacts are similar. There is an increase of impacts (here starting soon as 1890s). Thus, the way the impacts are stored in the database (by location or by municipality) affect the absolute values of impact per year but not the observed temporal changes over time.

Changes in exposure and vulnerability related to land use is another potential explanation of the rise in the number of impacts (Magnan et al., 2012; Garnier and Desarthe, 2013; Camuffo et al., 2020). In fact, as major population growth happened, especially in Bonneville and Chamonix, and lead to a strong and fast urban sprawl in the flood plain between the 1950's and the 2010's as shown by aerial photographs (Figure 6). They also show the vanishing of the alluvial forest (Dufour and Piégay, 2006) and cultivated fields to the benefit of urbanization in both towns. Upstream, in Chamonix, the demographic expansion dates back to the early 20<sup>th</sup> century with the flourish of mountain tourism. In downstream towns – e.g. Bonneville and Annemasse – the expansion starts in the 1950's because of the economic attractiveness of Geneva.



478  
 479 Figure 6. Aerial photographs highlighting changes in land use and urban sprawl growth in  
 480 Chamonix and Bonneville as well as plots stressing changes in impacts and population growth  
 481 from 1848 to 2011 (IGN, INSEE, HIFAVa).  
 482


483 Besides these numbers, the urban expansion and the growth in tourism entail the arrival of  
 484 new residents in the valley (INSEE, 2020), unaware of the local hazard history. The valley  
 485 narrowness, the demand for land and the loss of memory of past events have led to a rise of  
 486 constructions within historical flood-prone areas, resulting in an increased exposure. For  
 487 instance, in 1944 recently built houses in Chamonix were washed away by the Grépon River.  
 488 The same situation also happened during the 1968 flood that destroyed a new residential area  
 489 in Bonneville. To explore to what extent the urban growth may have resulted in an increased  
 490 exposure explaining the observed rise of impacts, we use the population growth data as a first-  
 491 order proxy of urban growth. In Chamonix, the number of impacts increase from year to year,  
 492 somehow mirroring the population growth (Figure 6). Therefore, population exposure might be  
 493 one explanation for the increased number of impacts. One can, however, notice the decrease  
 494 in impacts in the early 2000s due to the heightening of the dikes after the 1996 flood. In  
 495 Bonneville, the link between the number of impacts and the population growth is not as clear  
 496 as in Chamonix. This may partly result from the absence of major floods since 1968. Therefore,  
 497 an increasing exposure might locally explain the increase in impacts. Further studies should  
 498 however reconstruct diachronic maps of land use to assess in a finer way the urban growth in  
 499 flood-prone areas and its link with changes in impacts number. Overall, the potential role of an  
 500 increased exposure is not excluding the indirect sources affect (emergence and dominance of  
 501 the RTM source), but both factors can be combined and complement each other.  
 502

#### 503 **4.4. Nature of the flood impacts.**

504 The quantitative analysis of text content reveals the distribution of the impact categories  
 505 by river and illustrates the diversity of the catchment in terms of land use and economic  
 506 development (Figure 7). This analysis of text content is particularly relevant because it allows  
 507 to overcome the database scarcity of quantitative information on the severity of the flood.  
 508 Indeed, it is difficult to estimate the severity of a flood event as the flow rate and water height  
 509 are only mentioned in rare cases. However, according to Barriendos et al. (2019) a link can be  
 510 made between the nature of the impacts and the severity of the triggering phenomena. In order  
 511 to establish this link, we also need to consider the various dynamics of the flooding phenomena  
 512 as fast flooding generally affect smaller surfaces but in a more violent manner than slow floods

513 of the main rivers. Therefore, they may trigger a lesser number of impacts but the level of  
 514 destruction of the impacted element might be more important. Of course, the level of  
 515 destruction also depends on the sensitivity and physical vulnerability of the exposed element,  
 516 it is why categorizing impacts by their nature might help exploring the question of the flood  
 517 severity.

518  
 519 The categories used for the analysis are partly inspired from a recent paper from Diakakis  
 520 (2020) and refined based on textual analysis of the words used to describe the impacts in the  
 521 database. The number of occurrences of the words determining the category is here named  
 522 as the number of mentions. For the overall Arve catchment, damage to the transport network  
 523 is the most frequent (406 mentions), followed by damage to urbanized areas (286) and natural  
 524 environment (253). Impacts on transport network are in proportion equally distributed among  
 525 river types (all around 30%). Impacts on industrial facilities are mainly caused by the Arve and  
 526 the Giffre Rivers as their wider valleys allow the installation of these facilities. The Giffre and  
 527 the Borne Rivers have caused the least impacts specifically on urban areas, probably because  
 528 of less dense population and of an economy based on farming and tourism activities.  
 529




530  
 531 **Figure 7. Distribution of flood impacts categories according to the river types. The class “non**  
 532 **attributed” gathers the impact that could not be assigned to a river (e.g. overland flows).**  
 533

534 Most of the mentions of victims refer to impacts caused by tributaries (20 out of 28)  
 535 characterized by faster hydrological responses. They have caused 80% of the deadly impacts  
 536 registered since 1850 in the whole catchment, e.g. the 1892 glacial lake outburst and the 1987  
 537 flood of the Borne River in Grand-Bornand. The mentions of victims of the Borne River should  
 538 belong to the small tributaries class as the impacts occurred in the uppermost part of the  
 539 catchment. As shown by Jonkman et al. (2005) and Ruin et al. (2008) high mortality rates are  
 540 mostly due to the suddenness and violent responses of small catchments affecting people in  
 541 the open air (as for campers).

542  
 543 To track potential changes in the nature of impacts since 1850, they are scrutinized by  
 544 decades over the last 165 years (Figure 8). Impacts on transportation networks are present in  
 545 every decade since 1850 but they increase after 1930 as well as impacts on urbanized area  
 546 and natural environment categories. It was not until 1960 that impacts on the industrial,  
 547 commercial and recreational facilities category increased. Mentions about critical installation  
 548 (sewers and water pipes) are recorded for the first time in the 1960's. Mentions of victims are  
 549 present in almost every decade.  
 550 Nevertheless, no major evolution of the impacts' categories can be seen, except the  
 551 emergence of mentions of critical installations in the 1930's. A more in-depth analysis will be

552 conducted later on to define severity classes based on the nature of the impacts and also to  
 553 identify whether there is an evolution of the lexicon used to describe the impacts.  
 554  
 555



556  
 557 **Figure 8. Decennial histogram of the evolution of the categories of impacts and the number of**  
 558 **events.**  
 559

560 The data representation of Figure 8 does not allow to visualize the evolution of the absolute  
 561 values. For instance, at the catchment scale there is an increase in the number of mentions of  
 562 the victim category with 18 mentions during the period 1960-2015 compared to 10 for the  
 563 period 1850-1959 (Figure 8). This increase is not easily noticeable when looking at proportions  
 564 of impact categories because of the strong augmentation in the total number of mentions since  
 565 1930. Yet, apart from the disaster in the Grand-Bornand in 1987 (Borne River), since 1931 all  
 566 the mentions of victims refer to evacuations, recues or injuries.

567 The number of impacts has been almost multiplied by two since 1920. Mentions of impacts on  
 568 urbanized areas during 1960-2015 has been multiplied almost by four compared to the 1850-  
 569 1930 period. This agrees with the evolution of the land use due to the demographic growth,  
 570 i.e. the observed vanishing of forest areas and cultivated land to the benefit of urbanization  
 571 (Figure 6). The mentions of impacts to natural environment for the period 1850-1930 are more  
 572 than double compared to the period 1960-2015. During the first period, mentions of impacts to  
 573 natural environment refer mainly to forest, field and crops, while after 1960 there is no mention  
 574 of field or crops and most of the mentions are about gullying, deposition of sediments and  
 575 banks.

576  
 577

## 578 **5. Conclusions.**

579

580 This paper describes and analyses historical data documenting 916 flood impacts  
 581 associated to 327 flood events that occurred between 1850 and 2015 in the Arve valley, an  
 582 Alpine catchment characterized by a high hydrological and socioeconomic diversity. This  
 583 dataset is stored in the HIFAVa database fully described in this paper.

584

585 A first spatial and temporal analysis of this dataset allowed to identify three key findings  
 586 and research perspectives:

- 587 - The predominance of impacts due to torrential tributaries. There are two main types of  
 588 flood events causing impacts in the Arve catchment, e.g. floods related to the main  
 589 river and those related to the smaller mountain streams. Floods from these small  
 590 streams are characterized by sudden and fast hydrological responses and most of the  
 591 time by high volumes of sediment transport, making hazard management difficult. They  
 592 caused two third of the 916 recorded impacts with numerous casualties, such as the  
 593 Bon Nant (1892) and the Borne (1987) Rivers floods. In contrast, slow rising, day-long  
 594 floods of the Arve River affect larger areas and trigger large economic consequences

595 but no casualties. The 1968 flood event affecting a large part of the Arve catchment is  
596 an exemplary case of this flood type. As suggested by Ruin et al. (2008), the number  
597 of impacts caused by torrential streams being much higher than those triggered by the  
598 main rivers calls for a greater attention to flood risk assessment and management in  
599 small catchments.

600 - The rise in the number of impacts starting in 1920 and well-marked from 1960. This  
601 increase in impacts may be explained by various factors. Based on the available data,  
602 we discussed the potential source effect and changes in exposure. It appeared that the  
603 emergence and dominance of the RTM among the other sources as well as an  
604 increased exposure linked to urban expansion may play a role at some places.  
605 However, exposure was assessed through the population growth as a first-order proxy  
606 of urban expansion. A more detailed study of changes in land use based on e.g. old  
607 maps and aerial photography is necessary to confirm this preliminary result.

608 - The evolution of the impacts' nature (increase of impacts on urbanized area) mirroring  
609 the land use changes and probably due to the urban expansion linked to the large  
610 demographic growth in the catchment area. Further work is required to explore the  
611 evolution of the vocabulary used to describe those impacts across centuries in order to  
612 evaluate how this evolution might relate to changes in what the societies values and  
613 care for across history. Another path of future research concerns the identification of  
614 severity classes based on the description of the nature and extent of the damage in  
615 order to be able to characterize the level of impact on a given territory, to allow a  
616 classification of past events according to their intensity and to identify the most  
617 significant ones.

618 Moreover, the other drivers that may have induce the observe increase in impacts still need to  
619 be investigated. The lack of gauge data (available only for the main river and since 1950)  
620 precludes the study of potential links between the increased number of impacts and changes  
621 in flood occurrence and/or magnitude over the whole catchment. In contrast, data on the risk  
622 memory and its transmission within the society can be acquired and analyzed to explore the  
623 evolution of the territory's vulnerability through time.

624  
625  
626



627 **Appendixes**

628 Table A1. List of the historical records – or their origin – used to provide information about flood events  
 629 and impacts collected in the HIFAVa database. The date mentions the year of publication for books  
 630 and period covered by the newspapers.

<b>Manuscript materials</b>	<b>Printed materials</b>	<b>Newspapers</b>	<b>Other records</b>
ADHS	AM Cluses	ADHS	BDHI
AM Bonneville	Conard 1931	Le Messenger (1965-1968)	RTM (movie 1990)
AM Chamonix	Douvinet 2011	Le Dauphiné libéré (1963-2014)	SM3A
AM Cluses	Goy 2002	Journal de Genève (1855-1978)	
	Mestrallet 1986	La Croix 74 (1898)	
	Mougin 1914	L'Allobroge (1903-1940)	
	Parde 1931	Le Faucigny (1968)	
	Payot 1951	Le Figaro (1924)	
	PPR	Le Messenger (1940)	
	PPRI	Le Mont-Blanc républicain (1903)	
	Rannaud 1916	Le Progrès (1898)	
	Rougier	L'Industriel Savoisien (1910)	
	RTM	La Revue Savoisienne (1887-1889)	

631

632 Table A2. Presentation of the HIFAVa database showing its structure with each data entry.


id	[PK] serial	Primary key. ID number of an impact.
event	integer	Number of the event that triggers the impact. In case of an hydro-meteorological event, several impacts located at different places on different river are associated to the same event.
sources	integer	The different sources that provide information.
start_day	integer	Start day of event.
start_mont	integer	Start month of event.
start_year	integer	Start year of event.
start_date	date	Start date of event.
hour	text	Start hour of event.
time_unc	text	Uncertainty of the start date. H means that the start date is known at the hour scale, D at the day scale, M at the month scale and Y at the year scale. By default, when the day and/or the month is not known, "1" is attributed to start_day and/or start_month.
duration	real	Duration of event
river	text	River that trigger the flood (the cell may be empty if the impacts are not related to river flooding).
impact	text	Nature of impact.
municipality	text	Municipality where the impact is located.
imp_lat	real	Latitude of the impact in decimal degrees.
imp_long	real	Longitude of the impact in decimal degrees.
Space_unc		Describe the spatial scale of impact location. A means that the impact is located at the scale of a point on a map, B at the scale of a part of a city, C at the scale of a city, D at a coarser scale than the one of a city.
count	integer	Number of victims
hydrometeo_descript	text	Description of the hydrometeorological event according to the sources.
precipitation	real	Precipitation rate given in the sources (mm).
flood_descript	text	Description of the flood from the sources
river_water_level	real	Water level of the river (m).
water_level	real	Water level of flooded area (m).
discharge	real	Discharge of the river (m <sup>3</sup> /s)
impact_descript	text	Description of the impacts according to the sources.

633

634 Figure A1. The wiped-out thermal bath of Saint-Gervais-les-Bains after the debris flow of the 12th of  
 635 July 1892 (Thermal bath establishment of Saint-Gervais, via [www.thermes-saint-gervais.com](http://www.thermes-saint-gervais.com)).



636  
 637  
 638  
 639 Figure A2. Yearly occurrence of impacted municipalities and decennial moving averages of impacted  
 640 municipalities (orange curve) and associated impact events.



641  
 642  
 643 Acknowledgments: We would like to thank the *Syndicat Mixte d'Aménagement de l'Arve et de*  
 644 *ses Affluents* who shared its database of events and impacts. The HIFAVa database was partly  
 645 fed thanks to the funding of the projects Labex ITEM CrHistAI and OSUG2020 RHAAP. This  
 646 study was developed in the framework of *CDP-Trajectories* of the Grenoble-Alpes University  
 647 (France) funded by the *Agence Nationale de la Recherche* through the project  
 648 "*Investissements d'avenir*" (ANR-15-IDEX-02).

649 **Bibliography:**

650

651 ACTHYS-Diffusion: Etude pour la restauration des systèmes d'endiguement de l'Arve et du Borne.  
652 Fiches d'information historique (FIH) par système d'endiguement., 2017.

653 Antoine, J.-M.: Vulnérabilité et adaptation des sociétés montagnardes à la torrencialité au cours du  
654 Petit Âge Glaciaire dans les Pyrénées, *soe*, 53–66, <https://doi.org/10.4000/soe.685>, 2011.

655 Barriendos, M., Coeur, D., Lang, M., Llasat, M. C., Naullet, R., Lemaitre, F., and Barrera, A.: Stationarity  
656 analysis of historical flood series in France and Spain (14th–20th centuries), *Nat. Hazards Earth Syst.*  
657 *Sci.*, 3, 583–592, <https://doi.org/10.5194/nhess-3-583-2003>, 2003.

658 Barriendos, M., Ruiz-Bellet, J. L., Tuset, J., Mazón, J., Balasch, J. C., Pino, D., and Ayala, J. L.: The  
659 “Prediflood” database of historical floods in Catalonia (NE Iberian Peninsula) AD 1035-2013, and its  
660 potential applications in flood analysis, *Hydrol. Earth Syst. Sci.*, 18, 4807–4823,  
661 <https://doi.org/10.5194/hess-18-4807-2014>, 2014.

662 Barriendos, M., Gil-Guirado, S., Pino, D., Tuset, J., Pérez-Morales, A., Alberola, A., Costa, J., Balasch, J.  
663 C., Castellort, X., Mazón, J., and Ruiz-Bellet, J. L.: Climatic and social factors behind the Spanish  
664 Mediterranean flood event chronologies from documentary sources (14th–20th centuries), *Global and*  
665 *Planetary Change*, 182, 102997, <https://doi.org/10.1016/j.gloplacha.2019.102997>, 2019.

666 Beck, U.: *Risk Society : Towards a New Modernity*, Sage., London, 1992.

667 Bernard, C.: *Restauration et conservation des terrains en montagne : les terrains et les paysages*  
668 *torrentiels (Haute-Savoie)*, Impr. nationale (Paris), 60 pp., 1900.

669 Black, A. R. and Law, F. M.: Development and utilization of a national web-based chronology of  
670 hydrological events/Développement et utilisation sur internet d'une chronologie nationale  
671 d'événements hydrologiques, *Hydrological Sciences Journal*, 49, 3,  
672 <https://doi.org/10.1623/hysj.49.2.237.34835>, 2004.

673 Blöschl, G., Kiss, A., Viglione, A., Barriendos, M., Böhm, O., Brázdil, R., Coeur, D., Demarée, G., Llasat,  
674 M. C., Macdonald, N., Retsö, D., Roald, L., Schmockler-Fackel, P., Amorim, I., Bělínová, M., Benito, G.,  
675 Bertolin, C., Camuffo, D., Cornel, D., Doktor, R., Elleder, L., Enzi, S., Garcia, J. C., Glaser, R., Hall, J.,  
676 Haslinger, K., Hofstätter, M., Komma, J., Limanówka, D., Lun, D., Panin, A., Parajka, J., Petrić, H.,  
677 Rodrigo, F. S., Rohr, C., Schönbein, J., Schulte, L., Silva, L. P., Toonen, W. H. J., Valent, P., Waser, J., and  
678 Wetter, O.: Current European flood-rich period exceptional compared with past 500 years, *Nature*,  
679 583, 560–566, <https://doi.org/10.1038/s41586-020-2478-3>, 2020.

680 Boudou, M.: *Approche multidisciplinaire pour la caractérisation d'inondations remarquables :*  
681 *enseignements tirés de neuf évènements en France (1910-2010)*, *Geography*, Université Paul Valéry -  
682 Montpellier I, Lyon, 462 pp., 2015.

683 Camuffo, D., della Valle, A., and Becherini, F.: A critical analysis of the definitions of climate and  
684 hydrological extreme events, *Quaternary International*, 538, 5–13,  
685 <https://doi.org/10.1016/j.quaint.2018.10.008>, 2020.

686 Conseil départemental de la Haute-Savoie: *Archives départementales de Haute-Savoie*, n.d.

687 Cutter, S. L.: The Vulnerability of Science and the Science of Vulnerability, *Annals of the Association of*  
688 *American Geographers*, 93, 1–12, <https://doi.org/10.1111/1467-8306.93101>, 2003.

- 689 Diakakis, M., Deligiannakis, G., Antoniadis, Z., Melaki, M., Katsetsiadou, N. K., Andreadakis, E., Spyrou,  
690 N. I., and Gogou, M.: Proposal of a Flash Flood Impact Severity Scale for the classification and mapping  
691 of flash flood impacts., 2020.
- 692 Douvinet, J., Defossez, S., Anselme, A., and Denolle, A.-S.: Les maires face aux plans de prévention du  
693 risque inondation (PPRI), 40, 31, <https://doi.org/10.3917/eg.401.0031>, 2011.
- 694 Dufour, S. and Piégay, H.: Forêts riveraines des cours d'eau et ripisylves : spécificités, fonctions et  
695 gestion, *Rev. For. Fr.*, <https://doi.org/10.4267/2042/6704>, 2006.
- 696 Ferenczi, T.: L'invention du journalisme en France, Naissance de la presse moderne à la fin du XIXème  
697 siècle, Payot., Paris, 274 pp., 1996.
- 698 GADM: GADM, 2018.
- 699 Garambois, S., Legchenko, A., Vincent, C., and Thibert, E.: Ground-penetrating radar and surface  
700 nuclear magnetic resonance monitoring of an englacial water-filled cavity in the polythermal glacier of  
701 Tête Rousse, 81, 2016.
- 702 Garnier, E.: Genève face à la catastrophe 1350-1950. Un retour d'expérience pour une meilleure  
703 résilience urbaine., Slatkine, Genève, Suisse, 195 pp., 2016.
- 704 Garnier, E.: Xynthia, February 2010: Autopsy of a Foreseeable Catastrophe, in: Management of the  
705 Effects of Coastal Storms: Policy, Scientific and Historical Perspectives, Quevauviller, P., 111–148, 2017.
- 706 Garnier, E.: Lessons learned from the past for a better resilience to contemporary risks, 28, 778–794,  
707 2019.
- 708 Garnier, E. and Desarthe, J.: Cyclones and Societies in the Mascarene Islands 17th-20th Centuries,  
709 *AJCC*, 1–13, <https://doi.org/10.4236/ajcc.2013.21001>, 2013.
- 710 Garnier, E., Ciavola, P., Spencer, T., Ferreira, O., Armaroli, C., and Mclvor, A.: Historical analysis of storm  
711 events: Case studies in France, England, Portugal and Italy, 134, 10–23, 2018.
- 712 Gex, F.: La Haute-Savoie aujourd'hui et il y a 100 ans avec un tableau de la population par commune  
713 de 1801 à 1921., Librairie M. Dardel, Chambéry, 1924.
- 714 Giacona, F., Eckert, N., and Martin, B.: A 240-year history of avalanche risk in the Vosges Mountains  
715 based on non-conventional (re)sources, 17, 887–904, <https://doi.org/10.5194/nhess-17-887-2017>,  
716 2017.
- 717 Giacona, F., Martin, B., Furst, B., Glaser, R., Eckert, N., Himmelsbach, I., and Edelblutte, C.: Improving  
718 the understanding of flood risk in the Alsatian region by knowledge capitalization: the ORRION  
719 participative observatory, *Nat. Hazards Earth Syst. Sci.*, 19, 1653–1683, <https://doi.org/10.5194/nhess-19-1653-2019>, 2019.
- 721 Gil-Guirado, S., Espín-Sánchez, J.-A., and Del Rosario Prieto, M.: Can we learn from the past? Four  
722 hundred years of changes in adaptation to floods and droughts. Measuring the vulnerability in two  
723 Hispanic cities, *Climatic Change*, 139, 183–200, <https://doi.org/10.1007/s10584-016-1768-0>, 2016.
- 724 Gil-Guirado, S., Pérez-Morales, A., and Lopez-Martinez, F.: SMC-Flood database: a high-resolution  
725 press database on flood cases for the Spanish Mediterranean coast (1960–2015), *Nat. Hazards Earth  
726 Syst. Sci.*, 19, 1955–1971, <https://doi.org/10.5194/nhess-19-1955-2019>, 2019.

- 727 Giorgi, F., Zsolt Torma, C., Coppola, E., and Ban, N.: Enhanced summer convective rainfall at Alpine high  
728 elevations in response to climate warming, 9, 2016.
- 729 Gobiet, A., Kotlarski, S., Beniston, M., Heinrich, G., Rajczak, J., and Stoffel, M.: 21st century climate  
730 change in the European Alps—A review, *Science of The Total Environment*, 493, 1138–1151,  
731 <https://doi.org/10.1016/j.scitotenv.2013.07.050>, 2014.
- 732 Goy, J.: Autour de la catastrophe de l'établissement thermal de Saint-Gervais-les-Bains en 1892 :  
733 problèmes d'histoire des catastrophes naturelles., *Les pouvoirs publics face aux risques naturels dans*  
734 *l'histoire.*, Grenoble, 39–49, 2002.
- 735 Higuchi, K.: KH Coder, 2015.
- 736 IGN: GEOFLA, 2015.
- 737 IGN: BD Carthage (Base de Données sur la CARTographie THématique des AGences de l'eau et du  
738 ministère chargé de l'environnement), 2017.
- 739 INSEE: <https://www.insee.fr/fr/statistiques/3689656>, last access: 1 August 2020.
- 740 INSEE: Populations légales communales depuis 1968, 2019.
- 741 IPCC: Managing the risks of extreme events and disasters to advance climate change adaptation. A  
742 Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change,  
743 Intergovernmental Panel on Climate Change, 2012.
- 744 IPCC: The Ocean and Cryosphere in a Changing Climate, Intergovernmental Panel on Climate Change,  
745 Genève, Suisse, 2019.
- 746 Jonkman, S. N. and Kelman, I.: An Analysis of the Causes and Circumstances of Flood Disaster Deaths:  
747 An Analysis of the Causes and Circumstances of Flood Disaster Deaths, 29, 75–97,  
748 <https://doi.org/10.1111/j.0361-3666.2005.00275.x>, 2005.
- 749 Lastoria, B., Simonetti, M. R., Casaioli, M., Mariani, S., and Monacelli, G.: Socio-economic impacts of  
750 major floods in Italy from 1951 to 2003, *Adv. Geosci.*, 7, 223–229, [https://doi.org/10.5194/adgeo-7-](https://doi.org/10.5194/adgeo-7-223-2006)  
751 [223-2006](https://doi.org/10.5194/adgeo-7-223-2006), 2006.
- 752 Leone, F. and Vinet, F.: La vulnérabilité, un concept fondamental au cœur des méthodes d'évaluation  
753 des risques naturels, in: *La vulnérabilité des sociétés et des territoires face aux menaces naturelles*,  
754 GESTER, 2006.
- 755 Llasat, M. C., Llasat-Botija, M., and López, L.: A press database on natural risks and its application in  
756 the study of floods in Northeastern Spain, *Nat. Hazards Earth Syst. Sci.*, 9, 2049–2061,  
757 <https://doi.org/10.5194/nhess-9-2049-2009>, 2009.
- 758 Macdonald, N. and Sangster, H.: High-magnitude flooding across Britain since AD 1750, *Hydrol. Earth*  
759 *Syst. Sci.*, 21, 1631–1650, <https://doi.org/10.5194/hess-21-1631-2017>, 2017.
- 760 Magnan, A., Duvat, V., and Garnier, E.: Reconstituer les « trajectoires de vulnérabilité » pour penser  
761 différemment l'adaptation au changement climatique, *Nat. Sci. Soc.*, 20, 82–91,  
762 <https://doi.org/10.1051/nss/2012008>, 2012.

- 763 Mélo, A., Wilhelm, B., Giguet-Covex, C., and Arnaud, F.: Construire une chronique d'inondations :  
 764 évènements hydrologiques et histoire climatique dans le bassin de l'Arve (Alpes du Nord, France) entre  
 765 les XVIIIe et XXIe siècles., 40, 411–419, 2015.
- 766 Ménégos, M., Valla, E., Jourdain, N. C., Blanchet, J., Beaumet, J., Wilhelm, B., Gallée, H., Fettweis, X.,  
 767 Morin, S., and Anquetin, S.: Contrasting seasonal changes in total and intense precipitation in the  
 768 European Alps from 1903 to 2010, 1–37, <https://doi.org/10.5194/hess-2019-690>, 2020.
- 769 Merz, R. and Blöschl, G.: Regional flood risk—what are the driving processes?, 40–58, 2003.
- 770 Meunier, M.: La catastrophe du Grand Bornand : crue torrentielle du Borne le 14 juillet 1987, 78, 103–  
 771 114, 1990.
- 772 Min, S.-K., Zhang, X., Zwiers, F. W., and Hegerl, G. C.: Human contribution to more-intense precipitation  
 773 extremes, *Nature*, 470, 378–381, <https://doi.org/10.1038/nature09763>, 2011.
- 774 Ministère de la Transition écologique: Prévention des risques naturels, 2020.
- 775 Ministère de la Transition Ecologique: Base de Données Historiques sur les Inondations, n.d.
- 776 Mougin, P.: Les torrents de la Savoie, Société d'histoire de la Savoie, Grenoble, 1914.
- 777 Papagiannaki, K., Lagouvardos, K., and Kotroni, V.: A database of high-impact weather events in  
 778 Greece: a descriptive impact analysis for the period 2001–2011, *Nat. Hazards Earth Syst. Sci.*, 13, 727–  
 779 736, <https://doi.org/10.5194/nhess-13-727-2013>, 2013a.
- 780 Papagiannaki, K., Lagouvardos, K., and Kotroni, V.: A database of high-impact weather events in  
 781 Greece: a descriptive impact analysis for the period 2001–2011, *Nat. Hazards Earth Syst. Sci.*, 13, 727–  
 782 736, <https://doi.org/10.5194/nhess-13-727-2013>, 2013b.
- 783 Pardé, M.: Les crues de l'Arve en octobre 1930, *rga*, 19, 495–497,  
 784 <https://doi.org/10.3406/rga.1931.4782>, 1931.
- 785 Peiry, J. L. and Bravard, J. P.: Evolution naturelle d'un remplissage sédimentaire intramontagnard et  
 786 impacts des aménagements contemporains : L'exemple de la vallée de l'Arve (74), *La Houille Blanche*,  
 787 221–225, <https://doi.org/10.1051/lhb/1989018>, 1989.
- 788 Petrucci, O., Aceto, L., Bianchi, C., Bigot, V., Brázdil, R., Pereira, S., Kahraman, A., Kiliç, Ö., Kotroni, V.,  
 789 Llasat, M. C., Llasat-Botija, M., Papagiannaki, K., Pasqua, A. A., Řehoř, J., Rossello Geli, J., Salvati, P.,  
 790 Vinet, F., and Zêzere, J. L.: Flood Fatalities in Europe, 1980–2018: Variability, Features, and Lessons to  
 791 Learn, *Water*, 11, 1682, <https://doi.org/10.3390/w11081682>, 2019.
- 792 Préfecture de la Haute-Savoie: Données communales : plans de prévention des risques naturels  
 793 (PPRN), Les services de l'Etat en Haute-Savoie, n.d.
- 794 Rougier, H.: Les inondations du 24 juillet dans la haute vallée de l'Arve : faits et conséquences à tirer  
 795 pour l'aménagement du territoire., *Les pouvoirs publics face aux risques naturels dans l'histoire.*,  
 796 Grenoble, 51–64, 2001.
- 797 Rousset-Mestrallet, M.: Des torrents et des hommes. Trois siècles d'histoire à Samoëns., *Imprim'off 7*,  
 798 Marignier, 150 pp., 1986.
- 799 RTM and ONF: Base de Données RTM, 2012.

800 Ruin, I., Creutin, J.-D., Anquetin, S., and Lutoff, C.: Human exposure to flash floods – Relation between  
801 flood parameters and human vulnerability during a storm of September 2002 in Southern France,  
802 *Journal of Hydrology*, 361, 199–213, <https://doi.org/10.1016/j.jhydrol.2008.07.044>, 2008.

803 Schlögl, M., Fuchs, S., Scheidl, C., and Heiser, M.: Trends in torrential flooding in the Austrian Alps: A  
804 combination of climate change, exposure dynamics, and mitigation measures, *Climate Risk*  
805 *Management*, 32, <https://doi.org/10.1016/j.crm.2021.100294>, 2021.

806 SITG: GEOMOL-GEOPOTENTIELS, 2020.

807 Soanes, C. and Stevenson, A.: *Catastrophe*, Concise Oxford English Dictionary, 11 (revised), 2009.

808 Terti, G., Ruin, I., Anquetin, S., and Gourley, J. J.: Dynamic vulnerability factors for impact-based flash  
809 flood prediction, *Nat Hazards*, 79, 1481–1497, <https://doi.org/10.1007/s11069-015-1910-8>, 2015.

810 Thoumas, P.: Histoire des crues du Roubion (Drôme, France) depuis 1501 AD, une approche  
811 hydrologique des fluctuations climatiques sur cinq siècles, *physio-geo*, 14, 87–111,  
812 <https://doi.org/10.4000/physio-geo.8984>, 2019.

813 Viani, A., Condom, T., Sicart, J.-E., Rabatel, A., Gascoin, S., Ranzi, R., and Wimez, M.: Impact of the  
814 glacier retreat and snow melt on the seasonal cycle of streamflow of the Arve catchment since the  
815 1960s (Northern French Alps), *EGU General Assembly, Vienna, Austria*, 1, 2018.

816 Vincent, C., Descloitres, M., Garambois, S., Legchenko, A., Guyard, H., and Gilbert, A.: Detection of a  
817 subglacial lake in Glacier de Tête Rousse (Mont Blanc area, France), *Journal of Glaciology*, 58, 866–878,  
818 <https://doi.org/10.3189/2012JoG11J179>, 2012.

819 Wetter, O.: The potential of historical hydrology in Switzerland, *Hydrol. Earth Syst. Sci.*, 21, 5781–5803,  
820 <https://doi.org/10.5194/hess-21-5781-2017>, 2017.

821 Wilhelm, B., Ballesteros Canovas, J. A., Macdonald, N., Toonen, W., Baker, V., Barriendos, M., Benito,  
822 G., Brauer, A., Corella, J. P., Denniston, R., Glaser, R., Ionita, M., Kahle, M., Liu, T., Luetscher, M.,  
823 Macklin, M., Mudelsee, M., Munoz, S., Schulte, L., St. George, S., Stoffel, M., and Wetter, O.:  
824 Interpreting historical, botanical, and geological evidence to aid preparations for future floods, 6,  
825 <https://doi.org/10.1002/wat2.1318>, 2019.

826 Zgheib, T., Giacona, F., Granet-Abisset, A.-M., Morin, S., and Eckert, N.: One and a half century of  
827 avalanche risk to settlements in the upper Maurienne valley inferred from land cover and socio-  
828 environmental changes., *Global Environmental Change*, 65, 102149,  
829 <https://doi.org/10.1016/j.gloenvcha.2020.102149>, 2020.

830