

1 **Spatial variability in the relation between Drivers of extreme burnt**
2 **area in Portugal: fire weather and burned area: patterns and**
3 **drivers in Portugal**
4 **vegetation**

5 T. Calheiros (1), A. Benali (2), ~~J. M. N. Silva (2)~~, M.G. Pereira (3,4), J. M. N. Silva (2), J.P.
6 Nunes (1,5)

7 (1) cE3c: centre for Ecology, Evolution and Environmental changes, Faculdade de Ciências,
8 Universidade de Lisboa, Lisboa, Portugal

9 (2) Forest Research Centre, School of Agriculture, University of Lisbon, Tapada da Ajuda,
10 1349-017 Lisboa, Portugal

11 (3) Centro de Investigação e de Tecnologias Agro-Ambientais e Biológicas (CITAB),
12 Universidade de Trás-os-Montes e Alto Douro, Vila Real, Portugal

13 (4) IDL, Universidade de Lisboa, Lisboa, Portugal

14 (5) Soil Physics and Land Management Group, Wageningen University and Research,
15 Wageningen, Netherlands

16
17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32 ABSTRACT

33 Fire weather indices are used to assess the effect of weather conditions on wildfire behaviour
34 ~~and. Previous studies identified~~ the high Daily Severity Rating percentile (DSRp) ~~is~~ strongly
35 related to the total burned area (BA) in Portugal. The aims of this study were to: 1) assess if the
36 90th DSRp (DSR90p) threshold is adequate ~~for to estimate large BA in mainland~~ Portugal; 2)
37 identify and characterize regional variations of the DSRp threshold, at higher resolution, that
38 justifies the ~~bulk~~majority of BA; and, 3) analyse if vegetation cover can explain the DSRp
39 spatial variability.

40 We used ~~wildfire data~~, weather reanalysis data from ERA5-Land, ~~for the 2001—2019 period~~,
41 ~~wildfire and the land use map for Portugal. DSRp were data and from official Portuguese~~
42 ~~authorities for the 2001 – 2019 study period. We computed for DSRp and associated it to large~~
43 ~~wildfires (BA > 100 ha) that occurred in an extended summer period and combined with~~
44 ~~individual large wildfires. Cluster analysis was performed using the relationship between DSRp~~
45 ~~and BA, in each municipality.~~


46 (15th May to 31st October). Results revealed that the DSR90p is an adequate ~~threshold for~~
47 ~~Portugal~~indicator of extreme fire weather days and ~~is well related to large extreme BA in~~
48 ~~Portugal~~. However, the spatial pattern of the DSRp associated with the majority of total BA
49 shows some variability at the municipality scale, ~~differences appear between the DSRp linked~~
50 ~~to the majority of accumulated BA. Cluster analysis revealed that municipalities where large~~
51 Municipalities where large wildfires occur ~~in high DSRp present higher BA with extreme~~
52 weather conditions have burned areas mostly in forests and are located in coastal areas. In
53 contrast, ~~clusters with lower DSRp present greater BA in municipalities where large fires occur~~
54 with less extreme weather conditions are predominantly covered by shrublands and are situated

55 in eastern and inland regions. These findings can support better prevention and fire suppression
56 planning.

57

58 KEY WORDS: Wildfires, Cluster analysis, Fire weather, Land Use/Land Cover.

59



60 1. Introduction

61 Fire regime can be defined, in a strict sense, by the spatial and temporal patterns of wildfire
62 characteristics (e.g. occurrence, frequency, size, seasonality, etc), as well as, in a broad sense,
63 by vegetation characteristics, fire effects and fire weather in a given area or ecosystem, based
64 on fire histories at individual sites over long periods, generally resulting from the cumulative
65 interaction of fire, vegetation, climate, humans, and topography over time (Crutzen and
66 Goldammer, 1993; NCWG, 2011; Whitlock et al., 2010). **Wildfire**

67 One of the most important factors of fire regime is the wildfire incidence, that is defined as the
68 number of fire events and/or burnt area (BA). This factor depends on the weather and climate,
69 especially in regions with a Mediterranean-type of climate, where mild and rainy winters and
70 springs favour vegetation growth, while dry and hot summers promote thermal and hydric stress
71 of live fuels and dryness of dead fuels (Romano and Ursino, 2020). In the western
72 Mediterranean, the influence of climate variability on wildfire incidence became more evident
73 after the 1970s, following a fire regime change, from fuel-limited to drought-driven (Pausas
74 and Fernández-Muñoz, 2012). The main factor for this change was the increase in fuel load and
75 continuity due to rural depopulation and land abandonment (Moreira et al., 2011; Moreno et
76 al., 2014). These changes in landscape and population favoured the occurrence of large
77 wildfires (Ferreira-Leite et al., 2016), which tend to occur with severe fire weather conditions,
78 being rare in other meteorological conditions (Telesca and Pereira, 2010). Wildfires can also
79 modify the landscape in the Mediterranean region (e.g. Stamou et al. (2016)) influenced by
80 regeneration patterns, topography and local fire histories.

81 Land use interfaces, in particular those between forests and other land use types (shrublands,
82 agricultural and urban areas), have a significant effect on human-caused wildfire occurrence in
83 Mediterranean Europe, increasing fire risk due to human causes (Vilar et al., 2016). In the
84 Iberian Peninsula, shrublands and pine forests have registered larger burnt areas (Barros and

85 Pereira, 2014; Pausas and Vallejo, 1999). This fact can be explained by the increasing landscape
86 homogenization, due to shrublands expansion and agricultural abandonment, as observed by -
87 ~~The Iberian Peninsula is the European region with the highest wildfire incidence and~~
88 ~~consequently, suffers large property damage~~Lloret *et al.* (2002).

89 Heatwaves and droughts have a strong influence on fire incidence, as shown by several studies
90 in the last years in Mediterranean Europe (e.g., Duane and Brotons, 2018; Sutanto *et al.*, 2020).
91 The impacts of droughts on vegetation create favourable conditions for the ignition and spread
92 of wildfires, especially during ~~and fatalities (San-Miguel-Ayanz *et al.*, 2020).~~ ~~In particular,~~
93 ~~Portugal has been severely affected by wildfires in the last decades, especially in 2003, 2005~~
94 ~~and 2017 (Gouveia *et al.*, 2012; Trigo *et al.*, 2006; Turco *et al.*, 2019).~~

95 ~~The impacts of droughts on vegetation can create favourable conditions for the ignition and~~
96 ~~spread of wildfires, especially in~~ summer (Pausas and Fernández-Muñoz, 2012; Russo *et al.*,
97 2017), but also in winter (Amraoui *et al.*, 2015; Calheiros *et al.*, 2020). In addition, fire
98 incidence increased dramatically with the combined effect of prolonged drought and heatwaves,
99 as pointed out by Ruffault *et al.*, (2018). Wildfire incidence in Mediterranean Europe is
100 expected to increase in the future because of climate change, especially due to global warming
101 and changes in the precipitation regime (Sousa *et al.*, 2015; Turco *et al.*, 2018).

102 The Iberian Peninsula is the European region with the highest wildfire incidence which causes
103 large property damages and fatalities (San-Miguel-Ayanz *et al.*, 2020). In particular, Portugal
104 has been severely affected by wildfires in the last decades, especially in 2003, 2005 and 2017,
105 mainly as a consequence of anomalous atmospheric synoptic patterns and extreme weather
106 conditions (Gouveia *et al.*, 2012; Trigo *et al.*, 2006; Turco *et al.*, 2019).~~Heatwaves and droughts~~
107 ~~have a strong influence on fire incidence, as shown by several studies in the last years in~~
108 ~~Mediterranean Europe (Duane and Brotons, 2018; Sutanto *et al.*, 2020).~~ ~~In addition, fire~~
109 ~~incidence increased dramatically with the combined effect of prolonged drought and heatwaves~~

110 ~~in Mediterranean France, as pointed out by Ruffault *et al.*, (2018), or as occurred in the~~
111 ~~catastrophic fires of 2017 in Portugal (Turco *et al.*, 2019).~~ Other studies identified weather
112 types, most of them connected with heatwaves or droughts in the western Iberian Peninsula,
113 associated with the occurrence of large wildfires (Rodrigues *et al.*, 2020; Vieira *et al.*, 2020).

114 ~~In Western Mediterranean, the influence of climate variability on fire incidence became more~~
115 ~~evident after the 1970s, following a fire regime change, from fuel-limited to drought-driven~~
116 ~~(Pausas and Fernández-Muñoz, 2012). The main factor for this change was the increase of fuel~~
117 ~~load and continuity due to rural depopulation and land abandonment (Moreira *et al.*, 2011;~~
118 ~~Moreno *et al.*, 2014). These changes in landscape and population favoured the occurrence of~~
119 ~~large wildfires (Ferreira-Leite *et al.*, 2016), which tend to occur with severe fire weather~~
120 ~~conditions, being rare in other meteorological conditions (Teleseca and Pereira, 2010). The~~
121 ~~contribution of landscape-level fuel connectivity for wildfire size was evident in the~~
122 ~~1998–2008 period (Fernandes *et al.*, 2016). These changes in the landscape, together with~~
123 ~~socioeconomic changes, impact the fire regime (Pereira *et al.*, 2014; Parente and Pereira, 2016;~~
124 ~~Parente *et al.*, 2018). Future climate change will increase fire incidence in Mediterranean~~
125 ~~Europe (Sousa *et al.*, 2015; Turco *et al.*, 2018).~~

126 ~~Fire regime can be defined, in a strict sense, as a statistical concept described by the spatial and~~
127 ~~temporal patterns of wildfire characteristics (occurrence, frequency, size, seasonality, etc), as~~
128 ~~well as, in a broad sense, vegetation characteristics, fire effects and fire weather in a given area~~
129 ~~or ecosystem, based on fire histories at individual sites over long periods, generally result from~~
130 ~~the cumulative interaction of fire, vegetation, climate, humans, and topography over time~~
131 ~~(Crutzen and Goldammer, 1993; NCWG, 2011; Whitlock *et al.*, 2010). Cluster analysis for~~
132 ~~the Iberian Peninsula has identified several regions with similar fire regimes, using several~~
133 ~~variables related to fire, such as the intra-annual pattern of burnt area (BA) (Trigo *et al.*, 2016;~~
134 ~~Calheiros *et al.*, 2020; Calheiros *et al.*, 2021), fire activity and weather risk (Jimenez-Ruano *et*~~

135 ~~al., 2018), large fire weather typologies (Rodrigues et al., 2020) or BA spatio-temporal trends~~
136 ~~(Silva et al., 2019).~~

137 Fire weather danger indices are commonly used to assess the current and/or cumulative effect
138 of atmospheric conditions on fuel moisture and fire behaviour. The Canadian Forest Fire
139 Weather Index (FWI) System (CFFWIS) consists of six components that account for those
140 effects (Van Wagner, 1987), including the Daily Severity Rating (DSR). The 90th percentile of
141 the DSR (DSR90p) is often used as the threshold for severe fire weather that is associated with
142 large fires (Bedia et al., 2012; Carvalho et al., 2008; Fernandes, 2019; Silva et al., 2019). More
143 recently, the 95th percentile of DSR (DSR95p) was also identified as a good indicator of extreme
144 fire weather and well related to the BA in the Iberian Peninsula (Calheiros *et al.*, 2020; Calheiros
145 *et al.*, 2021). ~~BA and extreme fire weather days have a strong link, noticeable in the similar~~
146 ~~intra-annual variability pattern in the four pyro-regions of the Iberian Peninsula (Calheiros et~~
147 ~~al., 2020). This robust link was used to anticipate fire regime changes caused by future climate~~
148 ~~change, revealing the potential displacement of fire regimes to the north (Calheiros et al., 2021).~~

149 ~~Wildfires in Portugal were the subject of several studies that developed zoning approaches to~~
150 ~~identify regions with similar fire regimes using burnt area data (Kanevski and Pereira, 2017;~~
151 ~~Scotto et al., 2014), combined with fire weather indices (Calheiros et al., 2020; Calheiros et al.,~~
152 ~~2021) Wildfires in Portugal were the subject of several studies that developed zoning approaches~~
153 ~~to identify regions with similar fire regimes using solely burnt area data (Kanevski and Pereira,~~
154 ~~2017; Scotto et al., 2014; Silva et al., 2019) or combined with fire weather indices (Calheiros~~
155 ~~et al., 2020, 2021; Jimenez-Ruano et al., 2018), large fire-weather typologies (Rodrigues et al.,~~
156 ~~2020), population density, topography, land cover changes (Oliveira et al., 2017) and net~~
157 ~~primary production (Fernandes, 2019), or fire prevention policy decisions (Parente et al., 2016).~~

158 ~~Generally, clustering. Their~~ results indicate that Portugal can be divided into two (dividing the
159 north and south of Tajo River) or three main clusters (the north part further divided in western

160 and eastern). ~~Oliveira et al. (2017) added a fourth cluster in the central littoral region. Actually,~~
161 ~~the~~The spatial and temporal distribution of wildfires presents clustering patterns, suggesting
162 that small fires are more dependent on local topographic or human conditions, while large fires
163 are a consequence of infrequent causes or with shorter periods such as weather extreme events
164 (Pereira et al., 2015). The temporal pattern is characterized by periodicities and scaling regimes
165 (Telesca and Pereira, 2010) including a main summer fire season and a secondary spring peak,
166 both driven by the type of climate and the occurrence of extreme weather conditions (Amraoui
167 et al., 2015; Trigo et al., 2016; Calheiros et al., 2020).

168 ~~Another essential element for fire incidence is the vegetation and land use type.~~ There have
169 been important changes in land use since the 1960s in Portugal which are related to wildfire
170 occurrence. Arable cropland decreased from 40% to only 12% of the total area in 2006, at the
171 national level; and forest declined since the 1980s, as a result of forest fires, in Central Portugal
172 (Jones et al., 2011). ~~The contribution of landscape-level fuel connectivity for wildfire size was~~
173 ~~evident in the 1998 – 2008 period (Fernandes et al., 2016).~~ ~~The analysis of Corine Land Cover~~
174 ~~maps for 2000 and 2006 and EFFIS BA perimeters, from 2000 to 2013 in Portugal, revealed an~~
175 ~~increase in the area of shrublands and a decrease in forest areas , together with socioeconomic~~
176 ~~changes, impact the fire regime (Pereira et al., 2014; Parente and Pereira, 2016; Parente et al.,~~
177 ~~2018b).~~ ~~In Portugal, eucalyptus expansion has not modified the fire regime, but the rising~~
178 ~~undermanaged and abandoned forest plantations, especially after large-fire seasons, is a concern~~
179 ~~for the future (Fernandes et al., 2019).~~ ~~The analysis of Corine Land Cover maps for 2000 and~~
180 ~~2006 and EFFIS BA perimeters, from 2000 to 2013 in Portugal, revealed an increase in the area~~
181 ~~of shrublands, a decrease in forest areas, 51% of total BA in shrublands but a much higher~~
182 ~~wildfire proneness in shrublands than in forest areas (Pereira et al., 2014). Other studies have~~
183 ~~confirmed that shrublands~~

184 Shrublands are more susceptible to wildfires, whereas agricultural areas and agroforestry
185 systems are less likely to burn, as revealed by several studies (Carmo *et al.*, 2011; Nunes, 2012;
186 Meneses *et al.*, 2018, 2018a). Barros and Pereira, (2014) identified shrublands as the most
187 wildfire-prone land cover, followed by pine forests while, on the contrary, annual crops and
188 evergreen oak woodlands tend to be avoided by wildfire. Ferreira-Leite *et al.*, (2016) concluded
189 that uncultivated land (shrublands, grasslands, and other sparse vegetation) was the most
190 important factor affecting burnt areas, considering large wildfires, greater than 100 ha.
191 Topography and uncultivated land were significant factors determining burnt area, in a study
192 for the 1980-2014 period conducted at the municipal level (Nunes *et al.*, 2016). Additionally,
193 there is evidence of an extending urban-rural interface in Portugal, due to an increase in the
194 urban area since 1990, which contributes to an increase in fire incidence (Silva *et al.*, 2019),
195 especially in those regions (Tonini *et al.*, 2018).

196 A previous study, assessed the recent evolution of spatial and temporal patterns of BA and fire
197 weather risk in the Iberian Peninsula and concluded that the DSR95p is a good indicator of
198 extreme fire weather and is well related to the BA, noticeable in the similar intra-annual
199 variability pattern in four pyro-regions (Calheiros *et al.*, 2020). This robust link was used to
200 anticipate fire regime changes caused by future climate change, revealing the potential
201 displacement of fire regimes to the north (Calheiros *et al.*, 2021). Another essential element for
202 fire incidence is the vegetation and land use type. For example, land use interfaces, that are
203 generally between forests and other land use types (shrublands, agricultural and urban), have a
204 significant effect on human caused wildfire occurrence in Mediterranean Europe, showing that
205 larger interfaces have a larger risk of fire happen due to human causes (Vilar *et al.*, 2016). Fuel
206 removal can be a solution for the extending area of wildland-urban interfaces (Elia *et al.*, 2016).
207 However, previous studies did not look at additional factors such as landcover. Accordingly,
208 the objectives of this work were:

209 ~~Wildfires can also modify the landscape in the Mediterranean region (e.g. Stamou et al. (2016))~~
210 ~~influenced by regeneration patterns, topography and local fire histories. In the Iberian~~
211 ~~Peninsula, shrublands and pine forests have registered larger burnt areas (Barros and Pereira,~~
212 ~~2014; Pausas and Vallejo, 1999). This fact can be explained by the increasing landscape~~
213 ~~homogenization, due to shrublands expansion and agricultural abandonment, as observed by~~
214 ~~Lloret et al. (2002), in eastern Spain. In Portugal, eucalyptus expansion has not modified the~~
215 ~~fire regime, but the rising undermanaged and abandoned forest plantations, especially after~~
216 ~~large fire seasons, is a concern for the future (Fernandes et al., 2019).~~

217 ~~There is evidence of an extending urban-rural interface in Portugal, due to an increase of the~~
218 ~~urban area to double since 1990, which contributes to an increase in fire incidence (Silva et al.,~~
219 ~~2019). Results obtained for the entire territory of Continental Portugal in the 1990–2012 period~~
220 ~~reveal that the rural-urban interface increased by more than two thirds, the total BA decreased~~
221 ~~by one third, but the BA within the interface doubled (Tonini et al., 2018).~~

222 ~~Although the incidence of fire has several factors with variable influence, this study focuses on~~
223 ~~the relationship between extreme fire weather and high BA, resulting from large wildfires in~~
224 ~~Portugal. A previous study, assessed the recent evolution of spatial and temporal patterns of BA~~
225 ~~and fire weather risk in the Iberian Peninsula (Calheiros et al., 2020) and concluded that the~~
226 ~~DSR90p is a good indicator of extreme fire weather and is well related to the BA in the Iberian~~
227 ~~Peninsula.~~

228 ~~Given the role of extreme weather on BA resulting from large wildfires, the common use of~~
229 ~~DSR thresholds and the effect of other factors, namely land use/land cover, the objectives of~~
230 ~~this work were:~~

- 231 1) ~~to~~ assess if the DSR90p threshold is adequate ~~for~~ to estimate large BA in mainland
232 Portugal;

- 233 2) to identify and characterize regional variations of the DSRp threshold, at higher
234 resolution, that justifies the bulkmajority of BA, and;
235 3) to analyse if vegetation cover can explain the spatial variability of the DSRp.


236

237 2. Data and methodology

238 2.1 Study Area: Portugal

239 This study focuses on mainland Portugal, ~~a territory divided by 23 NUTS III provinces~~
240 ~~themselves subdivided into 278 municipalities and topographically~~ characterized by
241 mountainous areasranges in north and central regions and vast plains in the south ~~(Figure 1),~~
242 divided in 23 NUTS III regions which, in turn, are subdivided into 278 municipalities (Fig. 1).

243 The BA variability is mainly influenced by the precipitation anomaly in spring and the
244 occurrence of abnormal atmospheric patterns that generate very hot and dry days in the western
245 Iberian Peninsula during summer (Pereira *et al.*, 2005). In fact, 97% of the total number of
246 extreme wildfires (with BA ≥ 5000 ha) were active during heatwaves (Parente *et al.*,
247 20182018a) while almost 90% of extreme wildfires during the 1981–2017 period occurred
248 within a region affected by drought (Parente *et al.*, 2019). ~~Fire weather in Portugal has usually~~
249 ~~been characterized using the CFFWIS (Calheiros *et al.*, 2021; Calheiros *et al.*, 2020; Silva *et*~~
250 ~~*al.*, 2019; Nunes *et al.*, 2019; Pereira *et al.*, 2013; Carvalho *et al.*, 2008), which provides good~~
251 ~~results in comparison with other methods of fire danger evaluation (Viegas *et al.*, 1999).~~



252
 253 Figure 1: Mainland Portugal topography and NUTSIII provinces: *Alto Minho*(1), *Cávado*(2), *Ave*(3), *Alto*
 254 *Tâmega*(4), *Terras de Trás-os-Montes*(5), *Área Metropolitana do Porto*(6), *Tâmega e Sousa*(7), *Douro*(8), *Região*
 255 *de Aveiro*(9), *Viseu Dão-Lafões*(10), *Beiras e Serra da Estrela*(11), *Região de Coimbra*(12), *Região de Leiria*(13);
 256 *Médio Tejo*(14), *Beira Baixa*(15), *Oeste*(16), *Lezíria do Tejo*(17), *Alto Alentejo*(18), *Área Metropolitana de*
 257 *Lisboa*(19), *Alentejo Central*(20), *Alentejo Litoral*(21), *Baixo Alentejo*(22) and *Algarve*(23). Data from European
 258 Environment Agency (2021) and DGT (2010). Pyro-regions limits from Calheiros *et al.*, (2020), for comparison
 259 purposes, were also added, at blue: NW pyro-region is located in northwestern Portugal and SW pyro-region in
 260 southwestern and eastern of the country.

261
 262 **2.2 Fire Weather Index and.**




Figure 1: Mainland Portugal topography and administrative division based on NUTSIII provinces: *Alto Minho* (1), *Cávado* (2), *Ave* (3), *Alto Tâmega* (4), *Terras de Trás-os-Montes* (5), *Área Metropolitana do Porto* (6), *Tâmega e Sousa* (7), *Douro* (8), *Região de Aveiro* (9), *Viseu Dão-Lafões* (10), *Beiras e Serra da Estrela* (11), *Região de Coimbra* (12), *Região de Leiria* (13), *Médio-Tejo* (14), *Beira Baixa* (15), *Oeste* (16), *Lezíria do Tejo* (17), *Alto Alentejo* (18), *Área Metropolitana de Lisboa* (19), *Alentejo Central* (20), *Alentejo Litoral* (21), *Baixo Alentejo* (22) and *Algarve* (23). NUTSIII frontiers were loaded from the European Environment Agency (EEA, 2021) and altitude data from *Direção Geral do Território* (DGT, 2010). Borders (thick black line) of the pyro-regions found by *Calheiros et al.*, (2020), for comparison purposes, were also added: NW pyro-region is located in northwestern Portugal and SW pyro-region in southwestern and eastern of the country.

The territory of Continental Portugal is mostly covered by forests (39%), agricultural lands (26%), shrublands (12%) and agroforestry systems (8%), according to data from *Direção*

276 Geral do Território (DGT, 2019). The most common tree species are *Eucaliptus Globulus* (26%
277 of all forests), *Pinus Pinaster* (22%), both prevalent in the north and centre; and *Quercus*
278 *suber* (22%), with larger areas in the south, using forest data from *Instituto Nacional da*
279 *Conservação da Natureza e das Florestas* (ICNF, 2019). Pyro-regions shown in Fig. 1 are both
280 characterized by a high peak of BA centred in August and a much smaller one in March. The
281 main difference between the NW and SW pyro-region is the larger values of BA in the NW
282 pyro-region, compared with the SW, especially in August (Calheiros *et al.*, 2020).

284 **2.2 Meteorological Data and Fire Weather Indices**

285 We used the DSR which is ~~an additional component of the FWI system~~ more accurate to rate
286 ~~more accurately~~ the expected efforts required to suppression ~~or control the wildfire and is based~~
287 ~~on the FWI which, in turn, rates the fire intensity and is frequently used to inform the general~~
288 ~~public about fire weather danger conditions~~ a wildfire, being an additional component of the
289 FWI system (De Groot, 1987; Van Wagner, 1987).–The indices of the FWI system were
290 computed for the 2001 – 2019 study period with the equations provided by Van Wagner and
291 Pickett (1975) and daily values at 12h00UTC of air temperature and relative humidity (at 2
292 meters), wind speed (at 10 meters), and accumulated total precipitation.

293 ~~The~~ Data of the meteorological variables were obtained from the fifth generation of European
294 Centre for Medium-Range Weather Forecasts (ECMWF) atmospheric reanalyses of the global
295 climate (ERA5-Land). The ERA5-Land dataset was loaded from the Copernicus Climate
296 Change Service (Copernicus Climate Change Service (C3S), 2017) has (C3S, 2020), with a
297 much higher spatial resolution (0.1° lat × 0.1° long; the native resolution is 9 km) and temporal
298 (hourly) resolution than the previous reanalysis data service, that were widely used and with
299 good performances for different purposes, including FWI calculation in Portugal (Bedia *et al.*,
300 2012). The ERA5 is recognized as the best or one of the best global atmospheric reanalysis

301 datasets (Huai *et al.*, 2021; Muñoz-Sabater *et al.*, 2021; Urban *et al.*, 2021) and used worldwide
302 (Chinita *et al.*, 2021; Sianturi *et al.*, 2020). ~~Therefore, it is one of the most used meteorological~~
303 ~~datasets in the world.~~

305 2.3 Land use and wildfire data

306 Land use and land cover (LULC) map for 2018 (COS2018) and wildfire data, for the 2001 to
307 2019 period, were provided by ~~Portuguese national authorities, respectively, Direção Geral do~~
308 ~~Território (DGT, 2019) and the Instituto Nacional da Conservação da Natureza e das Florestas~~
309 ~~(ICNF, 2020). These datasets were the previously mentioned Portuguese national authorities~~
310 ~~(DGT, 2019; ICNF, 2020). These datasets were successfully~~ used in many other studies, by a
311 large number of authors for a wide variety of purposes (Bergonse *et al.*, 2021; Tarín-Carrasco
312 *et al.*, 2021). Only wildfires ~~larger than 100ha that with BA > 100 ha~~ occurred during the
313 extended summer season (~~here~~ defined between 15th May and 31st October), were investigated.
314 ~~When a given wildfire affected more than one municipality, the resulting BA extent was~~
315 ~~allocated considered in this study. It is important to each of the administrative units burned by~~
316 ~~the wildfire explain these methodological options.~~

317 ~~The starting and ending dates of each wildfire were fundamental information to attribute the~~
318 ~~DSR to each BA. This process was accomplished using MODIS satellite data, computed using~~
319 ~~the same method as in Benali *et al.* (2016), with start and end dates and ignition location~~
320 ~~estimated for circa 92% of the total BA, for large wildfires. Daily DSR was computed for the~~
321 ~~same period (2001–2019) and all ERA5 Land grid points within continental Portugal. The size~~
322 ~~of Portuguese municipalities is relatively small, so there are no major weather variations within.~~
323 ~~The DSR percentiles (DSRp) considered in the analysis carried out for the entire territory of~~
324 ~~mainland Portugal was the maximum value of DSR recorded during the duration of the wildfire.~~

325 In the case of the analysis performed based on the municipalities, the considered DSRp was the
326 maximum value of DSR during the duration of the wildfire in each municipality. Afterwards,
327 we computed the and assigned to the BA within the administrative unit.

328 BA data were normalized using both the decimal logarithm and fraction of the total burnt area
329 (FTBA), in percentage. Exploratory analysis showed that the BA extent of individual small fires
330 was poorly correlated with DSRp and, consequently, sorting was performed. Afterwards, BA
331 data for the entire mainland Portugal territory were sorted by assigned DSRp and the logarithm
332 of accumulated Burnt Area was plotted against DSRp to assess if this relationship is linear.
333 Subsequently, we analysed if a fixed threshold of DSR for extreme days – DSR90p – is adequate
334 to estimate extreme fire weather and is well related to large FTBA, for the entire territory. We
335 considered the correspondent 80% and 90% of FTBA as sufficient to classify DSRp as the
336 extreme threshold, justified by the results of Pereira *et al.*, (2005), which showed that 80% of
337 TBA occurs in 10% of summer days.

338 We selected 175 municipalities (from 278) affected by more than three individual wildfires and
339 a total BA > 500 ha in the studied period (2001–2019). Restricting the analysis to the
340 administrative units with sufficient data aims to increase the robustness of the results and to
341 prevent possible interpretation errors. Figures assessing the relation between DSRp and FTBA
342 were produced, for all the selected municipalities, concerning the second objective.

343 In each municipality, the selection of the maximum spatial value of DSR to associate with fires
344 is justified by the low spatial variability of the DSR, the small size of administrative units and
345 the native reanalysis data resolution (Copernicus Climate Change Service (C3S), 2017). The
346 BA division between municipalities can produce noise in the data. This procedure artificially
347 generates wildfires, some of them with relatively small size but high or very high DSRp. To
348 circumvent this difficulty, we decided to analyze BA percentages, which reduce the influence
349 of small wildfires on the final results.

~~It is important to address some methodological options. Only wildfires that occurred in the extended summer period, from~~ The focus on relatively large wildfires (here defined as wildfires with BA>100 ha) has two main reasons. First, mainland Portugal registers a huge number of small wildfires but they account only for a small amount of total BA (TBA). For example, wildfires with BA>100 ha are just about 1% of all wildfires but account for 75% of TBA (Pereira *et al.*, 2011). Second, wildfires in Portugal are mainly (99.4%) caused by humans, either by negligence (about one-quarter of the total number of wildfires with known cause) and intentionally (about three quarters), associated with the use of fire, accident and structural/land use (Parente *et al.*, 2018b), which means that small wildfires can occur with relatively low DSR.

~~The study only considered wildfires occurred during the 15th May to 31st October, were studied period~~ because of also two main reasons: (i) BA caused by large wildfires within this period accounts for 97.5% of TBA, assuming only large fires; and, (ii) the secondary peak of fire incidence in Portugal occurs in late winter \ /early spring, with low when DSR values is lower and depends much more on drought than on high air temperature (Amraoui *et al.*, 2015; Calheiros, *et al.*, 2020). Only large wildfires (BA>100 ha), similarly defined by the Portuguese forest authorities (ICNF), have been included also for two reasons. First, wildfires in Portugal are mainly (99.4%) caused by humans, by negligence (about one quarter of the total number of wildfires with known cause) and intentionally (about three quarters), associated with the use of fire, accident and structural/land use (Parente *et al.*, 2018), i.e., small wildfires can occur with relatively low DSR. Second, mainland Portugal registers a very large number of small wildfires but they account only for a small amount of TBA. For example, wildfires with BA>100 ha are just about 1% of all wildfires but account for 75% of total BA (Pereira *et al.*, 2011). The datasets and wildfire metrics used in this study are summarized in Table 1 and Table 2, respectively.

Table 1. Data sources, types, variables and methodology where it is used.

<u>Data source and type</u>	<u>Variables</u>	<u>Methodology</u>
<u>ERA5-Land. Meteorological data for 2001-2019</u> <u>Provided by the ECMWF</u>	<u>Temperature</u>	<u>To compute FWI indices, including DSR</u>
	<u>Relative Humidity</u>	
	<u>Wind speed</u>	
	<u>Precipitation</u>	
<u>COS2018. Land Use and Land Cover data.</u> <u>Provided by the <i>Direção Geral do Território</i></u>	<u>Forest</u>	<u>To assess burnable areas and the land cover type affected by each wildfire</u>
	<u>Shrublands</u>	
	<u>Agriculture</u>	
	<u>Agroforestry</u>	
	<u>Other burnable areas</u>	
<u>Wildfire data for 2001 – 2019.</u> <u>Provided by the <i>Instituto da Conservação da Natureza e das Florestas</i></u>	<u>Burnt area (BA) polygons for wildfires with BA > 100 ha</u>	<u>To compute burnt area metrics (Table 2)</u>

375

376 **2.4 Linking wildfires with weather and land use**

377 The relationship between wildfires, weather and land use was based on derived data, processed
378 as described in the following lines. The starting and ending dates of each wildfire were
379 fundamental to attribute the DSR to each BA. The dating process of the BA polygons relied on
380 MODIS satellite data and the methodology of Benali *et al.* (2016). It was possible to estimate
381 the starting and ending dates as well as ignition location for 2016 wildfire events, corresponding
382 to 92% of the initial total BA.

383 Daily DSR was computed for the study period and all ERA5-Land grid points within the
384 territory of Continental Portugal. In the case of the analysis carried out for the entire mainland
385 Portugal, the value of the DSR_p associated to each wildfire was the maximum value of DSR
386 registered in the area affected during the duration of the wildfire. When the analysis carried out
387 based on the municipalities, the procedure is similar with one exception: when a wildfire
388 affected more than one municipality, the BA in each municipality was allocated to this
389 administrative unit and analysed as single wildfire event. The division of the BA between
390 affected municipalities can introduce noise in the data since artificially generates BA with
391 relatively small size but high or very high DSR_p. To circumvent this potential problem, we
392 decided to analyze BA percentages, which reduce the influence of small wildfires on the final
393 results.

Table 2. Burnt area metrics used in the manuscript, including acronym, definition and spatial scale of application/use.

Burnt area metric	Definition	Scale
<u>Total Burnt Area (TBA)</u>	$TBA = \sum_{i=1}^n BA_i$ <i>n</i> =total number of wildfires	<u>National and Municipal</u>
<u>Log(accumulatedBA)</u>	$Log(accumulatedBA) = Log\left(\sum_{i=1}^n BA_i\right)$ <i>n</i> =total number of wildfires (sorted by correspondent DSR _p)	<u>National</u>
<u>Fraction of Total Burnt Area (FTBA)</u>	$FTBA = 100 - \left(\frac{\sum_{i=1}^m BA_i}{TBA} \times 100\%\right)$ <i>m</i> =number of sampled wildfires	<u>National and Municipal</u>
<u>DSR percentile associated to 90% of TBA (DSRp90TBA)</u>	$DSRp90TBA = DSRp(0.90 \times TBA)$	<u>National and Municipal</u>
<u>DSR percentile associated to 80% of TBA (DSRp80TBA)</u>	$DSRp80TBA = DSRp(0.80 \times TBA)$	<u>National and Municipal</u>
<u>Burnable Area (BNA)</u>	$BNA = \frac{\text{Area of burnable land cover type}}{\text{Total area}} \times 100\%$	<u>Municipal</u>
<i>BNAF/BNAS</i>	$\frac{\text{Area of forest}}{\text{Area of Shrubland}}$	<u>Municipal</u>
<i>TBAF/TBAS</i>	$\frac{\text{TBA in Forest}}{\text{TBA in Shrubland}}$	<u>Municipal</u>
<u>Burnt Area in Forest (BAF)</u>	$BAF = \sum_{i=1}^f BA_i \text{ in forest areas}$ <i>f</i> =number of wildfires occurred in forest	<u>Cluster</u>
<u>Burnt Area in Shrubland (BAS)</u>	$BAS = \sum_{i=1}^s BA_i \text{ in shrubland areas}$ <i>s</i> =number of wildfires occurred in shrubland	<u>Cluster</u>
<u>Burnt Area in Agriculture (BAA)</u>	$BAA = \sum_{i=1}^a BA_i \text{ in agricultural areas}$ <i>a</i> =number of wildfires occurred in agriculture	<u>Cluster</u>

We only selected (175) municipalities (from 278) affected by more than three wildfires and TBA > 500 ha. Restricting the analysis to the administrative units with sufficient data aims to increase results' robustness and prevent potential interpretation errors. The selection of the maximum value of DSR to associate with wildfires is justified by the low spatial variability of the DSR, the small size of administrative units and the native reanalysis data resolution (C3S, 2020).

403 To achieve the first objective, we start by making and analysing plots of BA metrics vs DSRp
404 (Table 2) for all the 2016 large wildfires occurred in mainland Portugal during the study period,
405 by this order:

406 1) We firstly compared the BA values with DSRp and analysed it.

407 2) Those results lead us to sort BA data by the respective DSRp, compute accumulated values
408 of BA, normalize it using the natural logarithm and plot against DSRp to assess if this
409 relationship is linear.

410 3) Subsequently, we analysed if a fixed threshold of DSR for extreme days - DSR90p - is
411 adequate to estimate extreme fire weather and is well related to large FTBA, for the entire
412 territory. It is important to note that FTBA was calculated as the difference between 100 and
413 the percentage of TBA correspondent to a certain DSRp (Table 2). This methodology was made
414 with the purpose to visualize the TBA that burns above a DSRp threshold. ~~LULC data can limit~~
415 ~~the analysis and affect the obtained results. LULC changed during the 19 years (2001—2019)~~
416 ~~of the study period in many locations, including in the BA polygons. Effectively, Meneses *et*~~
417 ~~*al.*, (2018) observed that the main land-use changes, for the 1990—2012 period, are related to~~
418 ~~reductions in forests and agricultural areas, together with increases in urban areas, with~~
419 ~~relatively small changes between 2000—2006 and 2006—2012 periods.~~ We considered the
420 correspondent 80% and 90% of FTBA as sufficient to classify DSRp as the extreme threshold,
421 justified by the results of Pereira *et al.*, (2005), which showed that 80% of TBA occurs in 10%
422 of summer days.

423 ~~Therefore, LULC changes do not significantly affect the findings, knowing that we only use~~
424 ~~LULC data for one year/inventory to assess wildfire selectivity. Understory vegetation is also~~
425 ~~a very important factor in fire vulnerability, spread and intensity (Espinosa *et al.*, 2019; Fonseca~~
426 ~~and Duarte, 2017). Consequently, wildfires only tend to occur and spread in managed forests~~

427 with very high DSR, higher than in unmanaged forests (Fernandes et al., 2019). However, land
 428 use data does not include forest management information. Despite the small fraction of managed
 429 forested areas, roughly 20%, as estimated by Beighley and Hyde, (2018), this lack of
 430 information can influence our results, particularly in the municipalities with a significant share
 431 of managed forest area.

433 2.35 Cluster Analysis

434 Potential clustering was assessed using the curves of FTBA vs. DSR_p for all the selected
 435 municipalities. The high number (278175) of these administrative regions ~~diffieult~~complicates
 436 the interpretation of the results. Therefore, cluster analysis was performed to identify the major
 437 macro-scale spatial patterns and to objectively and statistically assess the significant differences
 438 between the results obtained for different municipalities.

439 The following notation was adopted to describe the linkages (the distance between two clusters)
 440 used in the *complete* clustering method (The MathWorks Inc, 2021):

- 441 • Cluster r is formed from clusters p and q .
- 442 • n_r is the number of objects in cluster r .
- 443 • x_{ri} is the i th object in cluster r .
- 444 • *Complete linkage* (d), also called the *farthest neighbour*, which uses the largest distance
 445 between objects in the two clusters (Eq.1).

$$446 \quad d(r, s) = \max \left(\text{dist}(x_{ri}, x_{sj}) \right), i \in (1, \dots, n_r), j \in (1, \dots, n_s) \quad (1)$$

447 A distance metric is a function that defines the distance between two observations. The
 448 ~~Matlab~~MATLAB function *pdist* used in this study, which ~~computescan~~ compute the pairwise

449 distance between pairs of observations, ~~supports various distance with different~~ metrics. We
 450 used the correlation distance because it provides a more easily interpretable dendrogram.

451 Given an m -by- n data matrix X , which is treated as m (1-by- n) row vectors x_1, x_2, \dots, x_m , the
 452 correlation distance between the vector x_s and x_t are defined as in Eq.2:

$$453 \quad d_{st} = 1 - \frac{(x_s - \bar{x}_s)(x_t - \bar{x}_t)'}{\sqrt{(x_s - \bar{x}_s)(x_s - \bar{x}_s)'}\sqrt{(x_t - \bar{x}_t)(x_t - \bar{x}_t)'}} \quad (2)$$

454 where \bar{x}_s is described in Eq.3:

$$455 \quad \bar{x}_s = \frac{1}{n} \sum_j x_{sj} \quad \text{and} \quad \bar{x}_t = \frac{1}{n} \sum_j x_{tj}. \quad (3)$$

456 The selected ~~(1-R²)~~ threshold was 0.35, meaning that the coefficient of determination in
 457 the municipalities within the same cluster is higher than 0.65. This value was selected after a
 458 benchmarking analysis of the obtained dendrograms and results from an intended balance
 459 between the correlation between municipalities and the total number of clusters. For example,
 460 on one hand, if we have chosen 5 clusters, the correspondent correlation between municipalities
 461 within the same cluster will be larger than 0.5, a value that we considered too low for this
 462 analysis. On the other hand, for a higher correlation, for example, 0.75, which corresponds to
 463 ~~1-R²=0.25~~ $1 - R^2 = 0.25$, the number of clusters will be much higher, increasing the difficulty
 464 of interpreting ~~the~~ maps and dendrogram.

465 ~~Algorithms were processed with Matlab software.~~

466

467 **2.46 The influence of ~~the type of vegetation~~ on the fire-weather relationship**

468 The ~~LULC was related to BA to accomplish the third objective of the study by computing~~
 469 ~~several metrics (Table 2), namely: (i) the burnable area (BNA) in each municipality~~ was
 470 computed as the total burnable area (sum of the land cover types that are susceptible to burn
 471 based on the land cover map) in: (ii) the 2001–2019 period, divided by the total area of the

472 municipality, and presented in percentage. LULC was related to TBA by computing the TBA
 473 in the 5 classes of vegetation, namely: in forests; (BAF), shrublands; (BAS), agriculture; (BAA),
 474 agroforestry and others: other vegetation types; (iii) the ratio between forest and shrublands
 475 BNA (BNAF/BNAS) and TBA (TBAF/TBAS). Computations were made for each analysed
 476 municipality and cluster, ~~to accomplish the third objective. Two additional ratios were~~
 477 ~~computed for each municipality, the first between forest and shrublands BNA and the second~~
 478 ~~between forest and shrublands TBA.~~ Moreover, the spatial distribution of prevailing land-use
 479 types that were most affected by wildfires was investigated to identify which municipalities
 480 have a BA in forests larger than 50% or BA in shrublands larger than 40% of TBA. The adoption
 481 of different thresholds for BA in forests and shrublands is due to a much lower area of
 482 shrublands (12%) than of forests (39%) in continental Portugal (IGFDGT, 2019).

483 ~~A contingency~~Contingency table, accuracy metrics and statistical measures of association were
 484 used to analyse the influence of the type of vegetation cover on the relationship between DSRp
 485 and TBA. The contingency table contains the number of municipalities that ~~are characterized~~
 486 ~~by diverse~~belong to a different group of clusters, i.e., different DSRp thresholds at 90% of TBA
 487 (DSRp90TBA) and, ~~therefore, a different group of clusters. are characterized by BAF > 50%~~
 488 ~~or BAS + BAA > 40%.~~ The objective was to relate the municipalities (within the groups of
 489 clusters) with TBA in diverse vegetation cover types, ~~taking into consideration that pre-~~
 490 ~~conceived relationships must be made. These statistics. Statistical measures of association~~ were
 491 used for classification accuracy against a reference as, for example, municipalities with higher
 492 DSRp90TBA will have the largest TBA in forested areas, compared with other land use types;
 493 and accuracy metrics were computed according to this initial classification. ~~A contingency table~~
 494 ~~needs, at least, two rows and two columns and, therefore, two relationships.~~

495 The list of accuracy metrics includes: (i) the Overall Accuracy (OA), which represents the
 496 samples that were correctly classified and are the diagonal elements in the contingency table,


497 from top-left to bottom-right (Alberg *et al.*, 2004); (ii) the User's Accuracy (UA), or reliability,
498 that is indicative of the probability of a sample that was classified in one category belongs to
499 that category; and, (iii) the Producer's Accuracy (PA), represents the probability of a sample
500 being correctly classified (Congalton, 2001). Statistical measures are: the Chi-squared (χ^2) test
501 (Greenwood and Nikulin, 1996), which test the independence of two categorical variables; the
502 Phi-test (Φ) or phi coefficient (David and Cramer, 1947) is related to the chi-squared statistic
503 for a 2×2 contingency table, and the two variables are associated if $\Phi > 0$. Lastly, we computed
504 the Cohen's Kappa coefficient, firstly presented by Cohen (1960) and recently analysed by
505 McHugh (2012), that measures the interrater agreement of the two nominal variables. This
506 coefficient ranges from -1 to 1 and is interpreted as < 0 indicating no agreement to 1 as almost
507 perfect agreement.

508

509 3. Results

510 3.1 Patterns Linking wildfires with weather, at the national level

511 The scatter plot of ~~DSR vs BA does not reveal a simple robust relationship between these two~~
512 ~~variables, as visible in Figure a function of DSRp (Fig. 2, where-) reveals that most of large~~
513 ~~wildfires, including those with the logarithm of highest amounts of BA, were registered with~~
514 ~~the BA - Log(BA) - is plotted against highest values of DSRp. For low DSR values, e.g. below~~
515 ~~the percentiles of DSR. Effectively 80th percentile, the coefficient of determination, r^2 , is very~~
516 ~~low (0.04).~~




517

518 Figure 2: Scatterplot of vast majority of BA are the decimal logarithm of lowest in the burnt area (Log(BA))
 519 vs DSR percentile (DSRp), for each individual fire (blue circles), considering the fires with an area larger than 100
 520 ha that occurred between May 15 and October 31, in the 2001–2019 period. Best fit (red line) and r-square are
 521 also presented. 2016 sample values.

522


523 However In addition, the scatter plot of the decimal natural logarithm of the accumulated BA
 524 versus DSRp for the entire mainland Portugal territory (Figure (Fig. 3) showed presents a linear
 525 relationship, with a very high coefficient of determination ($R^2=0.94$) and p-value lower than
 526 the significance level. Nevertheless Furthermore, the increase logarithm of Log (accumulated
 527 BA) is exponential (with r^2 increases exponentially ($R^2=0.92$) for DSRp extreme values
 528 (DSR>DSR90p), meaning that BA rises suddenly with extreme meteorological conditions.



529

530 Figure 3: In summary, the results of these analysis reveal that: (i) wildfires can occur with a large
531 spectrum of DSRp values, in extended summer; and (ii) very large wildfires only occur with
532 high DSRp.

533




534

535

536


Figure 2. Scatterplot of the burnt area (BA) vs. DSR percentile (DSRp) for wildfires (blue circles) with BA > 100 ha that occurred between May 15 and October 31, in the 2001 – 2019 period.




537
 538 **Figure 3.** Scatterplot of the decimal logarithm of the accumulated burnt area (Log(accumulatedBA)) vs DSR
 539 percentile (DSRp), considering the fires with an area larger than 100 ha that occurred between May 15 and October
 540 31, in the 2001 – 2019 period. The blue circles represent each individual fire, with respective accumulated BA,
 541 after being sorted by the assigned DSRp. Best fit (red line), respective equation, r -square R -squared and p-value
 542 are also presented.


543 The analysis of the dependence of FTBA with DSRp in the entire mainland Portugal territory
 544 (Figure Fig. 4) revealed that most of the TBA occurred with very high DSRp values. For
 545 example, for days with $DSR > 50$ th DSRp (DSR50p) the FTBA is almost 100%, meaning that
 546 fires in days with lower DSR have a negligible impact on TBA. (please see Section 2.4). Fires
 547 in days with DSRp between 85 and 95 were responsible for more than 80% of TBA in the
 548 2001 – 2019 period, making this a good DSRp threshold for extreme days. This result justifies
 549 using the DSR90p at the national scale, which is widely used for a threshold of extreme values
 550 (Bedia *et al.*, 2012; Carvalho *et al.*, 2008; Fernandes, 2019; Silva *et al.*, 2019). However, if the

551 ~~analysis is performed at a higher spatial resolution, namely at the municipality level, some~~
552 ~~differences become apparent (Figure 5).~~






553



554

555 **Figure 4:** Fraction of total burnt area (FTBA) vs DSR percentile (DSRp), computed for mainland Portugal, in the
 556 2001 – 2019 period. The circle (cross) is the DSRp when the FTBA reaches 80% (90%).



557
 558 **Figure 5:** DSR percentile (DSRp) for 80% (left panel) and 90% (right panel) of the fraction of total burnt area
 559 (FTBA) in each municipality.


560

561 3.2 Linking wildfires with weather and land use, at the municipality level

562 However, if the analysis is performed at a higher spatial resolution, namely at the municipality
 563 level, some differences become apparent (Fig. 5). The spatial distribution of DSRp for
 564 FTBA=80% (DSRp80TBA) or FTBA=90% (DSRp90TBA) (Figure 5) in each municipality
 565 presents important differences between regions, together with more visible contrasts in
 566 DSRp90TBA than in DSRp80TBA. The much lower values of DSRp in the north-eastern (*Alto*
 567 *Tâmega, Terras de Trás-os-Montes, Douro* and northern *Beiras e Serra da Estrela*) and in the

568 southern interior regions (*Alentejo Central* and *Baixo Alentejo*) should be highlighted.

569 DSRp90TBA is ~~very high~~higher in most of the coastal and ~~in some of~~ central hinterland
570 municipalities (portions of *Área Metropolitana do Porto*, *Viseu Dão-Lafões*, *Região de*
571 *Coimbra*, *Beira Baixa* and *Região de Leiria*), reaching values similar to the mean country level
572 value (85 – 95). In some ~~NUTSIII provinces~~municipalities of the northern and central
573 hinterland, DSRp90TBA is between 60 and 70 ~~in most of the municipalities~~, particularly in
574 *Douro* and *Terras de Trás-os-Montes*. It is important to underline that DSRp80TBA >
575 DSRp90TBA which is a consequence of the adopted methodology to perform this analysis
576 (please see ~~section~~Section 2.24). This also helps understand why DSRp=50 is associated with
577 FTBA=100% (~~Figure 4). The spatial distribution of DSRp80TBA and DSRp90TBA suggests~~
578 ~~the existence of municipality clustering.~~Fig. 4).



579
 580 **Figure 5:** DSR percentile (DSRp) for 80% (left panel) and 90% (right panel) of the fraction of total burnt area
 581 (FTBA) in each municipality.

582


583 **3.2 Patterns at the municipality level**

584 WeThe spatial distribution of DSRp80TBA and DSRp90TBA suggests the existence of
 585 clustering. Therefore, we explored other features of wildfiresthe fire regime in mainland
 586 Portugal, tonamely BA metrics (Table 2) that could explain the similarities and differences
 587 observed in DSRptheir patterns at the municipality level. BurnableThe burnable area (BNA),
 588 the ratio of Forest/Shrublands BNA, andthe ratio of Forest/Shrublands TBA in each
 589 municipality were assessed and analysed (FigureFig. 6). Additionally, the number of wildfires
 590 and the TBA/BNA ratio in each municipality were also evaluated (see Appendix).

591 ~~Burnable area (Figure-~~The BNA (Fig. 6a) is much lower in coastal municipalities (except in
 592 *Algarve*) and in most of the northern and central hinterland, particularly in *Terras de Trás-os-*
 593 *Montes*, *Douro* and portions of *Beiras e Serra da Estrela*. ~~Those~~These relatively low values are
 594 explained by the high ~~density of~~population density and urban areas near the coastline ~~and~~or by
 595 agriculture patches in the countryside. On the other hand, higher ~~burnable areas~~BNA are
 596 presentfound in the mountain ranges, especially in the northwest (some municipalities located
 597 in *Alto Minho*, *Cávado* and *Alto Tâmega*) as well as in some specific highly forested regions in
 598 central hinterland (within *Área Metropolitana do Porto*, *Viseu Dão-Lafões*, *Região de Coimbra*,
 599 *Região de Leiria*, *Médio Tejo* and *Beira Baixa*) and one municipality in *Algarve*. These patterns
 600 are justified by low population density, low availability of land suitable for agriculture, and, in
 601 some regions, extensive forest plantations.

602 ~~Results (Figure-6b) also~~The Forest/Shrublands BNA (Fig. 6b) show that forest cover is
 603 prevalent in most of the analysed municipalities, ~~with special intensity ones~~especially near the
 604 west coast. Conversely, shrublands BNA is ~~more~~dominant in a few municipalities located in
 605 the northern hinterland, particularly ~~situated in~~*Alto Minho*, *Alto Tâmega*, *Douro* and *Beiras e*
 606 *Serra da Estrela*. ~~Results are considerably different analysing~~However, the spatial distribution
 607 of the Forest/Shrublands TBA (Figure-Fig. 6c), ~~with~~ present some considerable differences,
 608 namely an extensive ~~amount~~number of municipalities at the north, including coastal and inland,
 609 that have larger TBA in shrublands (a large number of municipalities are located in *Alto*
 610 *Tâmega*, *Tâmega e Sousa*, *Douro*, *Viseu Dão-Lafões* and *Beiras e Serra da Estrela*).
 611 Nevertheless, the municipalities with higher Forest/Shrubland BNA correspond with those with
 612 larger ratios of Forest/Shrubland TBA. Results of both maps are similar when analysing the
 613 southern provinces of the country (*Alto Alentejo*, *Alentejo Central*, *Alentejo Litoral*, *Baixo*
 614 *Alentejo* and *Algarve*), where almost all municipalities are characterized by higher forest BNA
 615 and TBA.

616



617

618 **Figure 6:** a) Burnable area (BNA), in percentage; b) Forest/Shrubland BNA and c) Forest/Shrubland total burnt
 619 area (TBA); all in the 2001 – 2019 period, for the selected municipalities.

620


621 *Other municipalities also highly affected by fires are located in the extreme northwest (Alto*
 622 *Minho), surrounding mountain ranges in the northwest (Área Metropolitana do Porto and*
 623 *Tâmega e Sousa), and in the south (Alto Alentejo and Algarve). By contrast, the lower BA*
 624 *percentages occur in most of the southern provinces (except Algarve) and the northeast (Terras*
 625 *de Trás-os-Montes). The largest TBA/BNA is observed in mountains ranges and forested*
 626 *regions of central hinterland, particularly in parts of Viseu-Dão-Lafões, Beiras e Serra da*
 627 *Estrela, Região de Coimbra, Região de Leiria, Médio Tejo and one municipality in Algarve. In*
 628 *some of these municipalities, this value is >100%, meaning that in the 19 years TBA is larger*
 629 *than BNA and, consequently, there were a large number of recurrent wildfires in those areas.*

630


631 **3.3 Cluster analysis patterns**

632 ~~Based on the relationship between TBA and DSRp~~The spatial distribution of DSRp80TBA and
633 ~~DSRp90TBA suggests the existence of clustering, which should also help explaining the feature~~
634 ~~similarities or differences between municipalities. Therefore,~~ the municipalities were grouped
635 in ten clusters. ~~However, the based on the relationship between TBA and DSRp. The obtained~~
636 dendrogram (Figure-Fig. 7) discloses that cluster 10 is ~~isolated, with only~~composed by just one
637 municipality; and, therefore, ~~can be eliminated~~was removed from further analysis. ~~Cluster~~
638 ~~numbers are sorted by descending order of the DSRp90TBA, i.e., 90% of TBA was registered~~
639 ~~with DSRp larger than this value. Cluster 2 includes the largest number of municipalities (23%~~
640 ~~of total) and highest TBA, almost 500,000 ha (26% of total). Generally, clusters group 13 or~~
641 ~~more municipalities, except for clusters 3 and 8, with only 5 and 6 municipalities, respectively.~~
642 ~~Each cluster represents between 8% and 16% of the total TBA for the study period, except for~~
643 ~~the two smaller clusters, where TBA is only 1% of the total.~~

644




645



646

647 **Figure 7:** Dendrogram results: cluster colours are the same as in [FigureFig. 6](#), for better identification. X axis
 648 numbers are the cluster numbers. Y axis is $(1-r^2R^2)$, where r is the correlation coefficient between FTBA and
 649 DSRp. The blue line is the clustering threshold, at 0.35. Each vertical line is a municipality.



650
 651 **Figure 8:** Clusters spatial distribution. Cluster colours are the same as in [Figure-Fig.7](#). Municipalities without
 652 colour were excluded from the cluster analysis, justifying only 5.2% of TBA.

653
 654 The spatial pattern of [FigureFig. 8](#) reveals a relatively homogeneous distribution of the
 655 municipalities of equivalent clusters, meaning that municipalities with similar DSRp are often
 656 neighbours. In general, patches of municipalities belonging to consecutive clusters are
 657 observed.

658 The FTBA vs. DSRp plots were produced for each cluster to illustrate and interpret the
 659 clustering results (Fig. 9). FTBA=100% occurs for DSR90p in cluster 1, confirming that large

660 wildfires in these municipalities only occurred with very extreme meteorological conditions.

661 The FTBA vs. DSRp curves for the first three clusters present a very steep slope for the highest

662 DSRp values (Figure 9), revealing that large wildfires take place in the municipalities of these

663 clusters in days with high DSRp (above 90). Moreover, the FTBA vs. DSRp plots for these

664 clusters present very low dispersion suggesting that the curves for the municipalities of each of

665 these clusters are very similar. These municipalities are located in north and central western

666 coastal areas, also include mountain ranges (predominantly in *Alto Minho*, *Cávado*, *Área*

667 *Metropolitana do Porto*, *Tâmega e Sousa*, *Região de Aveiro*, *Região de Coimbra* and *Alentejo*

668 *Litoral*), within some central and south hinterland regions (parts of *Viseu Dão-Lafões*, *Beiras e*

669 *Serra da Estrela*, *Médio-Tejo* and *Alto Alentejo*) and in the south coast (almost all of *Algarve*).

670 Clusters 4, 5 and 6 are prone to burn with less extreme conditions, where the median of DSR90p

671 corresponds to 85 – 90% of TBA. The slope of FTBA vs DSRp curves is less steep but the

672 dispersion is higher than the previous clusters, ~~and dispersion is higher in these clusters, with~~


673 ~~more municipalities where meaning that large~~ wildfires can occur with lower values of DSRp.

674 Both features suggest that in these clusters, wildfires tend to occur in a widest range of

675 meteorological conditions. These clusters are spread throughout the country and can be viewed

676 as a transition between the group of clusters with extreme (1, 2 and 3) and less extreme (7, 8


677 and 9) DSRp80TBA or DSRp90TBA.



678
 679 **Figure 9:** Fraction of total burnt area (FTBA) vs DSR percentile (DSRp), for the municipalities of each of the 9
 680 clusters. The black line is the median of all curves in each cluster. The shaded area is defined by the maximum and
 681 minimum curves in each cluster.

682 Clusters 7, 8 and 9 can be considered as the group of lower DSRp clusters, due to the relatively
 683 lower values of the DSRp80TBA or DSRp90TBA, which range from 70 to 80%. **Additionally,**
 684 **higher curve**Higher dispersion is also apparent, especially in cluster 9, which integrates
 685 municipalities where large wildfires can occur with lower values of DSRp (in some cases, below
 686 DSR50p). In this group of clusters, the slope of the FTBA vs DSRp curves, at higher values of
 687 DSRp is the lowest, especially in clusters 8 and 9. Nevertheless, the median curve of cluster 8
 688 has a different behaviour, compared to the other two clusters: the steeper interval is between
 689 70th and 80th percentile, meaning that a larger amount of BA occurs in less extreme conditions.
 690 The municipalities within these clusters are mostly located in the northern and central
 691 hinterland, particularly in *Alto-Tâmega*, *Terras de Trás-os-Montes*, *Douro*, *Beiras e Serra da*

692 *Estrela* and *Beira Baixa*. Additionally, a few municipalities within these clusters belong to
 693 *Alentejo Central* and *Baixo Alentejo*, two provinces with a scarce number of [fireslarge wildfires](#)
 694 and BA.




695
 696 **Figure 9:** Fraction of total burnt area (FTBA) vs DSR percentile (DSRp), for the municipalities of each of the 9
 697 clusters. The black line is the median of all curves in each cluster. The shaded area is defined by the maximum and
 698 minimum curves in each cluster.

699
 700 Box-plots of the DSRp80TBA and DSRp90TBA for the municipalities of each cluster
 701 ([FigureFig. 10](#)) are consistent with the previous results. Dispersion is considerably [much](#) higher
 702 in the latter than in the former case, especially in clusters 3, 7 and 8. In some municipalities of
 703 clusters 7 and 8, large wildfires, with the ability to exceed FTBA=10% ([FigureFig. 9](#)), start to
 704 occur with relatively low values of DSRp. Another notable difference is the boxplot medians:
 705 for DSRp90TBA they decrease with the ascending number of clusters as expectable, but not for

706 DSRp80TBA, where they increase between clusters 4 and 5, between 6 and 7, and between 8
 707 and 9.

708



709

710 **Figure 10:** Boxplots for DSRp80TBA (top panel) and DSRp90TBA (bottom panel), i.e., the DSR when the
 711 municipality curves reach associated to 80% (top) and 90% (bottom) BA of TBA, respectively, for the 9 clusters.

712 The central line is the median; the edges of the box are the 25th and 75th percentiles; and, the plus signs ~~represent~~
713 ~~the outliers~~represents outliers, defined as a value that is more than three scaled median absolute deviations away
714 from the median.

716 **3.4 Major drivers**


717 **3.4 The influence of vegetation on the fire-weather relationship**

718 The spatial distribution of the clusters resembles the general pattern of LULC in Portugal
719 (~~Figure~~Fig. 11, bottom panel). In general, municipalities with high DSRp90TBA are located in
720 regions of forests while municipalities with lower DSRp90TBA are located in regions where
721 shrublands tend to be predominant. Analysis of BA in LULC type ~~analysis~~, made for each
722 cluster, indicates that BA in forests (BAF) is notably higher than in shrublands (BAS), for the
723 first five clusters than for the last four clusters (~~Figure~~Fig. 11, top panel). This means that BAF
724 is higher for clusters with higher DSRp90TBA while BAS is higher for clusters with lower
725 DSRp90TBA. In addition, there is an increase in the fraction of BA in agricultural land
726 associated with the decrease of DSRp90TBA. This amount is ~~larger or very close to~~higher and
727 about 10% – 20% in clusters 6- ~~– 9~~and, but lower in clusters 1- ~~– 5~~.


728 Results show marked pieces of evidence between ~~most~~-coastal and ~~the~~ northern/north-~~eastern~~
729 hinterland municipalities, which present similar DSRp90TBA and, therefore, similar cluster
730 distribution. Highest BAF characterizes the majority of the municipalities with the observed
731 highest DSRp at 90% of TBA (generally above 85) while the territory with higher BAS is also
732 characterized by lower DSRp90TBA (below 85). These clusters (7-9) also present relatively
733 high percentages of BA in agriculture (mostly between 10 and 20%). It is also worth mentioning
734 that some municipalities present similar BAF and BAS, although being located in the coastal
735 regions, usually characterized by higher forest cover.

736 The land cover also helps to understand the DSRp80TBA and DSRp90TBA boxplots for each
 737 cluster, especially the higher dispersion in the latter in comparison with the former (FigureFig.
 738 10). These dissimilarities are especially evident in cluster 8, which is the cluster with the highest
 739 BA in shrublandsBAS and agricultureBAA (twice the value of clusters 1 – 5) and less in
 740 forestBAF (half the value of clusters 1 – 5). Additionally, cluster 8 is the one with a less
 741 burnable-areaBNA (not shown). ~~The combination of these factors could explain the high~~
 742 ~~dispersion: high BA in shrublands can occur~~

743



744



745
 746 **Figure 11.** Top: Burnt area in three land use types: forest, shrublands and agriculture; represented for each
 747 cluster, identified by the respective DSRp and also by letter c. Bottom: Municipalities with Burnt Area in
 748 Forest>50%, Shrublands>40% or other cases. Municipalities without colour were excluded from the cluster
 749 analysis.

750 ~~with~~ The combination of these factors could explain the high dispersion: high BAS can occur
 751 with low DSRp, high ~~BA in agricultural lands~~ BAA is much more likely to occur with high
 752 DSRp; and, finally, low ~~burnable areas~~ BNA prevent very large wildfires to occur, even with
 753 extreme DSRp.

754 A contingency table permitted to objectively and quantitatively assess the influence of
 755 vegetation cover in the spatial distribution of the clusters and, therefore, also in DSRp90TBA.
 756 Table 43 is based on the results ~~illustrated~~ depicted in ~~Figure~~ Fig. 11 and aims to assess if the
 757 differences in groups of clusters or DSRp90TBA can be explained by the BA prevailing in
 758 forested areas or shrubland+agricultural zones. Specifically, it purposes to assess if

759 municipalities of clusters 1 – 5, with $DSRp90TBA > 90$, have higher BAF ($BAF > 50\%$), and, on
 760 the contrary, clusters 7 – 9, with $DSRp90TBA < 90$, present higher BAS+BAA
 761 ($BAS+BAA > 50\%$).

762 Results reveal that the number of municipalities of clusters 1-5 and $BAF > 50\%$ is 4.6 times
 763 higher than the number of municipalities in clusters 7-9 and $BAF > 50\%$. However, the number
 764 of municipalities of clusters 7-9 and $BAS+BAA > 50\%$ is 1.3 higher than the number of
 765 municipalities of clusters 1-5 and $BAS+BAA > 50\%$. Consequently, the OA (71%), UA
 766 (71% – 70%) and PA (82% – 55%) reveal moderate to high accuracy. The $BAS+BAA > 50\%$
 767 threshold is probably a too demanding criterion for $DSRp90TBA = 90$ limit, as shrublands and
 768 agricultural land cover will also burn with higher $DSRp$ in a large number of municipalities.
 769 For forests ($BAF > 50\%$), the accuracy is better, i.e., this threshold has been accurate in more
 770 than four times of the municipalities that were incorrectly classified. The Cohen's Kappa test
 771 allows to conclude a fair agreement ($\kappa = 0.3828$) and reject the null hypothesis: observed
 772 agreement is not accidental (Landis and Koch, 1977). The Φ and C tests also corroborated that
 773 variables are dependent, with similar values, 0.39 and 0.36, meaning moderate correlation
 774 (Frey, 2018) and the existence of a relationship (De Espindola *et al.*, 2009), respectively.
 775 However, the χ^2 test results indicate that we can claim that the samples are independent (Frey,
 776 2018), with an error risk of about $4e-06$.

777 **Table 4.3.** Contingency tables and accuracy metrics to assess the role of vegetation BA assessed with $DSRp90BA$
 778 thresholds, for the municipalities used in cluster analysis. The contingency tables computed the number of
 779 municipalities (NM) for the following criteria: CLUST 1-5 (CLUST 7-9) and $BAF > 50\%$ ($BAS+BAA > 50\%$).
 780 Overall Accuracy (OA), User's Accuracy (UA) and Producer's Accuracy (PA) were the calculated accuracy
 781 metrics, together with the statistical tests Chi-squared (χ^2) test (with p-value), Phi coefficient (Φ), Contingency
 782 coefficient (C) and the Cohen's Kappa coefficient (κ).

NM	$BAF > 50\%$	$BAS+BAA > 50\%$
CLUSTERS 1-5	65	27

CLUSTERS 7-9	14	33
OA	71%	
UA	71%	70%
PA	82%	55%
χ^2	21.175 (4E-6)	
Φ	0.390	
C	0.363	
K	0.383	

783

784 Thus, three out of four computed statistics prove a dependent relationship and, consequently,
785 we can state that the cluster's spatial distribution patterns are correlated with vegetation type.

786

787 4. Discussion

788 The scatter plot of ~~BA vs DSR vs BA does not reveal a simple robust clearly~~ illustrate the
789 relationship between these two variables (Figure Fig. 2). ~~This fact can be explained by several~~
790 ~~reasons (e.g., firefighting activities, geographical/landscape features, fuel breaks, limitations of~~
791 ~~the Fire Weather Index System, etc.) but, in essence, the most important~~ one is that the
792 wildfire activity does not only depend on the weather. This means that: (i) ~~hand, large~~ wildfires
793 can occur in days with a wide range of relatively low values of DSR; (ii) ~~small wildfires can~~
794 ~~occur in days of high DSR, DSRp (<80)~~ due to several reasons including rapid fire-
795 suppression activities (e.g., firefighting) or ~~other~~fuel constraints (especially fuel). However, it
796 ~~is well known that~~ e.g., fuel breaks, geographical and landscape features). On the other hand,
797 extreme large wildfires only occur in days of extreme fire weather as pointed out by several
798 studies (Fernandes *et al.*, 2016). ~~These facts are validated by~~ According to our results, ~~revealing~~
799 ~~that~~ only 6% of the ~~Total Burnt Area (TBA)~~ occurs with DSRp<80 and 12% of TBA are
800 registered in wildfires with DSRp<90. ~~These reasons explain all the main features of Figure 2,~~
801 ~~namely: small wildfires are registered in days with almost all values of DSR, although the much~~

802 ~~small number of wildfires in the lower left quarter of the plot area, and the huge number of~~
803 ~~events near the right vertical axis, especially for $DSR > DSR_{90p}$. In effect, DSR seems to act as~~
804 ~~a limiting or conditioning factor of the maximum BA.~~

805 The ~~scatter~~ plots of Log (~~accumulatedBA~~accumulated BA) and FTBA ~~versusys.~~ DSRp
806 (~~FigureFig.~~ 3 and ~~FigureFig.~~ 4) suggest that DSR_{90p} is a suitable threshold to identify extreme
807 fire weather days for the entire territory of mainland Portugal which is in line with previous
808 studies (Bedia *et al.*, 2012; Carvalho *et al.*, 2008; Fernandes, 2019; Silva *et al.*, 2019). ~~The~~
809 ~~importance of extreme weather for the occurrence of large wildfires in Portugal has been already~~
810 ~~pointed out in several studies (Calheiros *et al.*, 2020, 2021; Parente *et al.*, 2018a, 2019; Trigo *et*
811 *al.*, 2006). Large wildfires ($BA > 100$ ha) are essentially dependent on the existence of extreme
812 fire weather and small and medium size wildfires are much more dependent on the daily and
813 annual (weather/vegetation) cycles (Telesca and Pereira, 2010).~~

814 However, analysis performed at a finer spatial scale (~~FigureFig.~~ 5) discloses interesting
815 deviations, namely differences between coastal areas and the hinterland municipalities. Large
816 wildfires/high BA can occur in most of the inland municipalities in the northeast and parts of
817 southern Portugal with $DSR_p < 80$, but can only occur in coastal and some mountainous
818 municipalities with higher DSR ($DSR > DSR_{90p}$).

819 Differences in DSRp throughout the territory are expected due to distinct characteristic factors,
820 including climate and landscape features. Mainland Portugal has two slightly different types of
821 temperate (group C) climate, namely Csb (dry and warm summer) in the north and Csa (dry and
822 hot summer) in the south, which promote different fire regimes in these two regions (~~Parente *et*~~
823 ~~*al.*, 2016). In fact, patterns of $DSR_p < 80$ TBA or $DSR_p > 90$ TBA (Figure 5) strongly resemble the~~
824 spatial distribution of the type of climates in Portugal (please see Fig. 1 of AEMET (2011)), in
825 the sense that regions with higher (lower) $DSR_p < 80$ TBA or $DSR_p > 90$ TBA present Csb (Csa) type
826 of climate.

827 [\(Parente et al., 2016\)](#). LULC is also an important wildfire factor in Portugal (Barros and Pereira,
828 2014; Leuenberger *et al.*, 2018; Parente and Pereira, 2016; Pereira et al., 2014; Tonini *et al.*,
829 2018). Therefore, it is not surprising the high similarity between the spatial patterns of
830 DSRp80TBA or DSRp90TBA and the LULC maps for Portugal (e.g., please see Figure 4 of
831 Parente and Pereira (2016)). Other wildfire-related [landscapevegetation](#) features were assessed
832 [\(Fig. 6\)](#) to explain the heterogeneity of DSRp80TBA and DSRp90TBA maps [\(Figure 6Fig. 5\)](#).
833 The ratio Forest/Shrublands BNA shows higher BNA in forests in most of the territory but the
834 ratio Forest/Shrublands TBA reveals higher TBA in shrublands, especially in regions of lower
835 DSRp80TBA and DSRp90TBA. ~~We did not analyse different types of forest or shrublands~~
836 ~~separately. Land~~[These findings are in line with the higher land](#) cover proneness to wildfires ~~is~~
837 ~~higher~~ for shrublands and pine forests than for annual crops, mixed forests and evergreen oak
838 woodlands (Barros and Pereira, 2014; Pereira *et al.*, 2014). ~~Those authors also observed that,~~
839 ~~as wildfire size increases, selectivity decreases for all land cover types. These findings may be~~
840 ~~a consequence of the different impacts of the fire weather on the different land cover types~~
841 ~~which motivates further research on the role of vegetation in the spatial distribution of DSRp~~
842 ~~associated with a larger fraction of TBA.~~
843 [The cluster analysis based on the DSRp vs FTBA curves aimed to find groups of municipalities](#)
844 [with similar fire-weather relationships](#). As expected, the spatial distribution of the clusters
845 [\(FigureFig. 8\)](#) is also very similar to the DSRp80TBA and DSRp90TBA maps [\(FigureFig. 5\)](#),
846 especially the marked differences between the coastal and hinterland municipalities of the
847 northeast and south-central.
848 The curves of DSRp_ vs_ FTBA for the clusters [\(FigureFig. 9\)](#) show decreasing derivatives and
849 increasing variability with the decrease in the DSR, which means a trend for large wildfires to
850 occur with less extreme weather conditions and greater variability between the municipalities
851 of each cluster.

852 ~~The cluster analysis based on the DSRp vs FTBA curves aimed to find groups of municipalities~~
853 ~~with similar fire weather relations. Contingency tables account for the municipalities of two~~
854 ~~distinct groups of clusters in terms of DSR.~~ Contingency tables, accuracy and statistical tests
855 led us to conclude that vegetation types, particularly forest and shrublands, influence the spatial
856 distribution of DSRp observed in Portugal.

857 ~~In addition to the type of climate, the~~The different vegetation cover ~~justifies~~ is able to explain
858 the spatial distribution of DSRp within mainland Portugal and, therefore, ~~explains~~ clusters'
859 dissimilarities (FigureFig. 11). On one hand, ~~extreme~~ DSR extremes are strongly influenced by
860 long-lasting severe droughts (~~before and not only~~ during ~~but before the~~ fire season), heatwaves
861 (during fire season), or both. Heat waves and droughts are important extreme weather/climate
862 events, promoting wildfires occurrence and spread, and, therefore, ~~for TBA~~high BA (Russo *et*
863 *al.*, 2017; Parente *et al.*, 20182018a; Parente *et al.*, 2019). On the other hand, shrublands are
864 more likely to suffer from droughts than forests. As observed by Gouveia *et al.*, (2012), during
865 drought shrublands presented higher levels of dryness, whereas broad-leaved forests exhibited
866 lower water stress. Coniferous forests are more resistant to short-term droughts than broad-
867 leaved forests, because of their decreased vulnerability to xylem cavitation (Allen *et al.*, 2010).
868 Consequently, forests tend to burn only under extreme DSR values, typically caused by
869 simultaneous drought and heatwave, while shrublands (and also agricultural areas) can burn
870 with lower DSRp. These facts can be additionally justified by biological features. In the
871 Mediterranean region, precipitation is the main constrain to photosynthesis and growth (Pereira
872 *et al.*, 2007). This is particularly critical for shallow-rooted species, like those of the herbaceous
873 vegetation and some shrub species, which are unable to access groundwater. It is less critical
874 for the deeply rooted species such as cork oak, and other drought-resistant Mediterranean
875 species (Cerasoli *et al.*, 2016).

876 LULC data can limit the analysis and affect the obtained results. LULC changed during the
877 19 years (2001 – 2019) of the study period in many locations, including in the BA polygons.
878 ~~Effectively, It is important to underline that this study is not about the relationship between~~
879 ~~LULC and weather and fire occurrence. In summary, this study is about the relationship~~
880 ~~between extreme fire weather and high BA resulting from large wildfires which is spatially~~
881 ~~affected due to LULC (among other factors). Additionally, while LULC, topography,~~
882 ~~population statistics, etc. are structural (essentially fixed or stationary) wildfire hazard factors,~~
883 ~~the meteorological conditions are conjunctural (essentially variable or dynamic) wildfire hazard~~
884 ~~factors. Despite a few space-time analyses (e.g., Parente et al., 2016; Pereira et al., 2015; Vega~~
885 ~~Orozeo et al., 2012), usually, and for obvious reasons, the influence of these two types of factors~~
886 ~~on the fire incidence is studied separately.~~

887 ~~However, it was precisely as a result of an in-depth analysis of the relationship between extreme~~
888 ~~fire weather (specifically DSRp) and fire incidence (specifically BA) that it was possible to~~
889 ~~conclude that LULC—a structural factor—influences the impacts of meteorological conditions~~
890 ~~—a conjectural factor of fire risk. As far as we know, this is the first study that identifies and~~
891 ~~establishes that the relationship between fire weather and fire incidence depends on LULC, for~~
892 ~~Portugal.~~

893 ~~It is also important to underline that, to establish this relationship~~Mencses *et al.*, (2018b)
894 ~~observed that the main land-use changes, for the 1990 – 2012 period, are related to reductions~~
895 ~~in forests and agricultural areas, together with increases in urban areas, with relatively small~~
896 ~~changes between 2000 – 2006 and 2006 – 2012 periods. Therefore, LULC changes do not~~
897 ~~significantly affect the findings, knowing that we only use LULC data for one year/inventory~~
898 ~~to assess wildfire selectivity. Understory vegetation is also a very important factor in fire~~
899 ~~vulnerability, spread and intensity (Espinosa et al., 2019; Fonseca and Duarte, 2017).~~
900 Consequently, wildfires only tend to occur and spread in managed forests with very high DSR,

901 higher than in unmanaged forests (Fernandes et al., 2019). However, land use data does not
902 include forest management information. Despite the small fraction of managed forested areas,
903 roughly 20%, as estimated by Beighley and Hyde, (2018), this lack of information can influence
904 our results, particularly in the municipalities with a significant share of managed forest area.

905 It is also important to underline that, to identify the drivers of extreme burnt area in Portugal,

906 we used objective methods and adequate statistics that ensure the robustness and statistical
907 significance of the results. The description of the study carried out also includes the chronology
908 of the performed analysis. In a previous study (Calheiros *et al.*, 2020), the relationship between
909 fire weather and fire incidence was ~~analyzed~~analysed in-depth for the entire Iberian Peninsula.

910 Among other results, they found that the DSR90p is a good indicator of extreme fire weather
911 and is well related to the BA in the Iberian Peninsula. In this study, we started by verifying
912 whether the relationship between DSRp and BA found, in general terms, for the Iberian
913 Peninsula, was also verified in mainland Portugal, at municipality level, and what is the spatial
914 variability of the extreme value of DSRp above which most of the burned area is registered. To

915 objectively interpret the obtained spatial patterns (~~Figure~~Fig. 5), we complemented and
916 deepened the analysis with the use of clustering algorithms, to classify the municipalities into
917 statistically different groups in terms of the relationship between FTBA and DSRp. The
918 emerging patterns showed that all of those most likely factors, such as topography, altitude

919 (~~Figure~~Fig. 1), slope (please see ~~Figure~~Fig. 5 of Parente and Pereira, 2016), population density
920 (please see ~~Figure~~Fig. 2 of Pereira *et al.*, 2011), rural and urban area type (please see ~~Figure~~Fig.
921 3 of Pereira *et al.*, 2011), road density/distance to the nearest road (please see ~~Figure~~Fig. 2a of

922 Parente *et al.*, 2018b) and climate type (please see ~~Figure~~Fig. 1a of Parente *et al.*, 2016) were
923 not able to explain the obtained spatial patterns. The only factor with a similar spatial pattern
924 was the LULC, which is the reason why we decide to explore this possibility more deeply, with

925 contingency tables and several accuracy metrics to assess the influence of the type of vegetation
926 cover on the relationship between DSRp and TBA.

927

928 5. Conclusions

929 ~~The relationship between DSR and BA was investigated, initially revealing low correlation but~~
930 ~~presenting the highest values of BA with extreme DSR. Those results lead us to differ the~~
931 ~~analysis to accumulated Log (BA) vs DSR, revealing that they are strongly correlated and the~~
932 ~~DSR90p is an adequate threshold for an extreme BA in mainland Portugal. Nevertheless, at~~
933 ~~higher resolution, relevant differences appear among DSRp thresholds that explain 90 and 80%~~
934 ~~of the TBA. Cluster analysis shows that these differences justified the existence of several~~
935 ~~statistically significant clusters. Generally, municipalities where large wildfires occur with high~~
936 ~~or very high DSRp values are located in the north and central coastal areas, central hinterland~~
937 ~~mountainous parts and in the extreme south. In contrast, clusters where large fires were~~
938 ~~registered with low DSRp values mostly appear in the north eastern. The type of climate and~~
939 ~~vegetation cover explain the clusters' distribution pattern and the relationship between DSRp~~
940 ~~and total BA. Large wildfires tend to occur mostly in forests with very high or extreme DSRp~~
941 ~~while, in shrublands, with relatively lower DSRp. This fact is explained by the different species~~
942 ~~features, which causes that shrublands are more suitable to dryness and heatwaves than forests.~~
943 ~~The relationship between vegetation cover and DSRp was statistically validated with the~~
944 ~~contingency tables and statistical tests. Results indicate an overall accuracy of 71% and a~~
945 ~~statistical relationship between dependent variables. BNA highest values are visible in the~~
946 ~~mountainous regions between the coastal and hinterland municipalities and, oppositely, lowest~~
947 ~~values are present in urban municipalities near the coast and some hinterland regions, due~~
948 ~~mostly to agricultural patches. BNA also can influence DSRp vs FTBA curve in the~~
949 ~~municipalities and explain the high variability in DSRp in the clusters.~~

950 ~~In summary, this work disclosed that the usual 90th percentile of DSR is a good indicator for~~
951 ~~the extreme BA in mainland Portugal.~~This work disclosed that the 90th percentile of DSR,
952 usually used to identify extreme fire weather days, is a good indicator for the extreme BA in
953 mainland Portugal. However, at higher resolution, this threshold presents regional variations
954 that should be considered, namely for landscape and wildfire management.


955 This analysis of the relationship between extreme fire weather (specifically DSRp) and fire
956 incidence (specifically BA) lead us to conclude that LULC – a structural factor – influences the
957 impacts of meteorological conditions – a conjectural factor of fire risk. As far as we know, this
958 is the first study that identifies and establishes that the relationship between fire weather and
959 fire incidence depends on LULC, in Portugal.

960 The role of vegetation cover on these regional variations is an important outlook of our results.

961 ~~Shrublands are more suitable to burn in less extreme conditions than forests. However, at higher~~
962 ~~resolution, this threshold presents regional variations that should be considered, namely for~~
963 ~~landscape and wildfire management.~~ These findings could help firefighters and civil protection
964 in prevention and combat planning, more importantly knowing the reputation and operational
965 use of DSR in Portugal. Climate type and vegetation cover explain the DSRp spatial distribution
966 dissimilarities, highlighting that landscape and forest management are key factors for the
967 adaptation to future climate change.

968
969
970
971
972 **Appendix**

973 In this section, we present the results that were important but not fundamental for this
974 manuscript. The ~~Number~~number of fires in Portugal (~~Figure~~Fig. 1), in each analysed
975 municipality, were assessed. The distribution of the number of wildfires, between 2001 and
976 2019, discloses a notable contrast between north and southern provinces (the last ones
977 considered as *Alto Alentejo*, *Alentejo Central*, *Alentejo Litoral*, *Baixo Alentejo* and *Algarve*).
978 Wildfires were more frequent in the extreme northwest (*Alto Minho* and *Alto Tâmega*) and some
979 municipalities located in *Beiras e Serra da Estrela*. Wildfire frequency is much lower in the
980 south and on most of the western coast.



981
982 Figure 1: Number of fires larger than 100 ha, all in the 2001 – 2019 period, for the selected municipalities

983
984 **Data availability:** This research was developed using three public data sources. The
985 meteorological variables were obtained from the fifth generation of ECMWF atmospheric

986 reanalyses of the global climate (ERA5-Land) dataset (Copernicus Climate Change Service
987 (C3S), ~~20172020~~). ~~Land use and land cover data were provided by Portuguese national~~
988 ~~authorities (DGT, 2019), and the wildfire database from the Portuguese Institute for the~~
989 ~~Conservation of Nature and Forests. Land use and land cover data were provided by Portuguese~~
990 ~~national authorities, respectively, *Direção Geral do Território* (DGT, 2019), and the wildfire~~
991 ~~database from the *Instituto Nacional da Conservação da Natureza e das Florestas* (ICNF,~~
992 2020).

993

994 **Author contribution:** TC developed the code to analyse the data, produced the results and
995 plots, and wrote the original draft of the manuscript. AB contributed to the supervision, the code
996 to analyse data and produce plots, and also to the writing. ~~JNS contributed to the supervision,~~
997 ~~methodology and writing.~~ MP contributed to the supervision, production of plots and writing.
998 JNS contributed to the supervision, methodology and writing. JPN contributed to the
999 supervision and writing. All authors contributed to the conceptualization and methodology of
1000 this research.

1001

1002 **Competing interests**

1003 The authors declare that they have no conflict of interest.

1004

1005 **Acknowledgments**

1006 This work was funded by the Portuguese Fundação para a Ciência e a Tecnologia through the
1007 PhD fellowship attributed to T Calheiros (PD/BD/128173/2016). Additional funding was
1008 obtained through the individual research grant attributed to JP Nunes (IF/00586/2015), research
1009 project FRISCO (PCIF/MPG/0044/2018), and research unit funding attributed to the CE3C,

1010 CITAB and Forest Research Centre (CEF) research centres (UIDB/00329/2020,
1011 UIDB/04033/2020 and IDB/00239/2020, respectively).

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022 **REFERENCES**

1023 Alberg, A. J., Park, J. W., Hager, B. W., Brock, M. V. and Diener-West, M.: The use of
1024 “overall accuracy” to evaluate the validity of screening or diagnostic tests, *J. Gen. Intern.*
1025 *Med.*, 19(5 PART 1), 460–465, doi:10.1111/j.1525-1497.2004.30091.x, 2004.

1026 Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M.,
1027 Kitzberger, T., Rigling, A., Breshears, D. D., Hogg, E. H. (Ted., Gonzalez, P., Fensham, R.,
1028 Zhang, Z., Castro, J., Demidova, N., Lim, J. H., Allard, G., Running, S. W., Semerci, A. and
1029 Cobb, N.: A global overview of drought and heat-induced tree mortality reveals emerging
1030 climate change risks for forests, *For. Ecol. Manage.*, 259(4), 660–684,
1031 doi:10.1016/j.foreco.2009.09.001, 2010.

1032 Amraoui, M., Pereira, M. G., Dacamara, C. C. and Calado, T. J.: Atmospheric conditions
1033 associated with extreme ~~fi-refire~~ activity in the Western Mediterranean region, *Sci. Total*
1034 *Environ.*, 524–525, 32–39, doi:10.1016/j.scitotenv.2015.04.032, 2015.

Formatou: Inglés (Estados Unidos)

- 1035 Instituto da Conservação da Natureza e das Florestas, [online] Available from:
1036 http://www2.icnf.pt/portal/florestas/dfci/inc/mapas_2020.
- 1037 Barros, A. M. G. and Pereira, J. M. C.: Wildfire selectivity for land cover type: Does size
1038 matter?, PLoS One, 9(1), doi:10.1371/journal.pone.0084760, 2014.
- 1039 Bedia, J., Herrera, S. and Guti, J. M.: Sensitivity of fire weather index to different reanalysis
1040 products in the Iberian Peninsula, Nat. Hazards Earth Syst. Sci., 699–708, doi:10.5194/nhess-
1041 12-699-2012, 2012.
- 1042 Beighley, M. and Hyde, A. C.: Portugal Wildfire Management in a New Era: Assessing Fire
1043 Risks , Resources and Reforms, ,Beighley & Hyde (February), 52, 2018.
- 1044 Benali, A., Russo, A., Sá, A. C. L., Pinto, R. M. S., Price, O., Koutsias, N. and Pereira, J. M.
1045 C.: Determining fire dates and locating ignition points with satellite data, Remote Sens., 8(4),
1046 doi:10.3390/rs8040326, 2016.
- 1047 Bergonse, R., Oliveira, S., Gonçalves, A., Nunes, S., DaCamara, C. and Zêzere, J. L.:
1048 Predicting burnt areas during the summer season in Portugal by combining wildfire
1049 susceptibility and spring meteorological conditions, Geomatics, Nat. Hazards Risk, 12(1),
1050 1039–1057, doi:10.1080/19475705.2021.1909664, 2021.
- 1051 Calheiros, T., Nunes, J. P. and Pereira, M. G.: Recent evolution of spatial and temporal
1052 patterns of burnt areas and fire weather risk in the Iberian Peninsula, Agric. For. Meteorol.,
1053 287, 107923, doi:10.1016/J.AGRFORMET.2020.107923, 2020.
- 1054 Calheiros, T., Pereira, M. G. and Nunes, J. P.: Assessing impacts of future climate change on
1055 extreme fire weather and pyro-regions in Iberian Peninsula, Sci. Total Environ., 754, 142233,
1056 doi:10.1016/j.scitotenv.2020.142233, 2021.
- 1057 Carmo, M., Moreira, F., Casimiro, P. and Vaz, P.: Land use and topography influences on
1058 wildfire occurrence in northern Portugal, Landsc. Urban Plan., 100(1–2), 169–176,
1059 doi:10.1016/j.landurbplan.2010.11.017, 2011.

- 1060 Carvalho, A., Flannigan, M. D., Logan, K., Miranda, A. I. and Borrego, C.: Fire activity in
1061 Portugal and its relationship to weather and the Canadian Fire Weather Index System, *Int. J.*
1062 *Wildl. Fire*, 17(3), 328–338, doi:10.1071/WF07014, 2008.
- 1063 Cerasoli, S., Costa e Silva, F. and Silva, J. M. N.: Temporal dynamics of spectral
1064 bioindicators evidence biological and ecological differences among functional types in a cork
1065 oak open woodland, *Int. J. Biometeorol.*, 60(6), 813–825, doi:10.1007/s00484-015-1075-x,
1066 2016.
- 1067 Chinita, M. J., Richardson, M., Teixeira, J. and Miranda, P. M. A.: Global mean frequency
1068 increases of daily and sub-daily heavy precipitation in ERA5, *Environ. Res. Lett.*, 16(7),
1069 doi:10.1088/1748-9326/ac0caa, 2021.
- 1070 Cohen, J.: A Coefficient of Agreement for Nominal Scales, *Educ. Psychol. Meas.*, 20(1), 37–
1071 46, doi:10.1177/001316446002000104, 1960.
- 1072 Congalton, R. G.: Accuracy assessment and validation of remotely sensed and other spatial
1073 information, *Int. J. Wildl. Fire*, 10(3–4), 321–328, doi:10.1071/wf01031, 2001.
- 1074 Copernicus Climate Change Service (C3S): ERA5: Fifth generation of ECMWF atmospheric
1075 reanalyses of the global climate, Copernicus Clim. Chang. Serv. Clim. Data Store [online]
1076 Available from:
1077 [https://eds.climate.copernicus.eu/edsapp#!/www.ecmwf.int/en/forecasts/dataset/ecmwf-](https://eds.climate.copernicus.eu/edsapp#!/www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-era5-land?tab=overview,2017v5)
1078 [reanalysis-era5-land?tab=overview,2017v5](https://eds.climate.copernicus.eu/edsapp#!/www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-era5-land?tab=overview,2017v5) (Last access: October 2020), 2020.
- 1079 Crutzen, P. J. and Goldammer, J. G.: Quantification of Fire Characteristics from Local to
1080 Global Scales., 1993.
- 1081 David, F. N. and Cramer, H.: Mathematical Methods of Statistics., *Biometrika*, 34(3/4), 374,
1082 doi:10.2307/2332454, 1947.
- 1083 [Direção Geral do Território \(DGT\): Modelo Digital do Terreno \(Resolução 50 m\) - Portugal](#)
1084 [Continental](#), [online] Available from:

1085 [https://snig.dgterritorio.gov.pt/rndg/srv/por/catalog.search#/metadata/ba3f114f-51e2-4eaa-](https://snig.dgterritorio.gov.pt/rndg/srv/por/catalog.search#/metadata/ba3f114f-51e2-4eaa-9f61-b8ade36b2378?tab=techinfo)
 1086 [9f61-b8ade36b2378?tab=techinfo](https://snig.dgterritorio.gov.pt/rndg/srv/por/catalog.search#/metadata/ba3f114f-51e2-4eaa-9f61-b8ade36b2378?tab=techinfo) (Last access: 18 January 2022), 2010.

1087 [Direção Geral do Território \(DGT\): Carta de Uso e Ocupação do Solo \(COS\) de Portugal](#)
 1088 [Continental para 2018, 2019.](#)

1089 Duane, A. and Brotons, L.: Synoptic weather conditions and changing fire regimes in a
 1090 Mediterranean environment, *Agric. For. Meteorol.*, 253–254(January), 190–202,
 1091 doi:10.1016/j.agrformet.2018.02.014, 2018.

1092 ~~Elia, M., Lovreglio, R., Ranieri, N. A., Sanesi, G. and Laforzezza, R.: Cost effectiveness of~~
 1093 ~~fuel removals in mediterraneanwildland-urban interfaces threatened by wildfires, *Forests,*~~
 1094 ~~7(149), 1–11, doi:10.3390/f7070149, 2016.~~

1095 De Espindola, R. S., Luciano, E. M. and Audy, J. L. N.: An overview of the adoption of IT
 1096 governance models and software process quality instruments at Brazil - Preliminary results of
 1097 a survey, *Proc. 42nd Annu. Hawaii Int. Conf. Syst. Sci. HICSS*, 1–9,
 1098 doi:10.1109/HICSS.2009.70, 2009.

1099 Espinosa, J., Palheiro, P., Loureiro, C., Ascoli, D., Esposito, A. and Fernandes, P. M.: Fire-
 1100 severity mitigation by prescribed burning assessed from fire-treatment encounters in maritime
 1101 pine stands, *Can. J. For. Res.*, 49(2), 205–211, doi:10.1139/cjfr-2018-0263, 2019.

1102 European Environment Agency: [\(EEA\): Copernicus Land Monitoring Service](#), Copernicus L.
 1103 Monit. Serv. - EU-DEM [online] Available from: [https://www.eea.europa.eu/data-and-](https://www.eea.europa.eu/data-and-maps/data/copernicus-land-monitoring-service-eu-dem)
 1104 [maps/data/copernicus-land-monitoring-service-eu-dem](https://www.eea.europa.eu/data-and-maps/data/copernicus-land-monitoring-service-eu-dem) (~~Acessed~~Last access: 17 March
 1105 2021), 2021.

1106 ~~Fernandes, P. M. Fernandes, P. M., Variation in the canadian fire weather index thresholds~~
 1107 ~~for increasingly larger fires in Portugal, *Forests*, 10(10), doi:10.3390/f10100838, 2019.~~

1108 ~~Fernandes, P. M.,~~ Monteiro-Henriques, T., Guiomar, N., Loureiro, C. and Barros, A. M. G.:
 1109 Bottom-Up Variables Govern Large-Fire Size in Portugal, *Ecosystems*, 19(8), 1362–1375,

- 1110 doi:10.1007/s10021-016-0010-2, 2016.
- 1111 Fernandes, P. M.: Variation in the canadian fire weather index thresholds for increasingly
1112 larger fires in Portugal, Forests, 10(10), doi:10.3390/f10100838, 2019.
- 1113 Fernandes, P. M., Guiomar, N. and Rossa, C. G.: Analysing eucalypt expansion in Portugal as
1114 a fire-regime modifier, *Sci. Total Environ.*, 666, 79–88, doi:10.1016/j.scitotenv.2019.02.237,
1115 2019.
- 1116 Ferreira-Leite, F., Bento-Gonçalves, A., Vieira, A., Nunes, A. and Lourenço, L.: Incidence
1117 and recurrence of large forest fires in mainland Portugal, *Nat. Hazards*, 84(2), 1035–1053,
1118 doi:10.1007/s11069-016-2474-y, 2016.
- 1119 Fonseca, T. F. and Duarte, J. C.: A silvicultural stand density model to control understory in
1120 maritime pine stands, *IForest*, 10(5), 829–836, doi:10.3832/ifor2173-010, 2017.
- 1121 Frey, B. B.: The SAGE Encyclopedia of Educational Research, Measurement, and Evaluation,
1122 SAGE Encycl. Educ. Res. Meas. Eval., (March), 1–4, doi:10.4135/9781506326139, 2018.
- 1123 Gouveia, C. M., Bastos, A., Trigo, R. M. and Dacamara, C. C.: Drought impacts on vegetation
1124 in the pre- and post-fire events over Iberian Peninsula, *Nat. Hazards Earth Syst. Sci.*, 12(10),
1125 3123–3137, doi:10.5194/nhess-12-3123-2012, 2012.
- 1126 Greenwood, P. E. and Nikulin, M. S.: A Guide to Chi-Squared Testing, , 1–2, 1996.
- 1127 Groot, W. J. De: Interpreting the Canadian Forest Fire Weather Index (FWI) System, Fourth
1128 Cent. Reg. Fire Weather Comm. Sci. Tech. Semin., ~~Proceeding, 3–14 [online] Available from:~~
1129 ~~[http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Interpreting+the+canadian+](http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Interpreting+the+canadian+forest+fire+weather+index+(FWI)+system#0Proceeding,3-14)~~
1130 ~~[forest+fire+weather+index+\(FWI\)+system#0Proceeding, 3–14](http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Interpreting+the+canadian+forest+fire+weather+index+(FWI)+system#0Proceeding,3-14)~~, 1987.
- 1131 Huai, B., Wang, J., Sun, W., Wang, Y. and Zhang, W.: Evaluation of the near-surface climate
1132 of the recent global atmospheric reanalysis for Qilian Mountains, Qinghai-Tibet Plateau,
1133 *Atmos. Res.*, 250(November 2020), 105401, doi:10.1016/j.atmosres.2020.105401, 2021.
- 1134 ~~Instituto de Meteorologia de Portugal and Agência Estatal de Meteorologia de Espanha: Atlas~~

- 1135 climático ibérico: Temperatura do ar e precipitação (1971-2000), 2011.
- 1136 Instituto da Conservação da Natureza e das Florestas (ICNF): 6.º Inventário Florestal
- 1137 Nacional (IFN6) - 2015 Relatório Final, , 284 [online] Available from:
- 1138 <http://www2.icnf.pt/portal/florestas/ifn/ifn6>, 2019.
- 1139 Instituto da Conservação da Natureza e das Florestas (ICNF), [online] Available from:
- 1140 <http://www2.icnf.pt/portal/florestas/dfci/inc/mapas> (last access: December 2020), 2020.
- 1141 Jimenez-Ruano, A., Rodrigues, M., Jolly, W. M. and de la Riva, J.: The role of short-term
- 1142 weather conditions in temporal dynamics of fire regime features in mainland Spain, J.
- 1143 Environ. Manage., In press(September), 1–12, doi:10.1016/j.jenvman.2018.09.107, 2018.
- 1144 Jones, N., de Graaff, J., Rodrigo, I. and Duarte, F.: Historical review of land use changes in
- 1145 Portugal (before and after EU integration in 1986) and their implications for land degradation
- 1146 and conservation, with a focus on Centro and Alentejo regions, Appl. Geogr., 31(3), 1036–
- 1147 1048, doi:10.1016/j.apgeog.2011.01.024, 2011.
- 1148 Kanevski, M. and Pereira, M. G.: Local fractality: The case of forest fires in Portugal, Phys. A
- 1149 Stat. Mech. its Appl., 479, 400–410, doi:10.1016/j.physa.2017.02.086, 2017.
- 1150 Landis, J. R. and Koch, G. G.: The Measurement of Observer Agreement for Categorical
- 1151 Data, Biometrics, 33(1), 159, doi:10.2307/2529310, 1977.
- 1152 Leuenberger, M., Parente, J., Tonini, M., Pereira, M. G. and Kanevski, M.: Wildfire
- 1153 susceptibility mapping: Deterministic vs. stochastic approaches, Environ. Model. Softw., 101,
- 1154 194–203, doi:10.1016/j.envsoft.2017.12.019, 2018.
- 1155 Lloret, F., Calvo, E., Pons, X. and Díaz-Delgado, R.: Wildfires and Landscape Patterns in the
- 1156 Eastern Iberian Peninsula, Landsc. Ecol., 17(8), 745–759,
- 1157 doi:<https://doi.org/10.1023/A:1022966930861>, 2002.
- 1158 McHugh, M. L.: Lessons in biostatistics interrater reliability : the kappa statistic, Biochem.
- 1159 Medica, 22(3), 276–282 [online] Available from: <https://hrcak.srce.hr/89395>, 2012.

- 1160 Meneses, B. M., Reis, E. and Reis, R.: Assessment of the recurrence interval of wildfires in
1161 mainland Portugal and the identification of affected land patterns, *J. Maps*, 14(2), 282–292,
1162 doi:10.1080/17445647.2018.1454351, 2018a.
- 1163 Meneses, B. M., Reis, E., Vale, M. J. and Reis, R.: Modelling land use and land cover
1164 changes in Portugal: A multi-scale and multi-temporal approach, *Finisterra*, 53(107), 3–26,
1165 doi:10.18055/finis12258, 2018b.
- 1166 Moreira, F., Viedma, O., Arianoutsou, M., Curt, T., Koutsias, N., Rigolot, E., Barbati, A.,
1167 Corona, P., Vaz, P., Xanthopoulos, G. and Mouillot, F.: Landscape ~~e-wild-fire~~ wildfire
1168 interactions in southern Europe-Europe: Implications for landscape management, *J. Environ.*
1169 *Manage.*, 92(10), 2389–2402, doi:10.1016/j.jenvman.2011.06.028, 2011.
- 1170 Moreno, M. V., Conedera, M., Chuvieco, E. and Pezzatti, G. B.: Fire regime changes and
1171 major driving forces in Spain from 1968 to 2010, *Environ. Sci. Policy*, 37, 11–22,
1172 doi:10.1016/j.envsci.2013.08.005, 2014.
- 1173 Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G.,
1174 Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles,
1175 M., Rodríguez-Fernández, N. J., Zsoter, E., Buontempo, C. and Thépaut, J. N.: ERA5-Land:
1176 A state-of-the-art global reanalysis dataset for land applications, *Earth Syst. Sci. Data*, 13(9),
1177 4349–4383, doi:10.5194/essd-13-4349-2021, 2021.
- 1178 National Wildfire Coordinating Group (NCWG), NCWG: Glossary of Wildland Fire
1179 Terminology, October, 2005(July), 189 [online] Available from:
1180 <http://www.nwccg.gov/pms/pubs/glossary/pms205.pdf>, 2011.
- 1181 Nunes, A. N.: Regional variability and driving forces behind forest fires in Portugal an
1182 overview of the last three decades (1980–2009), *Appl. Geogr.*, 34(March), 576–586,
1183 doi:10.1016/j.apgeog.2012.03.002, 2012.
- 1184 Nunes, A. N., Lourenço, L. and Meira, A. C. C.: Exploring spatial patterns and drivers of

- 1185 forest fires in Portugal (1980–2014), *Sci. Total Environ.*, 573, 1190–1202,
1186 doi:10.1016/j.scitotenv.2016.03.121, 2016.
- 1187 ~~Nunes, S. A., Dacamara, C. C., Turkman, K. F., Calado, T. J., Trigo, R. M. and Turkman, M.
1188 A. A.: Wildland fire potential outlooks for Portugal using meteorological indices of fire
1189 danger, *Nat. Hazards Earth Syst. Sci.*, 19(7), 1459–1470, doi:10.5194/nhess-19-1459-2019,
1190 2019.~~
- 1191 Oliveira, T. M., Guiomar, N., Baptista, F. O., Pereira, J. M. C. and Claro, J.: Is Portugal's
1192 forest transition going up in smoke?, *Land use policy*, 66(May), 214–226,
1193 doi:10.1016/j.landusepol.2017.04.046, 2017.
- 1194 Parente, J. and Pereira, M. G.: Structural fire risk : The case of Portugal, *Sci. Total Environ.*,
1195 573, 883–893, doi:10.1016/j.scitotenv.2016.08.164, 2016.
- 1196 Parente, J., Pereira, M. G. and Tonini, M.: Space-time clustering analysis of wildfires: The
1197 influence of dataset characteristics, fire prevention policy decisions, weather and climate, *Sci.*
1198 *Total Environ.*, 559, 151–165, doi:10.1016/j.scitotenv.2016.03.129, 2016.
- 1199 Parente, J., Pereira, M. G., Amraoui, M. and Fischer, E. M.: Heat waves in Portugal: Current
1200 regime, changes in future climate and impacts on extreme wildfires, *Sci. Total Environ.*, 631–
1201 632, 534–549, doi:10.1016/j.scitotenv.2018.03.044, 2018a.
- 1202 Parente, J., Pereira, M. G., Amraoui, M. and Tedim, F.: Negligent and intentional fires in
1203 Portugal : Spatial distribution characterization, *Sci. Total Environ.*, 624, 424–437,
1204 doi:10.1016/j.scitotenv.2017.12.013, 2018b.
- 1205 Parente, J., Amraouia, M., Menezes, I. and Pereira, M. G.: Drought in Portugal: Current
1206 regime, comparison of indices and impacts on extreme wildfires, *Sci. Total Environ.*, 685,
1207 150–173, doi:10.1016/j.scitotenv.2019.05.298, 2019.
- 1208 Pausas, J. G. and Vallejo, V. R.: The role of fire in European Mediterranean ecosystems, in
1209 Remote Sensing of Large Wildfires, pp. 3–16. Springer Berlin Heidelberg, Berlin,

- 1210 [Heidelberg, 1999.](#)
- 1211 [Pausas, J. G.](#) and Fernández-Muñoz, S.: Fire regime changes in the Western Mediterranean
1212 Basin: From fuel-limited to drought-driven fire regime, *Clim. Change*, 110(1–2), 215–226,
1213 doi:10.1007/s10584-011-0060-6, 2012.
- 1214 ~~[Pausas, J. G.](#) and [Vallejo, V. R.](#): The role of fire in European Mediterranean ecosystems, in
1215 [Remote Sensing of Large Wildfires](#), pp. 3–16, Springer Berlin Heidelberg, Berlin,
1216 Heidelberg, 1999.~~
- 1217 Pereira, J. S., Chaves, M. M., Caldeira, M. C. and Correia, A. V.: Water Availability and
1218 Productivity, *Plant Growth Clim. Chang.*, (December 2017), 118–145,
1219 doi:10.1002/9780470988695.ch6, 2007.
- 1220 ~~[Pereira, M.](#), [Calado, T.](#), [DaCamara, C.](#) and [Calheiros, T.](#): Effects of regional climate change
1221 on rural fires in Portugal, *Clim. Res.*, 57(3), 187–200, doi:10.3354/cr01176, 2013.~~
- 1222 Pereira, M. G., Trigo, R. M., da Camara, C. C., Pereira, J. M. C. and Leite, S. M.: Synoptic
1223 patterns associated with large summer forest fires in Portugal, *Agric. For. Meteorol.*, 129(1–
1224 2), 11–25, doi:10.1016/j.agrformet.2004.12.007, 2005.
- 1225 Pereira, M. G., Malamud, B. D., Trigo, R. M. and Alves, P. I.: The history and characteristics
1226 of the 1980 – 2005 Portuguese rural fire database, *Nat. Hazards Earth Syst. Sci.*, (Table 1),
1227 3343–3358, doi:10.5194/nhess-11-3343-2011, 2011.
- 1228 Pereira, M. G., Aranha, J. and Amraoui, M.: Land cover fire proneness in Europe, *For. Syst.*,
1229 23(3), 598–610, 2014.
- 1230 Pereira, M. G., Caramelo, L., Orozco, C. V., Costa, R. and Tonini, M.: Space-time clustering
1231 analysis performance of an aggregated dataset: The case of wildfires in Portugal, *Environ.*
1232 *Model. Softw.*, 72, 239–249, doi:10.1016/j.envsoft.2015.05.016, 2015.
- 1233 Rodrigues, M., Trigo, R. M., Vega-García, C. and Cardil, A.: Identifying large fire weather
1234 typologies in the Iberian Peninsula, *Agric. For. Meteorol.*, 280(November 2019), 107789,

- 1235 doi:10.1016/j.agrformet.2019.107789, 2020.
- 1236 Romano, N. and Ursino, N.: Forest fire regime in a mediterranean ecosystem: Unraveling the
1237 mutual interrelations between rainfall seasonality, soil moisture, drought persistence, and
1238 biomass dynamics, *Fire*, 3(3), 1–20, doi:10.3390/fire3030049, 2020.
- 1239 Ruffault, J., Curt, T., Martin-Stpaul, N. K., Moron, V. and Trigo, R. M.: Extreme wildfire
1240 events are linked to global-change-type droughts in the northern Mediterranean, *Nat. Hazards*
1241 *Earth Syst. Sci.*, 18(3), 847–856, doi:10.5194/nhess-18-847-2018, 2018.
- 1242 Russo, A., Gouveia, C. M., Páscoa, P., DaCamara, C. C., Sousa, P. M. and Trigo, R. M.:
1243 Assessing the role of drought events on wildfires in the Iberian Peninsula, *Agric. For.*
1244 *Meteorol.*, 237–238, 50–59, doi:10.1016/j.agrformet.2017.01.021, 2017.
- 1245 San-Miguel-Ayanz, J., Durrant, T., Boca, R., Maianti, P., Liberta`, G., Artes Vivancos, T.,
1246 Jacome Felix Oom, D., Branco, A., De Rigo, D., Ferrari, D., Pfeiffer, H., Grecchi, R., Nuijten,
1247 D. and Leray, T.: Forest Fires in Europe, Middle East and North Africa 2019, ,
1248 doi:10.2760/468688, 2020.
- 1249 Scotto, M. G., Gouveia, S., Carvalho, A., Monteiro, A., Martins, V., Flannigan, M. D., San-
1250 Miguel-Ayanz, J., Miranda, A. I. and Borrego, C.: Area burned in Portugal over recent
1251 decades: An extreme value analysis, *Int. J. Wildl. Fire*, 23(6), 812–824,
1252 doi:10.1071/WF13104, 2014.
- 1253 Sianturi, Y., Marjuki and Sartika, K.: Evaluation of ERA5 and MERRA2 reanalyses to
1254 estimate solar irradiance using ground observations over Indonesia region, *AIP Conf. Proc.*,
1255 2223(April), doi:10.1063/5.0000854, 2020.
- 1256 Silva, J. M. N., Moreno, M. V., Page, Y. Le, Oom, D., Bistinas, I. and Pereira, J. M. C.:
1257 Spatiotemporal trends of area burnt in the Iberian Peninsula , 1975 – 2013, *Reg. Environ.*
1258 *Chang.*, 515–527, 2019.
- 1259 Sousa, P. M., Trigo, R. M., Pereira, M. G., Bedia, J. and Gutiérrez, J. M.: Different

- 1260 approaches to model future burnt area in the Iberian Peninsula, *Agric. For. Meteorol.*, 202,
 1261 11–25, doi:10.1016/j.agrformet.2014.11.018, 2015.
- 1262 Stamou, Z., Xystrakis, F. and Koutsias, N.: The role of fire as a long-term landscape modifier:
 1263 Evidence from long-term fire observations (1922–2000) in Greece, *Appl. Geogr.*, 74, 47–55,
 1264 doi:10.1016/j.apgeog.2016.07.005, 2016.
- 1265 Sutanto, S. J., Vitolo, C., Di Napoli, C., D’Andrea, M. and Van Lanen, H. A. J.: Heatwaves,
 1266 droughts, and fires: Exploring compound and cascading dry hazards at the pan-European
 1267 scale, *Environ. Int.*, 134(January), 105276, doi:10.1016/j.envint.2019.105276, 2020.
- 1268 Tarín-Carrasco, P., Augusto, S., Palacios-Peña, L., Ratola, N. and Jiménez-Guerrero, P.:
 1269 Impact of large wildfires on PM10 levels and human mortality in Portugal, *Nat. Hazards Earth*
 1270 *Syst. Sci.*, 2018, 1–21, doi:10.5194/nhess-2021-38, 2021.
- 1271 Telesca, L. and Pereira, M. G.: Time-clustering investigation of fire temporal fluctuations in
 1272 Portugal, *Nat. Hazards Earth Syst. Sci.*, 10(4), 661–666, doi:10.5194/nhess-10-661-2010,
 1273 2010.
- 1274 ~~Território, D. G. do (DGT): Modelo Digital do Terreno (Resolução 50 m) – Portugal~~
 1275 ~~Continental, [online] Available from:~~
 1276 ~~[https://snig.dgterritorio.gov.pt/rndg/srv/por/catalog/search#/metadata/ba3f114f-51e2-4caa-](https://snig.dgterritorio.gov.pt/rndg/srv/por/catalog/search#/metadata/ba3f114f-51e2-4caa-9f61-b8ade36b2378?tab=techinfo)~~
 1277 ~~[9f61-b8ade36b2378?tab=techinfo](https://snig.dgterritorio.gov.pt/rndg/srv/por/catalog/search#/metadata/ba3f114f-51e2-4caa-9f61-b8ade36b2378?tab=techinfo) (Accessed 18 January 2022), 2010.~~
- 1278 ~~Território, D. G. do (DGT): Carta de Uso e Ocupação do Solo (COS) de Portugal Continental~~
 1279 ~~para 2018, 2019.~~
- 1280 The MathWorks Inc: Linkage. Agglomerative hierarchical cluster tree, Help Cent. [online]
 1281 Available from: <https://www.mathworks.com/help/stats/linkage.html> (~~Aeessed~~Last access:
 1282 15 November 2021), 2021.
- 1283 Tonini, M., Parente, J. and Pereira, M. G.: Global assessment of rural-urban interface in
 1284 Portugal related to land cover changes, *Nat. Hazards Earth Syst. Sci.*, 18(6), 1647–1664,

- 1285 doi:10.5194/nhess-18-1647-2018, 2018.
- 1286 Trigo, R. M., Pereira, J. M. C., Pereira, M. G., Mota, B., Calado, T. J., Dacamara, C. C. and
1287 Santo, F. E.: Atmospheric conditions associated with the exceptional fire season of 2003 in
1288 Portugal, *Int. J. Climatol.*, 26(13), 1741–1757, doi:10.1002/joc.1333, 2006.
- 1289 Trigo, R. M., Sousa, P. M., Pereira, M. G., Rasilla, D. and Gouveia, C. M.: Modelling wildfire
1290 activity in Iberia with different atmospheric circulation weather types, *Int. J. Climatol.*, 36(7),
1291 2761–2778, doi:10.1002/joc.3749, 2016.
- 1292 Turco, M., Rosa-Cánovas, J. J., Bedia, J., Jerez, S., Montávez, J. P., Llasat, M. C. and
1293 Provenzale, A.: Exacerbated fires in Mediterranean Europe due to anthropogenic warming
1294 projected with non-stationary climate–fire models, *Nat. Commun.*, 9(1), 1–9,
1295 doi:10.1038/s41467-018-06358-z, 2018.
- 1296 Turco, M., Jerez, S., Augusto, S., Tarín-Carrasco, P., Ratola, N., Jiménez-Guerrero, P. and
1297 Trigo, R. M.: Climate drivers of the 2017 devastating fires in Portugal, *Sci. Rep.*, 9(1),
1298 doi:10.1038/s41598-019-50281-2, 2019.
- 1299 Urban, A., Di Napoli, C., Cloke, H. L., Kysely, J., Pappenberger, F., Sera, F., Schneider, R.,
1300 Vicedo-Cabrera, A. M., Acquaotta, F., Ragetti, M. S., Íñiguez, C., Tobias, A., Indermitte, E.,
1301 Orru, H., Jaakkola, J. J. K., Rytí, N. R. I., Pascal, M., Huber, V., Schneider, A., de’ Donato,
1302 F., Michelozzi, P. and Gasparrini, A.: Evaluation of the ERA5 reanalysis-based Universal
1303 Thermal Climate Index on mortality data in Europe, *Environ. Res.*, 198(May),
1304 doi:10.1016/j.envres.2021.111227, 2021.
- 1305 ~~Vega Orozco, C., Tonini, M., Conedera, M. and Kanveski, M.: Cluster recognition in spatial-~~
1306 ~~temporal sequences: The case of forest fires, *Geoinformatica*, 16(4), 653–673,~~
1307 ~~doi:10.1007/s10707-012-0161-z, 2012.~~
- 1308 ~~Viegas, D. X., Bovio, G., Ferreira, A., Nosenzo, A. and Sol, B.: Comparative study of various~~
1309 ~~methods of fire danger evaluation in southern Europe, *Int. J. Wildl. Fire*, 9(4), 235,~~

1310 ~~doi:10.1071/WFE00015, 1999.~~
1311 Vieira, I., Russo, A. and Trigo, R. M.: Identifying local-scale weather forcing conditions
1312 favorable to generating Iberia’s largest fires, *Forests*, 11(5), 1–14, doi:10.3390/F11050547,
1313 2020.
1314 Vilar, L., Camia, A., San-Miguel-Ayanz, J. and Martín, M. P.: Modeling temporal changes in
1315 human-caused wildfires in Mediterranean Europe based on Land Use-Land Cover interfaces,
1316 *For. Ecol. Manage.*, 378, 68–78, doi:10.1016/j.foreco.2016.07.020, 2016.
1317 Van Wagner, C. . and Pickett, T. L.: Equations and fortran IV program for the 1976 metric
1318 version of the forest fire weather index, 1975.
1319 Van Wagner, C. E.: Development and structure of the Canadian Forest Fire Weather Index
1320 system., 1987.
1321 Whitlock, C., Higuera, P. E., McWethy, D. B. and Briles, C. E.: Paleocological Perspectives
1322 on Fire Ecology: Revisiting the Fire-Regime Concept ~~!2009-09-02 !2009-11-09 !2010-03-~~
1323 ~~05-!~~, *Open Ecol. J.*, 3(2), 6–23, doi:10.2174/1874213001003020006, 2010.

1324

Formatada: Sem controlo de linhas isoladas, Não ajustar o espaço entre texto asiático e texto em latim, Não ajustar o espaço entre texto asiático e números