

A Unified Probabilistic Framework for Volcanic Hazard and Eruption Forecasting

Warner Marzocchi¹, Jacopo Selva², Thomas H. Jordan³

¹ Department of Earth, Environmental, and Resources Sciences, University of Naples, Federico II, Complesso di Monte Sant'Angelo, Via Cupa Nuova Cintia, 21 - 80126 Napoli, Italy

² Istituto Nazionale di Geofisica e Vulcanologia, Via Donato Creti 12, 40128 Bologna, Italy

³ Department of Earth Sciences Southern California Earthquake Center, University of Southern California, Los Angeles, California 90089

10 *Correspondence to:* Warner Marzocchi (warner.marzocchi@unina.it)

Abstract. The main purpose of this article is to emphasize the importance of clarifying the probabilistic framework adopted for volcanic hazard and eruption forecasting. Eruption forecasting and volcanic hazard analysis seek to quantify the deep uncertainties that pervade the modeling of pre-, sin- and post-eruptive processes. These uncertainties can be differentiated into three fundamental types: (1) the natural variability of volcanic systems, usually represented as stochastic processes with parameterized distributions (*aleatory variability*); (2) the uncertainty in our knowledge of how volcanic systems operate and evolve, often represented as subjective probabilities based on expert opinion (*epistemic uncertainty*); and (3) the possibility that our forecasts are wrong owing to behaviors of volcanic processes about which we are completely ignorant and, hence, cannot quantify in terms of probabilities (*ontological error*). Here we put forward a probabilistic framework for hazard analysis recently proposed by Marzocchi & Jordan (2014), which unifies the treatment of all three types of uncertainty. Within this framework, an eruption forecasting or a volcanic hazard model is said to be complete only if it (a) fully characterizes the epistemic uncertainties in the model's representation of aleatory variability and (b) can be unconditionally tested (in principle) against observations to identify ontological errors. Unconditional testability, which is the key to model validation, hinges on an *experimental concept* that characterizes hazard events in terms of exchangeable data sequences with well-defined frequencies. We illustrate the application of this unified probabilistic framework by describing experimental concepts for the forecasting of tephra fall from Campi Flegrei. Eventually, this example may serve as a guide for the application of the same probabilistic framework to other natural hazards.

1 Introduction

Hazards associated with major eruptions and their consequences are highly uncertain owing to the stochastic behaviors of volcanic systems (aleatory variability) as well as our lack of knowledge about these behaviors (epistemic uncertainty). Because such complexity hampers the deterministic prediction of hazards, our goal is to describe them probabilistically (Sparks, 2003; Marzocchi and Bebbington, 2012; Poland and Anderson, 2020). Here we use the term probabilistic volcanic

hazard analysis (PVHA) to indicate the probabilistic forecast of any volcanological event of interest (eruption occurrence, ash fall loading, arrival of a pyroclastic flow, etc).

PVHA outcome may be generally described by the exceedance probability of a positive random variable X (Martin et al., 2004, Selva et al., 2018; Rougier and Beven, 2013; Sandri et al., 2016; Bear-Crozier et al., 2016)

$$f(x) = \Pr(X > x | \mathbf{H}), \quad x \in (0, \infty) \quad (1)$$

where \mathbf{H} is the PVHA model used, $f(x)$ is called the hazard curve (not to be confused with the hazard function commonly used to describe failure rate; from a statistical point of view $f(x)$ is a survival function). X is the (continuous or discrete) hazard intensity of interest in one specific time interval, for example, the tephra fall loading in one specific site, the dynamic pressure of a pyroclastic flow in one sector of the volcano, or the occurrence of an eruption, etc. Ideally, since we never know the true hazard curve, we may include an additional level of uncertainty considering a set of PVHA models; in this case, we have $\{f_i(x), \pi_i\}$ ($i = 1, \dots, N$), where π_i is the weight of the i^{th} PVHA model H_i , $f_i(x) \equiv f(x|H_i)$ (Rougier and Beven 2013).

The way in which we use the set of hazard curves $f_i(x)$ to estimate PVHA and the meaning of the weights depend on the probabilistic framework adopted. The importance of establishing the probabilistic framework to calculate PVHA cannot be overstated, because the framework implicitly defines the type of uncertainties, the meaning of probability, and consequently the possibility to validate (at least in principle) the PVHA model. Indeed, the qualitative difference between the objective probabilities of what we can see and count and the subjective probabilities of what we think we know has been recognized for 300 years, separating, for example, the frequentist and Bayesian approaches to probabilistic inference (Hacking 1965). Despite recent attempts to unify these two approaches (e.g., Box 1980; Rubin 1984; Bayarri and Berger 2004; Berger, 2004; Gelman and Shalizi 2013), controversy remains over how aleatory variability can be separated from epistemic uncertainty and whether such a dichotomy is actually useful in hazard analysis. For example, in the Bayesian view adopted by many researchers, the only type of uncertainty is epistemic (e.g., NRC, 1997; Bedford and Cooke, 2001; Lindley, 2000; Jaynes, 2003), whereas in the traditional frequentist framework, the only type of uncertainty is aleatory (Hacking, 1965).

Most practitioners recognize that distinguishing between different types of uncertainties can be useful in the interpretation of hazard estimates (e.g., Abrahamson and Bommer 2005; IAEA, 2012; Rougier 2013), but the confusion surrounding the topic is evident in the wide variety of schemes proposed for classifying uncertainties: shallow and deep (Stein and Stein, 2013), intra- and inter-model, external and internal (Rougier and Beven 2013), inter- and intra-model (Selva et al., 2013), value and structural uncertainty (e.g., Solomon et al., 2007), quantified measure of uncertainty and confidence on the validity of a finding (IPCC, 2013), model parameters and initial/boundary conditions, and many others (Reilly et al. 2021). It has been unclear whether these classifications are profound categories that must be reflected in the probabilistic framework or merely convenient, model-based divisions (e.g., NRC, 1997; Rougier and Beven, 2013).

The subject of this paper is to underline the importance of the probabilistic framework in PVHA and its practical implications. This framework must be able to (1) establish a coherent and clear hierarchy of different kinds of uncertainty

65 (aleatory variability, epistemic uncertainty, and ontological error), (2) assimilate subjective expert judgment into probabilistic models, and (3) unconditionally test complete probabilistic models against data. In the next sections, through a toy example and an application to the Campi Flegrei tephra-fall hazard, we describe the unified probabilistic framework developed by Marzocchi and Jordan (2014, 2017, 2018), which satisfies these requirements. To facilitate the reading and comprehension of the paper, we describe the notation and the new terminology both within the manuscript and in a glossary
70 at the end of the paper.

2 A tutorial example to describe a unified probabilistic framework for PVHA

We consider the case in which $f(x)$ is the annual probability to exceed a specific tephra fall threshold x in one site of specific interest around a volcano (the probability of exceedance, or PoE). The results can be generalized in a straightforward way to other time windows and other types of volcanic threats, as well as to eruption forecasting.

75 The unified probabilistic framework is rooted in the definition of an experimental concept, which allows us to define a hierarchy of uncertainties. Here we introduce it with a tutorial example, which may be easily generalized to more complex cases. We collect the sequence of annual observations for a particular value of the tephra fall loading x_0 , which we denote by the binary variable e_i . That is, $e_i = 0$ when there is no exceedance in the i^{th} year ($x < x_0$), and $e_i = 1$ when there is at least one exceedance ($x \geq x_0$). The experimental concept is the judgment of stochastic exchangeability of the sequence e_i , i.e.,
80 the joint probability distributions is invariant to data ordering when conditioned on a set of explanatory variables (Draper *et al.* 1993). This definition of the experimental concept has implications on the assumptions made about the eruptive process. With the above definition, we are assuming that, at the target volcano, eruptions (or the paroxysmal explosive phases) usually last less than one year (the time window used to define the experimental concept) and the inter-event times between consecutive eruptions (or paroxysmal explosive phases) are conditionally independent and mostly larger than one year. Of
85 course, if we are interested in volcanoes that behave differently, we have to define another more suitable experimental concept. This example has been chosen because it applies to volcanic systems like Campi Flegrei (the real example of the next section), and it presents similarities with seismic hazard to facilitate the comparison.

A theorem by de Finetti (1974) states that a set of events that is judged to be exchangeable (i.e., the events may come or not from different unknown distributions) can be modeled as identical and independently distributed random variables with a
90 well-defined frequency of occurrence, $\hat{\phi}$. The frequentist interpretation applied to a set of exchangeable events and the use of the Bayesian mathematical apparatus to handle uncertainties of different kind is the reason for use the term “unified” to characterize the probabilistic framework described in this paper.

In our example, $\hat{\phi} = \hat{f}(x_0)$, where $\hat{f}(\cdot)$ is the true hazard curve, and the unknown frequency $\hat{\phi}$ of the exchangeable sequence e_i is the aleatory variability. The estimation of the unknown true aleatory variability is the target of PVHA, and its estimation

95 is often possible with different models, $\{\phi_i, \pi_i\}$, where π_i is the weight of the i^{th} PVHA model $f_i(x)$, and $\phi_i = f_i(x_0)$. The
existence of such alternative models for the aleatory uncertainty reflects the existence of a known uncertainty on the true
frequency of the experimental concept (the aleatory variability), usually referred to as “epistemic uncertainty”. The definition
of the weight π_i has an unavoidably subjective nature. The weight of one model may be related in some ways to the hindcast
100 performance of that model and/or through expert judgments. In the usual application of the Bayesian framework to hazard
analysis, the set of models is considered complete and exhaustive, and the weight of a model is the probability to be the one
that should be used (Scherbaum and Kuehn, 2011); a similar interpretation is often adopted when using the Logic Trees
(Bommer and Scherbaum, 2008). In the unified framework the weight represents a measure of the forecasting skill of a
model with respect to the others (see Marzocchi and Jordan, 2017, for a deep description of this important issue). Then, from
the set $\{\phi_i, \pi_i\}$ we estimate the PoE distribution $p(\phi)$ (hereafter we use $p(\cdot)$ and $P(\cdot)$ to indicate the probability density
105 function and the cumulative distribution, respectively). This procedure is non-unique, it contains a degree of subjectivity, but
it is unavoidable; for example, choosing only one estimation ϕ_i , or the weighted average assumes that $p(\phi)$ follows a Dirac
distribution centered to this value. We call $p(\phi)$ the "Extended Experts' Distribution (EED)", which describes the full PVHA
(Marzocchi and Jordan, 2014).

In contrast to the Bayesian framework, for which all models are "wrong" and model validation is pointless (Lindley, 2000),
110 the unified framework allows model validation. Specifically, we can define an *ontological null hypothesis*, which states that
the true aleatory representation of future occurrence of natural events—the data generating process—mimics a sample from
the EED that describes the model’s epistemic uncertainty. According to the ontological null hypothesis, the true unknown
frequency $\hat{\phi}$ of the experimental concept cannot be distinguished from a realization of the EED, i.e., $\hat{\phi} \sim p(\phi)$. If the data
are inconsistent with the EED, the ontological null hypothesis can be rejected, which identifies the existence of an
115 *ontological error* (Marzocchi and Jordan, 2014). In other words, the “known unknowns” (epistemic uncertainty) do not
necessarily completely characterize the uncertainties, presumably due to effects not captured by the EED—“unknown
unknowns” associated with ontological errors.

In practice, collecting sufficient data for this kind of model validation is only feasible for specific sites surrounding active
volcanoes with a high frequency of eruptions. For a specific site near a high-risk volcano with a low eruptive frequency, the
120 data are usually insufficient for formal ontological testing. In probabilistic seismic hazard analysis (PSHA), the problem is
overcome trading time with space, i.e., considering many sites simultaneously for one or more time windows. This approach
requires that exceedances recorded at different sites can be considered statistically independent; although this may become
attainable in PSHA under some specific considerations, it clearly does not hold for sites surroundings a single volcano. In
volcanology, trading time with space is useful in validating models for global PVHA (e.g., Jenkins et al., 2015), which
125 consider the eruptive activity of all volcanoes of a specific type. In this case, it is possible to select sites that are far enough
to consider the observed exceedances conditionally independent one from each other, and the exchangeable sequence e_i ($i =$
 $1, \dots, N$) is given by the annual exceedances observed at different N sites. In summary, although the possibility to validate a

PVHA model is conceptually feasible through the unified probabilistic framework (whereas it is not in the Bayesian framework), it is currently often not practically feasible. In fact, the test of an ontological hypothesis (model validation) requires i) the definition of a proper experimental concept, ii) a complete PVHA including aleatory variability and epistemic uncertainty, and iii) enough independent data for testing. We are not aware of cases in which these three requirements are presently satisfied. But we argue that they could be achievable in future dedicated efforts, for example adopting procedures to standardize variables and group “analogue” and exchangeable volcanoes to increase the available data for testing (Tierz et al., 2019).

Future applications may also take advantage from the fact that the exchangeability judgment can be generalized beyond the stationarity of the process (implicit in our example) to more complex situations. For example, we may distinguish ash fall exceedances in the winter and summer seasons, because ash fall loading may be markedly affected by the seasonal dominant winds blowing in different directions. In this case, the data-generating process provides two sequences, $\{e_i^{(1)}: i = 1, \dots, N_1\}$ for the N_1 winter seasons, and $\{e_i^{(0)}: i = 1, \dots, N_0\}$ for the N_0 summer seasons. Both are judged to be Bernoulli sequences and they are observed to sum to k_0 and k_1 respectively. If the site is located downwind in the winter season, then the expected frequency of $\hat{\phi}^{(1)} = k_1/N_1$ might be greater than that of $\hat{\phi}^{(0)} = k_0/N_0$. As this example makes clear, it is neither the aleatory variability intrinsic to the model that matters in testing, nor the undisciplined randomness of the physical world, but rather the aleatory variability is defined by the exchangeability judgments of the experimental concept. In other words, aleatory variability is an observable behavior of the data-generating process conditioned by the experimental concept to have well-defined frequencies (Marzocchi and Jordan, 2014).

3 Accounting for epistemic uncertainty and aleatory variability: the unified framework applied to the tephra fall PVHA at Campi Flegrei

In this section we apply the probabilistic framework outlined in section 2 to the tephra fall PVHA at Campi Flegrei. Although the low eruption frequency of this volcano makes model validation unrealistic in the human time frame, the probabilistic framework has the advantage of providing a full description of the PVHA, accounting for all uncertainties; this may be of particular importance for decision-makers because, for example, they can immediately appreciate the level of uncertainty over the probabilistic assessment made by volcanologists.

Most (if not all) of the studies available for tephra fall PVHA at Campi Flegrei are based on Event Trees (ET; see Newhall and Hoblitt, 2002; Marzocchi et al., 2004; 2008; 2010; Marti et al., 2008; Sobradelo and Marti, 2010). The ET is a popular tree graph representation of events in which individual branches at each node point to different possible events, states, or conditions through increasingly specific subsequent events (intermediate outcomes) to final outcomes; in this way, an ET shows all relevant possible outcomes of volcanic unrest at progressively higher degrees of detail. The probability of each outcome is calculated combining the conditional probability of each branch belonging to the path from the first node to the

final outcome through classical probability theorems. By construction, ET is meant to describe only the intrinsic variability
160 of the process (aleatory variability) and not the epistemic uncertainty, hence it may produce only one single probabilistic
assessment $f(x)$. The ET has been generalized to account for uncertainties of different kind replacing the probability at each
node with a distribution of probability (Bayesian Even Tree, BET; Marzocchi et al., 2004; 2008; 2010; Neri et al., 2008;
Sobradelo and Marti, 2010), which aims at accounting for experts' judgment, different conceptual or physical models, data
from analog volcanoes, as well as data from the target or analog volcanoes (Marzocchi et al., 2004; Tierz et al., 2019).

165 The BET approach fits quite well the unified probabilistic framework that we advocate in this paper. If we consider an
experimental concept given by an exchangeable sequence e_i of annual tephra fall exceedances, the outcome of the BET code
may be seen as an EED, i.e., the distribution of probability $p(\phi)$ mimics where the true unknown frequency of the
exchangeable sequence may be. In the first paper that describes BET (Marzocchi et al., 2004), the use of a distribution of
probability at each node was generically advocated to account for the aleatory variability and epistemic uncertainty, using
170 the loose definition of irreducible randomness and limited knowledge of the process, respectively. This paper provides a
formal probabilistic background to justify the BET feature of considering the probability at each node as a distribution
($p(\phi)$) instead of a single number; at each node of the event tree, the central value of the distribution is the best guess of the
(unknown) long-term frequency of the experimental concept for that specific node (aleatory variability), and the dispersion
mimics the uncertainty over this unknown frequency.

175 The BET approach has been widely investigated for tephra fall PVHA at Campi Flegrei, adopting different choices,
hypotheses, and models (Selva et al., 2010; 2018; Sandri et al., 2016). For instance, Selva et al (2018) show the outcomes of
five different BET configurations (Figure 1) for one specific site inside the caldera (Figure 2), which differ in the
implementation of the tephra fall dispersion model (aggregation and granulometry). In Figure 3 we show the PoE
distribution $p(\phi)$ of each configuration for the tephra loading threshold of 300 kg/m². Each one of the five EED distributions
180 allows for a formal testing of the ontological hypothesis. Note that in this specific case, the EEDs do not account for seasonal
variabilities and are relative to the conditional PVHA (conditioned to the occurrence of an eruption). The generalization to
an unconditional PVHA can be straightforwardly made, for example, convolving all these distributions with the distribution
of the annual eruption probability.

When all EEDs are significantly overlapping (as for many points inside the Campi Flegrei caldera), it means that each BET
185 configuration describes the epistemic uncertainty in a consistent manner. Instead, for the site in Figure 2, we infer that
inconsistent BET outcomes (Figure 3) may be due to an underestimation of the epistemic uncertainty in each EED. For this
reason, in the example of Figure 3 we consider only the weighted average of each EEDs and then we build a new EED which
describes more satisfactorily the overall epistemic uncertainty given by the five BET configurations. This is equivalent to the
case of using alternative implementations of the classical ET (Newhall and Pallister, 2015), which produces a set of hazard
190 curves like in the upper panels of Figure 1. In the specific case of Figure 3, the reason for which the epistemic uncertainty is

underestimated in each EED may be due to the BET set up and/or to limitations of BET model to handle some sources of epistemic uncertainty.

The way in which we can build a single EED from a set of point forecasts $\{\phi_i, \pi_i\}$ (lower panels of Figure 1), or from a set of inconsistent EEDs (Figure 3), is what we call *ensemble modeling*. The terms ensemble modeling and ensemble forecasts
 195 have been used in many disciplines in different ways since early seventies (e.g., Leith, 1974). The book by Nate Silver (Silver, 2012) gives a wide range of successful applications and uses of ensemble modeling. The common feature across all these different flavors of ensemble modeling/forecasting is the attempt to account for the aleatory variability and the epistemic uncertainty by merging the forecasts of different models or parametrization of the same model in a proper way.

Standard methods are available for the induction of the EED $p(\phi)$ from the set $\{\phi_i, \pi_i\}$. Regardless the details of the
 200 procedure, we underline that the nonunique extrapolation of $\{\phi_i, \pi_i\}$ onto the continuous distribution $p(\phi)$ can contribute to ontological errors in the EED. Nonetheless, any reasonable procedure is not more subjective, and certainly less critical, than either not considering the epistemic uncertainty (for example, when using the weighted average $\bar{\phi}$), or assuming that the set of forecasts explore completely and exhaustively the epistemic uncertainty. In the first case, it is assumed that $p(\phi) \equiv \delta(\phi - \bar{\phi})$ is the Dirac distribution; in the latter case, it is assumed that the proper distribution for the epistemic uncertainty is
 205 fully described by the set of forecasts (Marzocchi et al., 2015).

Although ensemble modeling does not prescribe any specific procedure to estimate $p(\phi)$, in statistics random variables bounded in the range $[0,1]$ are often modeled by means of the Beta distribution (Gelman et al., 2003). Hence, we assume that $\Phi \sim \text{Beta}(\alpha, \beta)$, where the parameters α and β are related to the weighted average $E(\phi)$ and variance $\text{var}(\phi)$ of $\{\phi_i, \pi_i\}$ through:

$$210 \quad E(\phi) = \frac{\alpha}{(\alpha+\beta)} \quad (2)$$

and

$$\text{var}(\phi) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)} \quad (3)$$

Inverting equations 2 and 3, we can get the parameters of the Beta distribution $p(\phi)$, which describes the ontological hypothesis. Using the weights reported in Figure 1, we get the Beta distribution reported in Figure 4. As expected, this
 215 global EED is wider than each single EED of the BET configurations reported in Figure 3, and it aims at describing more realistically the complete uncertainty in PVHA. Although the choice of the Beta distribution is subjective, we advocate the use of a unimodal distribution (the Beta distribution is almost always unimodal), which describes more realistically, in most cases, the epistemic uncertainty over the true (unknown) frequency. (See the discussion of Figure 6 in Marzocchi et al., 2015.)

In this paper we have described a unified probabilistic framework which allows volcanologists to provide a complete description of PVHA, to define a clear taxonomy of uncertainties (aleatory variability, epistemic uncertainty, and ontological errors), and to account for experts' judgments preserving the possibility to unconditionally test PVHA against data, at least for high-frequency erupting volcanoes, or for global forecasting models. Although in this paper we focus entirely on PVHA, we think that this approach may potentially inspire other scientists working on different natural hazards.

One remarkable and distinctive feature of this probabilistic framework is that the mathematical description of PVHA is given by a distribution of probability (see, e.g., Marzocchi et al., 2004; 2008; 2010; Neri et al., 2008; Sobradelo and Marti, 2010; Bevilacqua et al., 2015), or, equivalently, through a bunch of hazard curves $f_i(x)$ ($i = 1, \dots, N$) (Rougier and Beven, 2013). The use of a distribution of probability instead of single numbers mark the main difference with probabilistic frameworks that are more commonly used in PVHA (Marzocchi and Bebbington, 2012), i.e., the frequentist (e.g., Bebbington, 2010; Deligne et al., 2010), and subjective Bayesian (Aspinall et al., 2003). Although these probabilistic frameworks are both legitimate because they are coherent with the Kolmogorov's axioms, they cannot provide a complete description of PVHA, because they cannot unambiguously distinguish and handle properly uncertainties of different kind, which are likely pervasive in natural systems (Marzocchi and Jordan, 2017; 2018).

Making explicit the probabilistic framework in PVHA is important. In the past, loose definitions of the probabilistic framework have provoked critiques of natural hazard analysis (see, e.g., Castanos and Lomnitz, 2002; Mulargia et al., 2017). For example, a vague definition of the nature of uncertainties and the role of subjective judgments brought some scientists to assert that (Stark, 2017) "what appears to be impressive 'science' is in fact an artificial amplification of the opinions and ad hoc choices built into the model, which has a heuristic basis rather than a tested (or even testable) scientific basis." This criticism is implicitly rooted in the (false) syllogism: science is objective, natural hazard analysis relies on subjective experts' judgment, hence natural hazard analysis is not science.

The unified probabilistic framework proposed here emphasizes the importance of model validation (at least in principle) as cornerstone of science; pure objectivity is a myth even in science and the presence of unavoidable subjectivity in PVHA cannot be used to dismiss its scientific nature. In an extreme case, experts' can behave like "models" expressing their subjective measure of the frequency of one defined experimental concept; mutatis mutandis, the same applies to the famous case of farmers who subjectively guess the weight of an ox (Galton, 1907), whose similarities and differences with natural hazard analysis has been discussed in Marzocchi and Jordan (2014). Conversely, the Bayesian framework, which is a full legitimate probabilistic framework to be used in PVHA, does not allow model validation; in this framework all models are wrong (hence, why waste time to validate them?), and we can only evaluate the relative forecasting performance of one model against the others (Lindley, 2000; Jaynes, 2003).

Besides the scientific aspects, the use of a PoE distribution has remarkable practical merits, because it shows to the decision makers both our best guess and the associated uncertainty. In plain words, if two PVHA have the same average, but with quite different variance, this may affect significantly the way in which PVHA could be used by decision-makers. For example, let us consider a case in which there is a critical threshold in PVHA that triggers a specific mitigation action when overcome (this is just a simplified example, because the decision-making has to be based on risk, not on hazard); both averages may be lower of such a critical threshold (hence, both suggesting no action), but, when considering the variance, one of the EED shows a significant part of the distribution above the threshold (suggesting to take action). In this case, the decision-makers may take into consideration the epistemic uncertainty deciding, for the sake of precautionary reasons, to use one specific high percentile of the EED, instead of the average; for example, the Ministry of Civil Defence & Emergency Management in New Zealand (MCDEM, 2008) uses 84th percentile of the tsunami hazard analysis as a threshold for taking actions.

As a final consideration, owing to the social implications, we think that only adopting a clear probabilistic framework to get a complete PVHA is the best way to defend probabilistic assessments against future scrutiny and criticism and to use these assessments in the most profitable way.

265 Glossary and notation

This glossary contains the statistical notation used in this paper, and the definitions of new and uncommon terms.

<i>Aleatory variability</i>	Intrinsic randomness of the data-generating process. In the uncertainty hierarchy, it is the event frequency defined by the experimental concept.
<i>Epistemic uncertainty</i>	Lack of knowledge about the data-generating process. In the <i>uncertainty hierarchy</i> , it is the modeling uncertainty in the event frequency, where the latter is defined by the <i>experimental concept</i> .
<i>Exchangeability</i>	A property that the joint probability distribution of a data set is invariant with respect to permutations in the data ordering.
<i>Experimental concept</i>	Collections of data, observed and not yet observed, that are judged to be exchangeable when conditioned on a set of explanatory variables. The experimental concept defines the uncertainty hierarchy.
<i>Extended experts' distribution (EED)</i>	The continuous probability distribution sampled by the discrete experts' distribution, used to set up the ontological hypothesis.

<i>Ontological error</i>	In the uncertainty hierarchy, an error in a model's quantification of the aleatory variability and epistemic uncertainty, where the dichotomy is defined by the experimental concept.
<i>Ontological hypothesis</i>	A statistical null hypothesis that the event frequency defined by the experimental concept is a sample from the extended experts' distribution. The rejection of the ontological hypothesis exposes an ontological error.
<i>PoE</i>	Probability of exceedance, i.e., the probability that one specific parameter of interest will be overcome in one specific time interval.
<i>PVHA</i>	Probabilistic volcanic hazard analysis; in this case it describes a forecast of the volcanic activity for any forecasting time window, and for different kind of events (eruption, magnitude, ash fall loading threshold, etc).
<i>Uncertainty hierarchy</i>	Levels of uncertainty, from aleatory variability to epistemic uncertainty to ontologic error, defined by the experimental concept.
$f(x)$	Hazard function or hazard curve; mathematically, it is a survivor function, as described by equation (1).
π_i	Weight of the i^{th} model.
Symbol ^	It indicates the true value of an unknown (e.g., $\hat{\phi}$ is the true frequency of the experimental concept), or the true probability distribution (e.g., $\hat{f}(\cdot)$ is the true hazard curve).
$z \sim p(z)$	The random variable z is distributed according to the distribution $p(z)$.
$p(\cdot)$ and $P(\cdot)$	Probability density function and the cumulative distribution, respectively
$E[z]$ and $\text{var}(z)$	Expectation and variance of the random variable z , respectively

Author contribution: All authors contributed, at different levels, to the conceptualization of the research. WM and JS carried out the formal analysis and numerical investigation. WM wrote the first draft of the paper with contributions from all the other authors. All authors reviewed and edited the final version.

Acknowledgments: TO BE WRITTEN [include: THJ was supported by a grant from the W.M. Keck Foundation.]

References

- Abrahamson, N. A., and Bommer, J. J.: Probability and uncertainty in seismic hazard analysis, *Earthq. Spectra*, 21, 603–607, 2015.
- 275 Aspinall, W.P., Woo, G., Voight, B., and Baxter, P.: Evidence-based volcanology: application to eruption crises, *J. Volcanol. Geoth. Res.*, 128, 273–285, 2003.
- Bayarri, M.J., and Berger, J.: P-values for composite null models, *J. Am. Stat. Assoc.*, 95, 1127–1142, 2000.
- Bear-Crozier, N., Miller, V., Newey, V., Horspool, N., and Weber, R.: Probabilistic Volcanic Ash Hazard Analysis (PVAHA) I: development of the VAPAH tool for emulating multi-scale volcanic ash fall analysis. *J. App. Volc.*, 5, 3, 2016.
- 280 Bebbington, M.: Trends and clustering in the onsets of volcanic eruptions, *J. Geophys. Res.*, 115, B01203, 2010.
- Bedford, T., and Cooke, R.: Probabilistic risk analysis: foundations and methods, Cambridge University Press, 2001.
- Berger, J.: The case for objective Bayesian analysis, *Bayesian Anal.*, 1, 1–17, 2004.
- Bevilacqua, A., Isaia, R., Neri, A., Vitale, S., Aspinall, W.P., Bisson, M., Flandoli, F., Baxter, P. J., Bertagnini, A., Esposti Ongaro, T., Iannuzzi, E., Pistolesi, M., and Rosi, M.: Quantifying volcanic hazard at Campi Flegrei caldera (Italy) with uncertainty assessment: I. Vent opening maps, *J. Geophys. Res. Solid Earth*, 120, 2309–2329, 2015.
- 285 Bommer, J. J., and F. Scherbaum: The use and misuse of logic trees in probabilistic seismic hazard analysis, *Earthq. Spectra* 24, 997-1009, 2008.
- Box, G.E.P.: Sampling and Bayes inference in scientific modelling and robustness, *Roy. Statist. Soc. Ser. A* 143, 383-430. 1980.
- 290 Castaños, H., and Lomnitz, C.: PSHA: Is it science? *Eng Geol*, 66, 315–317, 2002.
- de Finetti, B.: *Theory of Probability: A Critical Introductory Treatment*, John Wiley and Sons, London, 1974.
- Deligne, N. I., Coles, S. G., and Sparks, R. S. J.: Recurrence rates of large explosive volcanic eruptions. *J. Geophys. Res. Solid Earth*, 115, B06203, 2010.
- Draper, D., Hodges, J., Mallows, C., and Pregibon, D.: Exchangeability and data analysis (with discussion). *J. Roy. Statist. Soc, Ser. A*, 156, 9–37, 1993.
- 295 Galton, F.: *Vox populi*, *Nature*, 75, 450–451, 1907.
- Gelman, A., Carlin, J. B., Stern, H. S., and Rubin, D. B.: *Bayesian Data Analysis*, second edition, Chapman and Hall, London, 2003.
- 300 Gelman, A., and Shalizi C. R.: Philosophy and the practice of Bayesian statistics. *Br. J. Math. Stat. Psychol.*, 66, 8–38, 2013.

- Hacking, I.: *The Logic of Statistical Inference*, Cambridge University Press, Cambridge, 1965.
- IAEA: *Volcanic hazards in site evaluation for nuclear installations: specific safety guide*. Vienna: International Atomic Energy Agency, IAEA safety standards series, ISSN 1020–525X; no. SSG-21, 2012.
- 305 IPCC: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley, (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., 2013.
- Jaynes, E.T.: *Probability theory: the logic of Science*, Cambridge Univ. Press, NewYork, 2003.
- 310 Jenkins, S. F., Wilson, T. M., Magill, C., Miller, V., Stewart, C., Blong, R., Marzocchi, W., Boulton, M., Bonadonna, C., and Costa, A.: *Volcanic ash fall hazard and risk*. In: Susan Loughlin, Steve Sparks, Sarah Brown, Susanna Jenkins, Charlotte Vye-Brown (Eds.), *Global Volcanic Hazards and Risk*, pp. 173–222, 2015.
- Leith, C.E.: *Theoretical Skill of Monte Carlo Forecasts*, *Monthly Weather Rev.*, 102, 409–418, 1974.
- Lindley, D.V.: *The philosophy of statistics*, *Statistician* 49, 293-337, 2000.
- 315 Marti, J., Aspinall, W. P., Sobradelo, R., Felpeto, A., Geyer, A., Ortiz, R., Baxter, P., Cole, P., Pacheco, J., Blanco, M. J., and Lopez, C.: *A long-term volcanic hazard event tree for Teide-Pico Viejo stratovolcanoes (Tenerife, Canary Islands)*. *J. Volcanol. Geoth. Res.*, 178, 543–552, 2008.
- Martin, A.J., Umeda, K., Connor, C. B., Weller, J. N., Zhao, D., and Takahashi, M.: *Modeling long-term volcanic hazards through Bayesian inference: an example from the Tohoku volcanic arc, Japan*. *J. Geophys. Res.*, 109, B10208, 2004.
- 320 Marzocchi, W., Sandri, L., Gasparini, P., Newhall, C. N., and Boschi, E.: *Quantifying probabilities of volcanic events: the example of volcanic hazard at Mount Vesuvius*, *J. Geophys. Res.*, 109, B11201, 2004.
- Marzocchi, W., Sandri, L., and Selva, J.: *BET_EF: a probabilistic tool for long- and short-term eruption forecasting*, *Bull. Volcanol.*, 70 (5), 623–632, 2008.
- Marzocchi, W., Sandri, L., and Selva, J.: *BET_VH: a probabilistic tool for long-term volcanic hazard assessment*, *Bull. Volcanol.*, 72, 705–716, 2010.
- 325 Marzocchi, W., and Bebbington, M.: *Probabilistic eruption forecasting at short and long time scales*, *Bull. Volcanol.*, 74, 1777–1805, 2012.
- Marzocchi, W., and Jordan, T. H.: *Testing for Ontological Errors in Probabilistic Forecasting Models of Natural Systems*, *Proc. Nat. Acad. Sci.* 85, 955–959, 2014.
- 330 Marzocchi, W., Taroni, M., and Selva, J.: *Accounting for epistemic uncertainty in PSHA: logic tree and ensemble modeling*. *Bull. Seismol. Soc. Am.*, 105(4), 2151–2159, 2015.
- Marzocchi, W., and Jordan, T. H.: *A unified probabilistic framework for seismic hazard analysis*, *Bull. Seismol. Soc. Am.*, 107(6), 2738–2744, 2017.
- Marzocchi, W., and Jordan, T. H.: *Experimental concepts for testing probabilistic earthquake forecasting and seismic hazard models*, *Geophys. J. Int.*, 215, 780–798, 2018.
- 335 **MCDEM** [New Zealand Ministry of Civil Defence & Emergency Management] *Tsunami evacuation zones: director’s guideline for Civil Defence Emergency Management Groups*. (Ministry of Civil Defence & Emergency Management, 2008).
- Mulargia, F., Stark, P. B., and Geller, R. J.: *Why is probabilistic seismic hazard analysis (PSHA) still used?* *Phys. Earth Planet. Int.*, 264, 63–75, 2017.


- 340 Neri, A., Aspinall, W.P., Cioni, R., Bertagnini, A., Baxter, P. J., Zuccaro, G., Andronico, D., Barsotti, S., Cole, P. D., Esposti Ongaro, T., Hincks, T. K., Macedonio, G., Papale, P., Rosi, M., Santacroce, R., and Woo, G.: Developing an Event Tree for Probabilistic Hazard and Risk Assessment at Vesuvius, *J. Volcan. Geoth. Res.*, 178, 397–415, 2008.
- Newhall, C. G., and Hoblitt, R.: Constructing event trees for volcanic crises. *Bull. Volcanol.* 64, 3–20, 2002.
- Newhall, C. G., and Pallister, J. S.: Using Multiple Data Sets to Populate Probabilistic Volcanic Event Trees, in: *Volcanic Hazards, Risks and Disasters*, Ed(s): J. F. Shroder, P. Papale, Elsevier, 203–232, 2015.
- 345 NRC, National Research Council Panel on Seismic Hazard Evaluation: Review of Recommendations for Probabilistic Seismic Hazard Analysis: Guidance on Uncertainty and Use of Experts, National Academy of Sciences, Washington, D.C., 84 pp., ISBN:0-309-56207-4, 1997.
- Poland, M. P., and Anderson, K. R.: Partly cloudy with a chance of lava flows: Forecasting volcanic eruptions in the twenty-first century, *J. Geophys. Res. Solid Earth*, 125, e2018JB016974, 2020.
- 350 Reilly, A. C., Baroud, H., Flage, R., and Gerst, M. D.: Sources of uncertainty in interdependent infrastructure and their implications, *Reliability Engineering & System Safety*, 213, 107756, doi:10.1016/j.ress.2021.107756, 2021.
- Rougier, J. C.: Quantifying hazard losses. In: *Risk and uncertainty assessment for natural hazards*, eds. J. Rougier, S. Sparks, and L. Hill. Cambridge University press, Cambridge, 19–39, 2013.
- 355 Rougier, J. C., and Beven, K. J.: Model and data limitations: the sources and implications of epistemic uncertainty. In: *Risk and uncertainty assessment for natural hazards*, eds. J. Rougier, S. Sparks, and L. Hill. Cambridge University press, Cambridge, 40–63, 2013.
- Rubin, D.B.: Bayesian justifiable and relevant frequency calculations for the applied statistician, *Ann Stat*, 12, 1151–1172, 1984.
- 360 Sandri, L., Costa, A., Selva, J., Macedonio, G., Tonini, R., Folch, A., and Sulpizio, R.: Beyond eruptive scenarios: tephra fall from Neapolitan volcanoes, *Sci. Rep.*, 6, 24271, 2016.
- Scherbaum, F., and Kuehn N.M.: Logic tree branch weights and probabilities: Summing up to one is not enough, *Earthq. Spectra*, 27, 1237-1251, 2011.
- Selva, J., Costa, A., Marzocchi, W., and Sandri, L.: BET_VH: Exploring the influence of natural uncertainties on long-term hazard from tephra fallout at Campi Flegrei (Italy), *Bull. Volcanol.*, 72, 717–733, 2010.
- 365 Selva, J., Argyroudis, S., and Pitilakis, K.: Impact on loss/risk assessments of inter-model variability in vulnerability analysis, *Natural Hazards* 67, 723–746, 2013.
- Selva, J., Costa, A., De Natale, G., Di Vito, M. A., Isaia, R., and Macedonio, G.: Sensitivity test and ensemble hazard assessment for tephra fallout at Campi Flegrei, Italy, *J. Volcanol. Geotherm. Res.*, 351, 1–28, 2018.
- 370 Silver, N.: *The Signal and the Noise: Why So Many Predictions Fail-but Some Don't*, The Penguin Press, New York, New York, 2012.
- Sobradelo, R., and Marti, J.: Bayesian event tree for long-term volcanic hazard assessment: application to Teide-Pico Viejo stratovolcanoes, Tenerife, Canary Islands, *J. Geophys. Res.*, 115, B05206, 2010.
- Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L. eds.: *Climate change 2007: the physical science basis: contribution of Working Group I to the fourth assessment report of the intergovernmental panel on climate change*. Cambridge and New York, Cambridge University Press, 2007.
- 375 Sparks, R.S.J.: Forecasting volcanic eruptions, *Earth Plan. Sci. Lett.*, 210, 1–15, 2003.

Stark, P. B.: Pay no attention to the model behind the curtain, in *Significant Digits: Responsible Use of Quantitative Information*, A. Saltelli and Á. Guimarães Pereira (Editors), Megaloceros Press, University of Bergen, Bergen, Germany, 2017.


380 Stein, S., and Stein, J. L.: Shallow versus deep uncertainties in natural hazard assessments. *EOS, Tran. AGU*, 94, 133–140, 2013.

Tierz, P., Loughlin, S. C., and Calder, E. S.: VOLCANS: an objective, structured and reproducible method for identifying sets of analogue volcanoes, *Bull. Volcan.*, 81, 76, 2019.

385



390 **Figure 1: The left panel shows the mean tephra fall hazard curves relative to one point inside the Campi Flegrei (Figure 2) for each BET implementation described in Selva et al. (2018); colors indicate each implementation. The right panel shows the PoE relative to the tephra load threshold of 300 kg/m² (vertical dashed line in the upper panel), and the weight of each assessment.**



395

Figure 2: The location reference site for this study (red star) along with the hazard map relative the probability of 5% of exceedance, conditional upon the occurrence of one eruption of whatever size and from whatever vent at Campi Flegrei (mean of the epistemic uncertainty). The figure has been obtained modifying the Figure 11 of the corrigendum to the paper Selva et al., 2018 (available at doi.org/10.1016/j.jvolgeores.2018.07.008); the original maps have been obtained with the MATLAB software.

400




Figure 3: EED of each BET configuration relative to a tephra load threshold of 300 kg/m² for the site in Figure 2.

405




Figure 4: The overall EED obtained from the average of the five distributions of Figure 3. The EED is a Beta distribution with parameters given by equations 2 and 3. The histogram (right y-axis) shows the weight of each one of the five distributions.