

# ~~Compound~~ Characterizing multivariate coastal ~~flood risk~~ flooding events in a semi-arid ~~urbanized~~ region: The implications of copula choice, sampling, and infrastructure

Joseph T.D. Lucey<sup>1</sup> and Timu W. Gallien<sup>1</sup>

<sup>1</sup>Department of Civil and Environmental Engineering, University of California, Los Angeles, CA 90095, USA

**Correspondence:** Timu Gallien (tgallien@seas.ucla.edu)

**Abstract.** ~~Sea level rise will increase the frequency and severity of coastal flooding events. Compound coastal flooding~~ Multivariate coastal flooding is characterized by multiple flooding pathways (i.e., high offshore water levels, streamflow, energetic waves, precipitation) acting concurrently. This study explores the joint ~~flood~~ risks caused by the co-occurrence of high marine water levels and precipitation in a highly urbanized semi-arid, tidally dominated region. A novel structural function developed from the multivariate analysis is proposed to consider the implications of flood control infrastructure in ~~compound~~ multivariate coastal flood risk assessments. Univariate statistics are analyzed for individual sites and events. Conditional ~~;~~ and joint probabilities are developed using a range of copulas ~~and sampling methods. The Independent, and Cubic copulas produced poor results while the Fischer-Koek,~~ sampling methods, and hazard scenarios. The Nelsen, BB1, BB5, and Roch-Alegre were selected based on a Cramer-von Mises test and generally produced robust results across a range of sampling methods. The impacts of sampling are considered using annual maximum, annual coinciding, wet season monthly ~~coinciding~~ maximum, and wet season monthly ~~maximum coinciding~~ sampling. Although annual maximum sampling is commonly ~~recommended for characterizing compound~~ used for characterizing multivariate events, this work suggests annual maximum sampling ~~does not produce “worst-case” event pairs and substantially underestimates~~ may substantially underestimate marine water levels for extreme events. ~~Wet season coinciding water~~ Water level and precipitation combinations from wet season monthly coinciding sampling benefits from a dramatic increase in data pairs ~~;~~ improved goodness-of-fit statistics, and provide a range of physically realistic pairs. Wet season monthly coinciding sampling may provide a more accurate ~~compound multivariate~~ compound multivariate flooding risk characterization for long return periods in semi-arid regions. Univariate, conditional, and bivariate results emphasize the importance on proper event definition as this significantly influences the associated event risks.

## 1 Introduction

Coastal flooding is a significant human hazard (Leonard et al., 2014; Wahl et al., 2015) and is considered a primary health hazard by the U.S. Global Change Research Program (Bell et al., 2016). Coastal migration and utilization continues to increase (Nicholls et al., 2007; Nicholls, 2011). Over 600 million people populate coastal zones (Merkens et al., 2016). Climate change-induced sea level rise will substantially increase flood risk (Church et al., 2013; Horton et al., 2014), and negatively impact coastal populations (Bell et al., 2016). Even relatively modest sea level rise will significantly increase flood frequencies through

25 the US (e.g., Tebaldi et al., 2012; Taherkhani et al., 2020). Southern California is particularly vulnerable to sea level rise. Small changes in sea level ( $\sim 5$  cm) double the odds of the 50-year flooding event (Taherkhani et al., 2020) and the 100-year event is expected to become annual by 2050 (Tebaldi et al., 2012). Regional research has explored flood risks caused by sea level rise and coastal forcing (e.g., Heberger et al., 2011; Hanson et al., 2011; Gallien et al., 2015). However, accurately characterizing future, non-stationary coastal vulnerability requires considering the joint and potentially nonlinear impacts of  
30 compound (marine and hydrologic) events (Gallien et al., 2018).

Compound coastal flooding considers the combined impacts of marine and hydrologic ~~forcing~~forcings, typically within a physically relevant time window, and are considered multivariate events. Typical events, such as precipitation or high water levels, occurring simultaneously may combine to generate extreme events (Seneviratne et al., 2012). In urban coastal settings multiple flooding pathways (i.e., high marine water levels, wave runoff and overtopping, large fluvial flows, and pluvial  
35 flooding from precipitation) interact with infrastructure (e.g., sea walls, human-made dunes, and the storm system) potentially exacerbating hazards. Notably, ~~compound-multivariate~~ events that share a common return period may produce vastly different flooding outcomes. Traditionally, literature has focused on river discharge or storm surge dominated ~~compound-multivariate~~ events (Table 1).

From a flood risk perspective there are multiple methods to characterize events. A univariate approach is often used where  
40 a single variable (e.g., water level) is considered. For example, FEMA recommends characterizing ~~compound-multivariate~~ events by developing univariate water level and discharge statistics ~~, modeling each separately,~~ and then adopting ~~the more severe flooding~~a smooth, blended result for transitional areas (FEMA, 2011, 2016c). This can lead to underestimating flood risk because of the interplay between two flood pathways (i.e., a high tail water forces fluvial flooding upstream). Conditional probabilities represent an alternative where the ~~compound-multivariate~~ flood risk can be evaluated given available information  
45 on a primary variable (e.g., water level) to determine the exceedance probability of a secondary variable (e.g., precipitation) (Shiau, 2003; Karmakar and Simonovic, 2009; Zhang and Singh, 2012; Li et al., 2013; Mitková and Halmová, 2014; Serinaldi, 2015, 2016; Anandalekshmi et al., 2019). A third method uses copulas to analyze the dependence of multiple flood drivers and develop joint statistics.

Numerous studies have used a copula based approach to study floods manifested by various combinations of variables  
50 (Table 1). ~~Compound-Multivariate~~ flood risks can be described and quantified from previous copula studies (Salvadori, 2004; Salvadori and De Michele, 2004, 2007; Salvadori et al., 2011, 2013, 2016). Multivariate inland and coastal hydrology analysis have primarily focused on a small group of copulas: Archimedean (Clayton, Frank, and Gumbel), Student t, and Gaussian copulas. Alternative copulas may more accurately characterize urban coastal flooding (Jane et al., 2020). Specifically, hazard scenarios provide various perspectives on critical multivariate events (Salvadori et al., 2016). ~~Current studies are often limited to~~Initial studies were primarily focused on select hazard scenarios (Table 1 in Salvadori et al. (2016)).  
55

Data sampling methods in multivariate studies influence distribution fitting. Two primary sampling methods exist: peaks over threshold (Jarušková and Hanek, 2006) and block maxima (Engeland et al., 2004). Events selected using peaks over threshold sampling are above a predetermined threshold defining an “extreme” event. Block maxima sampling uses various block sizes (yearly, seasonal, semiannual, etc.) to separate and select the maximum event per block. Engeland et al. (2004) reports a

60 significantly different 1,000-year streamflow when using 12-month block sampling ( $160 \text{ m}^3\text{s}^{-1}$ ) compared to using a threshold  
of  $50 \text{ m}^3\text{s}^{-1}$  ( $120 \text{ m}^3\text{s}^{-1}$ ). Many studies utilize block maxima sampling with a yearly block size, i.e. the annual maximum  
sampling method (Baratti et al., 2012; Bezak et al., 2014; Wahl et al., 2015). This method is specifically recommended by  
FEMA (2016c) for evaluating coastal hazards. Alternatively, studies identify extremes in the primary variable using annual  
65 maximum sampling and select a secondary variable which co-occurs with the primary variable (Lian et al., 2013; Xu et al.,  
2014; Tu et al., 2018), creating a “coinciding” type sampling. Multivariate applications relying upon annual maximum sampling  
generate a “~~worse-case-scenario~~ events with severe flooding potentials” which may produce unrealistic variable combinations.  
Coinciding sampling draws from physically realistic event pairs. Although, multiple studies explore sampling effects on fitted  
distribution parameters and univariate return periods (Engeland et al., 2004; Jarušková and Hanek, 2006; Peng et al., 2019;  
Juma et al., 2020), sampling effects for multivariate coastal flooding events is unknown.

70 Coastal flooding studies primarily focus on locations defined by storm surge dominated oceanographic conditions with  
warm, humid (Wahl et al., 2012; Lian et al., 2013; Xu et al., 2014; Masina et al., 2015; Wahl et al., 2015; Mazas and Hamm,  
2017; Paprotny et al., 2018; Tu et al., 2018; Bevacqua et al., 2019; Didier et al., 2019; Xu et al., 2019; Yang et al., 2020),  
and monsoonal (Jane et al., 2020) climatic conditions (i.e. Köppen-Geiger system, Beck et al., 2018). In contrast, along the  
southern California coast typical tidal variability is 1.7 to 2.2 m (Flick, 2016) and storm surge rarely exceeds  $\sim 20$  cm (Flick,  
75 1998). Notably, during the wet season (October to March), when precipitation typically occurs, spring tide ranges are relatively  
large ( $\sim 2.6$  m). Critically, few studies consider areas where coastal flooding events are dominated by large tides and either  
precipitation or wave events (Masina et al., 2015; Mazas and Hamm, 2017; Didier et al., 2019; Jane et al., 2020). This study  
explores univariate and multivariate flooding events in a semi-arid, tidally dominated, highly urbanized region. Here, the  
dependency between observed water levels and precipitation, impacts of sampling methods and distribution fitting, and the  
80 most-likely-resulting flood values are explored.

## 2 Site description & data

This study considers observed water level and precipitation influences for coastal ~~compound~~ multivariate events at Santa  
Monica (SM), Sunset Beach (S), and LA Jolla (SD) areas in Los Angeles, Huntington Beach, and San Diego, California  
(Fig. 1); three semi-arid, tidally dominated sites in the US. All are low-lying estuarine or bay-backed highly urbanized beach  
85 communities requiring extensive coastal management to mitigate flooding events. For example, sea walls and artificial berms  
in Sunset Beach protect infrastructure from high embayment water levels, wave runup, and overtopping along the open coast.  
The storm drain network is managed to prevent back flooding during high tides. Notably, Gallien et al. (2014) suggested when  
tide valves are closed, the storm drain network cannot reduce pluvial flooding caused by alternative flooding pathways (e.g.,  
precipitation or wave overtopping). Pacific Coast Highway (PCH) is heavily utilized and is a primary transportation corridor  
90 along the southern California coastline. All locations are densely urbanized and highly impacted by flooding.

Observed water levels from the Los Angeles (Station ID: 9410660), La Jolla (Station ID: 9410230), and Santa Monica  
(Station ID: 9410840) tide gauges are available on NOAA’s Tides and Currents for daily high-low, hourly, or six minute

**Table 1.** [List A non-exhaustive list](#) of multivariate studies which utilized copulas to study the associated variables


Variable Pairs	References
Waves and water level	Masina et al. (2015); Mazas and Hamm (2017); Didier et al. (2019)
Waves and storm duration	De Michele et al. (2007); Salvadori et al. (2014, 2015)
Waves and storm surge	Wahl et al. (2012); Paprotny et al. (2018)
River discharge and water level	White (2007); Bray and McCuen (2014); Sadegh et al. (2018); Ganguli and Merz (2019a, b)
River discharge and storm surge	Paprotny et al. (2018); Ganguli et al. (2020)
River discharge and volume	Yue (2001a, b); Shiau (2003); Favre et al. (2004); De Michele et al. (2005); Poulin et al. (2007); Li et al. (2013); Salvadori et al. (2013); Requena et al. (2013); Aghakouchak (2014)
River discharge, rainfall, and water level	Bray and McCuen (2014); Jeong et al. (2014)
Multiple river discharges	Salvadori and De Michele (2010)
Rainfall and tide	Lian et al. (2013)*, Xu et al. (2014)*, Tu et al. (2018)*, Xu et al. (2019)*, Bevacqua et al. (2020), Yang et al. (2020)*
Rainfall and water levels	Jane et al. (2020)
Rainfall and storm surge	Wahl et al. (2015); Paprotny et al. (2018); Bevac- qua et al. (2019)
Rainfall intensity and depth	Yue (2000a, b, 2002); De Michele and Salvadori (2003)
Rainfall and groundwater	Anandalekshmi et al. (2019)
Rainfall and runoff	Zhang and Singh (2012); Hao and Singh (2020)
Rainfall and river discharge	Zhong et al. (2020)
Rainfall and temperature	Zhang et al. (2017)
Rainfall and duration	Salvadori and De Michele (2007)
Combinations of rainfall intensity, depth, and duration	Zheng et al. (2014)
Combinations of river discharge, vol- ume, and duration	Karmakar and Simonovic (2009); Reddy and Gan- guli (2012); Ganguli and Reddy (2013); Gräler et al. (2013); Mitková and Halmová (2014)

\*Note these studies use the term tide measurement but actually represent observed water level measurements. Please refer Section 2 for clarification.

intervals (NOAA, Accessed 2021d). Verified hourly water levels (m NAVD88) had the longest record length at all three stations and provided an additional 31-years of observations overlapping precipitation data for Los Angeles and La Jolla, and 6-years  
95 for Santa Monica. The resulting observations windows are November 22, 1973 to December 19, ~~2023~~ 2013 for Santa Monica, July 1, 1948 to December 1, 2012 for Sunset and July 1, 1948 to December 19, 2013 for San Diego (Table 2). It is worth noting, that within the body of ~~compound-multivariate~~ flooding literature, the terms tide and water level may be interchanged (e.g., Lian et al., 2013; Xu et al., 2014; Tu et al., 2018; Xu et al., 2019; Yang et al., 2020). This is a key distinction since compound event dependencies may change depending on water level selection. Recent efforts have been made to standardize language where  
100 tide represents only the astronomical changes in water levels and storm surge specifically excludes astronomical variability and consists only of the inverse barometric effects along with wind and wave setup (Gregory et al., 2019). In this study, the term observed water level (OWL) is adopted. OWL is the water level measured at the NOAA tide gauges which includes all tidal, storm, and climatic effects.

The U.S. Hourly Precipitation Data dataset provided by NOAA's National Centers for Environmental Information (NOAA,  
105 Accessed 2021c) at the Signal Hill (COOP:048230), Los Angeles International Airport (COOP:045114), and San Diego International Airport (COOP:047740) stations is used as the precipitation inputs. Observations do not contain trace amounts ( $< 0.25$  mm) and are provided as cumulative precipitation (mm) per event. Precipitation measurements were converted to a mm/hr rate by dividing the total event precipitation by the event ~~time-duration~~ to match the hourly OWL measurements. The final precipitation input is a 24-hour cumulative precipitation record made from the hourly observations. All data was transformed  
110 to UTC for analysis.

~~Compound~~ Multivariate flood probabilities are determined with combinations of sampling methods: Annual Maximum (AM), Annual Coinciding (AC), Wet Season Monthly Maximum (WMM), and Wet Season Monthly Coinciding (WMC). AM sampling pairs the single largest precipitation and OWL observations within a given year (without regard to co-occurrence), where AC sampling pairs the single largest precipitation observation within a given year to the largest OWL observation within  
115 its 24-hour accumulation period. Each sampling method samples from a unique probability space and therefore will provide varying perspectives for a return period. A summary of each sites' associated gauges, observation windows, and number of pairs is provided in Table 2. Southern California's wet season is defined between October to March and provides a majority of the total annual rainfall (Cayan and Roads, 1984; Conil and Hall, 2006). It is likely for extreme ~~compound-multivariate~~ events to occur during this period. ~~Wet-season-monthly-maximum~~ Maximum sampling pairs the single largest precipitation and  
120 OWL observations within each year or wet season month. A multivariate event created with the largest observed precipitation and OWL within a year or wet season month can result in an event with severe flooding potential. Although strictly speaking maximum pairings (annual or wet season) do not technically represent an observed ~~compound-events~~ since the co-occurrence of precipitation and water levels follows the FEMA guidance for considering a "worse case scenario" approach (FEMA, 2016e) ~~Wet-season-monthly-coinciding~~ multivariate event, they would represent a severe event and are consistent with the blended  
125 approach recommended by FEMA (2016c). Coinciding sampling pairs the single largest precipitation observation within each year or wet season month to the largest OWL observation within its 24-hour accumulation period, providing more realistic pairs compared to maximum sampling.



**Figure 1.** Map displaying (a) Santa Monica, (b) Sunset, and (c) San Diego sites along with locations of tide gauges (triangle) and precipitation stations (circle). The road drain (square) and boundary (yellow) at Sunset ( $\sim 2 \text{ km}^2$ ) is for the Structural scenario. Aerial imagery from NOAA (Accessed 2021a).

**Table 2.** Water level and precipitation observations at Santa Monica (SM), Sunset (S), and San Diego (SD) using annual maximum (AM), annual coinciding (AC), wet season monthly maximum (WMM), and wet season monthly coinciding (WMC) samplings

Site	Tide Gauge	Precip. Gauge	Observation Window	AM Pairs	AC Pairs	WMM Pairs	WMC Pairs
SM	9410840	045114	November 22, 1973 to December 19, 2013	40	38	193	191
S	9410660	048230	July 1, 1948 to December 1, 2012	63	63	257	258
SD	9410230	047740	July 1, 1948 to December, 19 2013	65	60	328	329


Distributions are fit with existing precipitation observations greater than zero consistent with previous studies (Swift Jr and Schreuder, 1981; Hanson and Vogel, 2008). Months with no OWL measurements were excluded. In the case of coinciding sampling, pairs that had three or more OWL measurements missing within the 24-hour window were manually reviewed and removed if their tidal peak was clearly missing. Specifically for WMM sampling, months with more than half their observations missing were also reviewed and removed if the tidal peak was missing. The resulting data pairs are shown in Fig. 2.

### 3 Methods

#### 3.1 Univariate, bivariate, & conditonal distributions

Potential flooding events are determined with three different probability definitions: univariate, conditional, and bivariate. Assuming  $X$  and  $Y$  are random variables,  $x$  and  $y$  are observations of these variables, and  $F_X$  and  $F_Y$  represent the variables' re-





**Figure 2.** Data pairs for each sampling method. ~~Annual Maximum~~annual maximum (AM,  $\times$ ), ~~Annual Coinciding~~annual coinciding (AC,  $+$ ), ~~Water Year Monthly Coinciding~~wet season monthly maximum (WMC,  $\triangle$ ), ~~Water Year Monthly Maximum~~and wet season monthly coinciding (WMM,  $\bullet$ ) at (a) Santa Monica, (b) Sunset, and (c) San Diego

spective cumulative distribution functions (CDF). Formulations for univariate ( $F_X(x), F_Y(y)$ ) and bivariate joint ( $F_{X,Y}(x,y)$ ) CDFs follow DeGroot and Schervish (2014) (Eq. (1) and (2)). Conditionals ( $F_{X|Y \geq y}(x|Y \geq y)$ ,  $F_{X|Y \leq y}(x|Y \leq y)$ , and  $F_{X|Y=y}(x|Y=y)$ ) are developed from Shiau (2003) (Eq. (3)) and Serinaldi (2015) (Eq. (4) and (5)). Conditionals 1 (C1), 2 (C2), and 3 (C3) represent Eq. (3), (4), and (5) going forward. Univariate statistics are developed using the appropriate continuous random variable distribution while conditional and bivariate CDFs are determined using copulas.

Copulas are functions that associate random variables' univariate CDFs to their joint CDF (e.g.,  $F_X$  and  $F_Y$  to  $F_{X,Y}(x,y)$ ) according to Sklar's theorem (Sklar, 1959; Salvadori, 2004). There is no requirement for the univariate distributions to be the same. This is particularly advantageous since the optimal univariate distributions may be used for each variable. Bivariate probabilities for different hazard scenarios, which represent various multivariate events, and conditional probabilities can be calculated using fitted copula functions.

$$F_X(x) = Pr(X \leq x) \quad (1)$$

$$F_{X,Y}(x,y) = Pr(X \leq x \text{ and } Y \leq y) \quad (2)$$

$$F_{X|Y \geq y}(x|Y \geq y) = Pr(X > x|Y \geq y) = \frac{F_X(x) - F_{XY}(xy)}{1 - F_Y(y)} \quad (3)$$

$$150 \quad F_{X|Y \leq y}(x|Y \leq y) = Pr(X > x|Y \leq y) = 1 - \frac{F_{X,Y}(x,y)}{F_Y(y)} \quad (4)$$

$$F_{X|Y=y}(x|Y=y) = Pr(X > x|Y=y) = 1 - \frac{\partial F_{XY}(xy)}{\partial y} \quad (5)$$

### 3.2 Hazard scenarios

Notation and definitions from Salvadori et al. (2016), unless otherwise stated, are used to define the upper set ( $S$ ) and scenario types. Salvadori et al. (2016) and Serinaldi (2015) present figures of each scenario's probability space. Further discussion of  
 155 hazard scenarios and copulas assume a bivariate situation.

#### 3.2.1 "OR"

"OR" scenario events have one or both random variables exceed a specified threshold. That is, what is the probability of a water level or precipitation event exceeding a given value? Standard univariate CDFs make up the associated copula.

$$\alpha_x^\vee = \mathbf{P}(\mathbf{X} \in S_x^\vee) = 1 - \mathbf{C}(\mathbf{F}_1(x_1), \dots, \mathbf{F}_d(x_d)) \quad (6)$$

#### 160 3.2.2 "AND"

"AND" scenario events have both random variables exceed a specified threshold. In this case the fundamental question is "what is the probability of a particular water level and precipitation rate exceeding specified values?". The survival copula ( $\hat{\mathbf{C}}(u, v)$ ) is comprised of univariate survival CDFs ( $\bar{\mathbf{F}}(x) = 1 - \mathbf{F}(x)$ ) and the provided equation can be found in Serinaldi (2015) and Salvadori and De Michele (2004).

$$165 \quad \alpha_x^\wedge = \mathbf{P}(\mathbf{X} \in S_x^\wedge) = \hat{\mathbf{C}}(\bar{\mathbf{F}}_1(x_1), \dots, \bar{\mathbf{F}}_d(x_d)) \quad (7)$$

$$\hat{\mathbf{C}}(u, v) = 1 - u - v + C(u, v) \quad (8)$$

#### 3.2.3 "Kendall"

The "Kendall" (K) scenario highlights an infinite set of OR events that separate the subcritical (i.e., "safe") and supercritical (i.e., "dangerous") statistical regions. In the OR scenario, events along an isoline ( $t$ ) share a common probability, but define



170 separate regions. Events along a Kendall  $t$  represent the same super critical region (Serinaldi, 2015) and provide a “safety lower bound” (Salvadori et al., 2011). Essentially the Kendall scenario considers the minimum OR events of concern.  $\mathbf{K}(t)$  is estimated by a method outlined in Salvadori et al. (2011).

$$\mathbf{K}(t) = \mathbf{P}(\mathbf{F}(X_1, \dots, X_d) \leq t) = \mathbf{P}(\mathbf{C}(F_1(X_1), \dots, F_d(X_d)) \leq t) \quad (9)$$

$$\alpha_t^{\mathbf{K}} = \mathbf{P}(\mathbf{X} \in S_t^{\mathbf{K}}) = 1 - \mathbf{K}(t) \quad (10)$$

### 175 3.2.4 “Survival Kendall”

“Survival Kendall” (SK) scenario highlights an infinite set of AND events which also separate safe and dangerous statistical spaces. AND events along a  $t$  also share a common probability, but define separate regions. Events along an SK  $t$  represent the same super critical region, but provide an “(upper) bounded safe region” (Salvadori et al., 2013). The Survival Kendall specifically considers the largest AND events of concern and is estimated by the method outlined in Salvadori et al. (2013).

$$180 \quad \hat{\mathbf{K}}(t) = \mathbf{P}(\bar{\mathbf{F}}(X_1, \dots, X_d) \leq t) = \mathbf{P}(\hat{\mathbf{C}}(\bar{F}_1(X_1), \dots, \bar{F}_d(X_d)) \leq t) \quad (11)$$

$$\alpha_t^{\hat{\mathbf{K}}} = \mathbf{P}(\mathbf{X} \in S_t^{\hat{\mathbf{K}}}) = 1 - \hat{\mathbf{K}}(t) = \hat{\mathbf{K}}(t) \quad (12)$$

### 3.2.5 “Structural”

The “Structural” scenario considers the probability of an output from a structural function,  $\Psi(\mathbf{X})$ , exceeding a design load or capacity ( $z$ ) (Salvadori et al., 2016). For example, De Michele et al. (2005) and Volpi and Fiori (2014) used a structural  
 185 function to evaluate a dam spillway while Salvadori et al. (2015) considers the preliminary design of rubble mound breakwater. In this work, the structural failure function focuses on the question “what is the probability of a water level forcing tide valve closure and subsequent flooding during a precipitation event?”.

$$\alpha_z^{\Psi} = \mathbf{P}(\mathbf{X} \in S_z^{\Psi}) = \mathbf{P}(\Psi(\mathbf{X}) > z) \quad (13)$$

## 3.3 MvCAT

190 The Multivariate Copula Analysis Toolbox (MvCAT) developed by Sadegh et al. (2017) is a publicly available MATLAB toolbox that fits 25 different copula functions to user data of two random variables. Copula parameters are optimized through a local optimization or with Markov Chain Monte Carlo methods (details in Sadegh et al. (2017)). The MvCAT framework is expanded to determine all scenarios and conditionals in this study. While all copulas have functional CDFs, the Cuadras-Auge,

Raftery, Shih-Louis, Linear-Spearman, Fischer-Hinzmann, Husler-Reiss, Cube, and Marshal-Olkin copulas do not have a PDF function. A PDF is required to determine the most likely value along an isoline, therefore it was decided to remove those copulas from the study. Copulas must also be computationally simple to derive or integrate to calculate Conditional 3 (Eq.(5)). Gaussian and Student t copulas' partial derivatives cannot be explicitly calculated, and estimates induce unrealistic errors (i.e., produce negative probabilities). Seventeen different copulas remain after eliminating those discussed above.

### 3.4 Return periods

Hydrologic events are commonly cast in the context of return periods (e.g., De Michele et al., 2005, 2007; FEMA, 2011; Wahl et al., 2012; USACE, 2013; Salvadori et al., 2014; Wahl et al., 2015; Salvadori et al., 2015). Return periods ( $T$ ) provide an easily interpreted metric describing the severity of an event and is the inverse of an event's probability of exceedance presented as  $F$  in Eq. (14) (Tu et al., 2018). In Eq. (15),  $N$  is the data's time window,  $n$  is the number of considered events within  $N$ , and  $N_e$  is the average number of events per unit of time (monthly, yearly, etc.). Therefore,  $N_e = 1$  when considering singular events within a year (Tu et al., 2018).

$$T = 1/(N_e * F) \quad (14)$$

$$N_e = n/N \quad (15)$$

### 3.5 Goodness of fit metrics

Multiple goodness of fit metrics and correlations serve to quantify the quality of distribution fits and dependencies between variables. Marginal and copula-fits are selected by Bayesian Information Criterion (BIC; Eq. (19)) and maximum likelihood (ML; Eq. (16)) values, respectively. must pass the Chi-square goodness-of-fit test at standard significance levels ( $\alpha = 0.05$ ), unless otherwise stated. Copulas are selected by BIC and must pass the Cramer-von Mises test (Genest et al., 2009; Couasnon et al., 2018; Sadegh Likelihood ( $\mathcal{L} \in [0, \infty)$ ) measures how well a distribution's estimated parameters fit the sample data with larger values suggesting a better fit. Log-likelihood ( $\ell \in (-\infty, \infty)$ ) is the log transformation of Eq. (16) used to calculate BIC. BIC ( $BIC \in (-\infty, \infty)$ ) is similar to the likelihood, but penalizes for the number of estimated parameters ( $D$ ) and the data's sample size ( $n$ ). Smaller BIC values represent a better fit. Equations and definitions can be found in Sadegh et al. (2017). Correlation measurements include Pearson's linear correlation, Kendall's tau, and Spearman's rho coefficients.

$$\mathcal{L}(\theta|\tilde{Y}) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi\tilde{\sigma}^2}} \exp\left\{-\frac{1}{2}\tilde{\sigma}^2[\tilde{y}_i - y_i(\theta)]^2\right\} \quad (16)$$

$$\tilde{\sigma}^2 = \frac{\sum_{i=1}^n [\tilde{y}_i - y_i(\theta)]^2}{n} \quad (17)$$

$$220 \quad \ell(\boldsymbol{\theta}|\tilde{\mathbf{Y}}) = -\frac{n}{2}\ln(2\pi) - \frac{n}{2}\ln\tilde{\sigma}^2 - \frac{1}{2}\tilde{\sigma}^2\sum_{i=1}^n[\tilde{y}_i - y_i(\boldsymbol{\theta})]^2 \quad (18)$$

$$BIC = D\ln(n) - 2\ell \quad (19)$$

## 4 Results


Univariate, conditional, and bivariate probabilities were developed using four sampling methods (~~AC, AM, WMC, WMMAM,~~  
AC, WMM, and WMC) and seventeen different copulas. Two marginal distributions do not pass the chi square test at the  
 225 standard 0.05 level of significance (San Diego AM OWL and Santa Monica WMM OWL). These distributions pass at reduced  
 significance levels of 0.01. Four copulas almost always passed (Nelsen, BB1, BB5, and Roch-Alegre) the Cramer-von Mises  
test and are used for analysis. It is noted the Roch-Alegre (Roch.) did not pass at Sunset for WMM sampling, and the BB1 and  
BB5 did not pass at San Diego for WMC sampling. Additionally, Santa Monica's AM data is slightly negatively correlated  
 (> -0.06). Copula and sampling effects differ significantly at low (i.e., low return period) and high (i.e., severe return period)  
 230 probabilities of non-exceedance. In the case of annual sampling, non-exceedance (exceedance) probabilities are 0.9 (0.1) and  
 0.99 (0.01) for the 10- and 100-year events, respectively. In wet season sampling, return period exceedance probabilities vary  
 depending on sampling type and location due to the average number of event observations per year ( $N_e$  from Eq. (15)). For  
 example, San Diego WMC sampling has 329 observations within the 65 year record (i.e.,  $N_e = 5.06$ ). Therefore, the exceedance  
 probabilities ( $F$  in Eq. (14)) associated to a 10- and 100-year event are 0.0198 and 0.0020 (non-exceedance probabilities at  
 235 0.9802 and 0.9980), respectively. Table 3 presents wet season (WMM and WMC) exceedance probabilities for all sites.

**Table 3.** Santa Monica, Sunset, and San Diego exceedance probabilities at the 10- and 100-year return periods for wet season monthly  
maximum (WMM) and wet season monthly coinciding (WMC) samplings.

	Santa Monica		Sunset		San Diego	
	10-year	100-year	10-year	100-year	10-year	100-year
WMM	0.0207	0.0021	0.0245	0.0025	0.0198	0.0020
WMC	0.0209	0.0021	0.0244	0.0024	0.0198	0.0020

### 4.1 Marginals

The selected marginal distributions (Fig. 3, Table 4) were tested and/or suggested fits in previous studies. Rainfall has been  
 widely fit with an Exponential distribution (refer to Table 2 in Salvadori and De Michele (2007)), but more recently been fit  
 using a variety of distributions including Gamma (Husak et al., 2007), Rayleigh (Pakoksung and Takagi, 2017; Esberto, 2018),  
 240 Generalized Pareto or Birnbaum-Saunders (Ayantobo et al., 2021). In the case of annual precipitation ~~sampling~~-(coinciding



**Figure 3.** (a) OWL and (b) precipitation marginals for Santa Monica (SM, solid lines), Sunset (S, dashed lines), and San Diego (SD, dotted lines) using AM-annual maximum (AM, blue), AC-annual coinciding (AC, green), WMM-wet season monthly maximum (WMM, black), and WMC-wet season monthly coinciding (WMC, red) samplings.

or maximum sampling) Santa Monica was well described by a Birnbaum-Sanders, Rayleigh best described Sunset data, and a Gamma was the best fit for San Diego data. Similarly, wet season precipitation data (maximum or coinciding sampling) was best described by the Exponential distribution for Santa Monica and Sunset while the Generalized Pareto best represented San Diego.

245 Historically, water levels have been described using a number of distributions including Normal (Hawkes et al., 2002), Generalized Pareto (Mazas and Hamm, 2017), Log Logistic and Nakagami (Sadegh et al., 2018), Birnbaum-Saunders (Sadegh et al., 2018; Didier et al., 2019; Jane et al., 2020), along with Gamma, Weibull, and Inverse Gaussian (Jane et al., 2020). Observed water levels did not exhibit site specific patterns and were described by a range of distributions (Table 4).

## 4.2 Copulas

250 The quality of a copula's fit is determined by its maximum likelihood (ML) values. Figure ?? presents ML values for the 17 tested copulas at each site (columns) per sampling method (rows). Fits produced similar ML statistics for a given site and sampling method, however, a few copulas produced particularly poor or unusual results. For example, the Independent and Cubic were consistently ranked as the lowest fit. Additionally, a number of copulas generally perform well (Fig. ??). For example, the Fischer-Koek (Fisc.), Tawn, and Roch-Alegre (Roch.) were amongst the better fitting copulas, ranking within the top five ML for nearly all datasets. The Tawn has previously been used in a OWL and precipitation analysis (Jane et al., 2020).

255

**Table 4.** Best fitting univariate distributions for each location and sampling method (annual maximum (AM), annual coinciding (AC), wet season monthly maximum (WMM), wet season monthly coinciding (WMC)).

Dataset	Variable	Santa Monica	Sunset	San Diego
AM	OWL	L	BS	GP
	Precip	BS	R	G
AC	OWL	N	GP	NA
	Precip	BS	R	G
WMM	OWL	N	W	GEV
	Precip	E	E	GP
WMC	OWL	G	IG	IG
	Precip	E	E	GP

BS - Birnbaum-Saunders; GP - Generalized Pareto; E - Exponential  
R - Rayleigh; N - Normal; L - Log logistic; G - Gamma  
W - Weibull; IG - Inverse Gaussian; NA - Nakagami


and in this case, is frequently the highest ranked. All other copula results were within the envelopes described by the five copulas. Unless otherwise noted, these focused copulas will be used to consider the impacts of copula choice and sampling.

ML Values per fitted copula for SM (left column), S (middle column), and SD (right column) (a), (b), (c) AM, (d), (e), (f) AC, (g), (h), (i) WMM, and (j), (k), (l) WMC

260 San Diego ~~WMC~~ wet season monthly coinciding conditional CDFs display individual copulas effects (Fig. ~~??4~~). The ~~Independent and Cubic~~ Nelsen, Roch., and BB1 copulas consistently suggest ~~lower~~ similar OWL (Fig. ~~??4a~~, e) and precipitation values (Fig. ~~??4b~~, f) while, in this example, the ~~Tawn suggests substantially~~ BB5 suggests higher OWL and precipitation values (~~dashed~~ solid black line, Fig. ~~??4a~~, b, e, f). C1's 100-year pair in Table 6 displays an example of the ~~Tawn~~ BB5's conservative nature. ~~The Roch-Alegre and Fischer-Koek provide very similar results for both precipitation and water level (black and~~  
265 ~~green lines, Fig. ??a, b, e, f).~~ Copula choice has nearly no effect on the Conditional 2 scenario (Fig. ~~??4c~~, d). Most probable OWL and precipitation values in Tables 5 and 6 further display the aforementioned behaviors. These conditional patterns generally persist at all locations with an additional note that the ~~Tawn, Roch-Alegre, and Fischer-Koek copulas typically~~ Nelsen can suggest lower OWL values and the BB5 can provide similar results ~~to the Roch. and BB1 copulas.~~

Figures ~~??~~ and ~~??~~

270 Figures 5 and 6 show the 10- and 100-year return periods, respectively, for the ~~five~~ four focused copulas using ~~WMC~~ sampling at SD. In all the hazard scenarios, the Independent and Cubic copulas suggest most probable events dominated by either large precipitation or OWL (Fig. ~~??~~ and ~~??~~). For example, AND 100-year OWL is unrealistically small compared to the other OWL values provided by alternative copulas (Table 6). Clearly the ~~Tawn~~ wet season monthly coinciding sampling at San Diego. Clearly the BB5 presents conservative results suggesting higher OWL and precipitation pairs in the AND ~~scenario and~~  
275 ~~SK scenarios~~ (Fig. ~~??a~~ and ??a5a, c and 6a, c), while the ~~Roch-Alegre and Fischer-Koek copulas present intermediate~~ BB1,



**Figure 4.** SD-WMC-San Diego wet season monthly coinciding OWL (left column) and precipitation (right column) (a), (b) C1, (c), (d) C2, and (e), (f) C3 CDFs using the Independent-Nelsen (blue), Cubic-Roch-Alegre (red), Roch-Alegre-BB1 (green), Fischer-Koek (black), and Tawn-BB5 (blackdashed) copulas. OWL/ Precipitation conditionals are conditioned on the occurrence of a 25-year precipitation/ OWL event.

Nelsen, and Roch-Alegre copulas present similar OWL and precipitation values. Similarly for the Survival Kendall scenario, the Tawn displays more conservative OWL and precipitation pairs while the Roch-Alegre and Fischer-Koek suggest the moderate OWL and precipitation levels (Fig. ??c and ??e). The OR and Kendall scenarios suggest quite similar isolines with the most probable precipitation varying between copulas between copulas, suggesting similar values, but the BB5 suggests less severe OWL and larger precipitation values according to the densest location along its isoline (Fig. ??5b, d, and ??6b, d). Tables 5 and 6 further display the bivariate patterns. Again, these bivariate patterns generally persist at all locations with an additional note that the Tawn, Roch-Alegre, and Fischer-Koek, copulas often generate similar values. the additional notes: the Nelsen may suggest lower OWL values in the AND and SK scenarios, the BB5 typically has similar results to the Roch. and BB1 copulas in the AND and SK scenarios, and the BB5 typically agreed with the other copula results outside of San Diego for the OR and K scenarios.


**Table 5.** SD-San Diego 10-year marginal (M), conditional (C), and bivariate OWL (m) and precipitation (mmday<sup>-1</sup>) values using WMC wet season monthly coinciding sampling. Conditionals are conditioned on a 25-year event occurring.

	Nelsen		Roch.		BB1		BB5	
	OWL	Precip	OWL	Precip	OWL	Precip	OWL	Precip
M	2.11	33.64	2.11	33.64	2.11	33.64	2.11	33.64
C1	2.11	33.64	1.94	36.27	2.16	35.57	2.59	48.57
C2	2.11	33.64	2.11	33.62	2.11	33.62	2.09	33.07
C3	2.11	33.64	1.94	36.24	2.16	35.56	2.40	44.73
AND	0.74	33.64	1.51	22.15	1.84	21.80	1.89	26.09
OR	2.11	55.32	2.11	38.29	2.20	37.06	2.14	39.10
K	1.97	48.34	1.97	55.32	2.06	31.79	2.06	31.80
SK	1.23	48.72	1.54	35.11	2.15	35.06	2.21	38.87


### 4.3 Sampling

San Diego conditional CDFs using the Fischer-Koek-BB1 copula clearly present sampling effects (i.e. maximum versus coinciding and annual versus wet season months). It should be noted, that each sampling method represents a unique probability space and accordingly results in alternative realizations of a given return period. Coinciding samplings exhibit similar OWL CDFs (green and red lines, Fig. ??7a, c, e), whereas wet season samplings exhibit similar precipitation CDFs (red and black lines Fig. ??7b, d, f). OWL values are generally can be larger for maximum sampling samplings at lower non-exceedance probabilities (i.e lower return periods) (Fig. ??7a, c, e blue and black lines, Table 7). However, the extended tail from WMC sampling produces larger OWL at higher return periods (red line Fig. ??7a, c, e; Table 8). Annual coinciding sampling displays significantly lower OWL values at low (Table 7) and high (Table 8) return periods. Only minimal differences between annual and wet season precipitation exist at the 10- and 100-year return periods (maximum difference of 1.79 and 3.08 1.06 and 4.56





**Figure 5.** ~~SD-WMC-San Diego wet season monthly coinciding~~ (a) AND, (b) OR, (c) SK, and (d) K hazard scenarios with the ~~Independent~~Nelsen, ~~Cubic~~, ~~Roch-Alegre (Roch)~~, ~~Fischer-Kock~~BB1, and ~~Tawn~~BB5 10-year isolines. Copula labels point to the mostly likely value on their respective isolines. ~~Most likely values outside the axis have a \* with their respective copula label.~~



**Figure 6.** ~~SD-WMC-San Diego wet season monthly coinciding~~ (a) AND, (b) OR, (c) SK, and (d) K hazard scenarios with the ~~Independent~~Nelsen, Cubic, Roch-Alegre (Roch), Fischer-Kock~~BB1~~, and Tawn~~BB5~~ 100-year isolines. Copula labels point to the mostly likely value on their respective isolines. ~~Most likely values outside the axis have a \* with their respective copula label.~~

**Table 6.** ~~SD-San Diego~~ 100-year marginal (M), conditional (C), and bivariate OWL (m) and precipitation (mmday<sup>-1</sup>) values using ~~WMC~~ wet season monthly coinciding sampling. Conditionals are conditioned on a 25-year event occurring.


	Nelsen		Roch.		BB1		BB5	
	OWL	Precip	OWL	Precip	OWL	Precip	OWL	Precip <del>OWL-Precip</del>
M	2.40	43.34	2.40	43.34	2.40	43.34	2.40	43.34 <del>2.40 43.34</del>
C1	<del>2.40</del> <u>2.47</u>	<del>43.34</del> <u>45.26</u>	<del>2.18 36.43</del> 2.46	44.95	2.44	<del>44.53</del> <u>44.48</u>	<del>2.85</del> <u>2.89</u>	52.72
C2	2.40	<del>43.34</del> <u>43.32</u>	2.40	<del>43.37 2.40</del> 43.32	2.40	43.34	<del>2.36</del> <u>2.35</u>	<del>42.07</del> <u>42.11</u>
C3	<del>2.40</del> <u>2.47</u>	<del>43.34</del> <u>45.26</u>	<del>2.20 36.88</del> 2.46	44.92	2.44	<del>44.51</del> <u>44.42</u>	<del>2.62</del> <u>2.68</u>	<del>49.38</del> <u>49.72</u>
AND	<del>0.74</del> <u>2.04</u>	<del>43.38</del> <u>30.77</u>	<del>1.53 41.02</del> 2.03	30.45	2.02	<del>30.06</del> <u>29.93</u>	<del>2.18</del> <u>2.20</u>	<del>38.04</del> <u>37.42</u>
OR	<del>2.39</del> <u>2.47</u>	<del>55.32</del> <u>45.53</u>	<del>2.39 55.32</del> 2.48	45.39	2.47	45.53	<del>2.43</del> <u>2.40</u>	<del>46.54</del> <u>49.17</u>
K	<del>2.13</del> <u>2.22</u>	<del>55.32</del> <u>31.79</u>	<del>2.15 42.63</del> 2.22	<del>37.67</del> <u>31.79</u>	2.17	<del>40.29</del> <u>40.13</u>	<del>2.17</del> <u>2.14</u>	<del>39.81</del> <u>38.59</u>
SK	<del>1.73</del> <u>2.31</u>	<del>51.49</del> <u>40.97</u>	<del>2.79 13.98</del> 2.33	41.53	2.32	41.26	<del>2.47</del> <u>2.46</u>	<del>46.35</del> <u>45.25</u>

mmday<sup>-1</sup> respectively; Tables 7 and 8). Annual (wet season) precipitation CDFs appear similar as OWL measurements are chosen subsequent to precipitation observations (Fig. ~~??7~~b, d, f).


~~Figures ?? and ??~~

~~Figures 8 and 9~~ present the 10- and 100-year return periods using the ~~Fischer-Koek-BB1~~ copula for all samplings (~~AC, AM, WMC, WMM~~AM, AC, WMM, and WMC). For the AND and K scenarios, AM sampling results in the largest OWL compared to the other sampling methods (Fig. ~~??8~~a, d and ~~??9~~a, d, Tables 7 and 8). Additionally for the AND and K scenarios (Fig. ~~??8~~a, d and ~~??9~~a, d), maximum samplings (AM and WMM) provide more conservative OWLs compared to WMC OWL values (Tables 7 and 8). When comparing maximum samplings (AM and WMM) to WMC sampling in the OR and SK scenarios, maximum samplings generally provide larger OWL values at lower return periods (Fig. ~~??8~~b, c; Table 7), but smaller or similar OWL at larger return periods (Fig. ~~??9~~b, c; Table 8). AC sampling generally results in the smallest OWL levels at all hazard scenarios. These behaviors persist across all locations,~~suggesting the~~ Given wet season monthly coinciding sampling results in larger OWL values for the marginal, conditional, OR, and Kendall scenarios, this suggests maximum type sampling may not accurately reflect OWL at extreme return periods~~for univariate and conditional situations~~.

Table 7 and 8 shows the most probable 10- and 100-year marginal and ~~compound-multivariate~~ event values. AM OWL exceed AC OWL across all probability types and sites, which is expected given the (nonphysical) paring of the two largest individual OWL and precipitation events without regard to co-occurrence. For example, in the 10-year return period ~~AM~~ annual maximum OWLs are at least 30 cm higher than AC (Table 7). In the 100-year return period ~~AM exceeds AC~~ annual maximum OWLs exceed annual coinciding OWLs by at least 17 cm (Table 8). Precipitation is generally consistent across ~~all conditionals, scenarios and~~ sampling types with only minor variations observed. ~~The AND scenario is the exception to this where substantially lower precipitation values are seen across all sampling and and scenarios.~~ AM and WMM sampling generally produced similar OWL results at both the 10- and 100-year return periods with maximum difference of 6 cm across all conditionals and copulas.




**Figure 7.** SD San Diego OWL (left column) and precipitation (right column) (a), (b) C1, (c), (d) C2, and (e), (f) C3 CDFs for AM-annual maximum (AM, blue), AC-annual coinciding (AC, green), WMM-wet season monthly maximum (WMM, black), and WMC-wet season monthly coinciding (WMC, red) samplings using the Fischer-Koek-BB1 copula. OWL/ Precipitation conditionals are conditioned on the occurrence of a 25-year precipitation/ OWL event.



**Figure 8.** SD-San Diego (a) AND, (b) OR, (c) SK, and (d) K hazard scenarios for AM-annual maximum (AM, cross), AC-annual coinciding (AC, plus), WMM-wet season monthly maximum (WMM, dot), WMC-wet season monthly coinciding (WMC, triangle) data and 10-year isolines using the Fischer-Koerk-BB1 copula. Sampling labels point to the mostly likely value on their respective isolines.

#### 4.4 Structural failure

A structural scenario is presented to consider flood severity along the Pacific Coast Highway (PCH) in Sunset Beach. PCH road elevation ranges from 1.7-2.4 m NAVD88 (Fig. 10), below typical spring tide ( $\sim 2.13$  m) and more extreme ( $\sim 2.3$  m)



**Figure 9.** SD-San Diego (a) AND, (b) OR, (c) SK, and (d) K hazard scenarios for AM-annual maximum (AM, cross), AC-annual coinciding (AC, plus), WMM-wet season monthly maximum (WMM, dot), WMC-wet season monthly coinciding (WMC, triangle) data and 100-year isolines using the Fischer-Kock-BB1 copula. Sampling labels point to the mostly likely value on their respective isolines.

water levels (NOAA, Accessed 2021b), requiring tide valves along PCH for flood prevention. Tide valve closures prevent back-flooding from high bay water levels coming up through subsurface storm drains that (normally) discharge to the bay.

**Table 7.** SD-San Diego 10-year marginal (M), conditional (C), and bivariate OWL (m) and precipitation (mmday<sup>-1</sup>) values using the Fischer-KoekBB1 with annual maximum (AM), annual coinciding (AC), wet season monthly maximum (WMM), and wet season monthly coinciding (WMC) samplings. Conditionals are conditioned on a 25-year event occurring.

	AM		AC		WMM		WMC	
	OWL	Precip	OWL	Precip	OWL	Precip	OWL	Precip
M	2.22	33.38	1.83	33.76	2.20	33.61	2.11	33.64
C1	<del>2.23</del> <u>2.24</u>	<del>33.78</del> <u>35.20</u>	1.86	<del>35.10</del> <u>35.03</u>	<del>2.20</del> <u>2.21</u>	<del>34.23</del> <u>34.39</u>	2.16	<del>35.57</del> <u>35.41</u>
C2	2.22	<del>33.37</del> <u>33.31</u>	1.83	33.70	2.20	33.60	2.11	33.62
C3	2.23	<del>33.77</del> <u>34.50</u>	1.86	<del>35.07</del> <u>34.99</u>	<del>2.20</del> <u>2.21</u>	<del>34.23</del> <u>34.31</u>	<del>2.16</del> <u>2.15</u>	<del>35.56</del> <u>35.37</u>
AND	2.14	<del>26.07</del> <u>26.13</u>	1.66	<del>26.77</del> <u>26.79</u>	2.09	20.89	1.84	<del>21.80</del> <u>21.65</u>
OR	2.25	<del>37.33</del> <u>37.03</u>	1.91	<del>37.49</del> <u>37.71</u>	2.22	<del>38.03</del> <u>38.48</u>	2.20	37.06
K	2.24	<del>36.03</del> <u>36.02</u>	1.89	36.44	2.18	<del>32.14</del> <u>32.10</u>	<del>2.06</del> <u>2.05</u>	<del>31.80</del> <u>32.52</u>
SK	2.25	<del>38.22</del> <u>38.45</u>	1.94	<del>38.78</del> <u>38.70</u>	2.21	<del>35.01</del> <u>35.06</u>	2.15	<del>35.06</del> <u>35.04</u>


**Table 8.** SD-San Diego 100-year marginal (M), conditional (C), and bivariate OWL (m) and precipitation (mmday<sup>-1</sup>) values using the Fischer-KoekBB1 with annual maximum (AM), annual coinciding (AC), wet season monthly maximum (WMM), and wet season monthly coinciding (WMC) samplings. Conditionals are conditioned on a 25-year event occurring.

	AM		AC		WMM		WMC	
	OWL	Precip	OWL	Precip	OWL	Precip	OWL	Precip
M	2.27	45.07	2.08	45.65	2.26	43.32	2.40	43.34
C1	2.27	<del>45.40</del> <u>48.54</u>	2.10	<del>46.78</del> <u>46.76</u>	<del>2.26</del> <u>2.27</u>	<del>43.70</del> <u>43.98</u>	2.44	<del>44.53</del> <u>44.48</u>
C2	2.27	<del>45.06</del> <u>44.90</u>	2.08	<del>45.59</del> <u>45.60</u>	2.26	<del>43.32</del> <u>43.30</u>	2.40	43.34
C3	2.27	<del>45.40</del> <u>46.35</u>	2.10	<del>46.75</del> <u>46.69</u>	<del>2.26</del> <u>2.27</u>	<del>43.70</del> <u>43.79</u>	2.44	<del>44.51</del> <u>44.42</u>
AND	2.23	<del>33.59</del> <u>34.01</u>	1.85	<del>34.43</del> <u>34.39</u>	<del>2.16</del> <u>2.17</u>	<del>29.27</del> <u>29.29</u>	2.02	<del>30.06</del> <u>29.93</u>
OR	2.27	<del>58.78</del> <u>48.15</u>	<del>2.09</del> <u>2.14</u>	<del>60.14</del> <u>48.83</u>	2.27	45.52	2.47	45.53
K	2.26	<del>43.02</del> <u>42.81</u>	2.02	42.45	2.22	<del>39.32</del> <u>39.68</u>	2.17	<del>40.29</del> <u>40.13</u>
SK	2.27	<del>45.05</del> <u>46.20</u>	2.08	<del>45.63</del> <u>45.65</u>	2.25	<del>40.76</del> <u>40.90</u>	2.32	41.26

Additionally, closed tide valves enable precipitation pooling since water cannot be drained to the bay. Severe pooling may result in a critical highway closure, which can further damage property and inhibit emergency service operations.

325     Areal precipitation flooding extent and depth can be estimated for water levels exceeding tide valve closure elevation. A water level equal to or greater than 1.68 m NAVD88 forces valve closures and frames the structural failure as a Conditional 1 type event. The local watershed is convex and drains an area of 94,897 m<sup>2</sup>. Water pools in the low elevation areas along PCH (Fig. 10). When pluvial water levels exceed the sea wall elevation, water overflows the sea wall and exits to the harbor. The maximum pool storage is 11,342 m<sup>3</sup>. The percent of flooding is then calculated with Eq. (20) as the structural function.






**Figure 10.** Elevations within the PCH Pacific Coast Highway boundary ranging from low (purple) to high (blue). Background imagery from NOAA (Accessed 2021a).

330 Structural scenario precipitation and percent flooding ( $\Psi$ ) values utilizing the Nelsen, Roch-Alegre, ~~Fischer-Koek, and Tawn~~  
and BB1 copulas are shown in Table 9. Rows and columns separate the utilized sampling methods and return periods of interest,  
 respectively. Fig. 11 shows  $\Psi$  as a function of precipitation for the Nelsen, Roch-Alegre, ~~Fischer-Koek, and Tawn~~ and BB1  
 copulas along with the 5- (square), 10- (circle), and 100-year (diamond) return periods. All copulas display similar values  
 across sampling methods ~~with the exception of the Tawn, which presents more conservative values using WMC sampling~~ (Fig.  
 335 11, Table 9). For example, the 10-year ~~WMC for the Tawn (Fig. 11e) is over 80 percent flooding compared to approximately~~  
~~70 percent with the other copulas~~ precipitation with wet season monthly coinciding sampling is 82.49, 81.95, and 81.09 for the  
Nelsen, Roch-Alegre, and BB1 copulas, respectively. Again, AC sampling severely underestimates precipitation and flooding.

$$\% \text{ flooding} = \frac{\text{precipitation} \times \text{area}}{\text{volume}} \times 100 \quad (20)$$

## 5 Discussion

340 Previous multivariate studies typically use a small, popular group of copulas (e.g., Clayton, Frank, Gumbel, Student t, and  
 Gaussian). Gaussian and Student t copulas were excluded from this study due to their lack of a computationally simple  
 derivative or integral while the Clayton, Frank, and Gumbel ~~generally provided average fits (Fig. ??). Tawn, Fischer-Koek,~~  
~~failed to pass the Cramer-con Mises test.~~ Nelsen, BB1, BB5, and Roch-Alegre copulas generally present similar values  
 with the ~~Tawn BB5~~ occasionally presenting more conservative pairs (Fig. ??). ~~Cubie and Independent copulas consistently~~  
 345 ~~present event pairs predominantly over- or underestimating OWL and precipitation (Fig. ??). The poor performance of the~~  
~~Independent copula further emphasizes the importance in accounting for dependencies between variables in multivariate~~



**Figure 11.** Structural scenario 5- (square), 10- (circle), and 100-year (diamond) return periods for AM-annual maximum (AM, blue), AC-annual coinciding (AC, green), WMM-wet season monthly maximum (WMM, black), and WMC-wet season monthly (WMC, red) data using the (a) Roch-AlegreNelsen, (b) Fischer-KoekRoch-Alegre, and (c) Tawn-BB1 copulas.

studies (Raynal-Villasenor and Salas, 1987; Yue and Rasmussen, 2002; De Michele et al., 2005). 6). Well fit copulas concentrate probabilities around more centralized intermediate OWL and precipitation values for compound multivariate events. This is most pronounced at higher (i.e., 100-year) return periods (Fig. ??6). Given that nearly all copulas, save the Cubic, Independent, and Tawn, exhibit similar ML the resulting copulas exhibit agreement between values (for a given sampling) suggests that choosing a reasonable copula may be sufficient to provide a robust characterization of considered compound multivariate flooding events.

The choice in sampling imparts a significant influence on flood event risk interpretation. Substantially lower ML values are observed in annual sampling (Fig. ??a-f) compared to wet season sampling (Fig. ?? g-i). This results from the the increased data availability when the entire wet season is used compared to a single annual observation. When maximum versus coinciding sampling is considered, maximum samplings (AM and WMM) tend to provide the largest OWL at low return periods (Fig. ??7a, c, e, Fig. ??8, and Table 7). At larger return periods, WMC wet season monthly coinciding then provides significantly larger OWLs (Fig. ??7a, c, e, Fig. ??9, and Table 8). This is observed in the conditionals and bivariate (minus the AND and K hazard scenarios which maximum samplings display the largest OWLs) at all sites. From a logical perspective, coinciding sampling provides a more realistic view of compound multivariate events (by definition these are pairs that have co-occurred

**Table 9.** Precipitation and percent flooding ( $\Psi$ ) associated to the 5-, 10-, and 100-year return periods ( $T$ ) using the [Nelsen](#), Roch-Alegre, [Fischer-Koek](#), and [Tawn-BB1](#) copulas to determine C1 values [with annual maximum \(AM\), annual coinciding \(AC\), wet season monthly maximum \(WMM\), and wet season monthly coinciding \(WMC\) samplings](#). Precipitation values are in  $\text{mmday}^{-1}$  and  $\Psi$  is a percentage. Values are based off a OWL of  $\geq 1.68$  m which forces tide valve closure.

$T$	5-yr.		10-yr.		100-yr.	
	Precip	$\Psi$	Precip	$\Psi$	Precip	$\Psi$
Nelsen						
AM	63.51	53.14	75.97	63.56	107.44	89.89
AC	<del>33.70</del> <a href="#">34.45</a>	<del>28.19</del> <a href="#">28.82</a>	<del>46.57</del> <a href="#">47.41</a>	<del>38.97</del> <a href="#">39.66</a>	<del>88.50</del> <a href="#">89.39</a>	<del>74.05</del> <a href="#">74.79</a>
WMM	<del>64.57</del> <a href="#">64.51</a>	<del>54.02</del> <a href="#">53.97</a>	<del>79.34</del> <a href="#">79.27</a>	<del>66.38</del> <a href="#">66.32</a>	<del>128.44</del> <a href="#">128.26</a>	100.00
WMC	<del>67.17</del> <a href="#">67.71</a>	<del>56.20</del> <a href="#">56.65</a>	<del>81.95</del> <a href="#">82.49</a>	<del>68.56</del> <a href="#">69.02</a>	<del>130.92</del> <a href="#">131.44</a>	100.00
Roch-Alegre						
AM	63.51	53.14	75.97	63.56	107.44	89.89
AC	<del>33.68</del> <a href="#">33.70</a>	<del>28.18</del> <a href="#">28.19</a>	<del>46.50</del> <a href="#">46.57</a>	<del>38.91</del> <a href="#">38.97</a>	<del>88.35</del> <a href="#">88.50</a>	<del>73.92</del> <a href="#">74.05</a>
WMM	<del>64.56</del> <a href="#">64.57</a>	54.02	<del>79.33</del> <a href="#">79.34</a>	66.38	<del>128.35</del> <a href="#">128.44</a>	100.00
WMC	<del>66.99</del> <a href="#">67.17</a>	<del>56.05</del> <a href="#">56.20</a>	<del>81.74</del> <a href="#">81.95</a>	<del>68.39</del> <a href="#">68.56</a>	<del>130.62</del> <a href="#">130.92</a>	100.00
BB1						
AM	63.51	53.14	75.97	63.56	107.44	89.89
AC	<del>33.42</del> <a href="#">33.30</a>	<del>29.64</del> <a href="#">27.86</a>	<del>49.15</del> <a href="#">46.41</a>	<del>41.12</del> <a href="#">38.83</a>	<del>92.98</del> <a href="#">89.44</a>	<del>77.79</del> <a href="#">74.83</a>
WMM	<del>65.12</del> <a href="#">64.60</a>	<del>54.48</del> <a href="#">54.05</a>	<del>79.88</del> <a href="#">79.38</a>	<del>66.83</del> <a href="#">66.42</a>	<del>128.88</del> <a href="#">128.44</a>	100.00
WMC	<del>83.39</del> <a href="#">66.34</a>	<del>69.77</del> <a href="#">55.50</a>	<del>98.10</del> <a href="#">81.09</a>	<del>82.08</del> <a href="#">67.84</a>	<del>146.99</del> <a href="#">130.03</a>	100.00

to produce a ~~compound-multivariate~~ flooding event). At long return periods ~~AC~~-annual coinciding sampling may require a long data record, which is often unavailable. Notably, in this study ~~AC~~-annual coinciding produced OWL samplings that were substantially lower than any of the other samplings. For example when comparing 100-year OWLs with ~~AC and WMC~~-annual and wet season monthly coinciding samplings, the marginal was 32 cm lower and the AND scenario was 17 cm lower (Table 8). Given that sea wall protected urban coastal areas are highly sensitive to even minor elevations differences (e.g., Gallien et al., 2011), this suggests with limited data records ~~AC~~-annual coinciding sampling should be avoided.

An important note is each probability type appropriately describes a unique event, characterized by OWL and precipitation. Serinaldi (2015) suggests inter-comparing univariate, multivariate, and conditional probabilities and return periods is misleading as each probability type describes its associated event. Events where only extreme OWL or precipitation is of concern, should simply utilize marginal statistics and follow current FEMA guidelines. ~~Compound-Multivariate~~ event analysis may utilize a variety of scenarios. Conditional type distributions become useful when future information on one variable is known (ex., predicted OWL levels). AND scenarios may be applied when both variables exceeding given limits is of concern. The Survival Kendall scenario is an alternative to the AND scenario using a more conservative approach to develop events of concern (Salvadori et al., 2013). An OR scenario should be applied when either ~~compounding-multivariate~~ variable exceeding a limit

375 is of concern, whereas the Kendall scenario provides minimum OR events of concern (Salvadori et al., 2011). The benefit of  
the Kendall and Survival Kendall is all the events along their isolines describe a similar probability space versus the AND and  
OR isolines describe events with similar probabilities of non-exceedance. ~~The~~ It is critical for practitioners and future studies  
to define concerning flood events within a region since the associated probability will result in varying event risk estimates as  
seen within this work. The selected probability type will have significant influence on flood risk studies and modeling efforts. A  
380 majority of previous studies focus on specific probability types and do not consider multiple flooding pathways. Only a single  
study explores all the probabilities associated to different extreme events (Serinaldi, 2016).

From a regulatory perspective, FEMA recommends individual (univariate) analysis to develop return periods for ~~compound~~  
multivariate coastal flooding applications (FEMA, 2011, 2016c, 2020), and blending the two hazard mapping results. Funda-  
mentally, this type of approach assumes (event) independence and may underestimate ~~compound-multivariate~~ flood hazards  
385 (e.g., Moftakhari et al., 2019; Muñoz et al., 2020). FEMA provides guidelines for coastal-riverine (FEMA, 2020), tide, surge,  
tide-surge (FEMA, 2016a), surge-riverine (FEMA, 2016c), and tropical storm (or hurricane) type flooding events (FEMA,  
2016b). Currently, FEMA does not ~~cover compound coastal flooding from~~ provide specific guidance for considering high  
marine water levels and precipitation. However, this work suggests, at high return periods the sampling method is critical to  
characterizing both univariate and joint probabilities.

390 Structural scenarios provide a quantitative context to frame flood vulnerability. In the structural failure context, ~~AC-annual~~  
coinciding sampling significantly underestimates flooding at all return periods, and ~~AM-annual maximum~~ sampling underesti-  
mates severe (i.e., 100-year) events, echoing previous ~~AC and AM samplings annual maximum and coinciding sampling~~ issues.  
Similar values between most copulas support the suggestion that choosing a reasonable copula will provide robust results in  
these types of applications. Precipitation events in the Structural scenario (Table 9) range between ~~33.70~~ 33.30  $\text{mmday}^{-1}$   
395 and ~~128.44~~ 131.44  $\text{mmday}^{-1}$ , resulting between ~~28.18~~ 27.86 % and complete (100 %) backshore flooding. This significant  
flooding at all return periods suggests severe flood vulnerability, which is validated by frequent closures of PCH. This struc-  
tural function provides a quick and simple alternative to poorly performing bathtub flood models (e.g., Bernatchez et al., 2011;  
Gallien et al., 2011, 2014; Gallien, 2016) to quantitatively explore flood severity while accounting for infrastructure and joint  
probabilities.

400 The maximum OWL and precipitation observations within the record are 2.33 m and  $118.93 \text{ mmday}^{-1}$  for Sunset, 2.27  
m and  $42.11 \text{ mmday}^{-1}$  for San Diego, and 2.45 m and  $76.83 \text{ mmday}^{-1}$  for Santa Monica. Most likely precipitation and  
OWL pairs in high return periods often exceed the current data record's maximums (e.g. Table 8). This study is limited to the  
available data records and sea level rise clearly imparts a non-stationary trend. Current water level values restricted to today's  
distribution tails, will become more frequent in the next century (Taherkhani et al., 2020). For example, Wahl et al. (2015)  
405 suggests a previously 100-year event in New York is now a 42-year event based on the increasing correlation between extreme  
precipitation and storm surge events. Similarly, our results suggest increasing precipitation and, particularly, OWL levels.


## 6 Conclusions

~~Univariate, conditional, and bivariate compound flood risk~~ Univariate and multivariate event risks from OWL and/or precipitation were explored at three sites in a tidally dominated, semi-arid region. Seventeen copulas were considered. Previous studies typically relied upon a small number of copulas (e.g. Clayton, Frank, Gumbel, Student t, and Gaussian) for ~~compound flooding multivariate flood~~ assessments. In this case, the ~~Fischer-Koek, Roch-Alegre, and Tawn copulas~~ Nelsen, BB1, BB5, and Roch-Alegre copulas passed the Cramer-von Misses test and produced similar, quality fits across all sampling methods. Although, in some cases, the ~~Tawn-BB5~~ produced conservative results. ~~Independent and Cubic copulas consistently produced poor fits which under-/overestimated values and favored either precipitation or water level dominated events. Multiple copulas exhibit similar goodness of fit values (Fig. ??) and~~ Copulas exhibit similar most probable pairs (e.g., Fig. ~~??, ??5~~, 6) suggesting a number of potential copulas may provide a robust multivariate analysis. The impact of sampling and distribution choice on uni- and multivariate return period values are presented, however uncertainties deserve further characterization. This work focused solely on exploring conditional and joint probabilities of OWL and precipitation in a ~~tidal~~ tidally and wave dominated semi-arid region and would not be applicable to regions experiencing multiple flooding seasons (e.g., Couasnon et al., 2022). Although wave impacts were not included in this assessment, they are fundamental to coastal flooding, particularly in regions subjected to long period swell. Joint probability methods explicitly including wave contributions to ~~compound flood risk characterization are needed~~, multivariate event risk characterizations are needed for future work.


The annual maximum method is widely recognized for hazard assessments (FEMA, 2011, 2016c), and is common practice in flood risk analysis (e.g., Baratti et al., 2012; Bezak et al., 2014; Wahl et al., 2015). Concerningly, this work suggests that ~~AM annual maximum sampling~~ does not characterize ~~the “worse case” scenarios~~ severe flooding potential for extreme events. Water levels are substantially underestimated as annual sampling neglects a large portion of observations (Table 8). Generally, maximum samplings produced larger values at minor return periods but significantly underestimated water levels at longer return periods than ~~WMC wet season monthly coinciding~~ sampling. Similarly, ~~AC annual coinciding~~ type sampling (Tables 7, 8) grossly underestimate OWLs. Wet season sampling quadruples data pairs (Table 2), providing additional historical joint event information. Further investigation into monthly coinciding and, where appropriate, water year coinciding are needed to develop optimal sampling strategies for given regional conditions.

*Data availability.* NOAA precipitation data is available for download at <https://www.ncei.noaa.gov/metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C00313/html#>. Tidal data is available for download on NOAA’s Tides & Currents website (<https://tidesandcurrents.noaa.gov>)

## 435 **Appendix A: Additional AICs and BICs**



**Figure A1.** Marginal OWL BIC values per fitted copula for **SM-Santa Monica** (left column), **S-Sunset** (middle column), and **SD-San Diego** (right column) **-Annual-Maximum-using annual maximum** ((a), (b), (c); **-Annual-Coinciding-**, **annual coinciding** ((d), (e), (f); **-Wet-Season Monthly-Maximum-**, **wet season monthly maximum** ((g), (h), (i); **-**, and **-Wet-Season Monthly-Coinciding-**, **wet season monthly coinciding** ((j), (k), (l)). The Y-axis is orientated to display best BIC (top) to **worst-worse** BIC (bottom).



**Figure A2.** Marginal precipitation BIC values per fitted copula for **SM** Santa Monica (left column), **S** Sunset (middle column), and **SD** San Diego (right column). ~~Annual-Maximum-using annual maximum~~ ((a), (b), (c)), ~~Annual-Coinciding~~, annual coinciding ((d), (e), (f)), ~~Wet-Season-Monthly-Maximum~~, wet season monthly maximum ((g), (h), (i)), and ~~Wet-Season-Monthly-Coinciding~~ wet season monthly coinciding ((j), (k), (l)). The Y-axis is orientated to display best BIC (top) to worst BIC (bottom).



Copula AIC values per fitted copula for SM (left column), S (middle column), and SD (right column). Annual Maximum (a), (b), (c); Annual Coinciding (d), (e), (f); Wet Season Monthly Maximum (g), (h), (i); and Wet Season Monthly Coinciding (j), (k), (l). The Y-axis is orientated to display best AIC (top) to worst AIC (bottom).

440 Copula BIC values per fitted copula for SM (left column), S (middle column), and SD (right column). Annual Maximum (a), (b), (c); Annual Coinciding (d), (e), (f); Wet Season Monthly Maximum (g), (h), (i); and Wet Season Monthly Coinciding (j), (k), (l). The Y-axis is orientated to display best BIC (top) to worst BIC (bottom).

*Author contributions.* JL conducted the primary analysis under the guidance and assistance of TG. Both authors wrote and edited the manuscript. TG conceived of and funded the work.

*Competing interests.* The authors declare no competing interest.

445 *Acknowledgements.* This work has been supported by the US Coastal Research Program under contract W912HZ-20-200-004, California Department of Parks and Recreation contact number C1670006, the National Science Foundation Graduate Research Fellowship Program grant number DGE-1650604, The National GEM Consortium Fellowship, and the UCLA Cota-Robles Fellowship. The US Coastal Research Program (USCRP) is administered by the US Army Corps of Engineers® (USACE), Department of Defense. The content of the information provided in this publication does not necessarily reflect the position or the policy of the government, and no official endorsement should be  
450 inferred. The authors' acknowledge the USACE and USCRP's support of their effort to strengthen coastal academic programs and address coastal community needs in the United States. Any opinions, findings, conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the agencies supporting the work. We would like to acknowledge Drs. Yeulwoo Kim and Nikos Kalligeris for their constructive feedback which strengthened this manuscript.

## References

- 455 Aghakouchak, A.: Entropy copula in hydrology and climatology, *Journal of Hydrometeorology*, 15, 2176–2189, 2014.
- Anandalekshmi, A., Panicker, S. T., Adarsh, S., Siddik, A. M., Aloysius, S., and Mehjabin, M.: Modeling the concurrent impact of extreme rainfall and reservoir storage on Kerala floods 2018: a Copula approach, *Modeling Earth Systems and Environment*, 5, 1283–1296, 2019.
- Ayantobo, O. O., Wei, J., and Wang, G.: Modeling Joint Relationship and Design Scenarios Between Precipitation, Surface Temperature, and Atmospheric Precipitable Water Over Mainland China, *Earth and Space Science*, 8, e2020EA001 513, 2021.
- 460 Baratti, E., Montanari, A., Castellarin, A., Salinas, J., Viglione, A., and Bezzi, A.: Estimating the flood frequency distribution at seasonal and annual time scales., *Hydrology & Earth System Sciences*, 16, 2012.
- Beck, H. E., Zimmermann, N. E., McVicar, T. R., Vergopolan, N., Berg, A., and Wood, E. F.: Present and future Köppen-Geiger climate classification maps at 1-km resolution, *Scientific data*, 5, 1–12, 2018.
- Bell, J. E., Herring, S. C., Jantarasami, L., Adrianopoli, C., Benedict, K., Conlon, K., Escobar, V., Hess, J., Luvall, J., Garcia-Pando, C.,  
465 et al.: Ch. 4: Impacts of extreme events on human health, Tech. rep., US Global Change Research Program, Washington, DC, 2016.
- Bernatchez, P., Fraser, C., Lefaivre, D., and Dugas, S.: Integrating anthropogenic factors, geomorphological indicators and local knowledge in the analysis of coastal flooding and erosion hazards, *Ocean & Coastal Management*, 54, 621–632, 2011.
- Bevacqua, E., Maraun, D., Vousdoukas, M. I., Voukouvalas, E., Vrac, M., Mentaschi, L., and Widmann, M.: Higher probability of compound flooding from precipitation and storm surge in Europe under anthropogenic climate change, *Science advances*, 5, eaaw5531, 2019.
- 470 Bevacqua, E., Vousdoukas, M. I., Zappa, G., Hodges, K., Shepherd, T. G., Maraun, D., Mentaschi, L., and Feyen, L.: More meteorological events that drive compound coastal flooding are projected under climate change, *Communications earth & environment*, 1, 1–11, 2020.
- Bezak, N., Brilly, M., and Šraj, M.: Comparison between the peaks-over-threshold method and the annual maximum method for flood frequency analysis, *Hydrological Sciences Journal*, 59, 959–977, 2014.
- Bray, S. N. and McCuen, R. H.: Importance of the assumption of independence or dependence among multiple flood sources, *Journal of*  
475 *Hydrologic Engineering*, 19, 1194–1202, 2014.
- Cayan, D. R. and Roads, J. O.: Local relationships between United States West Coast precipitation and monthly mean circulation parameters, *Monthly Weather Review*, 112, 1276–1282, 1984.
- Church, J., Clark, P., Cazenave, A., Gregory, J., Jevrejeva, S., Levermann, A., Merrifield, M., Milne, G., Nerem, R., Nunn, P., Payne, A., Pfeffer, W., Stammer, D., and Unnikrishnan, A.: Sea Level Change, in: *Climate Change 2013: The Physical Science Basis. Contribution*  
480 *of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., pp. 1137–1216, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Conil, S. and Hall, A.: Local regimes of atmospheric variability: A case study of Southern California, *Journal of Climate*, 19, 4308–4325, 2006.
- 485 Couasnon, A., Sebastian, A., and Morales-Nápoles, O.: A copula-based Bayesian network for modeling compound flood hazard from riverine and coastal interactions at the catchment scale: An application to the Houston Ship Channel, Texas, *Water*, 10, 1190, 2018.
- Couasnon, A., Scussolini, P., Tran, T., Eilander, D., Muis, S., Wang, H., Keesom, J., Dullaart, J., Xuan, Y., Nguyen, H., et al.: A flood risk framework capturing the seasonality of and dependence between rainfall and sea levels—an application to Ho Chi Minh City, Vietnam, *Water Resources Research*, p. e2021WR030002, 2022.

- 490 De Michele, C. and Salvadori, G.: A generalized Pareto intensity-duration model of storm rainfall exploiting 2-copulas, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- De Michele, C., Salvadori, G., Canossi, M., Petaccia, A., and Rosso, R.: Bivariate statistical approach to check adequacy of dam spillway, *Journal of Hydrologic Engineering*, 10, 50–57, 2005.
- De Michele, C., Salvadori, G., Passoni, G., and Vezzoli, R.: A multivariate model of sea storms using copulas, *Coastal Engineering*, 54, 734–751, 2007.
- 495 DeGroot, M. H. and Schervish, M. J.: *Probability and Statistics*, Pearson Education, 4 edn., 2014.
- Didier, D., Baudry, J., Bernatchez, P., Dumont, D., Sadegh, M., Bismuth, E., Bandet, M., Dugas, S., and Sévigny, C.: Multihazard simulation for coastal flood mapping: Bathtub versus numerical modelling in an open estuary, Eastern Canada, *Journal of Flood Risk Management*, 12, e12 505, 2019.
- 500 Engeland, K., Hisdal, H., and Frigessi, A.: Practical extreme value modelling of hydrological floods and droughts: a case study, *Extremes*, 7, 5–30, 2004.
- Esberto, M. D. P.: Probability Distribution Fitting of Rainfall Patterns in Philippine Regions for Effective Risk Management, *Environment and Ecology Research*, 6, 178–86, 2018.
- Favre, A.-C., El Adlouni, S., Perreault, L., Thiémonge, N., and Bobée, B.: Multivariate hydrological frequency analysis using copulas, *Water resources research*, 40, 2004.
- 505 FEMA: Coastal construction manual: Principles and practices of planning, siting, designing, constructing, and maintaining residential buildings in coastal areas, 2011.
- FEMA: Guidance for Flood Risk Analysis and Mapping: Coastal Water Levels, [https://www.fema.gov/sites/default/files/2020-02/Coastal\\_Wate\\_Levels\\_Guidance\\_May\\_2016.pdf](https://www.fema.gov/sites/default/files/2020-02/Coastal_Wate_Levels_Guidance_May_2016.pdf), 2016a.
- 510 FEMA: Guidance for Flood Risk Analysis and Mapping: Statistical Simulation Methods, [https://www.fema.gov/sites/default/files/documents/fema\\_coastal-statistical-simulation-methods\\_nov-2016.pdf](https://www.fema.gov/sites/default/files/documents/fema_coastal-statistical-simulation-methods_nov-2016.pdf), 2016b.
- FEMA: Guidance for Flood Risk Analysis and Mapping: Coastal Flood Frequency and Extreme Value Analysis, [https://www.fema.gov/sites/default/files/2020-02/Coastal\\_Flood\\_Frequency\\_and\\_Extreme\\_Value\\_Analysis\\_Guidance\\_Nov\\_2016.pdf](https://www.fema.gov/sites/default/files/2020-02/Coastal_Flood_Frequency_and_Extreme_Value_Analysis_Guidance_Nov_2016.pdf), 2016c.
- FEMA: Guidance for Flood Risk Analysis and Mapping: Combined Coastal and Riverine Floodplain, [https://www.fema.gov/sites/default/files/documents/coastal\\_riverine\\_guidance\\_dec\\_2020.pdf](https://www.fema.gov/sites/default/files/documents/coastal_riverine_guidance_dec_2020.pdf), 2020.
- 515 Flick, R. E.: A comparison of California tides, storm surges, and mean sea level during the El Niño winters of 1982-83 and 1997-98, *Shore & Beach*, 66, 7–11, 1998.
- Flick, R. E.: California tides, sea level, and waves—Winter 2015-2016, *Shore & Beach*, 84, 25–30, 2016.
- Gallien, T.: Validated coastal flood modeling at Imperial Beach, California: Comparing total water level, empirical and numerical overtopping methodologies, *Coastal Engineering*, 111, 95–104, 2016.
- 520 Gallien, T., Schubert, J., and Sanders, B.: Predicting tidal flooding of urbanized embayments: A modeling framework and data requirements, *Coastal Engineering*, 58, 567–577, 2011.
- Gallien, T., Sanders, B., and Flick, R.: Urban coastal flood prediction: Integrating wave overtopping, flood defenses and drainage, *Coastal Engineering*, 91, 18–28, 2014.
- 525 Gallien, T., O'Reilly, W., Flick, R., and Guza, R.: Geometric properties of anthropogenic flood control berms on southern California beaches, *Ocean & Coastal Management*, 105, 35–47, 2015.

- Gallien, T., Kalligeris, N., Delisle, M., Tang, B., Lucey, J., and Winters, M.: Coastal flood modeling challenges in defended urban backshores, *Geosciences*, 8, 450, 2018.
- 530 Ganguli, P. and Merz, B.: Trends in compound flooding in northwestern Europe during 1901–2014, *Geophysical Research Letters*, 46, 10 810–10 820, 2019a.
- Ganguli, P. and Merz, B.: Extreme coastal water levels exacerbate fluvial flood hazards in Northwestern Europe, *Scientific reports*, 9, 1–14, 2019b.
- Ganguli, P. and Reddy, M. J.: Probabilistic assessment of flood risks using trivariate copulas, *Theoretical and applied climatology*, 111, 341–360, 2013.
- 535 Ganguli, P., Paprotny, D., Hasan, M., Güntner, A., and Merz, B.: Projected changes in compound flood hazard from riverine and coastal floods in northwestern Europe, *Earth's Future*, 8, e2020EF001 752, 2020.
- Genest, C., Rémillard, B., and Beaudoin, D.: Goodness-of-fit tests for copulas: A review and a power study, *Insurance: Mathematics and economics*, 44, 199–213, 2009.
- Gräler, B., van den Berg, M., Vandenbergh, S., Petroselli, A., Grimaldi, S., De Baets, B., and Verhoest, N.: Multivariate return periods in hydrology: a critical and practical review focusing on synthetic design hydrograph estimation, *Hydrology and Earth System Sciences*, 17, 1281–1296, 2013.
- 540 Gregory, J. M., Griffies, S. M., Hughes, C. W., Lowe, J. A., Church, J. A., Fukimori, I., Gomez, N., Kopp, R. E., Landerer, F., Le Cozannet, G., et al.: Concepts and terminology for sea level: Mean, variability and change, both local and global, *Surveys in Geophysics*, 40, 1251–1289, 2019.
- 545 Hanson, L. S. and Vogel, R.: The probability distribution of daily rainfall in the United States, in: *World Environmental and Water Resources Congress 2008: Ahupua'a*, pp. 1–10, 2008.
- Hanson, S., Nicholls, R., Ranger, N., Hallegatte, S., Corfee-Morlot, J., Herweijer, C., and Chateau, J.: A global ranking of port cities with high exposure to climate extremes, *Climatic change*, 104, 89–111, 2011.
- Hao, Z. and Singh, V. P.: Compound Events under Global Warming: A Dependence Perspective, *Journal of Hydrologic Engineering*, 25, 03120 001, 2020.
- 550 Hawkes, P. J., Gouldby, B. P., Tawn, J. A., and Owen, M. W.: The joint probability of waves and water levels in coastal engineering design, *Journal of hydraulic research*, 40, 241–251, 2002.
- Heberger, M., Cooley, H., Herrera, P., Gleick, P. H., and Moore, E.: Potential impacts of increased coastal flooding in California due to sea-level rise, *Climatic Change*, 109, 229–249, 2011.
- 555 Horton, B. P., Rahmstorf, S., Engelhart, S. E., and Kemp, A. C.: Expert assessment of sea-level rise by AD 2100 and AD 2300, *Quaternary Science Reviews*, 84, 1–6, 2014.
- Husak, G. J., Michaelsen, J., and Funk, C.: Use of the gamma distribution to represent monthly rainfall in Africa for drought monitoring applications, *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 27, 935–944, 2007.
- Jane, R., Cadavid, L., Obeysekera, J., and Wahl, T.: Multivariate statistical modelling of the drivers of compound flood events in South Florida, *Natural Hazards and Earth System Sciences Discussions*, pp. 1–30, 2020.
- 560 Jarušková, D. and Hanek, M.: Peaks over threshold method in comparison with block-maxima method for estimating high return levels of several Northern Moravia precipitation and discharges series, *Journal of Hydrology and Hydromechanics*, 54, 309–319, 2006.
- Jeong, D. I., Sushama, L., Khaliq, M. N., and Roy, R.: A copula-based multivariate analysis of Canadian RCM projected changes to flood characteristics for northeastern Canada, *Climate dynamics*, 42, 2045–2066, 2014.

- 565 Juma, B., Olang, L. O., Hassan, M., Chasia, S., Bukachi, V., Shiundu, P., and Mulligan, J.: Analysis of rainfall extremes in the Ngong River Basin of Kenya: Towards integrated urban flood risk management, *Physics and Chemistry of the Earth, Parts A/B/C*, p. 102929, 2020.
- Karmakar, S. and Simonovic, S.: Bivariate flood frequency analysis. Part 2: a copula-based approach with mixed marginal distributions, *Journal of Flood Risk Management*, 2, 32–44, 2009.
- Leonard, M., Westra, S., Phatak, A., Lambert, M., van den Hurk, B., McInnes, K., Risbey, J., Schuster, S., Jakob, D., and Stafford-Smith, M.:  
570 A compound event framework for understanding extreme impacts, *Wiley Interdisciplinary Reviews: Climate Change*, 5, 113–128, 2014.
- Li, T., Guo, S., Chen, L., and Guo, J.: Bivariate flood frequency analysis with historical information based on copula, *Journal of Hydrologic Engineering*, 18, 1018–1030, 2013.
- Lian, J., Xu, K., and Ma, C.: Joint impact of rainfall and tidal level on flood risk in a coastal city with a complex river network: a case study of Fuzhou City, China, *Hydrology and Earth System Sciences*, 17, 679, 2013.
- 575 Masina, M., Lamberti, A., and Archetti, R.: Coastal flooding: A copula based approach for estimating the joint probability of water levels and waves, *Coastal Engineering*, 97, 37–52, 2015.
- Mazas, F. and Hamm, L.: An event-based approach for extreme joint probabilities of waves and sea levels, *Coastal Engineering*, 122, 44–59, 2017.
- Merkens, J.-L., Reimann, L., Hinkel, J., and Vafeidis, A. T.: Gridded population projections for the coastal zone under the Shared Socioeconomic Pathways, *Global and Planetary Change*, 145, 57–66, 2016.
- 580 Mitková, V. B. and Halmová, D.: Joint modeling of flood peak discharges, volume and duration: a case study of the Danube River in Bratislava, *Journal of Hydrology and Hydromechanics*, 62, 186–196, 2014.
- Moftakhari, H., Schubert, J. E., AghaKouchak, A., Matthew, R. A., and Sanders, B. F.: Linking statistical and hydrodynamic modeling for compound flood hazard assessment in tidal channels and estuaries, *Advances in Water Resources*, 128, 28–38, 2019.
- 585 Muñoz, D., Moftakhari, H., and Moradkhani, H.: Compound Effects of Flood Drivers and Wetland Elevation Correction on Coastal Flood Hazard Assessment, *Water Resources Research*, 56, e2020WR027544, 2020.
- Nicholls, R. J.: Planning for the impacts of sea level rise, *Oceanography*, 24, 144–157, 2011.
- Nicholls, R. J., Wong, P. P., Burkett, V., Codignotto, J., Hay, J., McLean, R., Ragoonaden, S., Woodroffe, C. D., Abuodha, P., Arblaster, J., et al.: Coastal systems and low-lying areas, 2007.
- 590 NOAA: 2018 NAIP 4-Band 8 Bit Imagery : California, Online, <https://coast.noaa.gov/dataviewer/#/imagery/search/-13145277.048600255,3989719.0716587296,-13141378.760157712,3992489.913934067/details/9159>, Accessed 2021a.
- NOAA: Extreme Water Levels, Online, [https://tidesandcurrents.noaa.gov/est/est\\_station.shtml?stnid=9410660](https://tidesandcurrents.noaa.gov/est/est_station.shtml?stnid=9410660), Accessed 2021b.
- NOAA: US Hourly Precipitation Data, Online, <https://www.ncei.noaa.gov/metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C00313/html#>, Accessed 2021c.
- 595 NOAA: Tides & Currents, Online, <https://tidesandcurrents.noaa.gov>, Accessed 2021d.
- Pakoksung, K. and Takagi, M.: Mixed Zero-Inflation Method and Probability Distribution in Fitting Daily Rainfall Data, *Engineering Journal*, 21, 63–80, 2017.
- Paprotny, D., Vousdoukas, M. I., Morales-Nápoles, O., Jonkman, S. N., and Feyen, L.: Compound flood potential in Europe, *Hydrology and Earth System Sciences Discussions*, pp. 1–34, 2018.
- 600 Peng, Y., Shi, Y., Yan, H., Chen, K., and Zhang, J.: Coincidence Risk Analysis of Floods Using Multivariate Copulas: Case Study of Jinsha River and Min River, China, *Journal of Hydrologic Engineering*, 24, 05018030, 2019.

- Poulin, A., Huard, D., Favre, A.-C., and Pugin, S.: Importance of tail dependence in bivariate frequency analysis, *Journal of Hydrologic Engineering*, 12, 394–403, 2007.
- Raynal-Villasenor, J. and Salas, J.: Multivariate extreme value distributions in hydrological analyses, IN: *Water for the Future: Hydrology in Perspective*. IAHS Publication, 1987.
- Reddy, M. J. and Ganguli, P.: Bivariate flood frequency analysis of upper Godavari River flows using Archimedean copulas, *Water Resources Management*, 26, 3995–4018, 2012.
- Requena, A., Mediero, L., and Garrote, L.: A bivariate return period based on copulas for hydrologic dam design: accounting for reservoir routing in risk estimation, *Hydrology and Earth System Sciences*, 17, 3023, 2013.
- Sadegh, M., Ragno, E., and AghaKouchak, A.: Multivariate Copula Analysis Toolbox (MvCAT): describing dependence and underlying uncertainty using a Bayesian framework, *Water Resources Research*, 53, 5166–5183, 2017.
- Sadegh, M., Moftakhari, H., Gupta, H. V., Ragno, E., Mazdiyasni, O., Sanders, B., Matthew, R., and AghaKouchak, A.: Multihazard scenarios for analysis of compound extreme events, *Geophysical Research Letters*, 45, 5470–5480, 2018.
- Salvadori, G.: Bivariate return periods via 2-copulas, *Statistical Methodology*, 1, 129–144, 2004.
- Salvadori, G. and De Michele, C.: Frequency analysis via copulas: Theoretical aspects and applications to hydrological events, *Water resources research*, 40, 2004.
- Salvadori, G. and De Michele, C.: On the use of copulas in hydrology: theory and practice, *Journal of Hydrologic Engineering*, 12, 369–380, 2007.
- Salvadori, G. and De Michele, C.: Multivariate multiparameter extreme value models and return periods: A copula approach, *Water resources research*, 46, 2010.
- Salvadori, G., Durante, F., and De Michele, C.: On the return period and design in a multivariate framework, *Hydrology and Earth System Sciences*, 15, 3293–3305, 2011.
- Salvadori, G., Durante, F., and De Michele, C.: Multivariate return period calculation via survival functions, *Water Resources Research*, 49, 2308–2311, 2013.
- Salvadori, G., Tomasicchio, G., and D'Alessandro, F.: Practical guidelines for multivariate analysis and design in coastal and off-shore engineering, *Coastal Engineering*, 88, 1–14, 2014.
- Salvadori, G., Durante, F., Tomasicchio, G., and D'alessandro, F.: Practical guidelines for the multivariate assessment of the structural risk in coastal and off-shore engineering, *Coastal Engineering*, 95, 77–83, 2015.
- Salvadori, G., Durante, F., De Michele, C., Bernardi, M., and Petrella, L.: A multivariate copula-based framework for dealing with hazard scenarios and failure probabilities, *Water Resources Research*, 52, 3701–3721, 2016.
- Seneviratne, S., Nicholls, N., Easterling, D., Goodess, C., Kanae, S., Kossin, J., Luo, Y., Marengo, J., McInnes, K., Rahimi, M., Reichstein, M., Sorteberg, A., Vera, C., and Zhang, X.: Changes in climate extremes and their impacts on the natural physical environment, in: *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation: A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change*, edited by Field, C., Barros, V., Stocker, T., Qin, D., Dokken, D., Ebi, K., Mastrandrea, M., Mach, K., Plattner, G.-K., Allen, S., Tignor, M., , and Midgley, P., pp. 109–230, Cambridge University Press, Cambridge, UK, and New York, NY, USA, 2012.
- Serinaldi, F.: Dismissing return periods!, *Stochastic Environmental Research and Risk Assessment*, 29, 1179–1189, 2015.
- Serinaldi, F.: Can we tell more than we can know? The limits of bivariate drought analyses in the United States, *Stochastic Environmental Research and Risk Assessment*, 30, 1691–1704, 2016.

- 640 Shiau, J.: Return period of bivariate distributed extreme hydrological events, *Stochastic environmental research and risk assessment*, 17, 42–57, 2003.
- Sklar, M.: Fonctions de repartition an dimensions et leurs marges, *Publ. inst. statist. univ. Paris*, 8, 229–231, 1959.
- Swift Jr, L. W. and Schreuder, H. T.: Fitting daily precipitation amounts using the SB distribution, *Monthly Weather Review*, 109, 2535–2540, 1981.
- 645 Taherkhani, M., Vitousek, S., Barnard, P. L., Frazer, N., Anderson, T. R., and Fletcher, C. H.: Sea-level rise exponentially increases coastal flood frequency, *Scientific reports*, 10, 1–17, 2020.
- Tebaldi, C., Strauss, B. H., and Zervas, C. E.: Modelling sea level rise impacts on storm surges along US coasts, *Environmental Research Letters*, 7, 014 032, 2012.
- Tu, X., Du, Y., Singh, V. P., and Chen, X.: Joint distribution of design precipitation and tide and impact of sampling in a coastal area, *International Journal of Climatology*, 38, e290–e302, 2018.
- 650 USACE: Coastal risk reduction and resilience, 2013.
- Volpi, E. and Fiori, A.: Hydraulic structures subject to bivariate hydrological loads: Return period, design, and risk assessment, *Water Resources Research*, 50, 885–897, 2014.
- Wahl, T., Mudersbach, C., and Jensen, J.: Assessing the hydrodynamic boundary conditions for risk analyses in coastal areas: a multivariate statistical aapproach based on Copula functions, *Natural Hazards and Earth System Science*, 12, 495–510, 2012.
- 655 Wahl, T., Jain, S., Bender, J., Meyers, S. D., and Luther, M. E.: Increasing risk of compound flooding from storm surge and rainfall for major US cities, *Nature Climate Change*, 5, 1093, 2015.
- Ward, P. J., Couasnon, A., Eilander, D., Haigh, I. D., Hendry, A., Muis, S., Veldkamp, T. I., Winsemius, H. C., and Wahl, T.: Dependence between high sea-level and high river discharge increases flood hazard in global deltas and estuaries, *Environmental Research Letters*, 13, 084 012, 2018.
- 660 White, C. J.: The use of joint probability analysis to predict flood frequency in estuaries and tidal rivers, Ph.D. thesis, University of Southampton, 2007.
- Xu, H., Xu, K., Lian, J., and Ma, C.: Compound effects of rainfall and storm tides on coastal flooding risk, *Stochastic Environmental Research and Risk Assessment*, 33, 1249–1261, 2019.
- 665 Xu, K., Ma, C., Lian, J., and Bin, L.: Joint probability analysis of extreme precipitation and storm tide in a coastal city under changing environment, *PLoS One*, 9, e109 341, 2014.
- Yang, X., Wang, J., and Weng, S.: Joint Probability Study of Destructive Factors Related to the “Triad” Phenomenon during Typhoon Events in the Coastal Regions: Taking Jiangsu Province as an Example, *Journal of Hydrologic Engineering*, 25, 05020 038, 2020.
- Yue, S.: Joint probability distribution of annual maximum storm peaks and amounts as represented by daily rainfalls, *Hydrological Sciences Journal*, 45, 315–326, 2000a.
- 670 Yue, S.: The Gumbel logistic model for representing a multivariate storm event, *Advances in Water Resources*, 24, 179–185, 2000b.
- Yue, S.: A bivariate gamma distribution for use in multivariate flood frequency analysis, *Hydrological Processes*, 15, 1033–1045, 2001a.
- Yue, S.: A bivariate extreme value distribution applied to flood frequency analysis, *Hydrology Research*, 32, 49–64, 2001b.
- Yue, S.: The bivariate lognormal distribution for describing joint statistical properties of a multivariate storm event, *Environmetrics: The official journal of the International Environmetrics Society*, 13, 811–819, 2002.
- 675 Yue, S. and Rasmussen, P.: Bivariate frequency analysis: discussion of some useful concepts in hydrological application, *Hydrological Processes*, 16, 2881–2898, 2002.



- Zhang, H., Wu, C., Chen, W., and Huang, G.: Assessing the impact of climate change on the waterlogging risk in coastal cities: A case study of Guangzhou, South China, *Journal of Hydrometeorology*, 18, 1549–1562, 2017.
- 680 Zhang, L. and Singh, V. P.: Bivariate rainfall and runoff analysis using entropy and copula theories, *Entropy*, 14, 1784–1812, 2012.
- Zheng, F., Westra, S., Leonard, M., and Sisson, S. A.: Modeling dependence between extreme rainfall and storm surge to estimate coastal flooding risk, *Water Resources Research*, 50, 2050–2071, 2014.
- Zhong, M., Wang, J., Jiang, T., Huang, Z., Chen, X., and Hong, Y.: Using the Apriori Algorithm and Copula Function for the Bivariate Analysis of Flash Flood Risk, *Water*, 12, 2223, 2020.