

# 1 **Lessons from the 2018-2019 European droughts: A collective need for**

## 2 **unifying drought risk management**

3 Veit Blauhut<sup>1</sup>, Michael Stoelzle<sup>1</sup>, Lauri Ahopelto<sup>2,3</sup>, Manuela I. Brunner<sup>1</sup>, Claudia Teutschbein<sup>4</sup>, Doris E  
4 Wendt<sup>5,6</sup>, Vytautas Akstinas<sup>7</sup>, Sigrid J. Bakke<sup>8</sup>, Lucy J. Barker<sup>9</sup>, Lenka Bartošová<sup>10</sup>, Agrita Briede<sup>11</sup>,  
5 Carmelo Cammalleri<sup>12</sup>, Ksenija Cindrić Kalin<sup>13</sup>, Lucia De Stefano<sup>14</sup>, Miriam Fendeková<sup>15</sup>, David C.  
6 Finger<sup>16, 17</sup>, Marijke Huysmans<sup>18</sup>, Mirjana Ivanov<sup>19</sup>, Jaak Jaagus<sup>20</sup>, Jiří Jakubínský<sup>10</sup>, Svitlana  
7 Krakovska<sup>21</sup>, Gregor Laaha<sup>22</sup>, Monika Lakatos<sup>23</sup>, Kiril Manevski<sup>24</sup>, Mathias Neumann Andersen<sup>24</sup>, Nina  
8 Nikolova<sup>25</sup>, Marzena Osuch<sup>26</sup>, Pieter van Oel<sup>27</sup>, Kalina Radeva<sup>25</sup>, Renata J. Romanowicz<sup>26</sup>, Elena Toth<sup>28</sup>,  
9 Mirek Trnka<sup>10</sup>, Marko Urošev<sup>29</sup>, Julia Urquijo Reguera<sup>30</sup>, Eric Sauquet<sup>31</sup>, Aleksandra Stevkov<sup>32</sup>, Lena M  
10 Tallaksen<sup>8</sup>, Iryna Trofimova<sup>21</sup>, Anne F. Van Loon<sup>33</sup>, Michelle T. H. van Vliet<sup>34</sup>, Jean-Philippe Vidal<sup>31</sup>,  
11 Niko Wanders<sup>34</sup>, Micha Werner<sup>35</sup>, Patrick Willems<sup>36</sup>, Nenad Živković<sup>37</sup>

12 <sup>1</sup> Environmental Hydrological Systems, Faculty of Environment and Natural Resources, University of Freiburg, Germany

13 <sup>2</sup> Water and Development Research Group, School of Engineering, Aalto University, Finland

14 <sup>3</sup> Freshwater Centre, Finnish Environment Institute, Finland

15 <sup>4</sup> Department of Earth Sciences, Program for Air, Water and Landscape Sciences; Hydrology, Uppsala University, Sweden

16 <sup>5</sup> School of Geography, Earth and Environmental Sciences, University of Birmingham, UK

17 <sup>6</sup> Department of Civil Engineering, University of Bristol, UK

18 <sup>7</sup> Laboratory of Hydrology, Lithuanian Energy Institute, Lithuania

19 <sup>8</sup> Department of Geosciences, University of Oslo, Oslo, Norway

20 <sup>9</sup> UK Centre for Ecology & Hydrology, Wallingford, UK

21 <sup>10</sup> Global Change Research Institute CAS, Brno, Czech Republic

22 <sup>11</sup> Faculty of Geography and Earth Sciences, University of Latvia, Latvia

23 <sup>12</sup> European Commission, Joint Research Centre, Ispra, Italy

24 <sup>13</sup> Croatian Meteorological and Hydrological Service, Zagreb, Croatia

25 <sup>14</sup> Facultad de Ciencias Geológicas, Universidad Complutense de Madrid, Spain

26 <sup>15</sup> Univerzita Komenského v Bratislave, Slovak Republic

27 <sup>16</sup> School of engineering, Reykjavik University, Reykjavik, Iceland

28 <sup>17</sup> Energieinstitut an der Johannes Kepler Universität, Linz, Austria

29 <sup>18</sup> Department of Hydrology and Hydraulic Engineering, Vrije Universiteit Brussel, Brussel, Belgium

30 <sup>19</sup> Institute of Hydrometeorology and Seismology, Montenegro

31 <sup>20</sup> Department of Geography, Institute of Ecology and Earth Sciences, University of Tartu, Estonia

32 <sup>21</sup> Laboratory of Applied Climatology, Ukrainian Hydrometeorological Institute, Kyiv, Ukraine

33 <sup>22</sup> Institute of Statistics, University of Natural Resources and Life Sciences, Vienna, Austria

34 <sup>23</sup> Hungarian Meteorological Service, Budapest, Hungary

35 <sup>24</sup> Department of Agroecology, Aarhus University, Tjele, Denmark

36 <sup>25</sup> Faculty of Geology and Geography, Sofia University “St. Kliment Ohridski”, Bulgaria

37 <sup>26</sup> Institute of Geophysics, Polish Academy of Sciences, Warsaw, Poland  
38 <sup>27</sup> Water Resources Management Group, Wageningen University, the Netherlands  
39 <sup>28</sup> Dept. of Civil, Chemical, Environmental and Materials Engineering, University of Bologna, Bologna, Italy  
40 <sup>29</sup> Geographical Institute “Jovan Cvijić”, Serbian Academy of Sciences and Arts, Serbia  
41 <sup>30</sup> Department of Agroforestry Engineering, Escuela Técnica Superior de Ingeniería Agronómica y de Biosistemas,  
42 Universidad Politécnica de Madrid, Madrid, Spain  
43 <sup>31</sup> INRAE, RiverLy, Villeurbanne, France  
44 <sup>32</sup> Department of Meteorology, National Hydrometeorological Service, Skopje, North Macedonia  
45 <sup>33</sup> Institute for Environmental Studies, Vrije Universiteit Amsterdam, the Netherlands  
46 <sup>34</sup> Department of Physical Geography, Utrecht University, the Netherlands  
47 <sup>35</sup> IHE Delft Institute for Water Education, Water Resources & Ecosystems Department, Delft, the Netherlands  
48 <sup>36</sup> Hydraulics and Geotechnics Section, Department of Civil Engineering, KU Leuven, Belgium  
49 <sup>37</sup> Faculty of Geography, University of Belgrade, Serbia

50

51 *Correspondence to:* Veit Blauhut (veit.blauhut@hydrology.uni-freiburg.de)

52 **Abstract.** Drought events and their impacts vary spatially and temporally due to diverse pedo-climatic and hydrologic  
53 conditions, as well as variations in exposure and vulnerability, such as demographics and response actions. While hazard  
54 severity and frequency of past drought events have been studied in detail, little is known about the effect of drought  
55 management strategies on the actual impacts, and how the hazard is perceived by relevant stakeholders. In a continental study,  
56 we characterised and assessed the impacts and the perceptions of two recent drought events (2018 and 2019) in Europe and  
57 examined the relationship between management strategies and drought perception, hazard and impact. The study was based  
58 on a pan-European survey involving national representatives from 28 countries and relevant stakeholders responding to a  
59 standard questionnaire. The survey focused on collecting information on stakeholders’ perceptions of drought, impacts on  
60 water resources and beyond, water availability and current drought management strategies on national and regional scale. The  
61 survey results were compared with the actual drought hazard information registered by the European Drought Observatory  
62 (EDO) for 2018 and 2019. The results highlighted high diversity in drought perception across different countries and in values  
63 of implemented drought management strategies to alleviate impacts by increasing national and sub-national awareness and  
64 resilience. The study identifies an urgent need to further reduce drought impacts by constructing and implementing a European  
65 macro-level drought governance approach, such as a directive, which would strengthen national drought management and  
66 mitigate damages to human and natural assets.

## 67 **1 Introduction**

### 68 **1.1 Drought impacts in Europe**

69 During recent decades, different parts of Europe have been affected by several severe, large-scale drought events, e.g. in 2003,  
70 2007, 2011, 2012, 2015, 2017, 2018, 2019 and 2020 (Baruth et al. 2020, Boergens 2020, 2017; Cindrić Kalin et al. 2015,

71 Garcia-Herrera et al. 2019; Hänsel et al. 2019; Ionita et al, 2017; Laaha et al., 2017). Each of these droughts was unique in  
72 terms of severity, spatio-temporal extent and associated direct and indirect impacts on human and natural resources (Stahl et  
73 al. 2015). Cammalleri et al. (2020) estimated drought-related losses in the European Union (EU) to be about 9 billion Euros  
74 annually. The largest share of these losses is typically seen in agriculture, energy and public water supply sectors (Cammalleri  
75 et al. 2020), triggered mainly by agricultural (soil moisture deficit) and hydrological drought (deficit in river flow and  
76 groundwater; Van Lanen et al. 2015). These sectoral losses likely represent only part of the actual drought impacts, as indirect,  
77 intangible or subtle impacts are more difficult to identify and quantify, such as adverse effects on ecosystem services and  
78 human health (UNDRR, 2021). According to the European Drought Impact report Inventory (EDII; Stahl et al. 2016), further  
79 impacts on aquaculture, ecosystems, humans and public safety, as well as conflicts between sectoral water users, have been  
80 reported. Herein, the occurrence and the composition of drought impacts are assumed to greatly vary with regional and national  
81 exposure, perception and vulnerability to droughts (e.g. Stahl et al., 2016).

## 82 **1.2 Drought Management in Europe**

83 A key element to mitigate drought impact is to respond promptly, i.e., implement drought management planning strategies and  
84 associated action plans (UNDRR 2019). However, a directive for drought risk management does not exist on near-continental  
85 scale such as in the EU (Hervás-Gámez et al. 2019), despite the identified potential for reducing emergency management costs  
86 through proactive management (Cammalleri et al 2020; Howarth et al. 2018). So far, “droughts have only been succinctly dealt  
87 with in the Water Framework Directive with no compulsory actions” (Hervás-Gámez et al. 2019). However, recommendations  
88 are not adopted in all relevant/major river basin districts (EC 2019). The “European Commission’s Communication on water  
89 scarcity and drought” and the ‘Blueprint to Safeguard Europe’s Water Resources’ (EC 2012) directly tackle drought and  
90 address current flaws and policy gaps. These documents have received a mixed response, ranging from “the Communication  
91 is still weak and lacks teeth in the policy landscape” (Stein et al. 2016) to “it is hoped to lead to an EU water policy development  
92 in a long term” (Hervás-Gámez et al. 2019). However, some countries have historically been and are more prone to drought  
93 compared to others due to their pedo-climatic settings, and although drought risk management does exist in these countries  
94 through national legislation, it mostly happens indirectly via policy-making regarding environmental protection, soil  
95 management, or water and climate adaptation (e.g. Caillet et al. 2019, Hanger-Kopp and Palka 2020). Moreover, a number of  
96 technical guidelines exist to support the development and the implementation of national drought resilience, adaptation and  
97 management plans (e.g. UNCCD 2019). In fact, different national legal approaches not being internationally coordinated can  
98 create conflicts, i.e. water scarcity in one region/country at the cost of another, such as the case of the Blue Nile between Egypt  
99 and Ethiopia (El Bastawesy et al., 2015) or the Danube between Hungary and Slovakia (Vuković et al., 2014). Therefore, a  
100 coordinated approach is required. Trnka et al. (2018) suggested to improve the understanding of triggers causing paradigm  
101 shifts from response-based to proactive drought management and policies as a priority research question.

### 102 **1.3 The 2018 & 2019 European droughts**

103 For several successive years, large parts of Europe were affected by severe and widespread droughts, which highlighted the  
104 vulnerability of its socio-economic and environmental systems. The 2018 event was special because of both rainfall deficit  
105 and high temperatures in many European countries (Rosner et al. 2019), with record-breaking high temperatures in several  
106 regions (Bakke et al., 2020), and reached otherwise cool and humid northern regions. This compound hot-dry event led to  
107 major impacts in north-central and north-eastern Europe, particularly affecting agriculture, livestock farming and forestry  
108 (Bakke et al. 2020, Beillouin et al. 2020, Rosner et al. 2019, Salmoral et al. 2020, Schuldt et al. 2020, Thompson et al. 2020)  
109 as reported for Sweden, Finland, Estonia, Lithuania, Latvia, Denmark, the Netherlands, Belgium, Germany, the United  
110 Kingdom, and eastern France (Moravec et al. 2021, Turner et al. 2021). The propagation of the meteorological drought resulted  
111 in low reservoir levels and river discharge, which impaired public water supply, leading to partial shut downs of nuclear power  
112 plants and triggering massive fish deaths in upstream watersheds (e.g. de Brito 2021). In contrast to central and northern  
113 Europe, the western Mediterranean countries experienced above-average wet conditions in 2018 after having experienced a  
114 very severe drought on the Iberian Peninsula in 2016–2017 and in Italy in 2017 (Garcia-Herrera et al. 2019; Rita et al. 2019),  
115 while the eastern Mediterranean experienced below average dry conditions (DriDanube-Watch 2018). Opposite to 2018, the  
116 2019 drought was centred on eastern Germany, Czech Republic and Poland before spreading westward (Boergens et al. 2020).  
117 The most affected regions were still suffering from large water balance deficits from the 2018 drought (Boergens et al. 2020)  
118 at the start of 2019. Hari et al. (2020) declared the period 2018–2019 in central Europe a two-year drought event unprecedented  
119 in severity in the last 250 years, whereas Büntgen et al. (2021) show an accumulation of drought signals in central Europe over  
120 five summers i.e. 2014-2018.

### 121 **1.4. Drought risk and perception**

122 The hydro-climatic aspects of past drought events have been studied in detail (e.g. Barker et al. 2019; Hisdal and Tallaksen  
123 2003; Dai 2013; Cheval et al. 2014; Jaagus et al. 2021; Laaha et al. 2017; Radeva et al. 2018; Spinoni et al. 2015, 2018),  
124 whereas knowledge of the relationship between drought management, perception and impacts remains limited (Blauhut 2020,  
125 Hagenlocher et al. 2019; Kreibich et al. 2019). Understanding how different stakeholders perceive a specific drought event  
126 and its potential impacts can contribute to defining and successfully implementing drought mitigation measures adapted to a  
127 site-specific context (Alduce et al. 2017). Only a few studies have analysed relationships between drought perceptions and  
128 impacts. For instance, Teutschbein et al. (2019) assessed the link between perceived drought severity, impacts, preparedness  
129 and management and measured hydrological drought impacts for two consecutive drought events (2017 and 2018) in Sweden.  
130 Although the authors did not find a significant relationship between the perceived level of drought impacts and the presence  
131 of a drought action plan, there was evidence that regions with a drought action plan applied significantly more measures in  
132 their drought response. Furthermore, the perceived drought severity did not match the observed severity of meteorological and  
133 hydrological droughts in Sweden: decision makers consistently overestimated the severity of mild drought events, while

134 underestimating more extreme drought conditions. In contrast, Blauhut et al. (2016) identified “drought awareness” and  
135 “drought management plans” as vulnerability factors driving drought risk for certain impact categories, such as agriculture and  
136 livestock farming, public water supply or freshwater ecosystems. The analysis of Blauhut et al. (2015b) suggested that while  
137 national and international water management policies and guidelines may have decreased vulnerability, they may also have  
138 increased awareness and recognition of environmental impacts, leading to an increased number of reported drought impacts.  
139 Hence, previous statements on the relationship between the existence of drought risk management plans and drought impacts  
140 cannot be generalised.

## 141 **1.5. Study aim**

142 The aim of this paper is to assess how monitored drought-hazard severity relates to drought perception and drought  
143 management strategies. We hypothesise that perceived drought impacts are not necessarily related to the severity of the drought  
144 hazard, but are strongly influenced by national awareness and drought management strategies. To verify this hypothesis, we  
145 investigated how the droughts of 2018 and 2019 in 28 European countries were related to a) the drought hazard as monitored  
146 by the European Drought Observatory (EDO; <https://edo.jrc.ec.europa.eu>), b) drought management actions taken in the  
147 different countries, c) drought perception by water managers and agencies and d) drought awareness. National drought  
148 perceptions, management and impacts were studied using a pan-European survey. On the basis of this survey, we discuss the  
149 potential benefits of a European drought directive, similar to the Floods Directive (2007/60/EC) with respect to reducing  
150 drought vulnerability and impacts by macro-level governance.

## 151 **2. Data**

152 In order to evaluate the hypothesis, two different types of spatial data were collected and compared: i) drought information as  
153 monitored by the EDO (<https://edo.jrc.ec.europa.eu>) and ii) information on drought impacts, perception and state of drought  
154 management plans collected through a pan-European survey targeting water managers and water agencies. Kosovo was not  
155 investigated disaggregated from Serbia (please see disclaimer)

156 The hydro-climatic situations in 2018 and 2019 were described using a set of drought indices compiled by EDO for a variety  
157 of drought types including meteorological drought (Standardised Precipitation Index (SPI) for 1, 3, 6, 9 and 12 month  
158 accumulation periods), soil moisture drought (soil moisture anomaly; SM), hydrological drought (Low Flow Index; LFI,  
159 representing the discharge anomaly with respect to a daily threshold), and vegetation drought (anomaly of Fraction of Absorbed  
160 Photosynthetically Active Radiation; fAPAR). The SPI is given at a monthly resolution, whereas the other indices are presented  
161 in 10-day non-overlapping intervals. To increase comparability of the four indices, the EDO data was further classified into  
162 categorical drought classes: no drought, moderate drought, severe drought and extreme drought. The standardised products  
163 SPI, fAPAR and SM are categorised following McKee et al. (1993) (Table 1), the Low-Flow Index (LFI) is computed from  
164 the daily streamflow values produced by the LISFLOOD hydrological model”. The drought classification scheme used for LFI

165 is taken from the European Drought Observatory ( Table 1). These drought classes are in operational use at the EDO.  
166 Furthermore, the fAPAR was restricted to the warm season in Europe from April to August and was not monitored for Iceland.  
167 Detailed information on the drought indices and drought classes applied herein can be found in the corresponding EDO  
168 indicator factsheets (<https://edo.jrc.ec.europa.eu/edov2/php/index.php?id=1101>).

169 In order to assess the country specific perception of drought, management and impacts with focus on the 2018 and 2019, a  
170 pan-European survey was designed by the International Association of Hydrological Science (IAHS) - Panta-Rhei “Drought  
171 in the Anthropocene” working group. National representatives of each country were selected and assigned responsibility to  
172 translate, distribute and evaluate the survey and all associated communication and feedback. The network of national  
173 representatives developed out of our active Panta Rhei- Drought in the Anthropocene group but also partly from the Euro-  
174 FRIEND low flow group. The idea was to have representatives affiliated to science or governmental agencies. Doing so, we  
175 expected a neutral point of view and comprehensive knowledge on the different aspects we were interested in. Furthermore,  
176 we expected such persons to be well networked and thus constitute a representative sample of stakeholders. The survey targeted  
177 representatives of water management organisations and water agencies. Survey respondents were selected by the national  
178 representatives aiming to provide a balanced view of national opinions and drought management practices (or actions), as well  
179 as local and regional knowledge within each country. The content of the survey was adapted from Teutschbein et al. (2019),  
180 who studied 290 Swedish municipalities to evaluate the relationship between perceived drought severity, impact, preparedness  
181 and management, aiming to compare stakeholder perception with hydrological drought indices. The perception of heat was  
182 not investigated.

183 The 26 questions of the survey covered the following themes:

- 184 • Respondent background and water resource(s) used/managed,
- 185 • General perception of drought and associated risks,
- 186 • Drought risk-related concepts and the drought management applied, and
- 187 • Perception of the 2018 and the 2019 drought events and their impacts.

188 The survey questions can be found in Table S1 in the supplements. The paper and the figures displayed in the main body  
189 present a synthesis and insights from the pan-European comparison of the responses. More detailed aspects of the individual  
190 country responses are shown in Figs. S1-7 of the supplements.

## 191 **3. Results and Discussion**

### 192 **3.1. The drought events of 2018 and 2019 – hydro-meteorological results**

193 The drought indices of the pan-European droughts in 2018 and 2019 are presented in Fig. 1, showing the specific hydro-  
194 meteorological conditions of a specific time in the year or the month with maximum proportion of area within each country  
195 affected by the hazard, as defined by severe or extreme category. The results on monthly and national resolution are shown in  
196 Fig. S1 in the supplements. Overall, the 2018 meteorological drought (as defined by SPI-3, SPI-6, SPI-9 and SPI-12) affected

197 mainly Central and Northern Europe. The Benelux countries, Germany, Denmark, Sweden and Finland showed an especially  
198 high spatial coverage of severe or extreme drought hazard. In early spring, rainfall deficits started in the North, i.e., Norway,  
199 Sweden and Finland, Lithuania and Latvia (Fig. S1) and accumulated over Central and Northern Europe, peaking in the  
200 summer with high shares of extreme drought hazard at short accumulation periods (SPI-1). Strong soil moisture deficits in the  
201 summer were detected in regions affected by strong precipitation deficits over multiple months (SPI-3, SPI-6; Fig. S1). At the  
202 European scale, soil moisture deficits were especially high in Northern Europe from June to August and the area under severe  
203 drought in central Europe peaked in October and November (Fig. 1).

204 The hydrological drought of 2018 followed a similar spatial pattern as the meteorological drought, with severe hazard levels  
205 in the Benelux countries, Germany, Czech Republic, Norway and Sweden. The maximum spatial coverage of severe or extreme  
206 low flows in Northern Europe occurred in June and July, in particular for countries where rainfall deficits continued and more  
207 intense deficits developed (SPI-9, SPI-12; Fig. S1). The maximum coverage of the 2018 hydrological drought in Central  
208 Europe occurred in October and November (Fig. 1).

209 Vegetation drought indicated by fAPAR was the most severe in Denmark and was contrasting with the drought signals of the  
210 other indices. For example, large parts of Belarus and France were under severe or extreme drought, while only small parts of  
211 Sweden and Finland were affected. In countries where precipitation deficits continued to accumulate over the 2018/2019 winter  
212 period, water deficits resulted in country-specific low flow conditions (Fig. 1). The multi-year drought 2018-2019 particularly  
213 affected Belgium, Belarus, the Czech Republic, Germany, Finland, Latvia, Luxemburg, Lithuania, the Netherlands, Poland,  
214 Sweden, Switzerland and the United Kingdom. Furthermore, Iceland also experienced an exceptionally long dry period in  
215 2019. However, the effects were not as intense as in mainland Europe, mainly due to the numerous icecaps that provided ice  
216 melt, an extensive snow cover during winter (Helmert et al. 2018), a subpolar climate, and warm and humid ocean winds, that  
217 could generate local rain events (Finger, 2018).

218 The 2019 drought was overall less severe compared to 2018, except for Iceland, who experienced an unusually long dry period.  
219 The centre of the meteorological drought moved eastwards with large areas under severe or extreme drought in Lithuania,  
220 Belarus and Ukraine. Despite the lower intensity of the 2019 meteorological drought, soil moisture deficits remained high in  
221 Central Europe, especially Poland, the Baltic and the Benelux countries. In Central Europe, soil moisture drought peaked in  
222 early February and March compared to a delayed peak in April in Poland and the Baltics and an even later peak in Ukraine  
223 and Moldova towards the end of 2019. The low flow situation in Central Europe and Scandinavia partly recovered, although  
224 severe hazard levels were still detected from July to September. The eastern European countries showed an overall increase in  
225 low flow severity peaking earlier in the year (April and May). In addition, fAPAR was less severe in 2019 for most months  
226 and most of Europe, while South-Eastern Europe and the Balkans showed increased hazard severity.

227 In southern and south-eastern Europe, the hydro-climatic conditions of 2018 and 2019 differed from the rest of Europe. In  
228 2018, Spain, Portugal and Italy had recovered from drought conditions, but deficits again developed in early 2019. In south-  
229 eastern Europe, winter 2018/2019 precipitation deficits were detected across much of the Balkan Peninsula, as well as in

230 Slovakia. In Ukraine, Moldova and Romania, the 2018 event was moderate in the second half of the year and further rainfall  
231 deficits accumulated during winter, which led to rising soil moisture deficits from summer 2019 to the end of 2020.

### 232 **3.2. The drought events of 2018 and 2019 - perception and management**

233 The online survey yielded contributions by 712 respondents from 28 European countries (Fig. 2a) with the number of responses  
234 varying by country, i.e., from a single expert (Romania) to over 100 replies (Sweden). The majority of the respondents were  
235 employees at governmental institutions (74%) at different administrative levels, with expertise related to water management,  
236 environment, meteorology and agriculture. Furthermore, private and public companies (operators of public water supply  
237 systems, hydropower plants; 13%), scientific institutions (4%) and other non-governmental organisations with a focus on  
238 environment and ecology (3%) also contributed to the survey.

239 The importance of the water resources as perceived by the participants (under normal conditions) ranked differently across the  
240 continent (Figs. 2b and c). The participants were asked to rank a selection of water resources, but were also free to add  
241 additional ones. The sources of “artificial recharge” were not specified. If the nationally averaged importance of water  
242 resources were ranked equally (e.g. regulated and individual groundwater use both ranked as second most important), their  
243 importance as rank #2 and #3 were also evaluated. Overall, the majority of the respondents selected groundwater as the most  
244 important resource (~35% of all participants), followed by surface water from rivers (22%), reservoirs (13%), individual wells  
245 (11%) and artificial groundwater recharge (11%). Further “Other” water resources such as rainfall collectors, ponds or water  
246 transfer systems were listed a few times (<1%). Specific spatial patterns of water resource importance were not apparent,  
247 although individual wells appeared to be more important in eastern Europe and artificial groundwater recharge was highlighted  
248 in Italy, Hungary and Bulgaria. In the case of Spain, the questionnaire was adapted to national specificities and resulted in less  
249 water-usage categories; here “regulated surface water” falls in the category of “surface water from reservoirs”. Accordingly  
250 water resources ranks are: #1 regulated surface water and #2 groundwater. A more detailed national breakdown of Fig.2 can  
251 be found in the supplements (Fig. S2).

252 The use of a drought definition to categorise drought hazard varied markedly across Europe (Fig. 3). About 40% of all  
253 participants did not have an operational drought definition in their public and private organisations, and a further 15% did not  
254 know whether there was one. In contrast, for Czech Republic, Spain, Italy and France all participants had an operational  
255 drought reporting system. With regard to the participants’ affiliation (see Table S2), about 60% of those working for  
256 governmental authorities did not have - or were not aware of - an operational drought definition, in contrast to private  
257 companies where around 30% were unaware of a drought definition. Overall, about 20% defined drought by a single drought  
258 type index (such as meteorological drought), 15% used two and 10% used three different drought indices. The majority of  
259 participants used meteorological and hydrological indices (30% each) and about 15% relied on soil moisture and vegetation  
260 conditions. Furthermore, drought-impact information such as vegetation activity (e.g. NDVI), crop yields or forest fire indices,  
261 but also media reports were used. In Spain, the ‘Special Drought Management Plans’ define two types of drought-related  
262 events: prolonged drought (meteorological) quantified by precipitation deficit over different time periods, and conjunctural



263 water scarcity identified through the assessment of available water resources. This question was not asked in Sweden and  
264 Poland, and in Latvia, drought definitions were not operationalised.

265 Following the drought definition question, respondents were asked whether an established governmental drought declaration  
266 system existed or the declaration of drought situations was based on case-specific decisions (Fig. S2). An operational  
267 declaration scheme is defined here as an official government implemented method of defining a drought situation, often  
268 including drought severity thresholds and pre-defined measures. Operational drought declaration schemes (at country or county  
269 level) were scarce across the continent, though these were found to be present in Spain, France, the Netherlands and the Czech  
270 Republic. In Czech Republic, drought declaration is based on the open national drought monitoring platform Intersucho (Trnka  
271 et al. 2020). The same platform is shared also by the Slovak Hydrometeorological Institute (Labudová et al. 2018).

272 In Spain, governmental drought declaration schemes are included in the Special Drought Management Plans approved at river  
273 basin level, whereas each one has adapted it to their context and characteristics. Outside of Spain, individual decisions on  
274 drought declarations are more commonly present in regions with a lack of fixed drought declaration schemes. In some  
275 countries, drought situations are declared by the “Emergency situations commission meetings”, for example in Lithuania, the  
276 Netherlands and the UK, where a national water management centre and drought committee advise the government. In Latvia,  
277 Estonia, Austria, Bulgaria and Denmark, more than half of the respondents did not know of the existence of any governmental  
278 drought declaration scheme or were not sure that one existed.

279 In over 25 countries, the majority of participants (>50%) responded positively to the question of whether future climate change  
280 may affect water resources. (Fig. 4 a). The majority of respondents expected the occurrence of droughts to “increase” or  
281 “strongly increase” in the (near-) future (Fig. 4 b). However, no relationship could be established between the expected future  
282 changes in drought hazard and the degree to which climate change is considered in policies. In addition, the responses about  
283 the expectation of “the need for more regulation of water distribution to fewer consumers due to shortages in the future” was  
284 neither linked to the expectation of future drought occurrence nor climate change consideration (Fig. 4 c). For example, in the  
285 UK, around 15% of respondents agreed that more regulation will be needed with a majority expecting an increase in drought  
286 occurrence, whereas in North Macedonia, about 85% agreed on a need for increased future regulation, having a similar share  
287 in future drought occurrence. Nevertheless, it appears that participants of northern European countries perceive less need for  
288 more regulation compared to the rest of Europe. As mentioned by the participants, future regulation is expected to take the  
289 form of an EU drought directive, ranking priorities, re-allocating water permits, technological enhancements to save water,  
290 water pricing and general water usage restrictions.

291 Few participants indicated that their countries had drought management action plans (~10%), although emergency action plans  
292 were more common (~25%), and both plans existed more in western compared to eastern Europe (Fig. 5 a). The UK, the  
293 Netherlands, Belgium, Spain, Switzerland, Italy and Montenegro were comparatively well prepared in this regard (>75% of  
294 the participants had an emergency action or management plan). The countries of Spain, Italy and Czech Republic had a high  
295 share of participants indicating drought risk management plans in operational mode. More than 150 participants of the entire  
296 survey indicated the intention to introduce new (or update existing) drought management plans. As indicated by the

297 participants, such management tools include strategies that range from the increase of water storage capacity or adapting  
298 farming practices, to the development of legally binding drought risk management actions.

299 To better understand the reasons for an absence of drought management plans, participants were asked for a possible  
300 explanation and answers were provided by the national experts as either pre-defined or free text options. At the country scale,  
301 “insufficient resources” and the perception that “drought is not seen as a risk” were the most frequent answers (Fig. 5b). For  
302 northern Europe and Austria, drought not seen as a risk was highlighted most often, whereas for the eastern European countries,  
303 the “lack of legal obligation- no European drought directive” and “Waiting for governmental advice” were selected by about  
304 15% of all participants. Further, a “Lack of knowledge” (on drought risk) was more prevalent in western Europe, whereas a  
305 “Lack of resources” (finance and capacity) were prominent in central and south-eastern Europe. Political issues (waiting for  
306 governmental advice, lack of forcing – no EU drought directive, and political lack of knowledge) were especially present in  
307 central and eastern Europe, but were less prominent in northern Europe.

308 With regard to communication and interaction during drought events, participants were asked whom they collaborate with to  
309 manage droughts (Fig. S3). On average, more than half of the participants collaborated with “other authorities” (e.g., county  
310 administrative boards or water authorities). About 45% interacted with “other departments or companies within the  
311 municipality” and about 20% with land owners and independent experts (such as universities). About 20% did not know about  
312 any collaboration and 5% of participants stated no existing collaboration.

### 313 **3.3. Survey-based perception and management of the 2018 and 2019 droughts in Europe**

314 The perception of the 2018 and 2019 drought events by the survey respondents showed country- and event-specific differences  
315 (Fig. 6). The participants could rank the hydroclimatic situation from extremely dry to wet; 2018 was mostly perceived as  
316 “drier than normal” and central and northern European countries in particular perceived “very dry” conditions (Fig. 6 a), with  
317 high proportions (25%) of “extremely dry” conditions in Czech Republic, Germany, Lithuania, North Macedonia and Norway  
318 (Fig. S4). The south-western European countries also perceived 2018 as “drier than normal”. In Iceland, 2018 was perceived  
319 as a wet year, in contrast to the dry conditions of 2019 which were perceived to be “one of the worst droughts” on record. For  
320 the rest of Europe, the 2019 drought was perceived as less severe than 2018 with the exception of France and Ukraine with  
321 high variations between countries (Fig. 6 b). The centre of the 2018 drought event shifted by the end of the year/ beginning of  
322 2019 from Central and northern Europe towards the East. Wetter conditions in northern Europe translated into perceptions of  
323 no or less severe drought in Scandinavia and the Baltics, respectively. The hydro-climatic situation in 2019 was still perceived  
324 as “very dry” (50%) in France, Belgium, Germany, Slovakia and Ukraine.

325 Drought management preparation in 2018 showed an east-to-west gradient, i.e., eastern, northern and central European  
326 countries felt overall more “prepared” while countries in western Europe perceived they were “not well” prepared (Fig. S5).  
327 The management of the 2018 drought event was generally perceived as being worse compared to 2019, except in some central  
328 and northern European countries. Most respondents thought that they were better prepared in 2019 due to the previous event  
329 that likely contributed to an earlier activation of emergency plans, if any. However, the perception of drought impacts only

330 shows minor differences between the two drought events (Fig. 6 c and d), with the exception of northern Europe. The  
331 Mediterranean and the Balkan countries perceived drought impacts as not severe or without impacts (Croatia) in both years,  
332 with a tendency towards a higher severity in 2019 for black sea countries. In Central Europe, participants perceived that they  
333 were severely affected in 2018 and this perception extended towards eastern Europe in 2019. Scandinavia and the Baltics were  
334 only slightly affected in 2018 with a lower perceived severity in 2019. For the majority of respondents, the drought of 2018  
335 played a crucial role in the perceived impacts of the 2019 drought event (Fig. S5). Most respondents perceived particularly  
336 negative consequences for agriculture, livestock farming, forestry and public water supply in 2018 and 2019 compared to  
337 relatively minor consequences regarding air pollution and conflicts (Fig. S6). At a first glance, the perception for these sectors  
338 differs only slightly between the events. However, soil moisture impacts, such as agricultural losses, impacts on freshwater  
339 aquaculture and fisheries or forest fires were less reported in 2019. Observing from 2018 to 2019, Denmark, Norway and the  
340 UK show stronger reductions in perceived impacts from 2018 to the 2019 drought event. Slightly more impacts were reported  
341 in 2019 for livestock farming in Ukraine, for forestry and terrestrial ecosystems in Belgium and Ukraine, for air quality in  
342 Bulgaria, Slovakia, North Macedonia and Ukraine, and for water quality in Austria, Czech Republic and North Macedonia. In  
343 contrast, Iceland was only affected in 2019 with strong effects on agriculture and water quality.

#### 344 **4. Discussion**

345 This is the first study that quantifies drought perception by water management related stakeholders at continental scale based  
346 on participatory survey. The survey analysis shows high diversity in perceived drought impacts (Fig. S6), which reflects  
347 Europe's pedo-climatic and socio-economic heterogeneity as is also shown by Stahl et al. (2016). The monitored and perceived  
348 drought hazard differed in some places as a result of the different drivers of drought impacts: hazard, exposure and vulnerability  
349 (IPCC 2014, UNDRR 2019). The diversity of impacted categories has been reported previously for similar drought events,  
350 e.g., in 1975, 1976 or 2003 (Stahl et al. 2016). Our findings corroborate those of Stahl et al. (2016) with different countries  
351 across Europe being affected by the hazard very differently. Large-scale weather patterns and differences in land surface  
352 properties play a crucial role in explaining this heterogeneity. For instance, Atlantic meridional dipole circulation anomalies  
353 have been found to be associated with Northern European droughts as represented by the SPI-6 and SPEI-6 (Standardised  
354 Precipitation-Evapotranspiration Index) indices (Kingston et al. 2015). The Scandinavian teleconnection pattern, which was  
355 unusually high in May and July 2018, resembles the large-scale atmospheric circulation pattern most associated with summer  
356 low flow in south- and eastern Scandinavia (Bakke et al. 2020). It should be also noted that the frequency of drought-related  
357 circulation patterns has been changing since the end of the 19th century with increasing frequencies over Central Europe (e.g.  
358 Lhotka et al., 2020; Trnka et al. 2009). The unique conditions of Iceland, where major drought events cannot be compared to  
359 the rest of Europe, was also shown by Spinoni et al. (2015), most likely attributed to its location influenced by warm humid  
360 winds and enhanced by the Gulf stream clashing with the cold Arctic winds from the North that generate frequent precipitation  
361 events (de Niet, 2020). Nevertheless, severe land degradation in Iceland has decreased the water holding capacity, making the

362 land susceptible to hydrological droughts (Finger et al. 2016, Keesstra et al. 2018). Furthermore, Spinoni et al. (2019) showed  
363 that major drought events as indicated by SPI and SPEI in central and northern Europe, north-eastern Europe, and southern  
364 Europe do not occur simultaneously, which was also evident in our results focusing on 2018 and 2019. The multi-year drought  
365 character of 2018 and 2019 became evident when focusing on the monitored hydrological drought conditions in Belgium,  
366 Switzerland, Czech Republic, Germany, Finland, Latvia, Luxemburg, Lithuania, Poland and Sweden.

367 In general, the hazard severity perceived by the surveyed stakeholders corresponded well with the hazard severity monitored  
368 by the EDO, though with some exceptions. For example, in 2018, large areas of Sweden and Finland were affected by severe  
369 (or extreme) hazard conditions according to the EDO, but the hydro-climatological situation was perceived as “dry”. In 2019,  
370 EDO reported a severe (or extreme) meteorological drought in Iceland and a severe (or extreme) soil moisture drought in the  
371 Baltic, but in both countries, the hazard severity was perceived as less severe and indexed as “drier than normal” to “dry”. In  
372 contrast, Norway’s participants perceived very dry conditions in 2018, but the proportion of monitored severe (or extreme)  
373 hazard conditions was low. These discrepancies could be attributed to the low awareness of the stakeholders for the drought  
374 conditions at a larger scale or the impact across different sectors, or the discrepancy between impact indicator and affected  
375 sector. Alternatively, standardised drought indices not effectively representing drought conditions everywhere may also  
376 contribute to these discrepancies. In higher latitude countries, a strong negative rainfall anomaly does not necessarily imply a  
377 deficit in water availability for e.g. plant water uptake or public water supply as storages are usually replenished after the snow  
378 melt period. (Cammalleri et al. 2016). As such, meteorological drought indices may not be appropriate to predict impacts and  
379 consequences for management of hydrological or agricultural (soil moisture) droughts. The wide range of drought definitions  
380 and associated high number of drought indices - combined with a widespread lack of operational declaration schemes -  
381 highlight the many obstacles when dealing with the complex inter- and transdisciplinary nature of drought impacts. A unique  
382 definition of drought that is valid across all regions and sectors is not possible in practice (Lloyd-Hughes 2014), especially if  
383 sectors, such as agriculture and water supply, are based on different laws and managed by different authorities. An effective  
384 implementation of macro-regional drought risk management requires a more holistic interdisciplinary view. Thus, drought  
385 cannot be declared by a single index only, the entire water cycle has to be considered, since droughts in different parts of the  
386 water cycle can lead to different impacts. Such a holistic view should start with initial meteorological drought (e.g. lower than  
387 normal precipitation often combined with higher than normal evaporation), causing a deficit in soil moisture, and if sustained  
388 for a sufficient time, may manifest itself as a hydrological drought (i.e. a deficit in streamflow and groundwater).

389 Our pan-European survey reflects the opinions of water professionals belonging to mostly the public sector and publicly owned  
390 companies. The perspectives of other citizens, local stakeholders, private companies and NGOs were less well represented.  
391 Nevertheless, the fraction of respondents’-affiliation differs among countries and can thus have had an influence of the herein  
392 generalised portrayal of drought risk. A statistical relation between affiliation and “other” replies (such as drought  
393 management, or reason for a lack of DRM) could not be found. Furthermore, sectoral and regional perceptions of drought risk  
394 might differ. For example, a hydropower production survey in southern Germany showed that legislative drought risk

395 regulation is not desired by reservoir operators, who would nevertheless support the development of drought risk management  
396 coupled with integrated river basin management (Siebert et al. 2021).

397 The preferential use of meteorological and hydrological indices to define drought by the participants was found similar to the  
398 findings of Bachmair et al. (2016). The absence of dedicated drought risk management strategies in many European countries  
399 is evident (Fig. 5 a) due to diverse and in some cases, contradictory reasons (Fig. 5 b). The country representatives were asked  
400 some broad questions on the state of national drought management and the potential for a European drought directive; the  
401 responses revealed an unsatisfactory state of national drought risk management in Europe (Fig. 7).

402 The existence of drought risk management plans or strategies tended to be higher in countries with more common water  
403 scarcity issues and more frequent drought events, such as those in the Mediterranean region (Tramblay et al. 2020). Moreover,  
404 only Spain's Special Drought Management Plans (updated in 2018; Hervás-Gómez & Delgado-Ramos, 2019) was considered  
405 as comprehensive and sufficient by national representatives. In addition, recent drought events may have forced governments  
406 to foster drought research and policy implementation, suggesting that a 'memory of recent disasters' improves disaster  
407 management and potentially mitigates drought impacts (DiBaldassarre et al. 2013; Kreibich, et al. 2017). Urquijo et al. (2016)  
408 stated that drought management is a combination of the history of water management and the frequency of drought, which is  
409 supported by our results from the Mediterranean countries and the Netherlands. Furthermore, case-specific effects of drought  
410 may also drive the need for risk management. In the Netherlands, for example, hydrological drought can increase salt water  
411 intrusion, increase land subsidence and structural instability of dikes. The resulting damages of these hydrological drought  
412 impacts decrease water security in the long-term, especially with regard to compound events. The engagement of non-  
413 governmental scientific groups also fosters drought risk management and particularly public and government awareness (e.g.  
414 Czech Republic).

415 The diversity of drought management approaches reflects the diversity of Europe's hydro-climatic conditions and governance  
416 contexts. However, droughts do not respect national borders and Europe has several shared river basins. In addition, climate  
417 change is estimated to increase drought severity and frequency globally and in Europe (UNDRR, 2021; Spinoni et al., 2018).  
418 The majority of the survey participants and all national representatives agreed that a pan-European drought management  
419 approach would support national and cross-boundary drought preparedness both now and in the future. While collaborations  
420 between water managers and agencies within countries are at least partly in place, as indicated in the survey, the difference  
421 between preparedness and proactive approaches to lower drought risk in Europe varies widely. Participants mainly in Central  
422 and Southern Europe indicated 'insufficient resources', 'lack of forcing', 'waiting for governmental advice' as a reason for not  
423 having a drought risk management plan. Across all national representatives and a majority of survey participants, there was a  
424 consensus that an EU Directive on drought risk management would be beneficial (whether or not countries are EU-member  
425 states). Similar to the Floods Directive (EU 2007), a common strategy should only set a coarse framework, delegating specific  
426 actions to the member states and especially regulating transboundary water management during drought. An EU Directive  
427 would be especially beneficial in countries where water resources management governance is not centralised, with wide  
428 procedural discrepancies among the different administrative regions and basin authorities. Recently, the Global Assessment

429 Report on Drought (GAR, 2021) highlighted that adaptive risk management and governance strategies are required as  
430 responses to complex risks such as drought, by means of actions, processes and institutions. A drought Directive, following  
431 the example of the European Floods Directive, would force member states and candidate countries to act and encourage  
432 cooperation across borders addressing the regional scale of drought hazard, secure resources and funding for drought risk  
433 research and most importantly initiate a common strategy to increase drought resilience. However, not all respondents of the  
434 survey fully shared this view, the main reason being that a pan-European approach would not be able to consider local  
435 specificities such as catchment physical characteristics, water infrastructure, water uses, and specific biodiversity needs.  
436 Accordingly, a pan-European approach should also be tailor-made to be trusted by the users. Therefore specific indicators and  
437 actions can be tailored to local situations and needs, but a general framework should be guiding the application of these. Thus,  
438 common action (e.g. a drought risk management strategy) may be conducted at a very general, broad and political level. At  
439 operational or local level, clear and common guidelines may be needed and the challenge is to be flexible enough to cover  
440 context-specific situations.

## 441 **5. Conclusions**

442 The pan-European survey on drought perception and management highlighted the heterogeneity in the perception of drought  
443 hazard, impacts and management across the European continent. The reflection on the drought events in the 2018-2019 period  
444 illustrated Europe's vulnerability to drought and the variable state of preparedness to withstand drought in many countries.  
445 Even though the awareness of a future increase of drought risk is prevalent, drought is often still not considered as a risk in  
446 Central, Northern and Eastern Europe. Here, we showed that drought hazard perception matched the observed or monitored  
447 drought hazard. In contrast, the occurrence of drought impacts does not always follow the pattern of hazard severity, and  
448 therefore requires assessment of drought beyond just the hazard. A relationship between national drought awareness and  
449 drought management strategies could not be established. Although a strong variability of drought risk management planning  
450 across the continent was evident, a common European strategy does not exist. As shown here, current national drought risk  
451 management practices range from a fundamental lack of legislation to country wide operational drought risk management  
452 plans. Future research might expand this survey to further explore and highlight potential benefits of a European drought  
453 directive. To foster national resilience to drought, drought management should be included in national legislation.

454 The key message of this study is that macro-governmental guidance by the EU is believed to be beneficial for national and  
455 international drought risk management. Such guidance should set a general framework which allows for regional flexibility of  
456 management strategies. To foster this kind of progress, sector specific databases on drought impacts, such as the EDII, are  
457 required to show and quantify the varied impacts of past droughts and increase public awareness in order to encourage political  
458 action. Going a step further, such information should be hosted by (inter-)national drought risk monitoring systems presenting  
459 sector specific drought risk.

460 As the first major steps towards a more unified drought risk management in Europe, we recommend:

- 461 1- The inclusion of a clear definition of drought in the Water Framework Directive, considering different types of  
462 drought, as well as their spatial and temporal occurrence,
- 463 2- The development of impact-driven, regional and sector-specific guidance on drought indices, and
- 464 3- The formation of an inter- and transdisciplinary collaborative EU-working group focusing on drought risk  
465 management and estimation of the potential benefits and downsides of a European Drought Directive.

#### 466 **Data availability**

467 Applied data can be accessed via <http://doi.org/10.34730/ae96ed78875c4caa9ee5c25c2e2f711a>.

#### 468 **Competing interests**

469 Some authors are members of the editorial board of NHSS. The peer-review process was guided by an independent editor,  
470 and the authors have no other competing interests to declare.

#### 471 **Disclaimer**

472 This study is based on the NUTS-level classification system (and GIS data) as provided by Eurostat for the year 2016.  
473 Accordingly, the pan-European survey followed the NUTS regions and countries as given.

#### 474 **Acknowledgments**

475 This research was carried out within the interdisciplinary research project DRiER. The project is supported by the  
476 Wassernetzwerk Baden-Württemberg (Water Research Network), which is funded by the Ministerium für Wissenschaft,  
477 Forschung und Kunst Baden-Württemberg (Ministry of Science, Research and the Arts of the State Baden-Württemberg)  
478 (grant no. AZ. 7532.21/2.1.6) and Maa- ja vesitekniikan tuki ry foundation. DW acknowledges her support as part of the  
479 NERC-funded Groundwater Drought Initiative (NE/R004994/1). LJB was supported by the Natural Environment Research  
480 Council (NE/R016429/1) as part of the UK-SCAPE programme delivering National Capability. MT, LB and JJ contributions  
481 have been supported by SustES—Adaptation strategies for sustainable ecosystem services and food security under adverse  
482 environmental conditions (CZ.02.1.01/0.0/0.0/16\_019/0000797). Special thanks go to the IAHS-Panta Rhei working group  
483 “Drought in the Anthropocene” who originally designed the study.

484 **References**

- 485 Aldunce, P., Araya, D., Sapiain, R., Ramos, I., Lillo, G., Urquiza, A., and Garreaud, R: Local perception of drought impacts  
486 in a changing climate: The mega-drought in central Chile. *Sustainability*, 9(11), 205, 2017.
- 487 Bachmair, S., Stahl, K., Collins, K., Hannaford, J., Acreman, M., Svoboda, M., Knutson, C., Smith, K.H., Wall, N., Fuchs, B.  
488 and Crossman, N.D.: Drought indicators revisited: the need for a wider consideration of environment and society. *Wiley*  
489 *Interdisciplinary Reviews: Water*, 3(4), pp.516-536, 2016.
- 490 Bakke, S. J., Ionita, M., and Tallaksen, L. M.: The 2018 northern European hydrological drought and its drivers in a historical  
491 perspective. *Hydrology and Earth System Sciences*, 24(11), pp.5621-5653. <https://doi.org/10.5194/hess-24-5621-2020>, 2020.
- 492 Barker, L. J., Hannaford, J., Parry, S., Smith, K. A., Tanguy, M., and Prudhomme, C.: Historic hydrological droughts 1891–  
493 2015: systematic characterisation for a diverse set of catchments across the UK, *Hydrol. Earth Syst. Sci.*, 23, 4583–4602,  
494 <https://doi.org/10.5194/hess-23-4583-2019>, 2019.
- 495 Baruth, B., Bassu, S., Bussay, A., Ceglar, A., Cerrani, I., Chemin, Y., De Palma, P., Fumagalli, D., Lecerf, R., Manfron, G.,  
496 Nisini Scacchiafichi, L., Panarello, L., Ronchetti, G., Seguini, L., Toreti, A., Van Den Berg, M., Zajac, Z., Zucchini, A. and  
497 Biavetti, I.: Crop monitoring in Europe, *JRC MARS Bulletin Vol. 28 No 9*, Van Den Berg, M., Baruth, B., Niemeyer, S., Van  
498 Der Velde, M. and Manfron, G. editor(s), Publications Office of the European Union, Luxembourg, 2020, JRC120746, 2020.
- 499 Beillouin, D., Schauburger, B., Bastos, A., Ciais, P., and Makowski, D.: Impact of extreme weather conditions on European  
500 crop production in 2018. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 375(1810),  
501 20190510. <https://doi.org/10.1098/rstb.2019.0510>, 2020.
- 502 Blauhut, V.: The triple complexity of drought risk analysis and its visualisation via mapping: a review across scales and sectors.  
503 *Earth-Science Reviews*, 210(September), 103345. <https://doi.org/10.1016/j.earscirev.2020.103345>, 2020.
- 504 Blauhut, V., Stahl, K., and Kohn, I.: The dynamics of vulnerability to drought from an impact perspective. In H. Andreu, J.,  
505 Solera, A., Paredes-Arquiola, J., Haro-Monteagudo, D., and van Lanen (Ed.), *Drought: Research and Science-Policy*  
506 *Interfacing*, e (p. 349–354.). London: CRC Press, 2015.
- 507 Blauhut, V., Stahl, K., Stagge, J. H., Tallaksen, L. M., Stefano, L. De, and Vogt, J.: Estimating drought risk across Europe  
508 from reported drought impacts, drought indices, and vulnerability factors. *Hydrology and Earth System Sciences*, 20(7), 2779–  
509 2800. <https://doi.org/10.5194/hess-20-2779-2016>, 2016.
- 510 Boergens, E., Güntner, A., Dobslaw, H., and Dahle, C.: Quantifying the Central European Droughts in 2018 and 2019 with  
511 GRACE Follow-On. *Geophysical Research Letters*, 47(14). <https://doi.org/10.1029/2020GL087285>, 2020.
- 512 Büntgen, U., Urban, O., Krusic, P.J. et al.: Recent European drought extremes beyond Common Era background variability.  
513 *Nat. Geosci.* 14, 190–196 (2021). <https://doi.org/10.1038/s41561-021-00698-0>, 2021.
- 514 Buras, A., Rammig, A., and S. Zang, C.: Quantifying impacts of the 2018 drought on European ecosystems in comparison to  
515 2003. *Biogeosciences*, 17(6), 1655–1672. <https://doi.org/10.5194/bg-17-1655-2020>, 2020.



516 Caillet, V.; Mager, U., and Zoth, P.: Herausforderung und Realität eines Dürremanagements in Baden-Württemberg.  
517 Verwaltungsblätter Baden-Württemberg (VBIBW) 2019, S. 133 ff.

518 Cammalleri, C., Naumann, G., Mentaschi, L., Formetta, G., Forzieri, G., Gosling, S., Bisselink, B., Roo, De A., and Feyen, L.  
519 (2020). Global warming and drought impacts in the EU. Publications Office of the European Union.  
520 <https://doi.org/10.2760/597045>, 2019.

521 Cammalleri, C., Micale, F., and Vogt, J.: A novel soil moisture-based drought severity index (DSI) combining water deficit  
522 magnitude and frequency. *Hydrological Processes*, 30(2), 289–301. <https://doi.org/10.1002/hyp.10578>

523 Cheval S, Busuioc A, Dumitrescu A, and Birsan M-V.: Spatiotemporal variability of meteorological drought in Romania using  
524 the standardized precipitation index (SPI). *Climate Research* 60. 235-248, 2014.

525 Cindrić Kalin, K., Prtenjak, M.T., Herceg-Bulić, I., Mihajlović, D. and Pasarić, Z.: Analysis of the extraordinary 2011/2012  
526 drought in Croatia. *Theoretical and applied climatology*, 123(3-4), pp.503-522, 2016.

527 Dai, A.: Increasing drought under global warming in observations and models. *Nat. Clim. Chang.*, 3, 52–58, 2013.

528 Di Baldassarre, G., Viglione, A., Carr, G., Kuil, L., Salinas, J. L., and Blöschl, G.: Socio-hydrology: conceptualising human-  
529 flood interactions, *Hydrol. Earth Syst. Sci.*, 17, 3295–3303, <https://doi.org/10.5194/hess-17-3295-2013>, 2013.

530 de Brito, M. M.: Compound and cascading drought impacts do not happen by chance: A proposal to quantify their relationships.  
531 *Science of the Total Environment*, 778(March). <https://doi.org/10.1016/j.scitotenv.2021.146236>, 2021.

532 de Niet, Julia; Finger, David D.C.; Bring, Arvid; Egilson, David; Gustafsson, David; and Kalantari, Zahra: Benefits of  
533 Combining Satellite-Derived Snow Cover Data and Discharge Data to Calibrate a Glaciated Catchment in Sub-Arctic Iceland,  
534 *Water* 12, no. 4: 975. <https://doi.org/10.3390/w12040975>, 2020.

535 Dessai, S., and Sims, C.: Public perception of drought and climate change in southeast England. *Environmental hazards*, 9(4),  
536 340-357, 2010.

537 DriDanube-Watch: Drought 2018 – DriDanube watch (Vol. 37). <https://drive.google.com/file/d/1m-qobw41JGauZdIANlS1NPaRe8SQ2Gp9/view>

538

539 EU 2007. 60/EC of the European Parliament and of the Council of 23 October 2007 on the assessment and management of  
540 flood risks. *Official Journal of the European Union L*, 288(27), p.2007, 2018.

541 European Commission: A Blueprint to Safeguard Europe’s Water Resources. COM (2012) 673 final. Brussels, 2012.

542 European Commission: Fitness Check of the Water Framework Directive and the Floods Directive, 1–184, 2019.

543 Falzoi, S., Gleeson, E., Lambkin, K., Zimmermann, J., Marwaha, R., O’hara, R., ... and Fratianni, S.: Analysis of the severe  
544 drought in Ireland in 2018. *Weather*, 74(Special Issue 11), 368–373. <https://doi.org/10.1002/wea.3587>, 2019.

545 Finger, D.C.: The value of satellite retrieved snow cover images to assess water resources and the theoretical hydropower  
546 potential in ungauged mountain catchments. *Jökull* 68, 47-66, 2018.

547 Finger D., Þórsson J., Pétursdóttir Þ., G. Halldórsson: Enhancing the resilience of water resources through land restoration in  
548 Rangárvellir, Iceland – an overview of the HydroResilience project, Extended abstract 10th European Conference on  
549 Ecological Restoration, Freising, Germany. SER Europe Knowledge Base ([www.ser.org/europe](http://www.ser.org/europe)), 5pp. ISSN2295-5704, 2016.

550 García-Herrera, R., Garrido-Perez, J. M., Barriopedro, D., Ordóñez, C., Vicente-Serrano, S. M., Nieto, R., ... Yiou, P. (2019).  
551 The European 2016/17 drought. *Journal of Climate*, 32(11), 3169–3187. <https://doi.org/10.1175/JCLI-D-18-0331.1>, 2019.

552 Hagenlocher, M., Meza, I., Anderson, C. C., Min, A., Renaud, F. G., Walz, Y., ... and Sebesvari, Z.: Drought vulnerability  
553 and risk assessments: state of the art, persistent gaps, and research agenda. *Environmental Research Letters*, 14(8), 83002.  
554 <https://doi.org/10.1088/1748-9326/ab225d>, 2019.

555 Hall, J. W., and Leng, G.: Can we calculate drought risk... and do we need to? *Wiley Interdisciplinary Reviews: Water*,  
556 (January), e1349. <https://doi.org/10.1002/wat2.1349>, 2019.

557 Hanger-Kopp, S., and Palka, M.: Exploring Drought Resilience Through a Drought Risk Management Lens in Austria, 115–  
558 138. <https://doi.org/10.1007/978-981-15-4320-3>, 2020.

559 Hari, V., Rakovec, O., Markonis, Y., Hanel, M., and Kumar, R.: Increased future occurrences of the exceptional 2018–2019  
560 Central European drought under global warming. *Scientific Reports*, 10(1). <https://doi.org/10.1038/s41598-020-68872-9>,  
561 2020.

562 Hänsel, S., Ustrul, Z., Łupikasza, E., and Skalak, P.: Assessing seasonal drought variations and trends over Central Europe.  
563 *Water Resources* 127, 53-57, 2019.

564 Helmert, J., Şensoy, Şorman A., Alvarado Montero, R., De Michele, C., De Rosnay, P., Dumont, M., Finger, D.C., Lange, M.,  
565 Picard, G., Potopová, V., Pullen, S., and Vikhamar-Schuler, D.: An Review of Snow Data Assimilation Methods for  
566 Hydrological, Land Surface, Meteorological and Climate Models: Results from a COST HarmoSnow Survey. *Geosciences*.  
567 8(12):489. <https://doi.org/10.3390/geosciences8120489>, 2018.

568 Hervás-Gámez, C., and Delgado-Ramos, F.: Drought management planning policy: From Europe to Spain. *Sustainability*  
569 (Switzerland), 11(7), 1–26. <https://doi.org/10.3390/su11071862>, 2019.

570 Hisdal H., Tallaksen L.M. (2003). Estimation of regional meteorological and hydrological drought characteristics: a case study  
571 for Denmark. *Journal of Hydrology* 281, 230–247. doi:10.1016/S0022-1694(03)00233-6, 2003.

572 Howarth, W.: Going with the flow: Integrated Water Resources Management, the EU Water Framework Directive and  
573 ecological flows, *LegalStudies*, 38, 298–319, <https://doi.org/10.1017/lst.2017.13>, 2018.

574 Ionita M, Tallaksen L. M., Kingston D. G., Stagge J. H., Laaha G, Van Lanen H. A. J., Scholz P., Chelcea S. M., and Haslinger  
575 K.: The European 2015 drought from a climatological perspective. *Hydrol. Earth Syst. Sci.* 21, 1397–1419, 2017.

576 IPCC: Climate Change 2014 Impact, Adaption, and Vulnerability Part A: Global and sectoral aspects. New York, USA.  
577 <https://doi.org/10.1017/CBO9781107415324.004>, 2014.

578 Jakubínský, J., Bláhová, M., Bartošová, L., Steinerová, K., Balek, J., Dížková, P., Semerádová, D., Alexandru, D., Bardarska,  
579 G., Bokal, S., Borojević, G., Bucur, A., Cindrić Kalin, K., Barbu Anda, C., Debre, B., Đorđević, M., Đurić, I., Mircea, B. F.,  
580 Gatarić, S., Gregorič, G., Hasenauer, S., Ivanov, M., Kiresi, A., Labudová, L., Turňa, M., Marinović, I., Marković, M.,  
581 Mateescu, E., Oblišar, G., Popescu, A., Srđević, Z., Savić-Šljivić, T., Supić, D., Sušnik, A., Pazin, N., Drljevic, M., Kuc, T.,  
582 Mitrovic, L., Micev, S., Wagner, W., Eitzinger, J., Daneu, V., Blauhut, V., Stahl, K. and Trnka, M. : Repository of Drought

583 Event Impacts Across the Danube Catchment Countries Between 1981 and 2016 Using Publicly Available Sources. *Acta*  
584 *Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 67(4): 925 – 938, 2019.

585 Jaagus, J., Aasa, A., Aniskevich, S., Boincean, B., Bojariu, R., Briede, A., Danilovich, I., Domínguez Castro, F., Dumitrescu,  
586 A., Labuda, M. and Labudová, L., Long-term changes in drought indices in eastern and central Europe. *International Journal*  
587 *of Climatology*, 2021.

588 Keesstra S., Nunes J., Novara A., Finger D., Avelar D., Kalantari Z. and A. Cerdà: The superior effect of nature based solutions  
589 in land management for enhancing ecosystem services, *Science of The Total Environment*, 610, 997-1009.doi:  
590 10.1016/j.scitotenv.2017.08.077, 2018.

591 Kingston, D. G., Stagge, J. H., Tallaksen, L. M., and Hannah, D. M.: European-Scale Drought: Understanding Connections  
592 between Atmospheric Circulation and Meteorological Drought Indices. *Journal of Climate*, 28(2), 505–516.  
593 <https://doi.org/10.1175/JCLI-D-14-00001.1>, 2015.

594 Kreibich, H., Blauhut, V., Aerts, J. C. J. H., Bouwer, L. M., Van Lanen, H. A. J., Mejia, A., ... Van Loon, A. F.: How to  
595 improve attribution of changes in drought and flood impacts. *Hydrological Sciences Journal*, 64(1), 1–18.  
596 <https://doi.org/10.1080/02626667.2018.1558367>, 2019.

597 Kreibich, H., Di Baldassarre, G., Vorogushyn, S., Aerts, J.C.J.H., Apel, H., Aronica, G.T., Arnbjerg-Nielsen, K., Bouwer,  
598 L.M., Bubeck, P., Caloiero, T., Chinh, D.T., Cortès, M., Gain, A.K., Giampá, V., Kuhlicke, C., Kundzewicz, Z.W., Llasat,  
599 M.C., Mård, J., Matczak, P., Mazzoleni, M., Molinari, D., Dung, N.V., Petrucci, O., Schröter, K., Slager, K., Thielen, A.H.,  
600 Ward, P.J. and Merz, B.: Adaptation to flood risk: Results of international paired flood event studies. *Earth's Future*, 5: 953-  
601 965. <https://doi.org/10.1002/2017EF000606>, 2017.

602 Laaha, G., Gauster, T., Tallaksen, L.M., Vidal, J.-P., Stahl, K., Prudhomme, C., Heudorfer, B., Vlnas, R., Ionita, M., Van  
603 Lanen, H.A.J., Adler, M.-J., Caillouet, L., Delus, C., Fendekova, M., Gailliez, S., Hannaford, J., Kingston, D., Van Loon, A.F.,  
604 McKee, T. B., Doesken, N. J., and Kleist, J.: The relationship of drought frequency and duration to time scales. In *Eight*  
605 *Conference on Applied Climatology I*. Journal Article, 1993.

606 Mediero, L., Osuch, M., Romanowicz, R., Sauquet, E., Stagge, J.H., and Wong, W.K.: The European 2015 drought from a  
607 hydrological perspective. *Hydrology and Earth System Sciences*, 21, 3001–3024. <https://doi.org/10.5194/hess-21-3001-2017>,  
608 2017.

609 Labudová, L., Ivaňáková, G., Trnka M., Bartošová, L.: Monitoring of drought impacts and the DriDanube project (Tromp  
610 Foundation Travel Award). *EMS Annual Meeting Abstracts*, Vol. 15, EMS2018-34-1, 2018.

611 Lloyd-Hughes, B.: The impracticality of a universal drought definition. *Theoretical and Applied Climatology*, 117, 607–611.  
612 <https://doi.org/10.1007/s00704-013-1025-7>, 2014.

613 Ministry for the Ecological Transition (MITECO). Instrucción Técnica para la Elaboración de los Planes Especiales de Sequía  
614 y la Definición del Sistema Global de Indicadores de Sequía Prolongada y de Escasez; Ministry for the Ecological Transition:  
615 Madrid, Spain; pp. 1–20, 2017.

616 Radeva K, Nikolova N, Gera M, Assessment of hydro-meteorological drought in the Danube Plain, Bulgaria. *Hrvatski*  
617 *geografski glasnik*, 80 (1), 7-25, 2018.

618 Rosner, B., Benedict, I., Heerwaarden, C. van, Weerts, A., Hazeleger, W., Bissolli, P., and Trachte, K.: The long heat wave  
619 and drought in Europe in 2018. *Bulletin of the American Meteorological Society*, (August), 2018–2020, 2019.

620 Salmoral, G., Ababio, B., and Holman, I. P.: Drought impacts, coping responses and adaptation in the UK outdoor livestock  
621 sector: Insights to increase drought resilience. *Land*, 9(6). <https://doi.org/10.3390/LAND9060202>, 2020.

622 Schuldt, B., Buras, A., Arend, M., Vitasse, Y., Beierkuhnlein, C., Damm, A., ... Kahmen, A.: A first assessment of the impact  
623 of the extreme 2018 summer drought on Central European forests. *Basic and Applied Ecology*, 45(June), 86–103.  
624 <https://doi.org/10.1016/j.baae.2020.04.003>, 2020.

625 Siebert, C., Blauhut, V., and Stahl, K.: Drought and its implications for the public water supply sector in Baden-Württemberg,  
626 Germany: impacts, management, perception. *WasserWirtschaft* 6/2021, 2021.

627 Spinoni, J., Barbosa, P., De Jager, A., McCormick, N., Naumann, G., Vogt, J.V., Magni, D., Masante, D. and Mazzeschi, M.:  
628 A new global database of meteorological drought events from 1951 to 2016. *Journal of Hydrology: Regional Studies*, 22,  
629 p.100593, 2019.

630 Spinoni J., Vogt J.V, Naumann G., Barbosa P., Dosio A.: Will drought events become more frequent and severe in Europe?  
631 *Int. J. Climatol.* 38, 1718–1736. doi: 10.1002/joc.5291

632 Spinoni, J., Naumann, G., Vogt, J. V., and Barbosa, P. (2015). The biggest drought events in Europe from 1950 to 2012.  
633 *Journal of Hydrology: Regional Studies*, 3, 509–524. <https://doi.org/10.1016/j.ejrh.2015.01.001>, 2018.

634 Stahl, K., Kohn, I., Blauhut, V., Urquijo, J., De Stefano, L., Acácio, V., ... Van Lanen, H. A. J.: Impacts of European drought  
635 events: Insights from an international database of text-based reports. *Natural Hazards and Earth System Sciences*, 16(3), 801–  
636 819. <https://doi.org/10.5194/nhess-16-801-2016>, 2016.

637 Stein, U., Özerol, G., Tröltzsch, J., Landgrebe, R., Szendrenyi, A., and Vidaurre, R.: European Drought and Water Scarcity  
638 Policies. In H. Bressers, N. Bressers, and C. Larrue (Eds.), *Governance for Drought Resilience: Land and Water Drought*  
639 *Management in Europe* (pp. 17–43). inbook, Cham: Springer International Publishing. [https://doi.org/10.1007/978-3-319-](https://doi.org/10.1007/978-3-319-29671-5_2)  
640 [29671-5\\_2](https://doi.org/10.1007/978-3-319-29671-5_2), 2016.

641 Teutschbein, C., Lundkvist, E., Albrecht, F., and Blicharska, M.: A culture of proactive drought management? Unraveling the  
642 perception and management of droughts in Swedish municipalities. In *Geophysical Research Abstracts* (Vol. 21), 2019.

643 Thompson, R. L., Broquet, G., Gerbig, C., Koch, T., Lang, M., Monteil, G., ... Rödenbeck, C.: Changes in net ecosystem  
644 exchange over Europe during the 2018 drought based on atmospheric observations. *Philosophical Transactions of the Royal*  
645 *Society of London. Series B, Biological Sciences*, 375(1810), 20190512. <https://doi.org/10.1098/rstb.2019.0512>, 2020.

646 Turner, S., Barker, L. J., Hannaford, J., Muchan, K., Parry, S., and Sefton, C.: The 2018/2019 drought in the UK: a  
647 hydrological appraisal. *Weather*, 1–6. <https://doi.org/10.1002/wea.4003>, 2021.

648 Trnka, M., Hlavinka, P., Možný, M., Semerádová, D., Štěpánek, P., Balek, J., ... Žalud, Z.: Czech Drought Monitor System  
649 for monitoring and forecasting agricultural drought and drought impacts. *International Journal of Climatology*, (May 2018),  
650 1–18. <https://doi.org/10.1002/joc.6557>, 2020.

651 Trambly, Y.; Koutroulis, A.; Samaniego, L.; Vicente-Serrano, S.M.; Volaire, F.; Boone, A.; Le Page, M.; Llasat, M.C.;  
652 Albergel, C.; Burak, S.; et al.: Challenges for Drought Assessment in the Mediterranean Region under Future Climate  
653 Scenarios. *Earth-Science Reviews*, 210, 103348, doi:10.1016/j.earscirev.2020.103348, 2020

654 Trnka M, Hayes M, Jurečka F, Bartošová L et al.: Priority questions in multidisciplinary drought research. *Clim Res* 75:241-  
655 260. <https://doi.org/10.3354/cr01509>, 2018.

656 Trnka M., Kyselý J., Možný M., Dubrovský M.: Changes in Central-European soil-moisture availability and circulation  
657 patterns in 1881–2005. *Int J Climatol* 29:655–672. <https://doi.org/10.1002/joc.1703>, 2009.

658 UNDRR- United Nations Office for Disaster Risk Reduction: GAR Special Report on Drought 2021. Geneva, 2021.

659 UNCCD: Drought Resilience, Adaptation and Management Policy Framework: Supporting Technical Guidelines, (July), 17,  
660 2019.

661 UNDRR- United Nations Office for Disaster Risk Reduction.: Global Assessment Report on Disaster Risk Reduction. Geneva,  
662 Switzerland: United Nations Office for Disaster Risk Reduction, 2019.

663 Urquijo, J., and De Stefano, L.: Perception of drought and local responses by farmers: a perspective from the Jucar River  
664 Basin, Spain. *Water resources management*, 30(2), 577-591, 2016.

665 Urquijo, J., Pereira, D., Dias, S., and De Stefano, L.: A methodology to assess drought management as applied to six European  
666 case studies. *International Journal of Water Resources Development*, 627(April), 1–24.  
667 <https://doi.org/10.1080/07900627.2016.1174106>, 2016.

668 Van Lanen, H. A. J., Laaha, G., Kingston, D. G., Gauster, T., Ionita, M., Vidal, J. P., ... and Van Loon, A. F.: Hydrology  
669 needed to manage droughts: the 2015 European case. *Hydrological Processes*, 30(17), 3097–3104.  
670 <https://doi.org/10.1002/hyp.10838>, 2016.

671

## 672 **Figures and tables**

673 **Table 1: Drought indices and their associated drought classes. SPI, fAPAR, SM and LFI are, respectively, Standardised Precipitation**  
674 **Index, fraction of Accumulated Photosynthetically Active Radiation, Soil Moisture and Low Flow Index.**


Indices	No drought	Moderate drought	Severe drought	Extreme drought
SPI, fAPAR, SM	> -1	-1 to -1.5	-1.5 to -2	< -2
LFI	0-0.25	0.25-0.5	0.5-0.75	0.75-1

675

676



678 Fig. 1. Drought hazard conditions for 2018 and 2019 across the European continent according to the European Drought  
 679 Observatory indicator factsheets (<https://edo.jrc.ec.europa.eu/edov2/php/index.php?id=1101>). Data are presented as the  
 680 proportion of the country's total area under severe (or extreme) drought hazard conditions. Standardised Precipitation Index  
 681 (SPI) is shown for 3, 6, 9 and 12 month accumulation periods for August (AUG), September (SEP), October (OCT) and December  
 682 (DEC). SM is Soil Moisture Anomaly/ Index, LFI is Low Flow Index and fAPAR is fraction of Photosynthetically Active  
 683 Radiation, all presented as for the day of the year in the corresponding colouring: 2018 as area and 2019 as lines.




685 Fig. 2. Water usage across Europe: a) number of survey participants by country, b) most important water resource by country, and  
 686 c) second most important water resource by country. Results are based on a pan-European survey designed by the International  
 687 Association of Hydrological Science (IAHS)- Panta-Rhei “Drought in the Anthropocene” working group and conducted in 28  
 688 countries.





689

690 Fig. 3. Major categories of drought indices used across Europe as a fraction of total replies per country (number of replies in  
 691 parentheses, total replies of 536). The mean index in each category (meteorology, soil moisture, hydrology and vegetation) is weighted  
 692 by the number of participants. Countries on the y-axis are sorted according to their mean index value, i.e., the highest for Czech  
 693 Republic, lowest for Slovakia. The category Others (n=37) comprise countries with less than 10 replies, namely Austria (9), Italy (8),  
 694 Belgium (6), Latvia (6), Iceland (4), Denmark (3) and Romania (1). Replies from Sweden and Poland are not considered here as  
 695 indices were not rated in these countries. Note that participants have different roles in their countries and thus might judge drought  
 696 indices differently. Results are based on a pan-European survey designed by the International Association of Hydrological Science  
 697 (IAHS) - Panta-Rhei “Drought in the Anthropocene” working group and conducted in 28 countries.



698

699


700

701

702

703

**Fig. 4.** Perception of climate change effect on drought management in Europe shown as percentage of participants responding to question: (a) whether future climate change will may affect water resources; (b) how droughts may change in future; (c) will drinking water providers in the future have to distribute water to fewer consumers due to shortages, e.g. 'rota cuts'. Results are based on a pan-European survey designed by the International Association of Hydrological Science (IAHS)- Panta-Rhei "Drought in the Anthropocene" working group and conducted in 28 countries.



704

705


706

707

708


709

*Fig. 5. Perception of drought risk management across Europe shown as percentage of participants in pie charts: a) distribution of drought risk management plans and emergency action plans by country; b) reasons for an absence of drought risk management by country and totals of selected reasons. Results are based on a pan-European survey designed by the International Association of Hydrological Science (IAHS)- Panta-Rhei “Drought in the Anthropocene” working group and conducted in 28 countries.*



710  
711  
712  
713  
714

**Fig. 6. Median perception of drought severity and impacts in 2018 and 2019 across Europe. Sweden participated only in 2018. Results are based on a pan-European survey designed by the International Association of Hydrological Science (IAHS) - Panta-Rhei “Drought in the Anthropocene” working group and conducted in 28 countries.**



715  
716  
717  
718

**Fig. 7. National representatives' joint opinion on a) the actual state of drought management in their country, b) the existence of a country-wide drought management plan, c) the existence of national recommendations for actions in order to minimise drought risk, and d) the benefit of an EU- drought directive for their country?, ordered by score (Very good=3; Not good = 0).**