

Brief communication:

Western Europe flood in 2021: mapping agriculture flood exposure from SAR

Kang He¹, Qing Yang², Xinyi Shen¹, Emmanouil N. Anagnostou¹

5

¹Department of Civil and Environmental Engineering, University of Connecticut, Storrs, CT 06269, USA

²College of Civil Engineering and Architecture, Guangxi University, Nanning, Guangxi, 530004, China

Correspondence to: Xinyi Shen (xinyi.shen@uconn.edu)

Abstract. In this communication, we present the exposure of agriculture lands to the flooding caused by extreme precipitation in western Europe from 12th to 15th of July 2021. Overlaying the flood inundation maps derived from the near-real-time Radar-Produced Inundation Diary (RAPID) system on the CORINE land cover map we estimate a ~~2470~~1920.26 km² area affected by the flooding, with ~~5764.4~~3635.9% representing agricultural land. Among the inundated agricultural land, ~~3333.7~~3635.9% of the area is pastures while ~~3333.7~~3333.7% is arable land. Most agricultural flood exposure is found in ~~south-eastern~~western France (~~~1680~~~1680 km²) along Rhône River, southern Netherlands along the Meuse River and western Germany along Rhine River and the coastal area of Marseille and Montpellier.

10
15

1. Introduction

The heavy precipitation between 12 and 15 July 2021 led to catastrophic floods in western European countries, including France, western Germany, Netherlands, Belgium, and Luxembourg. The flooding caused widespread power outages, infrastructure and crops damages in the affected areas. It is estimated that the loss from the flooding is up to €3 billion [Reinsurance News, 2021]. In addition, 46 people were confirmed dead in North Rhine-Westphalia state in Germany and in the neighbouring state of Rhineland-Palatinate 110 fatalities were confirmed. At least 20 people died following catastrophic flooding in Belgium. The Netherlands, Luxembourg and Switzerland are also affected. more than 200 people were killed, mostly in Germany and Belgium Thousands of people had been evacuated from their homes [CNN, 2021; FloodList, 2021]. In the same period, intensive floods occurred in China and the United States. Researchers highlighted that this is an effect of climate change and concluded that the frequency and intensity of such events will increase in a rapidly warming climate [World weather attribution, 2021].

20
25

Besides life loss, the flooding in western Europe have also taken a heavy toll on the agricultural sector according to European farmers' association COPA-COGECA. The European Union's crop monitoring unit stated that the exceptionally high rainfall and severe floods would reduce the grain quality in the affected countries [Successful farming, 2021] and had "effectively eliminated" any hope of a successful harvest in these areas [Euractiv, 2021]. Examples of crop damages include crops of grain, rapeseed and flax which have been washed away in Wallonia, Belgium and flood-affected fruit trees along the Meuse River

30

[Eurofruit, 2021]. In widespread crop loss scenarios like this one, damage assessment is an essential part of flood risk management and flood mitigation, which is also the basis of financial appraisals in the insurance sector [Tapia-Silva et al., 2011]. Even though the impact on the agriculture sector is expected to be severe, the magnitude of the damage is yet to be determined [Agence europe, 2021]. Therefore, it is important to have a quick assessment of the agriculture land exposure to flooding, which will inform crop loss estimates, especially for countries where agriculture plays an important role in the national economy, e.g., France and Germany. Near-real-time (NRT) flood mapping capability from satellite observations is vital to facilitate rapid assessment of flood loss and damage [Shen et al., 2019a].

In this brief communication, we use NRT inundation extents from the near-real-time RADar-Produced Inundation Diary (RAPID) system combined with CORINE land cover data to depict the flood-affected areas in western Europe, and particularly the agriculture land.

2. Methodology

We focus this communication on western Europe, which is mostly affected by the July 12-15 heavy precipitation event. The area extends from 1.5° E to 11.6° E, and 42.9° N to 53.1° N, and encompasses the Netherlands, Belgium, Luxembourg, Switzerland and portions of Germany, France, and Italy. This region is dominated by marine climate with abundant moisture supplemented by Atlantic Ocean. The weather is therefore moist and mild in winter, and moist and cool in summer.

We extract half hourly precipitation data of the event from the Integrated Multi-satellitE Retrievals for Global Precipitation Mission (IMERG) ~~Late~~ Final Precipitation L3 V06 product with 0.1-degree spatial resolution [https://disc.gsfc.nasa.gov/datasets/GPM_3IMERGHH_06/summary, Huffman et al., 2019]. ~~IMERG Late Run~~ The IMERG half-hourly Final Run product combines the multi-satellite data for the month with GPCC gauge analysis and thus provides the research-level products for precipitation estimation. ~~is computed about 14 hours after observation time, which integrates more data from sensors aboard on satellites to improve the accuracy.~~ We used IMERG data to calculate the maximum hourly precipitation rate and precipitation accumulation between 12 and 15 of July for each grid.

We generate inundation extents in NRT using the RAPID system and archive these maps on Amazon Web Services (AWS) [available at https://rapid-nrt-flood-maps.s3.amazonaws.com/index.html#Global_Flood_Event/Europe_Flood_2021/]. RAPID is a fully automated system delineating NRT inundation extents from high resolution (10 m) synthetic aperture radar (SAR) imagery [Yang et al., 2021; Shen et al., 2019a; Shen et al., 2019b]. Specifically, the RAPID system first segments water from non-water pixels by optimizing the threshold and probability density function (PDF) of the water class. Then, it runs a morphology-based procedure to reject false water bodies using rule sets defined at the body level instead of the pixel level. The morphological processing includes two sub modules, water source tracing (WST), and improved changed detection (ICD). WST traces water pixels from known water sources (e.g, rivers, lakes) indicated by a land use map. ICD rejects any water body that is disconnected from a known water source and does not have significantly increased area compared to the dry time. For dry reference, we use information from ground observation and satellite precipitation to determine non-flood period.

and image cover that period is select as dry reference. The RAPID requires approximately five dry reference images for each SAR image sensed on the flood day to reduce the error caused by noise-like speckle. In the third and last processing steps, RAPID uses multi-threshold compensation and machine learning to further reduce the speckles and strong scatter-caused false negatives. Figure 1 (a) presents an example of inundation delineation by RAPID system in Louhans, France. The CORINE land cover map, shown as Figure 1 (b). The corresponding SAR images sensed on July 16th, 2021 (Figure 1 (c), flooding period) and images sensed on July 10th, 2021 (Figure 1 (d), dry reference). To rule out false positives caused by glaciers and snow, we threshold the Height Above Nearest Drainage (HAND) data to mask out permafrost areas in Alps. The HAND used in this study is obtained from the Multi-Error-Removed Improved-Terrain (MERIT) Hydro Dataset [Yamazaki et al., 2019; Nobre et al., 2011]. Pixels over the Alps where HAND values are greater than 20 meters are removed from the inundated pixels. The threshold is determined by exploring the distribution of HAND for glaciers and perpetual snow recorded in CORINE land cover data and is large enough to avoid the removal of any true positives.

We obtain the latest land cover map over western Europe from Coordination of information on the environment (CORINE) Land Cover (CLC) inventory data [available at <https://land.copernicus.eu/pan-european/corine-land-cover/clc2018>]. CLC uses a Minimum Mapping Unit (MMU) of 25 hectares (ha) and a minimum width of 100 meter for linear elements. The standard CLC nomenclature includes 44 land cover classes, grouped in a three-level hierarchy. Five main categories used in this study are "artificial surfaces", "agricultural areas", "forest and semi-natural areas", "wetlands" and "water bodies". The detailed description of CORINE program and its nomenclature can be found in <https://www.eea.europa.eu/publications/COR0-part1>.

3. Results

The spatial pattern of the ~~maximum hourly precipitation and~~ accumulated precipitation from the July 12-15 heavy precipitation event are shown in Figure ~~1-2 (a) and (b)~~. Heavy precipitation (peak rate > 20 mm/hr) is observed in western Germany, ~~south~~north-eastern France, norther Luxembourg, south-western Netherlands, western Switzerland, and western Italy. The most ~~intense-intensive~~ precipitation (peak rate > 50 mm/hr) is found in ~~south France~~western Germany, as well as western Switzerland and Italy over the Alps. Heavier than ~~200-150~~ mm accumulated precipitation is found in eastern France (Châtel-de-Joux, Le Fied), north-eastern France (Plainfaing, Villers-la-Chèvre), mid-eastern Luxembourg (Echternach and Mersch), southern western Belgium (Liège), southern Netherlands (Limburg), western Germany (North Rhine-Westphalia, Rhineland-Palatinate), Switzerland and Italy, which represent an equivalent of two-month precipitation accumulation in these areas. Furthermore, accumulated precipitation is shown to exceed ~~250-200~~ mm in some parts of the region (e.g., western Switzerland ~~and Italy~~, south-north-eastern France, western Germany).

Figure ~~2-2 (b)~~ shows the inundation extents over western Europe. The RAPID inundation map shows high consistency with the precipitation map. The total inundated area determined from RAPID inundation map is around ~~2470-1920.26~~ km². We find extensive inundated areas in ~~south-eastern France, especially the coastal area, including Marseille and Montpellier. The the~~

upstream region of Rhône River where more than 120 mm precipitation fell in 72 -hours. The south-eastern France, especially the coastal area, exhibits extensive flood inundation as well, though the accumulated precipitation in these areas is around 10 mm. The flooded areas in south-eastern France, shown as Figure 3 (a), are mostly arable land. These areas exhibit clearly dampened backscattering from the dry date (Figure 3 (c), June 22) to the flood date (Figure 3 (b), July 16) while their backscattering on the flood date falls into the water category. These rainfed arable lands might be flooded by overflow from the Rhône River or the raised water table. The total inundated area over France is approximately ~~1680~~ 1318.28 km². In Germany, the main inundated area is found in the west which is caused by the intensive precipitation (120 mm), along the Rhine River (about ~~162~~ 162.02 km²). In the northern Netherlands where more than 100 mm precipitation is observed, regions near Markermeer and IJsselmeer, and regions around Hollands Diep and Meuse River are largely affected by the flood, which represents a total area of ~~245~~ 140.7 km². We compared the RAPID inundation maps and Landsat-based flood inundation maps (FIMs) for North Netherlands, presented as Figure 4 (a) and (b) respectively. The RAPID and Landsat based FIMs shows high consistency on the flooded areas according to the result of quantitatively -comparison, with precision, recall, F-1 score and Cohen kappa metrics are 0.8816, 0.8439, 0.8624, 0.8571, respectively. In Belgium and Luxembourg, the inundated areas are ~~116~~ 116.3 km² and ~~2~~ 1.79 km², mostly along Meuse River and Sauer River, respectively. In western Italy, an area of around ~~135~~ 50.38 km² along the Po River is affected by flooding. The flash floods in Switzerland also cause a ~~131~~ 130.79 km² inundation.

Figure ~~3-5~~ (a) shows the land use fraction in the inundated areas. Among them, ~~24~~ 24.2% (~~597~~ 463.94 km²) of the land is forested/semi-natural areas. For wetlands and artificial surfaces, the fractions are ~~11~~ 5.7% (~~269~~ 190.71 km²) and ~~85~~ 8% (~~192~~ 110.49 km²), respectively. The majority, nearly ~~57~~ 64.4% (~~1412~~ 1236.12 km²) of the flood inundated area is from agricultural land. Over inundated agricultural areas as Figure ~~3-5~~ (b) shows, ~~36~~ 35.9% (~~513~~ 443.64 km²) is pastures, ~~33~~ 33.6% (~~463~~ 416.07 km²) is arable land (including non-irrigated arable ~~land (382 km²) and land and rice fields, 339.09 km² and 76.98 km², respectively (81 km²)) and ~~24~~ 23.3% (~~342~~ 288.59 km²) is heterogeneous agricultural areas, which is the sum of complex cultivation patterns (~~272~~ 247.66 km²) and land principally occupied by agriculture, with significant areas of natural vegetation (~~70~~ 40.93 km²). The remaining ~~77~~ 7.1% (~~93~~ 87.82 km²) is permanent crops consisting of vineyards (~~71~~ 67.12 km²), fruit trees and berry plantations (~~21~~ 20.19 km²) and olive groves (~~1~~ 0.51 km²).~~

~~Figure 4 (a) shows inundated area of land use grouped by countries over western Europe.~~ Specifically, in France, ~~1085~~ 974.08 km² of agricultural land cover is affected by the flood. Among those inundated agricultural areas in France (~~Figure 4 (b)), 363~~ 317.5 km², ~~360~~ 337.3 km², ~~271~~ 233 km² and ~~91~~ 86.29 km² are pastures, arable land, heterogeneous agricultural areas, and permanent crops, respectively. Especially, the non-irrigated arable land in France is severely affected, the area is up to ~~283~~ 263.08 km² which is larger than the sum of inundated non-irrigated arable land in other countries. Besides, the rice fields and vineyards in France are also hit by flood. More than ~~70~~ 74 km² of rice fields and vineyards, mainly in the coastal areas, are inundated. In the Netherlands, ~~135~~ 98.97 km² of agricultural land is inundated, mostly are pastures (~~74~~ 50.41 km²), followed by heterogeneous agricultural areas (~~36~~ 28.29 km²). The inundated area of arable land (mostly is non-irrigated arable land) in Netherlands is ~~25~~ 20.14 km², while only ~~0.2~~ 13 km² of permanent crops (mainly fruit trees and berry plantations) are affected

130 by flood. In Germany, 88.33 km² of agricultural land is inundated with 59.32 km² and 25.06 km² of these areas being pastures
and non-irrigated arable land. The inundation over heterogeneous agricultural areas and permanent crops (including vineyards,
fruit trees and berry plantation) in Germany are estimated at 3.13 km² and 0.82 km², respectively. The total inundated areas in
Belgium and Italy are ~~both around 46-116.3 km² and 50.38 km², respectively.~~ In Belgium, the inundated areas of heterogeneous
135 agricultural land, pastures, and arable land were ~~19-20.21 km², 14-12.58 km² and 13.58 km², respectively~~, while nearly no
permanent crop is affected by flood. In Italy, most inundation among agricultural areas is arable land (~~30-12.22 km² of non-~~
irrigated arable land and ~~5-2.81 km² of rice field~~) and to a secondary effect heterogeneous agricultural area (~~9-1.42 km²~~). Only
~~0.2 km² of pastures in Italy are inundated while 0.2-0.6 km² of permanent crops (vineyards) are affected by flood.~~ In
Switzerland, the inundated areas of non-irrigated arable land, pastures and heterogeneous agricultural areas are ~~5-4.68 km²,~~
~~3.35 km² and 2.08 km², respectively.~~ ~~0.5-4.7 km² of permanent crops, mainly fruit trees and berry plantations, is also found to~~
140 be affected by flood in Switzerland. No permanent crop is inundated in Luxembourg, the total inundated area in Luxembourg
is ~~+1.79 km², with 0.4-4.2 km², 0.3-3.3 km² and 0.3-2.8 km² of them being heterogeneous agricultural areas, non-irrigated arable~~
land and pastures, respectively.

4. Closing remarks

~~The July 12-15 unprecedented precipitation and the associated catastrophic flood heavily impacts the western Europe with
145 more than 200 deaths and an estimated €3 billion of economic loss from infrastructure damages. The unprecedent precipitation
heavily damaged the western Europe with catastrophic flooding, causing damage to However, the impact that the flooding
across western Europe has on agriculture which is yet minimally quantified. In this communication, we analyze the inundated
area of agricultural land by overlaying the inundation extent derived from RAPID system with CORINE land cover data. The
results indicate that the total inundated area over western Europe is about ~~2470-1920.26 km²~~, of which ~~1680-1318.28 km²~~ is in
150 France. Around ~~5764.4%~~ of the flooded area is agricultural land. Because of the wide impact, we expect that the agricultural
productivity in western Europe will be significantly reduced. ~~The mid July when the extreme flood happened is the critical
growing season for crops like corn in Belgium, France, Luxembourg and Netherlands, and also the harvest season for wheat
in Belgium, France and Germany [Foreign Agricultural Service]. The oxygen supply is greatly reduced when a corn crop is
submerged in water, which greatly reduces or even stops critical plant functions such as nutrient and water uptake [Lauer
155 2008]. The quality and production of these crops would be severely damaged.~~ Besides the direct damage to livestock and
crops, the soil erosion and sedimentation due to the flood cause significant part of agricultural land be washed away or become
less fertile [Mst et al., 2019; Morris and Brewin, 2014,18]. In addition, extra costs are needed for pastures and cultivable land
to reconstruct and recover.~~

The limitation of this study is primarily inherited from the data sources. The RAPID system in Europe is triggered by IMERG
160 precipitation data, which is a satellite-based precipitation product found to systematically underestimate precipitation in
complex terrain areas, such as Alps [Navarro et al., 2019].

With the increasing flood observing capability brought by modern satellite constellations (for example, ICEYE [Ignatenko et al., 2020]), future directions of this study will include combining the NRT RAPID inundation estimates with developed flood models, crop data and other essential data (soil salinity, crop sensitivity, etc.) to predict flood-damaged cropland areas [Lazin et al., 2021] and associated socioeconomic impact [Gould et al., 2020].

Author contribution: KH: formal analysis, writing – original draft and editing. QY: software, formal analysis, data curation. XS and EA: conceptualization, project administration, writing – review and editing.

Competing interests: The authors declare that they have no conflict of interest.

Acknowledgements: This research was supported by National Science Foundation HDR award entitled “Collaborative Research: Near term forecast of Global Plant Distribution Community Structure, and Ecosystem Function.” [Kang He received the support of China Scholarship Council for four years' Ph.D. study in University of Connecticut \(under grant agreement no. 201906320068\).](#)

References

- 175 [Agence europe: Impact of flood disaster on EU agriculture is yet to be determined,
https://agenceurope.eu/en/bulletin/article/12766/19 , last access: 21 July 2021.](#)
- [CNN: Germany's deadly floods were up to 9 times more likely because of climate change, study estimates,
https://www.cnn.com/2021/08/23/europe/germany-floods-belgium-climate-change-intl/index.html , last access: 24 August 2021.](#)
- 180 [Euractiv:
https://www.euractiv.com/section/agriculture-food/news/eu-farmers-warn-harvest-will-fail-after-floods-plea-for-aid/, last access: 20 July 2021.](#)
- [Eurofruit: EU farmers warn harvest will fail after floods, plea for aid,
http://www.fruitnet.com/eurofruit/article/185840/europes-flood-damage-begins-to-emerge , last access: 22 July 2021.](#)
- [FloodList: Europe, https://floodlist.com/europe, last access: 16 July 2021.](#)
- 185 [Foreign Agricultural Service: Crop Calendars for Europe, https://ipad.fas.usda.gov/rssiws/al/crop_calendar/europe.aspx, last access: 1 March 2022](#)
- [Gould, I. J., Wright, I., Collison, M., Ruto, E., Bosworth, G., & Pearson, S. The impact of coastal flooding on agriculture: A case-study of Lincolnshire, United Kingdom. *Land Degradation & Development*, 31\(12\), 1545-1559, 2020.](#)

- 190 [Huffman, G.J., Stocker E.F., Bolvin D.T., Nelkin E.J., and Tan J.: GPM IMERG Late Precipitation L3 1 day 0.1 degree x 0.1 175 degree V06, Edited by Andrey Savtchenko, Greenbelt, MD, Goddard Earth Sciences Data and Information Services Center \(GES DISC\), 10.5067/GPM/IMERGDL/DAY/06, \(last access: September 2019\), 2019.](#)
- [Ignatenko, V., Laurila, P., Radius, A., Lamentowski, L., Antropov, O., & Muff, D. ICEYE Microsatellite SAR Constellation Status Update: Evaluation of first commercial imaging modes. In *IGARSS 2020-2020 IEEE International Geoscience and Remote Sensing Symposium* \(pp. 3581-3584\). IEEE, 2020.](#)
- 195 [Lazin, R., Shen, X., & Anagnostou, E. Estimation of flood-damaged cropland area using a convolutional neural network. *Environmental Research Letters*, 16\(5\), 054011, 2021.](#)
- [Lauer, Joe. "Flooding impacts on corn growth and yield." *Agronomy Advice*. University of Wisconsin. *Agronomy Department, Field Crops* 28 \(2008\): 49-56.](#)
- [Morris, J., & Brewin, P. The impact of seasonal flooding on agriculture: the spring 2012 floods in Somerset, England. *Journal of Flood Risk Management*, 7\(2\), 128-140, 2014.](#)
- 200 [Mst Jesmin Ara. Effect of floods on farmer's livelihood: a case study for building agriculture resilient to floods in Bangladesh. *International Journal of Science, Environment and and Technology*, 8\(2\), 334-344, 2019.](#)
- [Navarro, A., García-Ortega, E., Merino, A., Sánchez, J. L., Kummerow, C., & Tapiador, F. J. Assessment of IMERG precipitation estimates over Europe. *Remote Sensing*, 11\(21\), 2470, 2019.](#)
- 205 [Nobre, A. D., Cuartas, L. A., Hodnett, M., Rennó, C. D., Rodrigues, G., Silveira, A., & Saleska, S. Height Above the Nearest Drainage—a hydrologically relevant new terrain model. *Journal of Hydrology*, 404\(1-2\), 13-29, 2011.](#)
- [Reinsurance News: Berenberg says European floods to cost reinsurers up to €3bn, <https://www.reinsurancene.ws/berenberg-says-european-floods-to-cost-reinsurers-up-to-e3bn/>, last access: 20 July 2021.](#)
- [Shen, X., Anagnostou, E. N., Allen, G. H., Brakenridge, G. R., & Kettner, A. J. Near-real-time non-obstructed flood inundation mapping using synthetic aperture radar. *Remote Sensing of Environment*, 221, 302-315, 2019.](#)
- 210 [Shen, X., Dacheng W., Kebiao M., Anagnostou, E.N. and Hong Y., 2019b: Inundation Extent Mapping by Synthetic Aperture Radar: A Review, *Remote Sensing*, 11, 879.](#)
- [Successful farming: UPDATE 2-EU MONITOR TWEAKS 2021 CROP YIELD FORECASTS AS RAINS HIT QUALITY, <https://www.agriculture.com/markets/newswire/update-2-eu-monitor-tweaks-2021-crop-yield-forecasts-as-rains-hit-quality>, last access: 26 July 2021.](#)
- 215 [Tapia-Silva, F. O., Itzerott, S., Foerster, S., Kuhlmann, B., & Kreibich, H. Estimation of flood losses to agricultural crops using remote sensing. *Physics and Chemistry of the Earth, Parts A/B/C*, 36\(7-8\), 253-265, 2011.](#)
- [World weather attribution: Heavy rainfall which led to severe flooding in Western Europe made more likely by climate change, <https://www.worldweatherattribution.org/heavy-rainfall-which-led-to-severe-flooding-in-western-europe-made-more-likely-by-climate-change/>, last access: 23 August 2021.](#)
- 220 [Yamazaki, D., Ikeshima, D., Sosa, J., Bates, P. D., Allen, G. H., & Pavelsky, T. M. MERIT Hydro: A high-resolution global hydrography map based on latest topography dataset. *Water Resources Research*, 55\(6\), 5053-5073, 2019.](#)

- 225 Yang, Q., Shen, X., Anagnostou, E.N., Mo, C., Eggleston, J.R., Kettner, A.J., 2021. A High-Resolution Flood Inundation Archive (2016 – the Present) from Sentinel-1 SAR Imagery over CONUS. *Bulletin of the American Meteorological Society*, 1–40. <https://doi.org/10.1175/BAMS-D-19-0319.1>.
- Reinsurance News: Berenberg says European floods to cost reinsurers up to €3bn, <https://www.reinsurancene.ws/berenberg-says-european-floods-to-cost-reinsurers-up-to-e3bn/>, last access: 20 July 2021.
- CNN: Germany's deadly floods were up to 9 times more likely because of climate change, study estimates, <https://www.cnn.com/2021/08/23/europe/germany-floods-belgium-climate-change-intl/index.html>, last access: 24 August 230 2021.
- World weather attribution: Heavy rainfall which led to severe flooding in Western Europe made more likely by climate change, <https://www.worldweatherattribution.org/heavy-rainfall-which-led-to-severe-flooding-in-western-europe-made-more-likely-by-climate-change/>, last access: 23 August 2021.
- Successful farming: UPDATE 2 EU MONITOR TWEAKS 2021 CROP YIELD FORECASTS AS RAINS HIT QUALITY, <https://www.agriculture.com/markets/newswire/update-2-eu-monitor-tweaks-2021-crop-yield-forecasts-as-rains-hit-quality>, last access: 26 July 2021. 235
- Euractiv: <https://www.euractiv.com/section/agriculture-food/news/eu-farmers-warn-harvest-will-fail-after-floods-plea-for-aid/>, last access: 20 July 2021.
- Eurofruit: EU farmers warn harvest will fail after floods, plea for aid, <http://www.fruitnet.com/eurofruit/article/185840/europes-flood-damage-begins-to-emerge>, last access: 22 July 2021. 240
- Tapia-Silva, F. O., Itzerott, S., Foerster, S., Kuhlmann, B., & Kreibich, H. Estimation of flood losses to agricultural crops using remote sensing. *Physics and Chemistry of the Earth, Parts A/B/C*, 36(7-8), 253-265, 2011.
- Agence europe: Impact of flood disaster on EU agriculture is yet to be determined, <https://agenceeurope.eu/en/bulletin/article/12766/19>, last access: 21 July 2021.
- 245 Shen, X., Anagnostou, E. N., Allen, G. H., Brakenridge, G. R., & Kettner, A. J. Near real-time non-obstructed flood inundation mapping using synthetic aperture radar. *Remote Sensing of Environment*, 221, 302-315, 2019.
- Huffman, G.J., Stocker E.F., Bolvin D.T., Nelkin E.J., and Tan J.: GPM IMERG Late Precipitation L3 1-day 0.1-degree x 0.1-175-degree V06, Edited by Andrey Savtchenko, Greenbelt, MD, Goddard Earth Sciences Data and Information Services Center (GES DISC), 10.5067/GPM/IMERGDL/DAY/06, (last access: September 2019), 2019.
- 250 Yamazaki, D., Ikeshima, D., Sosa, J., Bates, P. D., Allen, G. H., & Pavelsky, T. M. MERIT Hydro: A high-resolution global hydrography map based on latest topography dataset. *Water Resources Research*, 55(6), 5053-5073, 2019.
- Nobre, A. D., Cuartas, L. A., Hodnett, M., Rennó, C. D., Rodrigues, G., Silveira, A., & Saleska, S. Height Above the Nearest Drainage – a hydrologically relevant new terrain model. *Journal of Hydrology*, 404(1-2), 13-29, 2011.
- Mst Jesmin Ara. Effect of floods on farmer's livelihood: a case study for building agriculture resilient to floods in Bangladesh. *International Journal of Science, Environment and and Technology*, 8(2), 334-344, 2019. 255

Morris, J., & Brewin, P. The impact of seasonal flooding on agriculture: the spring 2012 floods in Somerset, England. *Journal of Flood Risk Management*, 7(2), 128–140, 2014.

Navarro, A., García Ortega, E., Merino, A., Sánchez, J. L., Kummerow, C., & Tapiador, F. J. Assessment of IMERG precipitation estimates over Europe. *Remote Sensing*, 11(21), 2470, 2019.

260 Ignatenko, V., Laurila, P., Radius, A., Lamentowski, L., Antropov, O., & Muff, D. ICEYE Microsatellite SAR Constellation Status Update: Evaluation of first commercial imaging modes. In *IGARSS 2020–2020 IEEE International Geoscience and Remote Sensing Symposium* (pp. 3581–3584). IEEE, 2020.

Lazin, R., Shen, X., & Anagnostou, E. Estimation of flood-damaged cropland area using a convolutional neural network. *Environmental Research Letters*, 16(5), 054011, 2021.

265 Gould, I. J., Wright, I., Collison, M., Ruto, E., Bosworth, G., & Pearson, S. The impact of coastal flooding on agriculture: A case study of Lincolnshire, United Kingdom. *Land Degradation & Development*, 31(12), 1545–1559, 2020.






Figure 1. (a) RAPID flood map; (b) CORINE land cover map; (c) and (d) Sentinel-1 SAR image in VH polarization sensed on July 16th and 10th.

270



(a)



(b)

Figure 2. (a) Spatial pattern of precipitation accumulation over western Europe from 12th to 15th July, derived from IMERG half-hourly Final Run data. (b) Inundation extents over western Europe from 15th to 18th July, derived from RAPID system.

275




Figure 3. (a) RAPID flood map; (b) Sentinel-1 SAR image in VH polarization sensed on July 16th; (c) Sentinel-1 SAR image in VH polarization sensed on June 22.




Figure 4. Inundation extent from (a) Landsat-based flood map and (b) RAPID flood map on July 18 in northern Netherland.




Figure 5. Inundated area of land use grouped by countries over western Europe





Figure 1. Spatial pattern of (a) maximum hourly precipitation and (b) precipitation accumulation during flooding period (12 to 15 July) over western Europe.



285 Figure 2. Inundation extents over western Europe from 15th to 18th July, derived from RAPID system.




Figure 3. The land use fractions in inundated areas.




Figure 4. Inundated area of land use grouped by countries over western Europe