

# Characteristics and causes of natural and human-induced landslides in a tropical mountainous region: the Rift flank west of Lake Kivu (DR Congo)

Jean-Claude Maki Mateso<sup>1,2</sup>, Charles L. Bielders<sup>2</sup>, Elise Monsieurs<sup>3,4,5</sup>, Arthur Depicker<sup>6</sup>,  
5 Benoît Smets<sup>3,7</sup>, Théophile Tambala<sup>1</sup>, Luc Bagalwa Mateso<sup>1</sup> and Olivier Dewitte<sup>3</sup>

<sup>1</sup>Centre de Recherche en Sciences Naturelles, Department of Geophysics, Lwiro, DR Congo

<sup>2</sup>Université catholique de Louvain, Earth and Life Institute – Environmental Sciences, Louvain-La-Neuve,  
Belgium

<sup>3</sup>Royal Museum for Central Africa, Department of Earth Sciences, Tervuren, Belgium

10 <sup>4</sup>University of Liège, Department of Geography, Liège, Belgium

<sup>5</sup>F.R.S.-FNRS, Brussels, Belgium

<sup>6</sup>KU Leuven, Department of Earth and Environmental Sciences, Belgium

<sup>7</sup>Vrije Universiteit Brussel, Department of Geography, Brussels, Belgium

*Correspondence to:* Jean-Claude Maki Mateso (jean-claude.maki@uclouvain.be; makigeo2013@gmail.com)

## 15 **Abstract**

Tropical mountainous regions are often identified as landslide hotspots with growing population pressure. Anthropogenic factors are assumed to play a role in the occurrence of landslides in these populated regions, yet the relative importance of these human-induced factors remains poorly documented. In this work, we aim to explore the impact of forest cover dynamics, roads and mining activities on the characteristics and causes of  
20 landslides in the Rift flank west of Lake Kivu in the DR Congo. To do so, we compile a comprehensive multi-temporal inventory of 2730 landslides of different types that we group into five categories and that we analyze accordingly via frequency-area statistics, frequency ratio distribution and logistic regression susceptibility assessment. We find that natural factors contributing to the cause of recent (post 1950's) and old deep-seated  
25 landslides were either different or changed over time. Under similar topographic conditions, shallow landslides are more frequent, but of smaller size, in areas where deforestation has occurred since the 1950's. We attribute this size reduction to the decrease of regolith cohesion due to forest loss, which allows for a smaller minimum critical area for landsliding. In areas that were already deforested in 1950's, shallow landslides are less frequent, larger, and occur on less steep slopes. This suggests a combined role between regolith availability and soil  
30 management practices that influence erosion and water infiltration. Mining activities increase the odds of landsliding. Mining and road landslides are larger than shallow landslides but smaller than the recent deep-seated instabilities, and they are controlled by environmental factors that are not present under natural conditions. Our analysis demonstrates the role of human activities on the occurrence of landslides in the Lake Kivu region. Overall, it highlights the need to consider this context when studying hillslope instability characteristics and distribution patterns in regions under anthropogenic pressure. Our work also highlights the importance of considering the  
35 timing of landslides over a multi-decadal period of observation.

## 1 Introduction

Tropical mountainous regions are often identified as landslide hotspots with particularly vulnerable populations (Vanacker et al., 2003; Broeckx et al., 2018; Froude and Petley, 2018; Emberson et al., 2020). Nevertheless, the  
40 current knowledge on landslide processes in these regions remains limited as it is mostly derived from  
susceptibility models made at continental or global levels (Stanley and Kirschbaum, 2017; Broeckx et al., 2018).  
Because they are not based on detailed local inventories, such models do not allow to properly consider region-  
specific characteristics and causes of landslides (Depicker et al., 2020).

The growing demographic pressure and widespread land use and land cover (LULC) changes are expected to  
45 increase the frequency and impacts of landslides in tropical mountainous regions, especially in rural environments  
(Vanacker et al., 2003; Sidle et al., 2006; DeFries et al., 2010; Mugagga et al., 2012; Guns and Vanacker, 2014;  
Froude and Petley, 2018; Depicker et al., 2021a; Muñoz-Torrero Manchado et al., 2021). Deforestation and the  
associated loss of tree roots usually lower the slope stability by decreasing regolith cohesion and altering drainage  
50 patterns; whose effects are particularly pronounced on the occurrence of shallow landslides (Sidle and Bogaard,  
2016). Mining, quarrying and road construction alters the environment and commonly increases landslide activity  
(e.g. Sidle et al., 2006; Brenning et al., 2015; Arca et al., 2018; McAdoo et al., 2018; Vuillez et al., 2018; Muñoz-  
Torrero Manchado et al., 2021; Tanyaş et al., 2022). However, the impact of these anthropogenic factors on  
landslide processes (e.g. types, size, dynamics) depends on their timing and their legacy effect. It also depends on  
other environmental conditions such as slope angle and lithology (Depicker et al., 2021b). Developing further our  
55 understanding of landslides and their natural- and human-induced causes is therefore needed, especially in regions  
such as the tropics where the dearth of data is commonplace (Dewitte et al., 2022).

To achieve this, a detailed multi-temporal regional landslide inventory spanning several decades is essential  
(Guzzetti et al., 2012). New methodologies have been proposed in the past years to automatically map landslides  
with the use of, for example, Earth Observation data and machine learning (e.g. Prakash et al. 2021). However,  
60 such automatic approaches only perform well with recent landslides with a clear spectral signature. Furthermore,  
they are not always well adapted to an accurate understanding of the processes (Jones et al., 2021), especially when  
the landscapes are complex and highly influenced by human activities (Jacobs et al., 2018). The need for a visual  
identification of landslides is even more important when the movements that are studied are older and have  
occurred at an unknown period of time, much before the availability of satellite images (Pánek et al., 2021).

65 Historical aerial photographs offer the best opportunity at the regional level to work across several decades, both  
to compile a landslide inventory but also to reconstruct LULC changes (Glade, 2003; Guns and Vanacker, 2014;  
Shu et al., 2019). It is complementary to very high spatial resolution satellite images such as those available on ©  
Google Earth (Fisher et al., 2012), which are widely used in the identification of landslides in many environments  
(e.g. Broeckx et al., 2018; Pánek et al., 2021). Fieldwork is also essential in order to validate observations made  
70 from the different image sources, to discriminate between deep-seated and shallow processes, or to confirm depth  
estimates (Dewitte et al., 2021). Field surveys also help to understand the role of human activities on slope  
dynamics (Dewitte et al., 2021). Overall, sufficiently long and precise multi-decadal records of landslide activity  
and LULC are rare (e.g. Glade, 2003; Guns and Vanacker, 2014; Shu et al., 2019).

75 The aim of this work is to explore the role played by natural and human factors on the occurrence of landslides in  
a rural tropical mountainous region under high anthropogenic pressure. More specifically, we are interested in the  
Rift flank west of Lake Kivu, a region in the DR Congo where recent studies have shown that landslides are  
frequent and that recent deforestation has impacted the occurrence of shallow landslides (Maki Mateso and  
Dewitte, 2014; Depicker et al., 2020; Depicker et al., 2021b). We aim to: (1) further develop the existing landslide  
80 dataset and compile a comprehensive detailed multi-temporal regional landslide inventory spanning several  
decades; (2) describe the general characteristics of the landslides, and (3) analyze their causes according to  
different controlling factors, with special attention to multi-decadal forest cover dynamics. Historical aerial  
photographs and careful field surveys are key elements in this study.

## 2 Environmental settings and current knowledge of the landslide processes

85 The study is conducted in the Rift flank west of Lake Kivu in the DR Congo (Fig. 1a). It is one of the most seismic  
regions of the African continent, crossed by active faults and composed of six main rock types of varying age (Fig.  
1b) (Delvaux et al., 2017; Laghmouch et al., 2018). A significant portion of the study area is made of lithologies  
from the Archaen, the Mesoproterozoic and the Neoproterozoic, with various degrees of chemical weathering and  
fracturing (Kampunzu et al., 1998). Lastly formed rocks are the old Neogene basalts, highly weathered, that were  
90 deposited between 11-4 Ma years. The presence of mineral resources (gold and 3T minerals - tin, tantalum and  
tungsten) favours the proliferation of, often illegal, artisanal and small-scale mining and quarrying (Van Acker,  
2005; Geenen, 2012; Bashwira et al., 2014). Industrial mining is not present in the region and there is no new road  
construction associated with it (Bashwira et al., 2014).

The region has a tropical savannah/monsoon climate tempered by its elevation (Peel et al., 2007). The natural  
95 vegetation is mainly montane forest, still preserved in the Kahuzi-Biega National Park (Imani et al., 2017).  
However, between the 17th and 18th centuries, the region began to suffer the first strong effects of human influence  
through deforestation (Nzabandora and Roche, 2015). The roads built during the late 19<sup>th</sup> and first half of the 20<sup>th</sup>  
centuries 20<sup>th</sup> played a key role on further expanding this (Aleman et al., 2018). There has been significant  
deforestation and forest loss in recent decades as well (Basnet and Vodacek, 2015; Depicker et al., 2021a,b).  
100 Selective cutting is done for energy needs, house construction, furniture production and dugout canoes.  
Clearcutting, mostly small-scale, is associated with agriculture, mining and quarrying activities and road  
construction (Musumba Teso et al., 2019; Drake et al., 2019). After deforestation, the land is often permanently  
converted to agricultural land (cropland, grassland) or tree plantations (Depicker et al., 2021a). In some places,  
however, natural regeneration of the forest takes place (Masumbuko et al., 2012).

105




Figure 1: (a) Relief and (b) geology of the study area. The study area covers the districts of Kabare, Kalehe, Walikale, Masisi and Idjwi. Topography is derived from SRTM 1 arc second. Lithology and fault maps are from Laghmouch et al. (2018).

110 The study area (~ 5,700 km<sup>2</sup>) is one of the most densely populated regions of the DR Congo with more than 200 inhabitants/km<sup>2</sup> living mainly from agriculture, mining and quarrying activities (Linard et al., 2012; Michellier et al., 2016; Trefon, 2016). This region plays a key role in the supply of food and charcoal to the smaller rural centers and to the cities of Goma and Bukavu. Over the last decades, the population in both cities increased from a few tens of thousands to more than one million inhabitants (Michellier et al., 2016). The population growth in the study

115 area was partly caused by the influx of Rwandan refugees in 1994-1995, as well as the growing artisanal mining industry that offers job opportunities (Bashwira et al., 2014; Van Acker, 2005; Butsic et al., 2015;). The road network is relatively limited. Most roads are dirt roads and are poorly maintained, and there are no built-up walls (concrete, gabions) to stabilize the cut slopes.

Compiled from a limited number of very-high spatial resolution © Google Earth images partially covering the

120 study area, a first preliminary inventory of a few hundred landslides showed that the landslide processes are diverse

and that their impacts can be high (Maki Mateso and Dewitte, 2014). The inventory over the North Tanganyika-Kivu Rift region (hereafter called NTK Rift) of which our study area is a subregion was further expanded by Depicker et al. (2020) through the use of © Google Earth imagery with a search time limited per image. This inventory consisted of shallow and deep-seated landslides without paying attention to a further differentiation of the processes. Furthermore, it did not make a distinction between these two categories of processes in a susceptibility analysis. Depicker et al. (2020) showed that, in addition to slope angle, land cover is a key landslide predictor in the NTK Rift region. A more detailed investigation of the annual evolution of the forest cover over the last 20 years showed that deforestation increases landslide erosion 2-8 times during a period of approximately 15 years before it eventually falls back to a level similar to forest conditions (Depicker et al., 2021b). A catalogue of > 150 accurately dated landslide events, i.e. landslides that can be clearly associated with a common well-defined triggering rainfall event over the same area, was compiled for the NTK Rift for the last two decades. It allowed demonstrate the role of rainfall seasonality on the annual distribution of the occurrence of new landslides (Monsieurs et al., 2018b; Dewitte et al., 2021). Some landslide events consist of clusters of several hundreds of shallow slope failures. The spatial extent of such clustered events can be larger than 10 km<sup>2</sup>. A few events like these occur during each wet season (Depicker et al., 2020; Dewitte et al., 2021). They are commonly associated with particularly intense convective rainfall (Monsieurs et al., 2018b). None of the dated landslide events were triggered by earthquakes (Dewitte et al., 2021). This do not discard the role of earthquakes in triggering landslides in the region, but instead this reminds us that the return period of earthquakes with a magnitude large enough to trigger slope instabilities can be much longer than a few decades (Delvaux et al., 2017). Their potential impact, rather localized compared to that of climatic drivers, can be inexistent during a narrow time window of observation (Dewitte et al., 2021; Depicker et al., 2021b).

Landslides can also occur due to rock weathering and regolith formation (Dille et al., 2019). In other words, the long-term evolution of these preconditioning drivers alone can explain that a slope can also fail without any apparent trigger. This implies that the many landslides that occur in isolation of other events must be interpreted with care in terms of origin. For these features, it is not clear from a visual analysis of the satellite images whether they can be directly linked to a direct trigger. In addition, many landslides occur in isolation along roads (Dewitte et al., 2021). Some of the larger, historical, landslides (i.e. landslides that do not appear active in our oldest source of information) clearly occurred more than 10,000 years ago (Dewitte et al., 2021).

### **3 Material and methods**

#### **150 3.1 Landslide inventory**

The landslide inventory is a significant update of the inventory compiled by Depicker et al. (2020) who used only © Google Earth imagery for mapping the features whatever their type, age, and rainfall, seismic or non-triggered origin as explained in Section 2. Since the focus of Depicker et al., (2020) was to study landslides over a much larger region than the one of the present research, their inventory was not only built on a limited search-time on our study area but, also, without any field survey. Moreover, in our research, we differentiated between types (according to the updated Varnes' classification proposed by Hungr et al. (2014) and timing of landsliding. We strongly relied on three image products:

- A careful and detailed 3D (elevation exaggeration of 1) visual interpretation of © Google Earth images from 2005 to 2019, which provides a complete coverage of the region at a very high spatial resolution (~0.5 m), often multi-temporal;
- The interpretation of two hillshade images derived from a TanDEM-X digital elevation model (DEM) provided at 5 m resolution and covering most of the region (see Albino et al., (2015) and Dewitte et al., (2021) for technical explanation on the production of the DEM). The hillshade images were produced with a sun elevation angle of 30° and sun azimuth angle of 315° and 45°;
- The stereoscopic analysis of one single cover of historical panchromatic photographs acquired during the 1955-1958 period at the scale ~1/50,000 (i.e. about 1 m spatial resolution on the ground); the photographs are conserved at the Royal Museum for Central Africa (RMCA, Belgium).

The historical aerial photographs allowed to differentiate between old deep-seated landslides (i.e. landslides with an unknown time of origin and already present on the photographs) and recent deep-seated landslides that have occurred during the last 60 years (i.e. after the acquisition of the photographs). The aerial photographs were not used for mapping shallow landslides since this inventory would be biased. Indeed, the spatial resolution of the photographs is twice lower than that of the images in © Google Earth. Furthermore, the photographs provide a single temporal cover, whereas the multi-temporal © Google Earth images cover information for an imagery range of up to 13 years, i.e. the age difference between the oldest and youngest image. (e.g. Minova, Kalehe, Matanda in Fig. 1: ).

The estimation of the depth of a landslide is important when the role of LULC is to be considered; shallow landslides being much more sensitive to the vegetation characteristics than deep-seated landslides (Sidle and Bogaard, 2016). In the literature, a landslide is usually defined as shallow when the depth of its surface of rupture ranges between 2 to 5 m (Keefer, 1984; Bennett et al., 2016; Sidle and Bogaard, 2016). Here, landslides with a depth < 5 m were considered as shallow. This criteria is based on the numerous field observations in the region that show that regolith can easily develop over a depth of several meters and that trees often show deep rooting systems. Following the approach of Depicker et al. (2020) and Dewitte et al., (2021), the distinction between deep-seated and shallow landslides was made by visually estimating the relative landslide depth from © Google Earth and the 5 m resolution TanDEM-X hillshade images. Extensive in situ-field observations of several hundreds of recent landslides were then carried out to valid the assessment. The landslides occurring in mining and quarrying sites were all classified as mining landslides. A specific attention was also given to the landslides occurring along roads. Mining and road landslides are assumed to be related to important anthropogenic changes in the topography. Once they have occurred, field observations show that these landslides are commonly reworked and often further excavated. Therefore, for these two types of landslides, their depth was not assessed.

Six field surveys were conducted over the period 2016 to 2019 to validate the landslide inventory mapped from the images and get extra information on the landslide timing and their causes and triggers. Additional landslides identified only in the field were not considered in the analyses as they would bias the regional landslide distribution. The work was carried out by selecting representative areas with various types of landslides and areas with less or no landslides. These areas, that cover a total of ~20% of the region, were selected based on different landscape characteristics (lithology, slope, LULC), while taking into account accessibility and safety issues that

prevent to access many places (Jaillon, 2020). We also used information from media and grey literature (student theses, field reports from local research, and academic institutions and the civil protection).

200 The frequency of landslide surface area distributions were analyzed to check the completeness of the inventory and also enable comparison with other inventories in different environments. If the area frequency density can be properly fitted to an inverse  $\Gamma$  distribution, it is considered representative of the study area (Malamud et al., 2004). A bad fit could suggest that the inventory is biased and/or incomplete. Indeed, the use of several data sources in the inventory could bias the distribution of landslides, especially bearing in mind the limitations related to the interpretation of satellite images (Guzzetti et al., 2012). We performed this analysis separately for five categories  
205 of the inventory considered together or in isolation: all landslides, old and recent deep-seated landslides, shallow landslides, mining landslides and road landslides (see Section 3.1). The analysis of the frequency area distributions for the different shallow landslide populations defined according to the LULC and its dynamics was also used to infer about differences in environmental characteristics and slope failure mechanisms (Malamud et al., 2004; Van Den Eeckhaut et al., 2007; Guns and Vanacker, 2014; Tanyaş et al., 2018). Box-plots complemented the shallow  
210 landslide area analysis.

The extent of the study area is relatively small when considering regional climatic characteristics and the time window of the shallow landslide inventory built from Google Earth imagery is limited to a few years. Therefore, the location and spatial properties (areal extent, number of occurrences) of a rainfall-triggered landslide event forming a cluster of slope failures depends strongly on the stochastic nature (location, extent and magnitude) of  
215 the triggering rainfall event and less on local terrain conditions. The consideration of all landslides of such a cluster could bias the analysis by giving an excessive weight to the local terrain conditions (Depicker et al., 2020). Thus, for the shallow landslides susceptibility analysis (see Section 3.2), we retained a maximum of 30 landslides per cluster, randomly sampled in order to strengthen the statistical analysis and avoid overfitting. The choice of this selection is also guided by the concern to have at least the minimum of data required for training and validating  
220 the susceptibility models (Depicker et al., 2020). For the inverse  $\Gamma$  analysis, those landslides selected per cluster and other isolate landslides are called distributions *minus event*.

### 3.2 Multi-decadal forest dynamics


In the study area, the agricultural land use is complex (multiple cropping, multi-layer farming) and highly dynamic due to crop rotations and associations, shifting cultivation, and the bimodal annual rainfall pattern (Heri-Kazi and  
225 Biolders, 2021b). A detailed regional land use mapping serving as input variable in our susceptibility for shallow landslides and their distribution analysis (see Section 2.3) is therefore not feasible (e.g. Jacobs et al., 2018) which is an approach that differs from what can commonly be done in non-tropical environments (e.g. Chen et al., 2019; Shu et al., 2019). However, the dynamics of the forest can be better constrained. Here, to complement the year to year analysis conducted by Depicker et al. (2021b; see Section 2.1) that focused on the impact of deforestation on  
230 shallow landslides over the last 20 years, we reconstructed the forest dynamics over the last ~60 years (Fig. 2). We used the 1 m resolution orthomosaic generated from the RMCA's aerial photographs of the years 1955-1958 (Depicker et al., 2021a); these photographs being the only existing pre-satellite era source of information. The forest areas were delineated visually. The 2016 forest cover was extracted from the continental ESA CCI land cover model which is available at a 20 m resolution (ESA, 2016). This satellite-based product has an accuracy of



235 roughly 86 % in the region and has demonstrated its relevance in another study on landslides (Depicker et al.,  
2021b). Note also that between 2016 and 2019, i.e. the date that corresponds to the most recent images in Google  
Earth used for the inventory, very little forest cover changes were observed.

240 Knowing that the natural vegetation of the study area is forest (Section 2), in 1955-58, 42 % of the territory was  
already deforested (Fig. 2a). From 1955-58 to 2016, the loss of forest continued, the forest cover decreasing from  
58 % to 24 % of the study area. The area affected by the forest loss over the last 60 years is larger than the remaining  
permanent forest (Fig. 2b). The comparison of forest areas between 1955-58 and 2016 allows to consider four  
classes for the forest dynamics:

- Permanent forest corresponds to forest areas that are present at both dates.
- 245 • The forest loss class corresponds to forests present in 1955-58 that have disappeared in 2016. Since it is  
impossible to identify for each portion of the landscape the exact cause of forest loss, this class contains  
a mix of various forest management practices and other causes of forest cut/removal.
- The forest gain class represents the new forest that has appeared since 1955-58. Similarly, the causes  
associated with the occurrence of new forest are not exactly known; afforestation and natural forest  
250 regeneration being certainly drivers at play.
- Permanent anthropogenic environment (e.g. cropland, grassland, built-up land) means that the landscape  
was not forested in both dates and it is assumed that it remained so during that period.



255 Figure 2: Forest cover dynamics over the last 60 years. (a) Forest cover in 1955-58 and 2016; (b) Areas of forest  
cover change between 1955-58 and 2016. Details for the images used in this figure are in Table 1.



### 3.3 Landslide susceptibility and distribution analysis

Landslide susceptibility approaches are commonly used to determine the factors that control the occurrence of landslides. There are numerous approaches which are more or less complex in terms of modelling implementation, data needs, and result interpretability (Reichenbach et al., 2018). Since our study does not aim to develop a new methodology nor to show the ability to use complex methods; we relied on a logistic regression approach (Hosmer and Lemeshow, 2000) to determine the predictor variables related to the occurrence of the different types of landslides. Logistic regression is a straightforward and relatively low-data demanding method that has been widely used (Reichenbach et al., 2018) and that allows a rather easy interpretation of the results (e.g. Jacobs et al., 2018; Depicker et al., 2020).

Frequency ratio (Lee and Pradhan, 2007) models were used as a more simple but complementary approach to better understand the role of each variable in the contribution of the landslide occurrence in terms of process characterization. For example, when slope angle is highlighted by a logistic regression model as a significant variable, we still remain unaware of the types of slopes that actually influence the occurrence of landslides.

The analysis was carried out according to the five categories of landslides defined in Section 3.1. The analysis was done at the scale of one point (pixel) per landslide to avoid spatial autocorrelation (e.g. Jacobs et al., 2018; Kubwimana et al., 2021). The point is manually positioned in the central region of the visually delineated landslide's source area to represent as close to reality as possible the conditions that caused its occurrence. In doing so we also avoid the selection of the highest point of the landslide that rarely corresponds to its initiation point (Dille et al., 2019). As stressed by Tanyaş et al., (2018), landslides growth with time. Therefore, considering one pixel per landslide instead of its whole source area allows to avoid a temporal-induced bias. The digital elevation model used for the analysis (see Table 1) is posterior to the occurrence of the old deep-seated landslides. Therefore, for deep-seated landslides, a point outside the source area where topography does not appear to have been disturbed by the instability is visually determined for the calculation of the slope associated with the landslide origin. Calculating the slope values at the level of the landslide source for this type of landslide would give values that are the consequences of landslides rather than the causes of their origin.

#### 3.3.1 Predictor variables and landslide causes

The purpose of this research is to examine the predictor variables (See supplementary Figure 1 for the predictor variables not displayed in the main manuscript) that contribute to the susceptibility of the different landslide categories. As such we mainly investigate the causes of the landslides. Nevertheless, the predictors highlighted by the susceptibility analysis may also help to discuss triggering conditions since the tectonic, landscape and climate of a region are commonly interlinked (Whipple, 2009; Whittaker, 2012).

We used eight predictors that can be considered as natural factors that cause landslide occurrence (Table 1): elevation, slope angle, planar curvature, profile curvature, topographic wetness index (TWI), slope aspect, lithology, and distance to faults. Although these predictors are commonly used (Reichenbach et al., 2018), it is worth specifying that, here, elevation is used as proxy for climatic conditions, namely orographic rainfall and the probability of convective rainfall/thunderstorms, as the resolution of regional-climate derived products is too low (at least 2.8 km) to accurately capture at the scale of our study area the effect of elevation on rainfall (Monsieurs

et al., 2018a; Van de Walle et al., 2020). Distance to fault is used to determine the possible contribution of seismic activity in the occurrence of deep-seated landslides not only as a triggering factor (e.g. Keefer, 1984), but also as a rock weathering factor (e.g. Vanmaercke et al., 2017). Using the fault pattern is the most appropriate option to tackle the seismic zonation context since the most detailed seismic hazard assessment for this part of the continent is at a spatial resolution of 2.2 km; i.e. at a resolution that is too coarse for our study (Delvaux et al., 2017).

Table 1. Landside predictor variables used for the susceptibility and frequency ratio analyses and the ancillary data from which they are derived.

Variable	Type	Source
- Elevation (m)	Continuous	Nasa Shuttle Radar Topography Mission (SRTM) Version 3.0 Global 1 arc second Data (Temporal Extent 2000-02-11 to 2000-02-21)  <a href="https://lpdaac.usgs.gov/products/srtm_g1v003/">https://lpdaac.usgs.gov/products/srtm_g1v003/</a>
- Slope angle (°)	Continuous	
- Profile curvature (m <sup>-1</sup> )	Continuous	
- Plan curvature (m <sup>-1</sup> )	Continuous	
- Topographic wetness index	Continuous	
- Slope aspect (°)	Categorical	
• north	Dummy	
• northeast	Dummy	
• east	Dummy	
• southeast	Dummy	
• south	Dummy	
• southwest	Dummy	
• west	Reference*	
• northwest	Dummy	
- Lithology	Categorical	Geological map of the Kivu at scale 1/500,000 (Laghmouch et al., 2018)
• Old basalts	Dummy	
• Black shales and tillite	Dummy	
• Granites (mica and leuco-granites)	Dummy	
• Granitic rocks (rhyolite)	Reference*	
• Pelites and quartzopelites	Dummy	
• Gneiss and micaschists	Dummy	
- Distance to faults (m)	Continuous	
Distance to roads (m)	Continuous	<a href="https://www.openstreetmap.org/history#map=9/-2.0475/28.5535">https://www.openstreetmap.org/history#map=9/-2.0475/28.5535</a>
- Forest dynamics between 1955-58 and 2016	Categorical	Forest cover in 2016: (ESA, 2016: <a href="http://2016africalandcover20m.esrin.esa.int/viewer.php">http://2016africalandcover20m.esrin.esa.int/viewer.php</a> )
• Permanent forest	Reference*	Forest cover in 1955-58: Historical aerial photographs and derived orthomosaics from RMCA (see Section 2.1)
• Forest loss	Dummy	
• Forest gain	Dummy	
• Permanent anthropogenic environment	Dummy	

\* Each dummy variable is compared with the reference group.

Besides the natural factors, we identified two anthropogenic predictors (Table 1): forest dynamics and distance to roads. For the forest dynamics, we considered the four classes identified in Fig. 2. The main roads were retrieved from OpenStreetMap. Good knowledge of the study area and the analysis of the very high-resolution Google Earth images allowed us to verify the high accuracy of the road network proposed by OpenStreetMap. Using the historical photographs, we observe that the main roads date back to the colonial times and that no major changes in the network have occurred over the last 60 years. The few recent landslides that are observed in the field along these roads confirm the assumption that the direct impact of the main roads on the occurrence of recent landslides is currently limited. These landslides are clearly linked to the road cut topography, i.e. topographic conditions that cannot be constrained at the resolution of the SRTM elevation data (1" or roughly 30 m). They are often of very limited size, i.e. at a size that is too small to be features that can be identified in © Google Earth in a consistent manner. For our study, the distance to roads is taken as a proxy for human settlement, trail density, and intensity and diversity of agricultural practices. Since motorized transportation means are very limited in the region, the population growth, the expansion of villages and the agricultural activities are indeed highly associated with the main road networks.

Prior to analysis, the non topographically-derived predictor variables were resampled at the resolution of the SRTM DEM data; a resolution that is commonly used in many susceptibility analyses (Reichenbach et al., 2018) and that provided the best results in similar regions (e.g. Jacobs et al., 2018). The association between the dependent variable and each predictor variable was tested using the Pearson  $\chi^2$  test at a 95 % level of confidence (e.g. Van Den Eeckhaut et al., 2006). The predictors were tested for multicollinearity, variables with variance inflation factor (VIF) > 2 being excluded from the analysis (e.g. Van Den Eeckhaut et al., 2006). The flat areas (slope angle < 1°) that are spread across the region were not excluded from the analysis since their total extent is limited and their impact on the inflation of susceptibility model performance would be minor (Brenning, 2012).

For the analysis of deep-seated landslides, the predictor variables associated with anthropogenic activities were excluded. For the shallow landslides, the 'distance to faults' variable was also excluded. As explained earlier, the shallow landslide inventory represents a narrow time window of observation. As such, the spatial distribution of the shallow landslides could be biased by the stochastic pattern of the recent heavy rainfall events and anthropogenic disturbances rather than being the reflect of the longer-term impact of weathering conditions associated with seismicity.

### 3.3.2 Logistic regression

Logistic regression is used to describe the relationship between a binary dependent variable (the presence or absence of landslides) and one or more independent predictor variables (Hosmer and Lemeshow, 2000). Hence, the logistic regression does not only require landslide data, but also non-landslide data. We sampled this non-landslide data by generating a number of random points that is equal to the number of landslides in the inventory in order to avoid prevalence (Hosmer and Lemeshow, 2000). Non-landslide points were randomly generated outside a 40 m buffer zone around landslide areas. The basic equation for logistic regression is:

$$\log\left(\frac{p}{1-p}\right) = \alpha + \sum_{i=1}^n \beta_i X_i \quad (1)$$

where  $P$  is the likelihood of landslide occurrence and takes values between 0 and 1,  $\alpha$  is the intercept of the model,  $X_i$  represents  $i$ -th of  $n$  predictors, and  $\beta_i$  the accompanying coefficient that has to be fitted to the data.

Calculations were performed in an RStudio environment version 1.4.1717 with LAND-SE software (Rossi and Reichenbach, 2016). In order to be considered in the final logistic regression equation, continuous variable coefficients needed to be significant at the 95 % level of confidence (e.g. Jacobs et al., 2018). For categorical variables, as soon as one dummy variable was significant, all other dummy variables were included in the model (e.g. Depicker et al., 2020). The quality of the models was judged by (i) the prediction rate (e.g. Depicker et al., 2020), (ii) a visual plausibility inspection of the susceptibility maps after reclassifying each map into four classes of increasing susceptibility that cover 40 %, 30 %, 20 %, and 10 % of the study area, and (iii) considering the area under the curve (AUC) of the receiver-operating-characteristics curve (ROC). The AUC values vary between 0 and 1 and can be interpreted as the model's capacity of differentiating between landslide and non-landslide locations. An AUC = 0.5 shows that the model performance is equivalent to random classification, while an AUC = 1 indicates a perfect classification (Hosmer and Lemeshow, 2000). Training and validation datasets were taken in the proportions of 70 % and 30 %, respectively (Broeckx et al., 2018; Fang et al., 2020).

We assessed the importance of each individual predictor for the logistic regression in two ways. First, we calculated the AUC for landslide susceptibility models that only relied on the considered predictor, to assess the extent to which this predictor can be used to differentiate between landslide and non-landslide locations. Although this is quite a straightforward approach that does not consider the possible interplay among predictor variables, this allow to have a first quantitative insight on the importance of each variable to the susceptibility models (Depicker et al., 2020). A second way to determine the impact of the predictors was the analysis of the odds ratio (OR). The OR of a predictor expresses how a change of a predictor value translates into an increase/decrease in the odds of landsliding, whereby the odds of landsliding is calculated as  $\frac{P}{1-P}$  (see Eq. (1)). The  $OR_i$  of predictor  $i$  is calculated as:

$$OR_i = e^{\beta_i \delta_i}, \quad (2)$$

whereby  $\beta_i$  is the coefficient of predictor  $i$ , and  $\delta_i$  is the increase in predictor  $i$ . For continuous variables an arbitrary but realistic value for  $\delta_i$  is chosen. For the dummy variables,  $\delta_i$  equals 1. For the categorical variables, the OR for each dummy reflects an increase or decrease relative to the reference variable (Kleinbaum and Klein, 2010).

### 3.3.3 Frequency ratio

The frequency ratio model considers each landslide predictor variable individually and classifies its values into a set of bins to indicate for each bin of the predictor variable the probability of occurrence of a landslide (Lee and Pradhan, 2007; Lee et al., 2007; Kirschbaum et al., 2012). The frequency ratio is calculated as:

$$Fr_{cb} = \frac{a_{cb}/a_T}{A_{cb}/A_T}, \quad (3)$$

where  $Fr_{cb}$  is the frequency ratio value for a bin  $b = (1, 2, \dots, n)$  of a predictor variable  $c = (1, 2, \dots, m)$ ,  $a_{cb}$  is the cumulative landslide area within bin  $b$  of predictor  $c$ ,  $a_T$  is the cumulative landslide area in the entire study area,  $A_{cb}$  is the area attributed to bin  $b$  of predictor  $c$ , and  $A_T$  is the total extent of the study area.

## 4 Results

### 4.1 Landslide inventory

Overall, we mapped 2730 landslides (Fig. 3a; Table 2). The landslides are diverse in terms of size, age and type (Fig. 4). The inventoried landslides cover ~3 % of the study area. The largest landslide is an old and deep-seated complex movement (426 ha), while the smallest detected landslide is a shallow debris avalanche (16 m<sup>2</sup>). The landslides are grouped into five categories (Fig. 3a; Table 2):

- Old deep-seated landslides represent 45,5 % of the inventoried landslides and cover 93 % of the total landslide affected area. Most of these landslides are of the rock slide type. Rock avalanches, although much less frequent, are also present. Rockfalls can be associated with the presence of the main scarps of these old landslides. However, they have not been considered in the inventory and the subsequent analysis;
- Shallow landslides represent 40.4 % of inventoried landslides, but represent only 2.7 % of the total affected area. Most of these landslides are of the debris avalanche type. These landslides are all recent and clearly associated with rainfall. The landslides clustered events all fall in this category;
- Recent deep-seated landslides represent a small percentage of landslides (5.8 %) but cover an area (2.9 %) similar to shallow landslides. Most of the landslides are of the slide type. Their trigger, when identified, is associated with rainfall;
- Mining landslides (that also include quarrying landslides) represent 5.6 % of the inventoried landslides and cover 1.2 % of the total landslide affected area;
- Road landslides: the inventory shows that 115 landslides are located within 50 meters of roads. 60 of these landslides are shallow, 13 recent and deep-seated, 35 old and deep-seated, and 7 are mining landslides. Only the shallow and recent deep-seated landslides were classified as road landslides; i.e. a total of 73 landslides. The old deep-seated landslides located close to roads were retained in the old deep-seated landslide category because their timing is likely to precede road construction. The mining landslides were also retained in their respective category.




Figure 3: (a) Landslide inventory obtained from the image analysis and extent of the forest cover in 2016 (after ESA, 2016). Numbers represent clusters of shallow landslides that are associated with heavy rainfall events dated in ascending order from oldest to most recent. (b) Additional landslides identified only in the field.

405 We identified several shallow landslides clustered events. One of the events is related to the Kalehe rainstorm of October 2014 (Fig. 3a: event 2; Fig. 4a) reported by Maki Mateso and Dewitte (2014). This rainfall triggered 634 shallow landslides, 346 of them being connected to talwegs and providing materials to 17 debris flows. Ten debris flows were particularly destructive and deadly when they reached villages on the shores of Lake Kivu (Maki Mateso and Dewitte, 2014). In this area, 14 shallow landslides present on © Google Earth images before this event

410 were reactivated. Field observations and interviews with local populations confirmed that the shallow landslides that are not associated with these clustered events are also rainfall-triggered.

Table 2: Typology, size properties, and identification methods of the inventoried landslides (LS). The percentages of landslides linked to the TanDEM-X hillshade images (% of LS in TanDEM-X) represent landslides that could not be very well identified in © Google Earth alone.

Landslide type	Number of LS	% of LS	% of LS area	Max area (ha)	Min area (m <sup>2</sup> )	Average area (ha)	Standard deviation (ha)	% of LS in ©Google Earth	% of LS in TanDEM-X
Deep-seated (old)	1243	45.5	93.0	426.4	604	12.6	26.8	94.9	5.1
Deep-seated (recent)	159	5.8	2.9	28.9	210	3.1	5.4	97.5	2.5
Shallow	1103	40.4	2.7	53.8	16	0.4	2.4	100	0
Mining	152	5.6	1.2	13.4	99	1.4	1.9	100	0
Road	73	2.7	0.1	2.0	149	0.3	0.3	100	0
All landslides	2730	100	100			6.2		97.5	2.5

415

Table 3: Field-based validation of the landslides (LS) inventoried from the image analysis. True Positive (TP) = landslides that were mapped in the images and validated in the field. False Positive (FP) = landslides that were mapped in the images but not validated in the field. False Negative (FN) = landslides that were identified solely in the field. Precision = TP / (TP+TN)

Landslide type	Number of LS mapped in the images and checked in the field	TP	FP	FN	Precision (%)	Total number of LS viewed in the field
Deep-seated (old)	248	239	9	60	96	308
Deep-seated (recent)	47	44	3	4	94	51
Shallow	426	420	6	55	99	481
Mining	15	9	6	2	60	17
Road	50	45	5	5	90	55
Total	786	757	29	126	96	912

420


Landslide mapping was largely done using © Google Earth; the TanDEM-X hillshades being useful to confirm the identification of about one fifth of the old deep-seated landslides (Table 2). Fieldwork carried out to validate 786 landslides (25% of the inventory) showed that they were identified with a precision of 96 % (Table 3). Old deep-seated landslides and shallow landslides were mapped with the highest precision. Mining landslides were mapped with a lower precision due to the difficulty of differentiating between landslide processes and anthropogenic soil disturbance in © Google Earth imagery. The field validation allowed to also map an extra 126 landslides (Fig. 3b) that could only be identified in the field (Table 3). For the old deep-seated landslides, this

425



represents an extra 24% of observations (Table 3: see column FN). Nevertheless, landslides identified only in the field were not considered in the analysis to avoid biases due to overrepresentation.

430 Each debris flow is connected to up to hundreds of shallow landslides that act as source areas. A clear distinction was made between these sources and the debris flow path and deposition areas (Fig. 4a). Out of a total of the 184 debris flows identified from the images, 90 with a length-to-width ratio  $> 50$  were excluded from the analysis since they show greater similarities to debris-rich floods than to the other landslides present in the region (Malamud et al., 2004). Nevertheless, the shallow landslides acting as source areas were kept in the analysis. Also, 22 very large, 435 old, deep-seated landslides were excluded from the analysis because they have complex main scarps where it is difficult to determine the pixels that best represent the natural conditions of occurrence. Overall, from the 2730 landslides identified from the images, 2618 landslides were used for the subsequent analysis.



440 Figure 4: Examples of landslide types (according to Varnes' new classification – Hungr et al., 2014). (a) Cluster of recent debris avalanches and debris flows triggered during an intense rainfall event (25/10/2014) in the vicinity of Kalehe (-2.041°S, 28.874°E); the image illustrates a part of the landslides clustered event 2 shown on Fig. 3a). The source areas of these shallow landslides are identified. (b) Old earthflow (-2.053°S, 28.660°E). (c) Old rock slides/rock avalanches/ with path-dependent rockfalls (-2.007°S, 28.708°E). (d) Recent deep-seated rotational slide that occurred in 2002 (-1.530°S, 28.708°E). (e) Recent deep planar slide that occurred in 1994 and created a

445 dammed lake (-1.521°S, 28.977°E). (f) Recent slides, flows and avalanches associated with mining activities that occurred from 2013 onwards (-1.563°S, 28.885°E).

Except for the recent deep-seated and mining landslides, the inverse  $\Gamma$  distribution fits well the distributions for the other categories of the inventory, (Fig. 5a,c); which supports their use for further susceptibility analysis. The Wilcoxon rank comparison test confirms significant statistical differences ( $p$ -value  $< 0.05$ ) among the area distributions (Fig. 5b).

450




Figure 5: Landslide (LS) area characteristics. (a, c) Landslide frequency-area distributions for each landslide category. (b) Boxplots showing the distribution of landslide area for each landslide category. Boxplots show the lower and upper quartiles and median. The whiskers of each box represent 1.5 times the interquartile range. The average area of the landslides (red dots) is provided for each boxplot and the outliers beyond whiskers are shown as dots. The number of landslides in each category is shown in brackets.

455




Figure 6: Shallow landslide characteristics and forest cover dynamics. (a, c) Boxplots showing the distribution of landslide area and landslide slope, respectively, for each land cover class. A detailed description of boxplots is provided in Figure 5. (b, d) Shallow landslide frequency-area distributions for each land cover class.

A majority (72 %) of the shallow landslides is found in areas of forest loss (Fig. 6). The landslides in the permanent anthropogenic environment have the largest mean area, followed by the landslides in permanent forest, and the landslides in areas of forest loss. In forest gain zones, landslides are on average the smallest. The Wilcoxon rank comparison test confirms significant statistical differences ( $p$ -value  $< 0.05$ ) among the landslide area distributions. The same differences are also confirmed for the landslide slope distribution (Fig. 6b). In permanent forest areas, shallow landslides occur on steeper slopes compared to shallow landslides in anthropogenic environments (Fig. 6b). The analysis of the completeness of the inventory (Fig. 6b,d) shows that an acceptable distribution emerges for each category of shallow landslides except for the landslide inventory in *permanent forest minus event* (Fig. 6b).


#### 4.2 Landslide susceptibility and distribution analysis

The Pearson  $\chi^2$  tests confirm the association between the dependent variable and each predictor variable at a 95 % level of confidence. There was not multicollinearity between the predictors ( $VIF < 2$ ) retained for this study.

Depicker et al. (2020) assessed the impacts of the size of the landslide training dataset to calibrate a landslide susceptibility model. They showed that the quality of a susceptibility assessment is questionable if the number of



landslides is too small. In view of the low number of recent deep-seated, mining, and road landslides in the present study (Table 3), we did not calibrate susceptibility models from these three types of landslides. Instead, we tested these inventories against the two susceptibility models computed from the shallow and/or old deep-seated landslide datasets, from which we could derive prediction rates (Fig. 7).



480

Figure 7: Landslide susceptibility models and prediction rates. (a) shallow landslides (AUC: 0.78); (b) old deep-seated landslides (AUC: 0.82); (c) prediction rate curves for shallow, mining, and road landslides; (d) prediction rate curves for old deep-seated, recent deep-seated, mining and road landslides. The red highlight (c, d) represents the 10 % of the region with the highest landslide susceptibility values.

485

The univariate AUC values are all above 0.5 (Table, 4). All predictors considered for both categories of landslides where thus considered in the multivariate logistic regression models. The two susceptibility models of shallow and old deep-seated landslides show similar AUC and prediction rates (Fig. 7). At first sight, both models have spatial similarities of high susceptibility on the eastern part of the region; while the entire western part is weakly susceptible (Fig. 7a,b). However, when we go into detail, the spatial patterns of the susceptibility values of the two

490

models are quite different as it reflects the differences in the importance of the predictors included in the assessment (Table 4, Table 5).

Table 4: Relative importance of the predictors of the logistic regression models for shallow and old deep-seated landslides based on AUC<sub>i</sub> (ranked in descending order).

Shallow landslides		Old deep-seated landslides	
Predictor	AUC <sub>i</sub>	Predictor	AUC <sub>i</sub>
Forest loss	63	Profile curvature.	65.7
Elevation	61.5	Elevation	65.3
Slope angle	60.1	Distance to faults	64.2
Distance to roads	59.7	Slope angle	64
Pelites and quartzopelites	58.9	TWI	63.8
Permanent anthropogenic environment	55.9	Plan curvature.	59.5
TWI	55.3	Pelites and quartzopelites	54.5
Plan curvature.	53.1	South	52.4
East	52.2	North-east	52
South-east	52.2	North	51.6
Black shales and tillite	51.8	East	51.1
Old basalts	51.8	South-east	50.7
North	51.8	Granites (mica and leuco-granites)	50.7
Granites (mica and leuco-granites)	50.8	Old basalts	50.5
South-west	50.7	Gneiss and micaschists	50.5
Gneiss and micaschists	50.6	South-west	50.3
South	50.6	Black shales and tillite	50.3
Profile curvature	50.4	North-west	50.2
North-east	50.4		
North-west	50.1		
Forest gain	*		

495 \* Only four landslides are present in this category.

Table 5. Results of the logistic regression models for shallow landslides and old deep-seated landslides.

		Shallow landslides		Old deep-seated landslides		Step
AUC		0.78		0.82		
Predictor variable		LR coef.	Odds ratio	LR coef.	Odds ratio	$\delta_i$
(Intercept)		-3.560 ***		-1.661 ***		
Elevation		0.001 ***	1.857	0.002 ***	2.535	500
Slope aspect	Northwest	0.842 *	2.321	-0.366	0.694	1

	West	Ref.	-	Ref.	-		
	Southwest		0.674 *	1.962	-0.232	0.793	1
	South		0.630 *	1.878	0.032	1.033	1
	Southeast		0.599	1.820	-0.345	0.708	1
	East		0.513	1.670	-0.578 **	0.561	1
	Northeast		0.622	1.863	-0.897 ***	0.408	1
	North		0.481	1.618	-0.831 ***	0.436	1
	Plan curvature		-0.272 *	0.580	0.166 ***	1.394	2
	Profile curvature		-0.190	0.999	-0.463 ***	0.998	0.005
	Slope angle		0.050 ***	1.649	0.033 ***	1.391	10
	Topographic wetness index		0.093	1.000	-0.281 ***	1.000	0.001
Lithology	Old basalts		-0.753 -	0.471	0.201	1.223	1
	Black shales and tillite		-1.207 ***	0.299	-1.358 ***	0.257	1
	Granite coarse grain		-17.026 -	0.000	-2.126 ***	0.119	1
	Granitic rocks (rhyolite)	Ref.			Ref.		
	Pelites and quartzopelites		-1.274 ***	0.280	0.155	1.168	1
	Gneiss and micaschists		0.506	1.659	-0.468	0.626	1
	Distance to roads		0.000 ***	0.931	no	-	500
	Distance to faults	no	-		0.000 ***	0.914	500
Forest cover dynamics	Permanent forest	Ref.	-		no	-	
	Forest loss		0.922 ***	2.514			1
	Gain forest	no	-		no	-	
	Permanent anthropogenic environment		-0.159	0.853	no	-	1

No = variable not included in the logistic regression model

Ref. = reference category of the dummy variable

500 Coefficient included in the logistic regression model = \*p-value < 0.05, \*\* p-value < 0.01, \*\*\* p-value < 0.001

505 Forest loss has a large influence on the occurrence of shallow landslides as deforestation increases the odds of landsliding by a factor 2.5 (Table 4, Table 5). However, anthropogenic environments appear to be less landslide-prone than permanent forest. Elevation and slope angle are similarly important for the prediction of both types of landslides (Table 4) but have a slightly larger impact on the odds of deep-seated landsliding than on the odds of shallow landsliding (Table 5). Slope aspect has a greater impact on the occurrence of shallow landslides than for old deep-seated landslides. It appears that the plan curvature reduces the occurrence of shallow landslides while it favours the occurrence of old deep-seated landslides. The effect of lithology is also different for shallow and deep-seated landslides. For shallow landslides, the gneiss and micaschists are most landslide-prone and the lowest susceptibility is associated with black shales, tillite and old basalts. For deep-seated landslides, black shales, tillite and old basalts favour landslides while gneiss and micaschists do not. 'Distance to roads' and 'distance to faults' have a significant but rather limited impact on shallow and old deep-seated landslides, respectively.

Mining and road landslides are poorly predicted using the shallow landslide model (Fig. 7c). The prediction of road and mining landslides using the deep-seated model is also poor, although less problematic for the mining landslides (Fig. 7d). Recent deep-seated landslides are reasonably well predicted using the old deep-seated landslide model, which validates to some extent the multi-temporal predicting performance of the assessment.




Figure 8: Frequency distribution for shallow and deep-seated landslides in function of different predictor variables. The corresponding frequency ratio is shown for each class above the vertical bars. The green, orange and red curves indicate the proportion of forest cover, forest loss and slope > 25°, respectively, in the different classes of the predictor variables.

Slope angle is an important driver for shallow and old deep-seated landslides (Fig. 8a,b). Both types of landslides are favoured by slopes angles > 20-25°. We observe a trend in the landscape of increasing slopes and forest loss and decreasing forest cover with increasing elevation (Fig. 8c). The decrease in forest cover at high elevations is also associated with a natural change of the vegetation: bamboo vegetation is found at 2300-2600 m asl and subalpine vegetation such as ferns occur at 2400-3300 m asl (Mokoso et al., 2013; Cirimwami et al., 2019). At higher elevations (> 2000 m), shallow landslides occur more frequently, and this can probably be explained by a cumulative effect of forest loss, steeper slopes and increased orographic rainfall associated with these elevations



(Fig. 8c). The positive frequency ratio in the 1400-1700 m elevation class is related to the area of permanent anthropogenic environment. This zone is characterized by low forest cover and relatively low slopes (Fig. 8c). Deep-seated landslides are also favoured by steeper slopes and higher elevations. Regarding the dynamics of forest cover (Fig. 8e), the occurrence of shallow landslides is favoured in the deforested areas.

535

## 5 Discussion

### 5.1 Landslide types and completeness of the inventory

Despite its high precision, and the fact that with more than 2700 mapped landslides we have identified more than three times as many features as in the inventory of Depicker et al. (2020), we are aware that the dataset is still incomplete. This is particularly the case for the shallow landslides because their inventory covers a maximum period of 13 years. Furthermore, their scars can quickly be altered by natural vegetation regrowth, land reclamation and erosion (Malamud et al., 2004; Van Den Eeckhaut et al., 2007; Kubwimana et al., 2021; Dewitte et al., 2022); although, here, since we have used several image covers from Google Earth, this issue should be nuanced. In addition, small landslides frequently happen unnoticed at the resolution of the satellite images (Guzzetti et al., 2012). Finally, field validation showed that a significant proportion of old deep-seated landslides can be missed from image analysis (Table 3). This is because identifying the exact limits of the failed mass may not be easy for old deep-seated landslides, particularly in forest areas (Malamud et al., 2004). While building the inventory, we remained conservative and mapped only the features for which we had high confidence. As the protocol for landslide identification over the whole region was uniform and the number of identified landslides relatively important, we trust that the inventory is reliable and representative enough for the analysis.

540

545

550

555

560

565

The frequency area distributions of all landslides categories (Fig. 5a,c), with the exception of recent deep-seated and mining landslides, are similar to what has been observed in other parts of the world (e.g., Malamud et al., 2004; Guns & Vanacker, 2014; Jacobs et al., 2017 ). For the recent deep-seated landslides, an overrepresentation is noticed at the level of the smallest landslides and the rollover is absent. Since the spectral signature of these landslides is pronounced, we cannot invoke here a problem of subjectivity in the mapping. Additionally, we can give a high trust in the completeness of the inventory as evidenced by field validation that showed that almost no landslides were missed (Table 3). Therefore, we posit that this divergence in size is related to a lower influence of successive slope failure in the increase of landslide area through time; in other words, recent landslides did not have the time to grow (Tanyaş et al., 2018). This process of successive failures has been well documented for the Ikoma landslide, south of Bukavu (Figure 1b; Dille et al., 2019). The distribution of the mining landslides is irregular and different from what is typically observed, with a rollover that is flattened and a sudden increase in the frequency of the smallest slope failures. Similarly, to the inventory of the recent deep-seated landslides, the completeness and the reliability of the mapped features cannot be much questioned. We suggest that this unusual area distribution is the result of the human-induced alteration of the environmental conditions (see Section 4.4). To our knowledge, there are no similar studies that have been carried out on artificial mining slopes. Further investigations on other cases would be needed to verify our hypothesis.

The presence of a rollover in the frequency-area distribution of the shallow landslides in the anthropogenic environment (Fig. 6b,d) is in opposition to what we could have expected considering the study by Van Den

570 Eeckhaut et al. (2007). This study was also conducted in a populated rural environment and also relied on an  
inventory that is not associated with one single landslide triggering event. They did not find a positive power-law  
relation for the smaller landslides which is separated from the larger landslides by a rollover. This difference  
probably lies in the fact that our study area is much more landslide-prone. The research by Van Den Eeckhaut et  
al. (2007) was indeed carried out in a hilly region of Belgium where the temperate climate is much less favourable  
575 than that of Van Den Eeckhaut et al. (2007), that our region is not altered by mechanized farming, and that human  
activities such as works associated with building and road construction and drainage systems are much less present,  
i.e. factors that are highlighted as causes of landslides in Belgium, are issues that can also be invoked to explain  
this divergence in the frequency area distribution of shallow landslides.

580 Under permanent forest, we do not observe a rollover point in the shallow landslide distribution, (Fig. 6b). We  
hypothesize that the smallest landslides may be hidden under the canopy and therefore less visible on satellite  
images. A second explanation is that the presence of trees and their roots increases slope stability and therefore  
the minimal critical area for landsliding (Milledge et al., 2014).

## 5.2 Drivers of deep-seated landslides

585 The old deep-seated landslide susceptibility model is the first model proposed for the region that focuses only on  
deep-seated processes. The model shows a good quantitative prediction performance, both in terms of AUC and  
prediction rate. The model shows that terrain morphology and seismic activity seem to play a dominant role in  
deep-seated landslide distribution in the study area. The frequency ratio analysis (Fig. 8b,d) further supports this  
as it highlights the association of landslides with steep slopes and higher elevations, i.e. in topographic contexts  
590 nearer to the ridge crests that are known to amplify seismic shaking (Meunier et al., 2008). The role of elevation  
as a driver of more humid conditions should, however, not be ignored as rainfall is also known to trigger deep-  
seated landslides (LaHusen et al., 2020). Also, the role of the long-term weathering of the landscape and the  
occurrence of non-triggered landslides should not be underestimated (Dille et al., 2019). Lithology is of lesser  
importance in our study area; which is in agreement with the findings of Depicker et al. (2021b) that show that the  
595 various lithologies in the region have similar rock strength properties. As we also show that the topography and  
the presence of faults play a role, it is another factor that can explain that the influence of lithology is somehow  
limited.

600 The lower prediction rate of the recent deep-seated landslides using the old deep-seated landslide model could be  
related to the fact that the observations are made on a period that is too short to apprehend the full panel of  
environmental conditions that led to old deep-seated landslides. For example, no earthquake-induced recent deep-  
seated landslides were observed (Dewitte et al., 2021), whereas seismicity is an important component of the old  
deep-seated landslide model. In addition, the climatic and seismic conditions have evolved over the past tens of  
thousands of years (Felton et al., 2007; Wassmer et al., 2013; Ross et al., 2014; Smets et al., 2016). For example,  
605 the region experienced an abrupt shift from drier conditions to more humid conditions around 13,000 BP (Felton  
et al., 2007; Wassmer et al., 2013). In addition, about 10,000 BP, Lake Kivu water highstands were ~100 m above  
the current level, which could have triggered few large landslides (Ross et al., 2014; Dewitte et al., 2021). This

change in the lake level was not only due to a shift in the climatic conditions but also to the formation of the Virunga Volcano Province that created a dam on the upstream part of the Rift basin that used to drain northwards (Figure 1b; Haberyan and Hecky, 1987). During that period of volcano formation, the regional geodynamics and the seismicity pattern were different (Smets et al., 2016). Hence a large part of the old deep-seated landslides may have been triggered under different conditions.

Old and recent deep-seated landslides differ also in terms of size (Fig. 4). There have not been any major events during the past 60 years that caused large landslides comparable to the largest old deep-seated landslides (of area  $10^6 \text{ m}^2$ ). We identify five possible factors to explain this difference. First, our window of observation is too narrow to apprehend the impact of forcing events of high-magnitude such as large earthquakes (Marc et al., 2019). Second, the past environmental conditions may have been more favourable to large slope failures. A third factor explaining the size difference between old and recent deep-seated processes is that larger landslides are less frequent but have a longer-lived morphology legacy; therefore smaller old deep-seated landslides may no longer be visible. The fourth factor is that old landslides have a size that is the legacy of a history of phases of slope deformation, and not one single slope failure (Tanyaş et al., 2018). Fifth, amalgamation must not be excluded (Marc and Hovius, 2015), especially for the eldest features. Overall, our current knowledge does not allow to give more credit to one factor in particular. The common sense is certainly to assume that the difference in landslide size is the reflection of a combination of factors.

### 5.3 Drivers of shallow landslides

Rainfall is the trigger of the shallow landslides that we have identified in this study, which is in agreement with the other studies in the region (Dewitte et al., 2021; Kubwimana et al., 2021). The spatial distribution of shallow landslides differs from the distribution of deep-seated landslides. This is mainly due to the anthropogenic factors such as deforestation that influence shallow processes (Table 4). The regional susceptibility model also indicates that deforestation is the most important factor in their occurrence (Table 5). Similarly, the analysis of frequency ratios shows that landslides disproportionately occur within areas that were deforested in the past 60 years, demonstrating the role of the forest in slope stabilization (Grima et al., 2020).

Shallow landslides in forest loss areas (Fig. 6a,b) have, on average, a smaller size compared to landslides in forest. This observation is in line with the findings of Depicker et al. (2021b) and is attributed to the decrease of regolith cohesion by reduced root cohesion and evapotranspiration due to forest loss (Glade, 2003; Masi et al., 2021), which allows for a smaller minimum critical area for landsliding (Milledge et al., 2014). In short, human-induced land cover change is associated with an increase in the number of landslides and a shift of the frequency-area distribution towards smaller landslides (Guns and Vanacker, 2014).

In permanent anthropogenic environments (Fig. 6a,c), shallow landslides are less frequent, larger, and occur on less steep slopes as compared to shallow landslides in forest. Firstly, the steepest slopes in the anthropogenic environments have been subject to increased landslide erosion the first few years after the original forest cover was removed (prior to 1955-1958) (Depicker et al., 2021b). As a result, we can assume that steep slopes in

anthropogenic environments have less regolith available for landsliding compared to steep slopes in permanent forest areas. This process of regolith depletion is further exacerbated in cropland. Wilken et al. (2021) have measured in the region that erosion in cropland sites can reach up to about 40 cm in 55 years. Similarly, Heri-Kazi and Biielders (2021a) measured mean erosion rates of the order of 11 mm/year on cropland. Regolith erosion has therefore the consequence of reducing the spatial extent of areas where landslides can occur. A second process that may explain the landslide pattern in the anthropogenic environments is that, in parallel to regolith erosion, one also has sedimentation and the formation of colluvium (Wilken et al., 2021); which results in local accumulation of material. The material forms a loose sedimentary deposit usually in places with lower slope angles. This could be extra material available for the formation of landslides; the colluvium supply and a minimum depth of material being recognized as playing a key role in the occurrence of shallow landslides (Parker et al., 2016). Hence, we have less areas available for landslides, but a concentration of the susceptible places. A third explanation is probably related to soil management practices that influence erosion and water infiltration. In the region, usually on the less steep terrain, drainage ditches that favour water infiltration and hence an increase in pore-water pressure are widely applied by farmers (Heri-Kazi and Biielders, 2021b).

#### **5.4 Drivers of mining landslides and road landslides**

The poor prediction rates of mining and road landslides when compared to the two shallow and deep-seated susceptibility models (Fig.7) shows that they respond to different environmental factors. Road construction and mining activities are commonly associated with the presence of slope cuts and an increase of slope angle. These altered local topographic conditions cannot be constrained in the covariates derived from the SRTM or similar available products. In addition, the disturbances induced by roads and mining activities are not limited to the sole change of slope angle conditions. For example, this also implies changes in water runoff and infiltration, debutting, presence of fills and eventual overloading, excess stress from engine/digging, i.e., conditions that can influence the size and frequency characteristics of landslides (Brenning et al., 2015; Arca et al., 2018; Froude and Petley, 2018; McAdoo et al., 2018; Vuillez et al., 2018; Tanyaş et al., 2022).

Road landslides are mostly shallow. While it is obvious that roads create favourable conditions for the initiation of landslides, as observed not only in the region (e.g. Kubwimana et al., 2021), but also worldwide (Froude and Petley, 2018; Sidle et al., 2006; Brenning et al., 2015; Arca et al., 2018; McAdoo et al., 2018; Vuillez et al., 2018; Muñoz-Torrero Manchado et al., 2021; Tanyaş et al., 2022); an accurate spatio-temporal regional pattern of these human-induced slope failures cannot be assessed here. A substantial proportion of road landslides can only be observed in the field (Table 3). In addition, landslides along roads can easily disappear due to maintenance works. Furthermore, many of the main roads were already present in the 1950's, their current impact therefore being altered.

Overall, mining conditions seem to lead to landslides whose smallest features are more frequent than what would occur under natural conditions as attested in the frequency area distribution (see Section 4.2). The area of mining landslides is significantly larger than that of road landslides and their regional distribution is slightly more in agreement with the characteristics of deep-seated landslides (Fig. 7d), which is logical as mining activities are related to the lithological characteristics of the landscape; i.e. a cause that typically has more influence on deeper processes (Migoñ, 2013; Dille et al., 2019).

Considering the recent development of the mining activities in the region (Butsic et al., 2015; Tyukavina et al., 2018; Musumba Teso et al., 2019), we can assume with confidence that the associated landslides represent slope instabilities that have occurred over a period of about 20 years whereas the recent deep-seated landslides represent slope failures that have occurred over the last 60 years. The distribution of the mining landslides is also restricted spatially to some lithologies. With these specificities in mind and the fact that the number of inventoried mining and recent deep-seated landslides is relatively similar, respectively 152 and 159 (Table 2), this study confirms that mining activities increase the odds of landsliding. It has implication not only in terms of hazard assessment but also in assessing the population at risk, knowing that mined sites are populated. This is to be put in parallel with the findings of Depicker et al. (2021a) that show that the risk of shallow landslides has increased significantly in the region during the last decades in the places where mining activities are found due, notably, to an increase in population.

## 6 Conclusions

Our study improves the understanding of landslide processes and the human impact thereon in tropical rural mountainous environments. The use of several sources of data allowed to build a very detailed and comprehensive landslide inventory in time and space for the region; a source of information unprecedented in such environments. This inventory enabled the grouping of landslides into five categories: old and recent deep-seated landslides, shallow landslides, mining landslides and road landslides. Among deep-seated landslides, historical aerial photographs from the 1950's were an added value in the sense that they were used for differentiating between old and recent slope processes. We deduce the differences in the driving factors and area distribution for old and recent deep-seated landslides, suggesting that factors of landslide occurrence are either different or change over time depending on geodynamic and/or climatic conditions. The role of anthropogenic factors has been established in the occurrence of shallow landslides. Deforestation initially increases landsliding, but in the long term, when forest is permanently converted into agricultural land, landslide frequency appears to be lower compared to permanent forest lands. The impact of forest, forest cover changes and soil management practices depends on topographic conditions and regolith availability. The factors of occurrence of mining landslides significantly increase landsliding in areas that, under natural conditions, would be less prone to slope failures. Our analysis shows that the importance of human activities must be considered when investigating landslide occurrence in regions under anthropogenic pressure. This is particularly needed when one sees that the changing spatio-temporal patterns of landslides associated with these activities tend to further exacerbate the risks that the population face. On a more technical/methodological note, our study also demonstrates the importance of considering the timing of landslides in susceptibility and distribution assessments.

## Author contribution

J-C.M.M C.B and O.D conceived the study. J-C.M.M processed and analysed the data and created the figures with inputs from C.B, O.D and A.D. J-C.M.M wrote the manuscript, with main inputs from C.B and O.D and contributions from A.D and B.S. J-C.M.M and O.D compiled the landslide inventory. J-C.M.M, O.D, E.M, T.T and L.B.M participated in the fieldwork data acquisition and interpretation. B.S. generated the orthomosaics from

aerial photographs. All the authors contributed to the final version of the paper. C.B. and O.D. obtained funding  
725 for this work.

### **Declaration of Competing Interest**

The authors declare that they have no conflict of interest.

### **Acknowledgements**

730 Jean-Claude Maki Mateso was supported by the Université catholique de Louvain (doctoral scholarship from the  
Administration of International Relations) and the Development Cooperation programme of the Royal Museum  
for Central Africa with support of the Directorate-General Development Cooperation and Humanitarian Aid of  
Belgium (RMCA-DGD). Arthur Depicker and Benoît Smets were supported by the PASTECA project (BELSPO  
BRAIN-be Programme, Contract BR/165/A3/PASTECA; <http://pasteca.africamuseum.be/>). Elise Monsieurs was  
735 supported by a F.R.S. – FNRS PhD scholarship. The fieldwork was supported by the GeoRisCA (BELSPO SSD  
Programme, Contract SD/RI/02A; <http://georisca.africamuseum.be/>), RESIST (BELSPO STEREO-III  
Programme, Contract SR/00/305; <http://resist.africamuseum.be/>) and HARISSA (RMCA-DGD 2019-2024;  
<https://georiska.africamuseum.be/en/projects/harissa>) projects. We wish to thank B. Delvaux, J. Poesen and V.  
Vanacker for their insightful discussions and recommendations regarding this research. A special thank goes to  
740 François Kervyn for his constant support to conduct research in the region.

### **References**

- Albino, F., Smets, B., d'Oreye, N., and Kervyn, F.: High-resolution TanDEM-X DEM: An accurate method to  
estimate lava flow volumes at Nyamulagira Volcano (D. R. Congo), *J. Geophys. Res. Solid Earth*, 120, 4189–  
745 4207, <https://doi.org/10.1002/2015JB011988>, 2015.
- Aleman, J. C., Jarzyna, M. A., and Staver, A. C.: Forest extent and deforestation in tropical Africa since 1900,  
*Nat. Ecol. Evol.*, 2, 26–33, <https://doi.org/10.1038/s41559-017-0406-1>, 2018.
- Arca, D., Kutoğlu, H. Ş., and Becek, K.: Landslide susceptibility mapping in an area of underground mining  
using the multicriteria decision analysis method., *Environ. Monit. Assess.*, 190, 725,  
750 <https://doi.org/10.1007/s10661-018-7085-5>, 2018.
- Bashwira, M.-R., Cuvelier, J., Hilhorst, D., and van der Haar, G.: Not only a man's world: Women's  
involvement in artisanal mining in eastern DRC, *Resour. Policy*, 40, 109–116,  
<https://doi.org/10.1016/j.resourpol.2013.11.002>, 2014.
- Basnet, B. and Vodacek, A.: Tracking Land Use/Land Cover Dynamics in Cloud Prone Areas Using Moderate  
755 Resolution Satellite Data: A Case Study in Central Africa, *Remote Sens.*, 7, 6683–6709,  
<https://doi.org/10.3390/rs70606683>, 2015.
- Bennett, G. L., Miller, S. R., Roering, J. J., and Schmidt, D. A.: Landslides, threshold slopes, and the survival of  
relict terrain in the wake of the Mendocino Triple Junction, *Geology*, 44, 363–366,  
<https://doi.org/10.1130/G37530.1>, 2016.

- 760 Brenning, A.: Improved spatial analysis and prediction of landslide susceptibility: Practical recommendations, in: *Landslides and Engineered Slopes: Protecting Society Through Improved Understanding*, Chapter Improved Spatial Analysis and Prediction of Landslide Susceptibility: Practical Recommendations, Taylor & Francis, Alberta, 789–794, 2012.
- Brenning, A., Schwinn, M., Ruiz-Páez, A. P., and Muenchow, J.: Landslide susceptibility near highways is increased by 1 order of magnitude in the Andes of southern Ecuador, Loja province, *Nat. Hazards Earth Syst. Sci.*, 15, 45–57, <https://doi.org/10.5194/nhess-15-45-2015>, 2015.
- 765 Broeckx, J., Vanmaercke, M., Duchateau, R., and Poesen, J.: A data-based landslide susceptibility map of Africa, *Earth-Science Rev.*, 185, 102–121, <https://doi.org/10.1016/j.earscirev.2018.05.002>, 2018.
- Butsic, V., Baumann, M., Shortland, A., Walker, S., and Kuemmerle, T.: Conservation and conflict in the Democratic Republic of Congo: The impacts of warfare, mining, and protected areas on deforestation, *Biol. Conserv.*, 191, 266–273, <https://doi.org/10.1016/j.biocon.2015.06.037>, 2015.
- 770 Chen, L., Guo, Z., Yin, K., Shrestha, D. P., and Jin, S.: The influence of land use and land cover change on landslide susceptibility: a case study in Zhushan Town, Xuan’ en County (Hubei, China), *Nat. Hazards Earth Syst. Sci.*, 19, 2207–2228, <https://doi.org/10.5194/nhess-19-2207-2019>, 2019.
- 775 Cirimwami, L., Doumenge, C., Kahindo, J.-M., and Amani, C.: The effect of elevation on species richness in tropical forests depends on the considered lifeform: results from an East African mountain forest, *Trop. Ecol.*, 60, 473–484, <https://doi.org/10.1007/s42965-019-00050-z>, 2019.
- DeFries, R. S., Rudel, T., Uriarte, M., and Hansen, M.: Deforestation driven by urban population growth and agricultural trade in the twenty-first century, *Nat. Geosci.*, 3, 178–181, <https://doi.org/10.1038/ngeo756>, 2010.
- 780 Delvaux, D., Mulumba, J.-L., Sebagenzi, M. N. S., Bondo, S. F., Kervyn, F., and Havenith, H.-B.: Seismic hazard assessment of the Kivu rift segment based on a new seismotectonic zonation model (western branch, East African Rift system), *J. African Earth Sci.*, 134, 831–855, <https://doi.org/10.1016/j.jafrearsci.2016.10.004>, 2017.
- Depicker, A., Jacobs, L., Delvaux, D., Havenith, H.-B., Maki Mateso, J.-C., Govers, G., and Dewitte, O.: The added value of a regional landslide susceptibility assessment: The western branch of the East African Rift, *Geomorphology*, 353, 106886, <https://doi.org/10.1016/j.geomorph.2019.106886>, 2020.
- 785 Depicker, A., Jacobs, L., Mboga, N., Smets, B., Van Rompaey, A., Lennert, M., Wolff, E., Kervyn, F., Michellier, C., Dewitte, O., and Govers, G.: Historical dynamics of landslide risk from population and forest-cover changes in the Kivu Rift, *Nat. Sustain.*, 4, 965–974, <https://doi.org/10.1038/s41893-021-00757-9>, 2021a.
- Depicker, A., Govers, G., Jacobs, L., Campforts, B., Uwihirwe, J., and Dewitte, O.: Interactions between deforestation, landscape rejuvenation, and shallow landslides in the North Tanganyika–Kivu rift region, Africa, *Earth Surf. Dyn.*, 9, 445–462, <https://doi.org/10.5194/esurf-9-445-2021>, 2021b.
- 790 Dewitte, O., Dille, A., Depicker, A., Kubwimana, D., Maki Mateso, J.-C., Mugaruka Bibentyo, T., Uwihirwe, J., and Monsieurs, E.: Constraining landslide timing in a data-scarce context: from recent to very old processes in the tropical environment of the North Tanganyika-Kivu Rift region, *Landslides*, 18, 161–177,



- 795 <https://doi.org/10.1007/s10346-020-01452-0>, 2021.
- Dewitte, O., Depicker, A., Moeyersons, J., and Dille, A.: Mass Movements in Tropical Climates, in: *Treatise on Geomorphology*, vol. 5, edited by: Shroder, J. J. F., Elsevier, 338–349, <https://doi.org/10.1016/B978-0-12-818234-5.00118-8>, 2022.
- Dille, A., Kervyn, F., Mugaruka Bibentyo, T., Delvaux, D., Ganza, G. B., Ilombe Mawe, G., Kalikone Buzera, C., Safari Nakito, E., Moeyersons, J., Monsieurs, E., Nzolang, C., Smets, B., Kervyn, M., and Dewitte, O.: Causes and triggers of deep-seated hillslope instability in the tropics – Insights from a 60-year record of Ikoma landslide (DR Congo), *Geomorphology*, 345, 106835, <https://doi.org/10.1016/j.geomorph.2019.106835>, 2019.
- 800 Drake, T. W., Van Oost, K., Barthel, M., Bauters, M., Hoyt, A. M., Podgorski, D. C., Six, J., Boeckx, P., Trumbore, S. E., Cizungu Ntaboba, L., and Spencer, R. G. M.: Mobilization of aged and biolabile soil carbon by tropical deforestation, *Nat. Geosci.*, 12, 541–546, <https://doi.org/10.1038/s41561-019-0384-9>, 2019.
- 805 Emberson, R., Kirschbaum, D., and Stanley, T.: New global characterisation of landslide exposure, *Nat. Hazards Earth Syst. Sci.*, 20, 3413–3424, <https://doi.org/10.5194/nhess-20-3413-2020>, 2020.
- ESA: ESA Climate Change Initiative – Africa Land Cover Project 2017. 20 m Resolution, Eur. Sp. Agency, 2016.
- 810 Fang, Z., Wang, Y., Peng, L., and Hong, H.: Integration of convolutional neural network and conventional machine learning classifiers for landslide susceptibility mapping, *Comput. Geosci.*, 139, 104470, <https://doi.org/10.1016/j.cageo.2020.104470>, 2020.
- Felton, A. A., Russell, J. M., Cohen, A. S., Baker, M. E., Chesley, J. T., Lezzar, K. E., McGlue, M. M., Pigati, J. S., Quade, J., Curt Stager, J., and Tiercelin, J. J.: Paleolimnological evidence for the onset and termination of glacial aridity from Lake Tanganyika, Tropical East Africa, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 252, 405–423, <https://doi.org/10.1016/j.palaeo.2007.04.003>, 2007.
- 815 Fisher, G. B., Amos, C. B., Bookhagen, B., Burbank, D. W., and Godard, V.: Channel widths, landslides, faults, and beyond: The new world order of high-spatial resolution Google Earth imagery in the study of earth surface processes, in: *Google Earth and Virtual Visualizations in Geoscience Education and Research*, vol. 492, Geological Society of America, 1–22, [https://doi.org/10.1130/2012.2492\(01\)](https://doi.org/10.1130/2012.2492(01)), 2012.
- 820 Froude, M. J. and Petley, D. N.: Global fatal landslide occurrence from 2004 to 2016, *Nat. Hazards Earth Syst. Sci.*, 18, 2161–2181, <https://doi.org/10.5194/nhess-18-2161-2018>, 2018.
- Geenen, S.: A dangerous bet: The challenges of formalizing artisanal mining in the Democratic Republic of Congo, *Resour. Policy*, 37, 322–330, <https://doi.org/10.1016/j.resourpol.2012.02.004>, 2012.
- 825 Glade, T.: Landslide occurrence as a response to land use change: a review of evidence from New Zealand, *Catena*, 51, 297–314, [https://doi.org/10.1016/S0341-8162\(02\)00170-4](https://doi.org/10.1016/S0341-8162(02)00170-4), 2003.
- Grima, N., Edwards, D., Edwards, F., Petley, D., and Fisher, B.: Landslides in the Andes: Forests can provide cost-effective landslide regulation services, *Sci. Total Environ.*, 745, 141128,

<https://doi.org/10.1016/j.scitotenv.2020.141128>, 2020.

830 Guns, M. and Vanacker, V.: Shifts in landslide frequency–area distribution after forest conversion in the tropical Andes, *Anthropocene*, 6, 75–85, <https://doi.org/10.1016/j.ancene.2014.08.001>, 2014.

Guzzetti, F., Mondini, A. C., Cardinali, M., Fiorucci, F., Santangelo, M., and Chang, K.: Landslide inventory maps: New tools for an old problem, *Earth-Science Rev.*, 112, 42–66, <https://doi.org/10.1016/j.earscirev.2012.02.001>, 2012.

835 Haberyan, K. A. and Hecky, R. E.: The late Pleistocene and Holocene stratigraphy and paleolimnology of Lakes Kivu and Tanganyika, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 61, 169–197, [https://doi.org/10.1016/0031-0182\(87\)90048-4](https://doi.org/10.1016/0031-0182(87)90048-4), 1987.

Heri-Kazi, A. B. and Biielders, C. L.: “Cropland characteristics and extent of soil loss by rill and gully erosion in smallholder farms in the KIVU highlands, D.R. Congo,” *Geoderma Reg.*, 26, e00404,

840 <https://doi.org/10.1016/j.geodrs.2021.e00404>, 2021a.

Heri-Kazi, A. B. and Biielders, C. L.: Erosion and soil and water conservation in South-Kivu (eastern DR Congo): The farmers’ view, *L. Degrad. Dev.*, 32, 699–713, <https://doi.org/10.1002/ldr.3755>, 2021b.

Hosmer, D. W. and Lemeshow, S.: *Applied Logistic Regression*, Second Edi., Wiley, New York, 392 pp., 2000.

Hungr, O., Leroueil, S., and Picarelli, L.: The Varnes classification of landslide types, an update, *Landslides*, 11, 167–194, <https://doi.org/10.1007/s10346-013-0436-y>, 2014.

845 Imani, G., Boyemba, F., Lewis, S., Nabahungu, N. L., Calders, K., Zapfack, L., Riera, B., Balegamire, C., and Cuni-Sanchez, A.: Height-diameter allometry and above ground biomass in tropical montane forests: Insights from the Albertine Rift in Africa, *PLoS One*, 12, e0179653, <https://doi.org/10.1371/journal.pone.0179653>, 2017.

Jacobs, L., Dewitte, O., Poesen, J., Maes, J., Mertens, K., Sekajugo, J., and Kervyn, M.: Landslide characteristics and spatial distribution in the Rwenzori Mountains, Uganda, *J. African Earth Sci.*, 134, 917–930, 850 <https://doi.org/10.1016/j.jafrearsci.2016.05.013>, 2017.

Jacobs, L., Dewitte, O., Poesen, J., Sekajugo, J., Nobile, A., Rossi, M., Thiery, W., and Kervyn, M.: Field-based landslide susceptibility assessment in a data-scarce environment: the populated areas of the Rwenzori Mountains, *Nat. Hazards Earth Syst. Sci.*, 18, 105–124, <https://doi.org/10.5194/nhess-18-105-2018>, 2018.

855 Jaillon, A.: Maps of conflict minerals in eastern DRC - A0 posters (International Peace Information Service, Accessed September 25, 2020), 2020.

Jones, J. N., Boulton, S. J., Stokes, M., Bennett, G. L., and Whitworth, M. R. Z.: 30-year record of Himalaya mass-wasting reveals landscape perturbations by extreme events, *Nat. Commun.*, 12, 6701, <https://doi.org/10.1038/s41467-021-26964-8>, 2021.

860 Kampunzu, A. B., Bonhomme, M. G., and Kanika, M.: Geochronology of volcanic rocks and evolution of the Cenozoic western branch of the East African Rift system, *J. African Earth Sci.*, 26, 441–461, [https://doi.org/10.1016/S0899-5362\(98\)00025-6](https://doi.org/10.1016/S0899-5362(98)00025-6), 1998.

- Keefner, D. K.: Landslides caused by earthquakes., *Geol. Soc. Am. Bull.*, 95, 406–421, [https://doi.org/https://doi.org/10.1130/0016-7606\(1984\)95](https://doi.org/https://doi.org/10.1130/0016-7606(1984)95), 1984.
- 865 Kirschbaum, D. B., Adler, R., Hong, Y., Kumar, S., Peters-Lidard, C., and Lerner-Lam, A.: Advances in landslide nowcasting: evaluation of a global and regional modeling approach, *Environ. Earth Sci.*, 66, 1683–1696, <https://doi.org/10.1007/s12665-011-0990-3>, 2012.
- Kleinbaum, D. G. and Klein, M.: *Logistic Regression*, edited by: Intergovernmental Panel on Climate Change, Springer New York, New York, NY, 167-73 pp., <https://doi.org/10.1007/978-1-4419-1742-3>, 2010.
- 870 Kubwimana, D., Ait Brahim, L., Nkurunziza, P., Dille, A., Depicker, A., Nahimana, L., Abdelouafi, A., and Dewitte, O.: Characteristics and Distribution of Landslides in the Populated Hillslopes of Bujumbura, Burundi, *Geosciences*, 11, 259, <https://doi.org/10.3390/geosciences11060259>, 2021.
- Laghmouch, M., Kalikone, C. B., Ganza, G. B., Delvaux, D., Bachinyaga, J., Wazi, N., Nzolang, C., Fernandez, M., Nimpagaritse, G., Tack, L., Dewaele, S., and Kervyn, F.: *Carte géologique du Kivu au 1/500 000*, 2018.
- 875 LaHusen, S. R., Duvall, A. R., Booth, A. M., Grant, A., Mishkin, B. A., Montgomery, D. R., Struble, W., Roering, J. J., and Wartman, J.: Rainfall triggers more deep-seated landslides than Cascadia earthquakes in the Oregon Coast Range, USA, *Sci. Adv.*, 6, eaba6790, <https://doi.org/10.1126/sciadv.aba6790>, 2020.
- Lee, S. and Pradhan, B.: Landslide hazard mapping at Selangor, Malaysia using frequency ratio and logistic regression models, *Landslides*, 4, 33–41, <https://doi.org/10.1007/s10346-006-0047-y>, 2007.
- 880 Lee, S., Ryu, J.-H., and Kim, I.-S.: Landslide susceptibility analysis and its verification using likelihood ratio, logistic regression, and artificial neural network models: case study of Youngin, Korea, *Landslides*, 4, 327–338, <https://doi.org/10.1007/s10346-007-0088-x>, 2007.
- Linard, C., Gilbert, M., Snow, R. W., Noor, A. M., and Tatem, A. J.: Population Distribution, Settlement Patterns and Accessibility across Africa in 2010, *PLoS One*, 7, e31743, <https://doi.org/10.1371/journal.pone.0031743>, 2012.
- 885 Maki Mateso, J.-C. and Dewitte, O.: Towards an inventory of landslide processes and the elements at risk on the Rift flanks West of Lake Kivu (DRC), *Geo. Eco. Trop.*, 38, 137–154, 2014.
- Malamud, B. D., Turcotte, D. L., Guzzetti, F., and Reichenbach, P.: Landslide inventories and their statistical properties, *Earth Surf. Process. Landforms*, 29, 687–711, <https://doi.org/10.1002/esp.1064>, 2004.
- 890 Marc, O. and Hovius, N.: Amalgamation in landslide maps: Effects and automatic detection, *Nat. Hazards Earth Syst. Sci.*, 15, 723–733, <https://doi.org/10.5194/nhess-15-723-2015>, 2015.
- Marc, O., Behling, R., Andermann, C., Turowski, J. M., Illien, L., Roessner, S., and Hovius, N.: Long-term erosion of the Nepal Himalayas by bedrock landsliding: the role of monsoons, earthquakes and giant landslides, *Earth Surf. Dyn.*, 7, 107–128, <https://doi.org/10.5194/esurf-7-107-2019>, 2019.
- 895 Masi, E. B., Segoni, S., and Tofani, V.: Root Reinforcement in Slope Stability Models: A Review, *Geosciences*, 11, 212, <https://doi.org/10.3390/geosciences11050212>, 2021.

- Masumbuko, N. C., Habiyaremye, M. F., and Lejoly, J.: Woody climbing plants influence the structure of the mountain forest in the Kahuzi-Biega National Park, DR Congo, *Reg. Environ. Chang.*, 12, 951–959, <https://doi.org/10.1007/s10113-012-0309-2>, 2012.
- 900 McAdoo, B. G., Quak, M., Gnyawali, K. R., Adhikari, B. R., Devkota, S., Rajbhandari, P. L., and Sudmeier-Rieux, K.: Roads and landslides in Nepal: how development affects environmental risk, *Nat. Hazards Earth Syst. Sci.*, 18, 3203–3210, <https://doi.org/10.5194/nhess-18-3203-2018>, 2018.
- Meunier, P., Hovius, N., and Haines, J. A.: Topographic site effects and the location of earthquake induced landslides, *Earth Planet. Sci. Lett.*, 275, 221–232, <https://doi.org/10.1016/j.epsl.2008.07.020>, 2008.
- 905 Michellier, C., Pigeon, P., Kervyn, F., and Wolff, E.: Contextualizing vulnerability assessment: a support to geo-risk management in central Africa, *Nat. Hazards*, 82, 27–42, <https://doi.org/10.1007/s11069-016-2295-z>, 2016.
- Migoñ, P.: 4.10 Weathering and Hillslope Development, in: *Treatise on Geomorphology*, vol. 4, edited by: Shroder, J. F., Elsevier, San Diego, 159–178, <https://doi.org/10.1016/B978-0-12-374739-6.00075-0>, 2013.
- Milledge, D. G., Bellugi, D., McKean, J. A., Densmore, A. L., and Dietrich, W. E.: A multidimensional stability model for predicting shallow landslide size and shape across landscapes, *J. Geophys. Res. Earth Surf.*, 119, 2481–2504, <https://doi.org/10.1002/2014JF003135>, 2014.
- 910 Mokoso, J. M., Habiyaremye, F. M., Janssen, T., van Diggelen, R., Robbrecht, E., and Habimana, H. N.: Diversité des Fougères et leurs alliées le long du gradient altitudinal au sein de l'écosystème forestier des montagnes du Parc National de Kahuzi-Biega (RD CONGO), *Int. J. Environ. Stud.*, 70, 259–283, <https://doi.org/10.1080/00207233.2013.778007>, 2013.
- 915 Monsieurs, E., Kirschbaum, D. B., Tan, J., Maki Mateso, J.-C., Jacobs, L., Plisnier, P.-D., Thiery, W., Umutoni, A., Musoni, D., Mugaruka Bibentyo, T., Balumezi Ganza, G., Ilombe Mawe, G., Bagalwa Mateso, L., Kankurize, C., Michellier, C., Stanley, T., Kervyn, F., Kervyn, M., Demoulin, A., and Dewitte, O.: Evaluating TMPA Rainfall over the Sparsely Gauged East African Rift, *J. Hydrometeorol.*, 19, 1507–1528, <https://doi.org/10.1175/JHM-D-18-0103.1>, 2018a.
- 920 Monsieurs, E., Jacobs, L., Michellier, C., Basimike Tchanganaboba, J., Ganza, G. B., Kervyn, F., Maki Mateso, J.-C., Mugaruka Bibentyo, T., Kalikone Buzera, C., Nahimana, L., Ndayisenga, A., Nkurunziza, P., Thiery, W., Demoulin, A., Kervyn, M., and Dewitte, O.: Landslide inventory for hazard assessment in a data-poor context: a regional-scale approach in a tropical African environment, *Landslides*, 15, 2195–2209, <https://doi.org/10.1007/s10346-018-1008-y>, 2018b.
- 925 Mugagga, F., Kakembo, V., and Buyinza, M.: A characterisation of the physical properties of soil and the implications for landslide occurrence on the slopes of Mount Elgon, Eastern Uganda, *Nat. Hazards*, 60, 1113–1131, <https://doi.org/10.1007/s11069-011-9896-3>, 2012.
- Muñoz-Torrero Manchado, A., Allen, S., Ballesteros-Cánovas, J. A., Dhakal, A., Dhital, M. R., and Stoffel, M.: 930 Three decades of landslide activity in western Nepal: new insights into trends and climate drivers, *Landslides*, 18, 2001–2015, <https://doi.org/10.1007/s10346-021-01632-6>, 2021.

- Musumba Teso, P., Kavira, M., and Katcho, K.: Key Factors Driving Deforestation in North-Kivu Province, Eastern DR-Congo Using GIS and Remote Sensing, *Am. J. Geogr. Inf. Syst.*, 8, 11–25, <https://doi.org/10.5923/j.ajgis.20190801.02>, 2019.
- 935 Nzabandora, C. K. and Roche, E.: Six millennia of environment evolving on the western ridge of the Kivu Lake in the Mount Kahuzi area (D.R.Congo) Palynological analysis of the Ngushu sedimentary sequence, *Geo. Eco. Trop.*, 39, 1–26, 2015.
- Pánek, T., Břežný, M., Kilnar, J., and Winocur, D.: Complex causes of landslides after ice sheet retreat: Post-LGM mass movements in the Northern Patagonian Icefield region, *Sci. Total Environ.*, 758, 143684, <https://doi.org/10.1016/j.scitotenv.2020.143684>, 2021.
- 940 Parker, R. N., Hales, T. C., Mudd, S. M., Grieve, S. W. D., and Constantine, J. A.: Colluvium supply in humid regions limits the frequency of storm-triggered landslides, *Sci. Rep.*, 6, 34438, <https://doi.org/10.1038/srep34438>, 2016.
- Peel, M. C., Finlayson, B. L., and McMahon, T. A.: Updated world map of the Köppen-Geiger climate classification, *Hydrol. Earth Syst. Sci.*, 11, 1633–1644, <https://doi.org/10.5194/hess-11-1633-2007>, 2007.
- 945 Prakash, N., Manconi, A., and Loew, S.: A new strategy to map landslides with a generalized convolutional neural network, *Sci. Rep.*, 11, 9722, <https://doi.org/10.1038/s41598-021-89015-8>, 2021.
- Reichenbach, P., Rossi, M., Malamud, B. D., Mihir, M., and Guzzetti, F.: A review of statistically-based landslide susceptibility models, *Earth-Science Rev.*, 180, 60–91, <https://doi.org/10.1016/j.earscirev.2018.03.001>, 2018.
- 950 Ross, K. A., Smets, B., De Batist, M., Hilbe, M., Schmid, M., and Anselmetti, F. S.: Lake-level rise in the late Pleistocene and active subaquatic volcanism since the Holocene in Lake Kivu, East African Rift, *Geomorphology*, 221, 274–285, <https://doi.org/10.1016/j.geomorph.2014.05.010>, 2014.
- Rossi, M. and Reichenbach, P.: LAND-SE : a software for statistically based landslide, *Geosci. Model Dev.*, 3533–3543, <https://doi.org/10.5194/gmd-9-3533-2016>, 2016.
- 955 Shu, H., Hürlimann, M., Molowny-Horas, R., González, M., Pinyol, J., Abancó, C., and Ma, J.: Relation between land cover and landslide susceptibility in Val d’Aran, Pyrenees (Spain): Historical aspects, present situation and forward prediction, *Sci. Total Environ.*, 693, 133557, <https://doi.org/10.1016/j.scitotenv.2019.07.363>, 2019.
- 960 Sidle, R. C. and Bogaard, T. A.: Dynamic earth system and ecological controls of rainfall-initiated landslides, *Earth-Science Rev.*, 159, 275–291, <https://doi.org/10.1016/j.earscirev.2016.05.013>, 2016.
- Sidle, R. C., Ziegler, A. D., Negishi, J. N., Nik, A. R., Siew, R., and Turkelboom, F.: Erosion processes in steep terrain—Truths, myths, and uncertainties related to forest management in Southeast Asia, *For. Ecol. Manage.*, 224, 199–225, <https://doi.org/10.1016/j.foreco.2005.12.019>, 2006.
- 965 Smets, B., Delvaux, D., Ross, K. A., Poppe, S., Kervyn, M., d’Oreye, N., and Kervyn, F.: The role of inherited

- crustal structures and magmatism in the development of rift segments: Insights from the Kivu basin, western branch of the East African Rift, *Tectonophysics*, 683, 62–76, <https://doi.org/10.1016/j.tecto.2016.06.022>, 2016.
- Stanley, T. and Kirschbaum, D. B.: A heuristic approach to global landslide susceptibility mapping, *Nat. Hazards*, 87, 145–164, <https://doi.org/10.1007/s11069-017-2757-y>, 2017.
- 970 Tanyaş, H., Allstadt, K. E., and van Westen, C. J.: An updated method for estimating landslide-event magnitude, *Earth Surf. Process. Landforms*, 43, 1836–1847, <https://doi.org/10.1002/esp.4359>, 2018.
- Tanyaş, H., Görüm, T., Kirschbaum, D., and Lombardo, L.: Could road constructions be more hazardous than an earthquake in terms of mass movement?, *Nat. Hazards*, 112, 639–663, <https://doi.org/10.1007/s11069-021-05199-2>, 2022.
- 975 Trefon, T.: *Congo’s Environmental Paradox: potential and predation in a land of plenty*, Zed Books Ltd., London, 194 pp., 2016.
- Tyukavina, A., Hansen, M. C., Potapov, P., Parker, D., Okpa, C., Stehman, S. V., Kommareddy, I., and Turubanova, S.: Congo Basin forest loss dominated by increasing smallholder clearing, *Sci. Adv.*, 4, eaat2993, <https://doi.org/10.1126/sciadv.aat2993>, 2018.
- 980 Van Acker, F.: Where did all the land go? Enclosure & social struggle in Kivu (D.R.Congo), *Rev. Afr. Polit. Econ.*, 32, 79–98, <https://doi.org/10.1080/03056240500120984>, 2005.
- Van Den Eeckhaut, M., Vanwalleghem, T., Poesen, J., Govers, G., Verstraeten, G., and Vandekerckhove, L.: Prediction of landslide susceptibility using rare events logistic regression: A case-study in the Flemish Ardennes (Belgium), *Geomorphology*, 76, 392–410, <https://doi.org/10.1016/j.geomorph.2005.12.003>, 2006.
- 985 Van Den Eeckhaut, M., Poesen, J., Govers, G., Verstraeten, G., and Demoulin, A.: Characteristics of the size distribution of recent and historical landslides in a populated hilly region, *Earth Planet. Sci. Lett.*, 256, 588–603, <https://doi.org/10.1016/j.epsl.2007.01.040>, 2007.
- Van de Walle, J., Thiery, W., Brousse, O., Souverijns, N., Demuzere, M., and van Lipzig, N. P. M.: A convection-permitting model for the Lake Victoria Basin: evaluation and insight into the mesoscale versus synoptic atmospheric dynamics, *Clim. Dyn.*, 54, 1779–1799, <https://doi.org/10.1007/s00382-019-05088-2>, 2020.
- 990 Vanacker, V., Vanderschaeghe, M., Govers, G., Willems, E., Poesen, J., Deckers, J., and De Bievre, B.: Linking hydrological, infinite slope stability and land-use change models through GIS for assessing the impact of deforestation on slope stability in high Andean watersheds, *Geomorphology*, 52, 299–315, [https://doi.org/10.1016/S0169-555X\(02\)00263-5](https://doi.org/10.1016/S0169-555X(02)00263-5), 2003.
- 995 Vanmaercke, M., Ardizzone, F., Rossi, M., and Guzzetti, F.: Exploring the effects of seismicity on landslides and catchment sediment yield: An Italian case study, *Geomorphology*, 278, 171–183, <https://doi.org/10.1016/j.geomorph.2016.11.010>, 2017.
- Vuillez, C., Tonini, M., Sudmeier-Rieux, K., Devkota, S., Derron, M.-H., and Jaboyedoff, M.: Land use changes, landslides and roads in the Phewa Watershed, Western Nepal from 1979 to 2016, *Appl. Geogr.*, 94, 30–40,

1000 <https://doi.org/10.1016/j.apgeog.2018.03.003>, 2018.

Wassmer, P., Schwartz, D., Gomez, C., Ward, S., and Barrere, P.: Geomorphology and sedimentary structures of upper pleistocene to holocene alluvium within the nyabarongo valley (Rwanda). palaeo-climate and palaeo-environmental implications, *Geogr. Fis. e Din. Quat.*, 36, 199–210, <https://doi.org/10.4461/GFDQ.2013.36.17>, 2013.

1005 Whipple, K. X.: The influence of climate on the tectonic evolution of mountain belts, *Nat. Geosci.*, 2, 97–104, <https://doi.org/10.1038/ngeo413>, 2009.

Whittaker, A. C.: How do landscapes record tectonics and climate?, *Lithosphere*, 4, 160–164, <https://doi.org/10.1130/RF.L003.1>, 2012.

1010 Wilken, F., Fiener, P., Ketterer, M., Meusburger, K., Muhindo, D. I., van Oost, K., and Doetterl, S.: Assessing soil redistribution of forest and cropland sites in wet tropical Africa using  $^{239+240}\text{Pu}$  fallout radionuclides, *Soil*, 7, 399–414, <https://doi.org/10.5194/soil-7-399-2021>, 2021.

1015

1020

1025